

HEPATITIS A VIRUS

THE ORGANISM/TOXIN

- Hepatitis A virus (HAV) is a small, single-stranded RNA virus belonging to the *Picornaviridae* family and the only member of the genus *Hepatovirus* (1, 2, 3).
- There is a single HAV serotype and multiple genotypes. Genotypes I, II and III, further divided into subtypes A and B, infect humans. HAV subtypes are further divided into clusters or strains which have been associated with particular risk factors (2, 3, 4, 5, 6).
- HAV is the main cause of acute self-limiting hepatitis in humans worldwide. The virus is primarily transmitted by the faecal/oral route following exposure to infected people, or via contact with HAV-contaminated food, water, or surfaces (2, 3).

GROWTH AND ITS CONTROL

Growth:

- HAV cannot grow in food, water or the environment. Wild -type strains grow poorly in cell culture however, cell culture-adapted strains can be cultured easily and used for survival and inactivation studies (7).

Survival:

Temperature	<ul style="list-style-type: none"> HAV is stable in the environment and can persist for prolonged periods (weeks or months) under normal, domestic food storage conditions (8, 9). The virus remains intact and infectious for several years under freezing conditions (8, 10). HAV can survive for 90 days at -20°C on blueberries, raspberries, strawberries, basil and parsley (11). HAV is stable under storage at 4°C in shellfish and produce with only 1.7 log₁₀ reduction after 4 weeks (12) and 2 log₁₀ reduction after 9 days respectively (13).
Drying	<ul style="list-style-type: none"> The half-life of HAV ranges from more than 7 days at low humidity (25%) (5°C) to 2 hours at high humidity (95%) (35°C), demonstrating a greater resistance to inactivation at low humidity and temperatures (14).
pH	<ul style="list-style-type: none"> Stable at acid pH. In tissue culture experiments, HAV remained infectious for up to 5 hours at pH 1.0 (25°C) (15). Acidic fruits have been associated with HAV outbreaks due to the acid-resistance of this virus (10).

Inactivation:

Thermal inactivation:	
Temperature	<ul style="list-style-type: none"> The Ministry for Primary Industries and Food Standards Australia New Zealand recommend cooking food to 85°C for 1 minute to inactivate HAV (16). The extent of virus inactivation is influenced by the food matrix. For example, complete inactivation of HAV has been observed in

	<p>shellfish when heating to 85°C for 3 minutes (16, 17) however, heat stability is increased in foods (eg dairy) with a higher fat content. Heating cream or milk to 85°C for 30 seconds is sufficient to cause only a 5 log₁₀ reduction in HAV titre (9, 16, 18).</p> <ul style="list-style-type: none"> HAV reduced by 6 log₁₀ in natural berries after 4-6 minutes at 75°C. Addition of increasing concentrations of sucrose extended the time taken to achieve a similar reduction in virus titre (16).
D values	<ul style="list-style-type: none"> A literature review suggests the following time/temperature (°C/minutes) in “shellfish”: D₅₀ 54.2; D₅₆ 9.3; D₆₀ 3.3-6.5, “meat”:D₅₀ 42.1; D₅₆ 20.6; D₆₀ 5.9 and “spinach”: D₅₀ 34.4; D₅₆ 8.4; D₆₀ 4.6 (19).
Non-thermal inactivation:	
Sanitisers /disinfectants	<ul style="list-style-type: none"> Commercial disinfectants used in the food industry containing 2% glutaraldehyde, 0.4% quaternary ammonium compound and 6% sodium hypochlorite reduced virus particles on stainless steel by more than 4 log₁₀ after one minute of contact time. Sodium hypochlorite at 3,000 ppm (5 minutes contact time at 22°C) reduced HAV on surfaces by > 5log₁₀ (8). Many chemical disinfectants used in institutional and domestic environments, including ethanol, chloroform, Freon, Arklone and perchloroacetic acid, do not effectively inactivate the virus on hands or surfaces (8). Chlorine disinfection of ready-to-eat (RTE) food reduced HAV on fresh produce including strawberries, cherry tomatoes and head lettuce (1.7 to 2.4 log reduction when exposed to 20 ppm solution of free chlorine for 5-10 minutes) (20).
Radiation	<ul style="list-style-type: none"> Extremely resistant to ionising radiation compared with bacteria. Nearly 3 kGy of gamma irradiation produced a 1 log₁₀ reduction of HAV on lettuce and strawberries (21) and 3 kGy dose reduced HAV by less than 2 log₁₀ in oysters and clams (10, 22).
Preservatives and other non-thermal processing technologies	<ul style="list-style-type: none"> High pressure processing (350 MPa, 21°C, 5 minutes) reduced HAV load in pureed strawberries and green onions by 4 log₁₀ and in Mediterranean mussels (400 MPa, room temperature, 5 minutes) by 2.9 log₁₀ (18, 23). Modified atmosphere packaging is not a suitable strategy for reduction of HAV on food (10).

THE ILLNESS

Incubation: Range 10-50 days (average 28) (3, 8).

Symptoms:

- Most HAV infections are mild and self-limiting lasting up to two months. Many infections, particularly in children, are asymptomatic.
- When disease is evident, HAV usually presents as four

classical phases: 1) hepatocyte viral replication in the absence of symptoms; 2) anorexia, nausea, vomiting, headache, alterations in taste, abdominal discomfort and fatigue lasting 5 to 7 days; 3) darkening of the urine, light-coloured stools, jaundice and right upper quadrant pain with enlargement of the liver, jaundice and liver enzyme elevation. Faecal shedding and viraemia are maximal during this stage which occurs in fewer than 10% of children under 6, 40-50% of older children and 70-80% of adults; 4) symptoms resolve and liver enzymes return to normal (3, 8).

Condition: Hepatitis A

At Risk Groups:

- All age groups. The disease is milder in children under 6, compared to older children and adults. Complications are more likely to occur in older and immunocompromised people (8).
- In developing/endemic countries, primary infection at a young age confers life-long immunity, hence outbreaks are unusual. In developed countries, HAV infections are less common, due to increased standards of living, but the majority of adults remain susceptible to infection leading to the occurrence of outbreaks amongst the general population (2, 3, 24).
- Risk factors include consumption of contaminated RTE food and/or water, travel to HAV- endemic areas and contact with infected persons. Increased risk is linked with exposure to blood or blood products (for example, injection-drug use) (2).

Long-term effects:

- Long or recurring illness, lasting up to 6 months, can occur in approximately 3-20% of clinical cases (8).
- Extra-hepatic disorders rarely occur but may include haemolysis, acute renal failure, reactive arthritis, pancreatitis, fulminant hepatitis and neurological syndromes (8, 25).

Dose: Highly infectious. As few as 10-100 virus particles appear to be able to cause disease (26).

Incidence:

- New Zealand incidence rate of 0.7/100,000 (2016). Notifications for HAV have remained relatively stable at 0.6-2.1 during the last 10 years. There have been fluctuations due to outbreaks in 2008, 2012 and 2013 (27). Updates can be found on the ESR website https://surv.esr.cri.nz/surveillance/annual_surveillance.php
- New Zealand incidence rates are slightly lower than the EU (2.6/100,000 (2010)) (28), slightly higher than the USA (0.43/100,000 (2015)) (29) and similar to Australia (1/100,000 (2014)) (30).

Treatment:

- Treatment rarely required. Supportive care with treatment for specific complications as appropriate (31).

Prevention and control strategies:

- Most effective way to prevent HAV infection is to improve sanitation, food safety and immunisation practices (3, 31).
- HAV is a vaccine-preventable disease (inactivated HAV vaccines are approved for use in New Zealand although are not routinely recommended for children).

The vaccine is recommended for certain at-risk groups including transplant patients, people with chronic liver disease, close contacts of hepatitis A cases, travellers, health care workers and food handlers when situations warrant it (3, 31, 32, 33).

- Post-exposure prophylaxis (within 2 weeks) and prophylactic administration of human immunoglobulins may be recommended for short-term protection against HAV for certain groups including day-care centre, military and food industry staff (31, 33, 34).

SOURCES

Human:

- Humans are the major reservoir of natural HAV infection (3).

Animal:

- No animal vectors. Chimpanzees and several small, non-human primates are also susceptible to HAV but the disease is milder (3).

Food:

- Contaminated shellfish, salads, fruits and vegetables and any manually-prepared RTE foods (2).
- Poor hygiene practices, poor sanitation and contamination of fresh produce, with e.g. HAV-contaminated irrigation water during production, are major risk factors for the transmission of HAV to humans (2).

Environment/Water:

- HAV can survive, but not multiply, in the environment. Many factors will influence virus stability including relative humidity, temperature, type of suspending medium and type of surface contaminated (8).
- Survival of HAV in environmental sources (fresh and seawater, wastewater, marine sediments, soils and shellfish) is prolonged (>12 weeks) at 25°C (2, 8).
- Faecal pollution, from infected individuals, can contaminate shellfish beds, recreational, irrigation and drinking water (2).

Transmission Routes:

- Directly, via person-to-person contact, or indirectly via consumption of contaminated food or water or contact with fomites (2, 8).
- On rare occasions, HAV may also be transmitted via blood transfusion or blood products originating from infected donors (35).

OUTBREAKS AND INCIDENTS

Outbreaks:

New Zealand

- Since 2008, the number of outbreaks of HAV has ranged from 0 to 5. The number of associated cases has ranged from 2 to 54.
- The number of foodborne outbreaks has remained at 1 or less for the past 10 years (36).
- The remainder of the annual outbreaks/cases may be sporadic or can be attributed to other sources including person-to-person and travel (36).

New Zealand

Notable foodborne outbreaks in recent years are included below ⁽³⁶⁾.

Year	Foodborne outbreaks (cases)	Suspected foods
2008	1 (2)	No identifiable source
2009	0 (0)	N/A
2010	1 (3)	No identifiable source
2011	0 (0)	N/A
2012	0 (0)	N/A
2013	0 (0)	N/A
2014	0 (0)	N/A
2015	1 (7)	Fruit (imported frozen, mixed berries)

Worldwide

Notable foodborne outbreaks in recent years are included below.

Year	Cases (deaths)	Suspected foods	Country	Control measure failure
2004	269	Raw beef	Belgium	HAV-infected food handler ⁽³⁷⁾ .
2009	562	Semi-dried tomatoes	Australian (multistate)	Contamination of raw ingredients and manufacturing environment ⁽³⁸⁾ .
2012-2013	107	Strawberries/mango	EU (multistate)	Contaminated Egyptian produce. Cases among EU travellers returning from Egypt associated with exposure to strawberries or mango. Transmission likely to have occurred on hotel premises ⁽³⁹⁾ .
2013	165	Frozen pomegranate arils	USA (multistate)	Poor hygiene during food production for exportation the likely cause ⁽⁴⁰⁾ .
2013-2014	1589 (2)	Imported mixed, frozen berries	EU (multistate)	Consumption of various products containing contaminated berries. No point source of contamination identified ⁽⁴¹⁾ .
2016	143	Frozen strawberries	USA (multistate)	Contaminated produce imported from Egypt ⁽⁴²⁾ .

REFERENCES

- 1- Cohen, J.I., Ticehurst, J.R., Purcell, R.H., Buckler-White, A. and Baroudy, B.M. (1987). Complete nucleotide sequence of wild-type hepatitis A virus: comparison with different strains of Hepatitis A virus and other picornaviruses. *Journal of Virology*. **61**: 50-59.
- 2- Vaughan, G., Rossi, L.M.G., Forbi, J.C., de Paula, V.S., Purdy, M.A., Xia, G. and Khudyakov, Y.E. (2014). Hepatitis A virus: Host interactions, molecular epidemiology and evolution. *Infection, Genetics and Evolution* **21**: 227-243.
- 3- Lemon, S.M., Ott, J.J., Van Damme, P. and Shouval, D. (2017). Type A viral Hepatitis: A summary and update on the molecular virology, epidemiology, pathogenesis and prevention. *Journal of Hepatology* **68**: 167-184.
- 4- Robertson, B.H., Jansen, R.W., Khanna, B., Totsuka, A., Nainan, O.V., Siegl, G., Widell, A., Margolis, H.S., Isomura, S., Ito, K., Ihsizu, T., Moritzugu, Y. and Lemon, S.M. (1992). Genetic relatedness of Hepatitis A virus strains recovered from different geographical regions. *Journal of General Virology*. **73**: (pt 6). 1365-1377.
- 5- Nainan, O.V., Armstrong, G.L., Han, X.H., Williams, I., Bell, B.P. and Margolis, H.S. (2005). Hepatitis A molecular epidemiology in the United States 1996-1997: sources of infection and implications of vaccination policy. *Journal of Infectious Diseases*. **191**: 975-963.
- 6- Costa-Mattioli, M., Ferre, V., Casane, D., Perez-Bercoff, R., Coste-Burrel, M., Imbert-Marcille, B.M., Andre, E.C., Bressoliette-Bodin, C., Billaudel, S. and Cristina, S. (2013). Evidence of recombination in natural populations of Hepatitis A virus. *Virology*. **311**: 51-59.
- 7- The International Commission on Microbiological Specifications for Foods (1996). *Viruses*. In *Micro-organisms in Foods 5 Microbiological Specifications of Food Pathogens*. Blackie Academic and Professional, London.
- 8- D'Souza, D.H., Moe, C.L. and Jaukus, L. (2007) *Foodborne viral Pathogens*. In *Food Microbiology, fundamentals and Frontiers*, 3rd Edition. Edited by M.P. Doyle and L.R. Beuchat. ASM Press, Washington DC.
- 9- Sattar, S.A., Tetro, J., Bidawid, S. and Farber, J. (2000). Foodborne spread of Hepatitis A: Recent studies on virus survival, transfer and inactivation. *Canadian Journal of Infectious Diseases*. **11**: 159-163.
- 10- Baert, L., Debevere, J. and Uyttendaele, M. (2009). The efficacy of preservation methods to inactivate foodborne viruses. *International Journal of Food Microbiology*. **31**: 83-94.
- 11- Butot, S., Putallaz, T. and Sanchez, G. (2008). Effects of sanitation, freezing and frozen storage on enteric viruses in berries and herbs. *International Journal of Food Microbiology*. **126**: 30-35.
- 12- Hewitt, J. and Greening, G.E. (2004). Survival and persistence of norovirus, Hepatitis A virus and feline calicivirus in marinated mussels. *Journal of Food Protection* **67**: 1743-1750.
- 13- Croci, L., De Medici, D., Scalfaro, C., Fiore, A. and Toti, L. (2002). The survival of Hepatitis A virus in fresh produce. *International Journal of Food Microbiology* **73**: 29-34.
- 14- Mbithi, J.N., Springthorpe, V.S. and Sattar, S.A. (1991). Effect of relative humidity and air temperature on survival of Hepatitis A virus on environmental surfaces. *Applied and Environmental Microbiology*. **57**: 1394-1339.
- 15- Scholz, E., Heinrich, U. and Flehmig, B. (1989). Acid stability of Hepatitis A virus. *Journal of General Virology*. **70**: 2481-2485.
- 16- Food Standards Australia New Zealand (FSANZ). (2015). *Guidance for the thermal inactivation of Hepatitis A virus in berries*. Prepared by FSANZ Risk Assessment (microbiology) and the Ministry for Primary Industries Food Risk Assessment.
- 17- Millard, J., Appleton, H. and Parry, J.V. (1987). Studies on heat inactivation of Hepatitis A virus with special reference to shellfish. *Epidemiology and Infection*. **98**: 397-414.
- 18- Baert, L. (2013). *Foodborne virus inactivation by thermal and non-thermal processes*. Book chapter: viruses in food and water, Chapter 12. Pgs 237-260. Published by Woodhead Publishing Limited.
- 19- Horn, B. and Hewitt, J. (2016). *Review of microbial pathogen inactivation at sous vide cooking temperatures*. Client Report FW15038. Prepared for the Ministry for Primary Industries under project MFS/15/2 Microbiological Food Safety.
- 20- Casteel, M.J., Schmidt, C.E. and Sobsey, M.D. (2008). Chlorine disinfection of produce to inactivate Hepatitis A virus and coliphage MS2. *International Journal of Food Microbiology*. **125**: 267-273.

- 21- Bidawid, S., Farber, J.M. and Sattar, S.A. (2000). Inactivation of Hepatitis A virus (HAV) in fruits and vegetables by gamma irradiation. *International Journal of Food Microbiology*. **57**: 91-97.
- 22- Mallett, J.C., Beghian, L.E., Metcalf, T.G. and Kaylor, J.D. (1991). Potential of irradiation technology for improved shellfish sanitation. *Journal of Food Safety*. **11**: 231-245.
- 23- Kingsley, D.H. (2013). High pressure processing and its application to the challenge of virus-contaminated foods. *Food Environmental Virology* **5**: 1-12.
- 24- Sanchez, G., Bosch, A. and Pinto, R.M. (2007). Hepatitis A virus detection in food: current and future prospects. *Letters in Applied Microbiology* **45**: 1-5.
- 25- Gilkson, M., Galun, E., Oren, R., Tur-Kaspa, R. and Shouval, D. (1992). Relapsing Hepatitis A. Review of 14 literature cases and literature survey. *Medicine* **71**: 14-23.
- 26- Todd, E.C.D., Greig, J.D., Bartleson, C.A. and Michaels, B.C. (2008). Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 4. Infective doses and pathogen carriage. *Journal of Food Protection* **71**: 2339-2373.
- 27- ESR (2016) Notifiable diseases in New Zealand. Annual Report 2015. ESR, Porirua. https://surv.esr.cri.nz/PDF_surveillance/AnnualRpt/AnnualSurv/2016/2016AnnualNDReportFinal.pdf (accessed January 2018).
- 28- European Centre for Diseases Prevention and Control. The European Union Summary Report on trends and sources of Zoonoses, zoonotic agents and food-borne outbreaks in 2013. EFSA, Parma, Italy and ECDC, Stockholm, Sweden <http://ecdc.europa.eu/en/publications/Publications/EU-summary-report-trends-sources-zoonoses-2013.pdf> (accessed January 2018).
- 29- Centers for Disease Control and Prevention. Summary of Notifiable Infectious Diseases and Conditions – United States, 2014. https://www.cdc.gov/mmwr/mmwr_nd/index.html (accessed January 2018).
- 30- Annual Report of the National Notifiable Diseases Surveillance System. Australia's Notifiable Disease Status (2014): NNDS Annual Report Working Group. [https://health.gov.au/internet/main/publishing.nsf/Content/cda-cdi4001-pdf-cnt.htm/\\$FILE/cdi4001e.pdf](https://health.gov.au/internet/main/publishing.nsf/Content/cda-cdi4001-pdf-cnt.htm/$FILE/cdi4001e.pdf) (accessed January 2018).
- 31- Lorio, N. and John, S. (2017). Hepatitis A. From StatPearls Publishing LLC (October 23, 2017).
- 32- WHO position paper on Hepatitis A vaccines: June 2012 recommendations. (2013). *Vaccine* **31**: 285-286.
- 33- Ogholikhan, S. and Schwartz, K.B. (2016). Hepatitis vaccines. *Vaccines* **4(1)**: 6.
- 34- Hollinger, F.B. and Emerson, S.U. (2001). Hepatitis A virus. pgs 799-840. In D.M. Knipe, P.M. Howley, D.E. Griffin, R.A. Lamb, M.A. Martin, B. Roizman and S.E. Straus (ed). *Fields Virology*. Lippincott, Williams and Wilkins, Philadelphia, Pa.
- 35- De Paula, V.S. (2012). Laboratory diagnosis of Hepatitis A. *Future Virology*. **7**: 11.
- 36- ESR (2015) Summary of outbreaks in New Zealand 2015. Annual Report (2015). ESR, Porirua. https://surv.esr.cri.nz/surveillance/annual_outbreak.php?we_objectID=4480 (accessed January 2018).
- 37- Robesyn, E., de Schrijver, K., Wollants, E., Top, G., Verbeeck, J. and Van Ranst, M. (2009). An outbreak of Hepatitis A associated with the consumption of raw beef. *Journal of Clinical Virology*. **44(3)**: 207-210.
- 38- Donnan, E.J., Fielding, J.E., Gregory, J.E., Lalor, K., Rowe, S., Goldsmith, P., Antoniou, M., Fullerton, K.E., Knope, K., Copland, J.G., Scott Bowden, D., Tracy, S.L., Hogg, G.G., Tan, A., Adamopoulos, J., Gaston, J. and Vally, H. (2012). A multistate outbreak of Hepatitis A associated with semidried tomatoes in Australia, 2009. *Clinical Infectious Diseases*. **6**: 775-81.
- 39- Sane, J., MacDonald, E., Vold, L., Gossner, C. and Severi, E. (2015). Multistate foodborne Hepatitis A outbreak among European tourists returning from Egypt – need for reinforced vaccination recommendations, November 2012 to April 2013. *European Surveillance. Surveillance and outbreak reports* Jan 29; **20(4)** pii 21018.
- 40- Centers for Disease Control and Prevention (CDC). (2013). Multistate outbreak of Hepatitis A infection linked to pomegranate seeds from Turkey. *Advanced Molecular Detection (AMD)*. <https://www.cdc.gov/amd/stories/hepatitis-a.html> (accessed January 2018).
- 41- Severi, E. *et al* (2015). Large and prolonged food-borne multistate Hepatitis A outbreak in Europe associated with consumption of frozen berries, 2013 to 2014. *European Surveillance* **20(29)**:pii=21192. <http://www.eurosurveillance.org/images/dynamic/EE/V20N29/art21192.pdf> (accessed January 2018).
- 42- Centers for Disease Control and Prevention (CDC). (2016). Multistate outbreak of Hepatitis A linked to frozen strawberries (final update). <https://www.cdc.gov/hepatitis/outbreaks/2016/hav-strawberries.htm> (accessed January 2018).