



## Catches, size, and age structure of the 2015–16 hoki fishery, and a summary of input data used for the 2017 stock assessment

New Zealand Fisheries Assessment Report 2017/31

S.L. Ballara,  
R.L. O'Driscoll

ISSN 1179-5352 (online)  
ISBN 978-1-77665-606-6 (online)

June 2017



Requests for further copies should be directed to:

Publications Logistics Officer  
Ministry for Primary Industries  
PO Box 2526  
WELLINGTON 6140

Email: [brand@mpi.govt.nz](mailto:brand@mpi.govt.nz)  
Telephone: 0800 00 83 33  
Facsimile: 04-894 0300

This publication is also available on the Ministry for Primary Industries websites at:  
<http://www.mpi.govt.nz/news-and-resources/publications>  
<http://fs.fish.govt.nz> go to Document library/Research reports

**© Crown Copyright - Ministry for Primary Industries**

## CONTENTS

EXECUTIVE SUMMARY .....	1
1. INTRODUCTION .....	2
1.1 Stock structure .....	2
1.2 Description of the hoki fishery .....	3
1.3 Catch history .....	4
1.4 Recent hoki research .....	5
2. HOKI FISHERY, 2015–16 .....	5
2.1 Catch and effort information .....	5
2.1.1 Total Allowable Commercial Catch (TACC) and other management controls .....	5
2.1.2 Catch .....	6
2.1.3 CPUE analysis .....	7
2.1.4 Bycatch .....	9
2.2 Size and age composition of commercial catches .....	10
2.2.1 Size and age composition in spawning fisheries .....	11
2.2.2 Size and age composition in non-spawning fisheries .....	13
2.2.3 Comparison of size and age composition between main areas .....	14
3. HOKI RESEARCH .....	14
3.1 Resource surveys .....	14
3.1.1 Trawl surveys .....	14
3.1.2 Acoustic surveys .....	15
4. CONCLUSIONS .....	15
5. ACKNOWLEDGMENTS .....	16
6. REFERENCES .....	16
TABLES .....	19
FIGURES .....	35
APPENDICES .....	99



## EXECUTIVE SUMMARY

**Ballara, S.L.; O’Driscoll, R.L. (2017). Catches, size, and age structure of the 2015–16 hoki fishery, and a summary of input data used for the 2017 stock assessment.**

*New Zealand Fisheries Assessment Report 2017/31. 119 p.*

This report summarises catches by area and presents the length and age structure of hoki caught commercially during the 2015–16 fishing year. Length frequency and catch-at-age data from spawning and non-spawning fisheries are compared with those from previous years. Biomass indices from research surveys and results from other research on hoki in the last year are also briefly described. Data in this report were incorporated in the hoki stock assessment in 2017.

The total reported hoki catch in 2015–16 was 136 719 t, about 13 000 t lower than the TACC of 150 000 t, and 25 000 t lower than the catch in 2014–15. Catches in 2015–16 decreased in all areas except for the east coast South Island (ECSI) and east coast North Island (ECNI). The spawning fishery catch on the west coast South Island (WCSI) decreased by 10 000 t to 68 900 t, but was the largest hoki fishery for the sixth consecutive season. The non-spawning fishery on the Chatham Rise was the second largest hoki fishery, with 36 700 t taken in 2015–16, about 3000 t less than in 2014–15. The spawning fishery catch from Cook Strait decreased by 1900 t to 18 400 t, and the non-spawning catch from the Sub-Antarctic fishery decreased by 10 000 t to 6600 t in 2015–16. Catches from Puysegur decreased to 800 t, whereas catches from ECNI and ECSI increased by 100 and 500 t respectively. Overall, about 76 600 t of the total catch in 2015–16 was taken from western stock areas, well below the industry-agreed catch limit of 90 000 t, and about 60 100 t came from the eastern stock areas.

Length and age frequency distributions from the commercial fishery show that most of the catch in 2015–16 was fish of 50–90 cm total length. The 2011 year class (68–75 cm) was important in all areas as age 4+ years on the Chatham Rise and Sub-Antarctic, and as age 5 in Cook Strait (both sexes) and on the WCSI (mainly as males). The 2010 year class (age 5+) was also strongly represented in the Chatham Rise and in the Sub-Antarctic catches. There were few hoki from the 2012 year class in all areas. The 2014 year class also appeared in all main fishery areas. The percentage of small fish in the catch decreased in the non-spawning areas, mainly due to the presence of the strong 2010 and 2011 year class in these fisheries. Most fish caught on the Chatham Rise in 2015–16 were less than 90 cm, and the largest hoki came from the Cook Strait, ECSI, and WCSI fisheries.

Only one fishery independent research survey for hoki has been carried out since the 2016 stock assessment. The relative biomass index for hoki from the core strata in the November–December 2016 Sub-Antarctic trawl survey increased by 21% from 2014. The 2016 survey index was adjusted to account for strata that could not be completed due to rough weather. Several modes were present in the hoki scaled length frequency distribution including some 1+ fish (2015 year-class), relatively few 2+ fish (2014 year-class), and a broad mode of fish aged 5–12 years. A trawl survey was also carried out on the WCSI in August 2016, but trawl estimates from this survey are not considered to provide reliable indices of hoki abundance.

## 1. INTRODUCTION

This report provides biological data relevant to the 2017 hoki stock assessment. Catch statistics and data from commercial sampling carried out during the 2015–16 fishing year are presented, and results from other research programmes carried out since March 2016 are summarised, including results of the trawl survey of the Sub-Antarctic in December 2016. Details of model structure, results, and yield estimates for the hoki stock assessment carried out in 2017 are published separately.

### 1.1 Project objectives

This report fulfils the final reporting requirement for 3 Objectives in research projects DEE2016-08HOK, DEE2016-19, and DEE2016-20.

DEE2016-08HOK Objective 1: To provide a descriptive analysis of the hoki fishery in the 2015–16 year.

DEE2016-19 Objective 1: To collect otolith samples in the fish processing sheds of the commercial landings of hoki from Cook Strait and inside the line on the WCSI during winter 2016.

DEE2016-20 Objective 1: To determine the catch-at-age for commercial catches and resource surveys of specified middle depth and deepwater fishstocks. For hoki these include calculating WCSI spawning, Cook Strait spawning, Eastern non-spawning, and Western non-spawning catch-at-age.

### 1.2 Stock structure

The hoki catch is currently managed under a single TACC which can be caught in all areas of the EEZ excluding QMA 10 (Fishstock HOK 1). However, since 1990 the Hoki Working Group has assessed hoki as two stocks, “eastern” and “western” (Annala (1990) and subsequent Plenary Reports). Hoki on the west coast of the North and South Islands and in the area south of New Zealand, including Puysegur Bank, Snares Shelf, and Campbell Plateau, are assumed to be one stock unit, the “western stock”. The east coast of the South Island, Mernoo Bank, Chatham Rise, Cook Strait, and the east coast of the North Island up to North Cape are assumed to contain the “eastern stock”. Immature hoki (2–4 years old) from both “stocks” occur together on the Chatham Rise.

Livingston (1997) reviewed the two-stock hypothesis originally adopted in 1990 (Livingston 1990) with respect to data collected in 1990–97, and concluded that this hypothesis was still a valid interpretation for hoki. Morphometric and ageing studies (Horn & Sullivan 1996, Livingston & Schofield 1996) found consistent differences between adult hoki from the two main dispersed areas (Chatham Rise and Southern Plateau), and from the two main spawning grounds in Cook Strait and west coast South Island (WCSI), which suggested that there were two sub-populations of hoki. It is not known if differences between the two sub-populations, are the result of genetic, environmental, or some other factor. The chemistry of otoliths from the WCSI and Cook Strait stocks was similar (Kalish et al. 1996), and no genetic differences were detected between spawning stocks (Smith et al. 1981, 1996).

The hoki stock assessment model from 2006 to 2007 (Francis 2007, 2008) and since 2012 (McKenzie 2013, 2015a, 2015b, 2016, 2017), had two variants which were associated with different stock structure hypotheses. The ‘base case’ hypothesis assumed natal fidelity: a fish that was spawned in one area will grow up to spawn in the same area (i.e., a fish is ‘eastern’ or ‘western’ from birth). The alternative hypothesis does not assume natal fidelity, so fish spawned in one area can spawn in another area (i.e., a fish chooses to be ‘eastern’ or ‘western’ when it matures). Under both hypotheses, it was assumed that once a fish has spawned it shows site fidelity – it cannot later change spawning grounds. All model

runs from 2008–11 assumed natal fidelity because of technical problems concerning the definition of unfished biomass without this assumption (Francis 2009, McKenzie 2013). These problems were resolved and model runs which do not assume natal fidelity were included as sensitivity runs from then on (McKenzie 2015a, 2015b, 2016, 2017).

Francis et al. (2011) described a pilot study, aimed at determining whether analyses of stable isotopes and trace elements in otoliths could be used to test the stock structure hypothesis and the question of natal fidelity. However, none of the six trace elements or two stable isotopes considered, unambiguously differentiated the two hoki stocks. Two earlier pilot studies appeared to provide weak support for the hypothesis of natal fidelity for the western and eastern spawning stocks. Smith et al. (2001) found significant differences in gill raker counts, and Hicks & Gilbert (2002) found significant differences in measurements of otolith zones between samples of 3 year-old hoki from the 1997 year-class caught on the WCSI and in Cook Strait. However, when additional year-classes were sampled, differences were not always detected (Hicks et al. 2003).

Horn (2011) reviewed the published literature on natal fidelity in relationship to management of hoki. He concluded that, because hoki are an off-shore species, widely dispersed in the non-spawning season, with multiple diffuse spawning areas, it is unlikely that hoki exhibit 100% natal fidelity. Even if natal fidelity is the preferred option for hoki from an evolutionary perspective, it is likely that some proportion of the population would stray routinely. An independent review of the hoki assessment model, commissioned by Ministry for Primary Industries in February 2014, noted that “the extents of natal fidelity are important to identify”, and recommended exploration of a range of model structures (Butterworth et al. 2014).

### **1.3 Description of the hoki fishery**

Since the 1980s the main fishery for hoki operated from late June to late August on the WCSI, where hoki aggregate to spawn. The spawning aggregations begin to concentrate at depths of 300–700 m around the Hokitika Canyon from late June, and further north off Westport later in the season. Fishing in these areas continues into September in some years. In 1988 another fishery developed on large spawning aggregations of hoki in Cook Strait. The spawning season in Cook Strait runs from late June to mid-September, peaking in July and August. Small catches of spawning hoki are taken from other grounds off the east coast of South Island (ECSI), and late in the season at Puysegur Bank. There are also anecdotal reports of spawning hoki being caught near the Snares Islands, Chatham Islands, and several other locations off the east coast of North Island (ECNI).

Outside the spawning season, when hoki disperse to their feeding grounds, substantial fisheries have developed since the early 1990s on the Chatham Rise and in the Sub-Antarctic. These fisheries usually operate at depths of 300–800 m. The Chatham Rise fishery generally has similar catches over all months except in July–September, when catches are lower due to the fishery moving to the spawning grounds. In the Sub-Antarctic, catches have typically peaked in April–June. Out-of-season catches are also taken from Cook Strait and ECNI, but these are small compared to spawning season catches.

From 1986 to 1990 surimi vessels dominated the catches and took about 60% of the annual WCSI catch. However, since 1991, the surimi component of catches has decreased and processing to head and gut or to fillet product has increased, as has “fresher” catch for shore processing. The hoki fishery now operates throughout the year, producing high quality fillet product from both spawning and non-spawning fisheries. Twin-trawl rigs were used in some hoki fisheries since 1998, and trawls made of spectra twine (a high strength twine with reduced diameter resulting in reduced drag and improved fuel efficiencies) were introduced to some vessels in 2007–08. Since 2012–13, precision seafood harvest (PSH) technology has been tested in the hoki fishery. This is a prototype trawl system that aims to enable fish to be landed in much better condition than traditional trawls. The use of PSH in the hoki fishery is moving towards becoming “routine” although its use in high volume spawn aggregations is still very limited.

The Hoki Fishery Management Company introduced a Code of Practice for hoki target trawling in 2001 with the aim of protecting small fish (less than 60 cm). The Code of Practice was replaced by Operational Procedures for Hoki Fisheries, implemented by the Deepwater Group from 1 October 2009. The Operational Procedures aim to manage and monitor fishing effort within four industry management areas, where there are thought to be high abundance of juvenile hoki (Narrows Basin of Cook Strait, Canterbury Banks, Mernoo Bank, and Puysegur Bank). These areas are closed to hoki target trawling by vessels larger than 28 m, with increased monitoring when targeting species other than hoki. There is also a general recommendation that vessels move from areas where catches of juvenile hoki (defined as less than 55 cm total length) comprise more than 20% of the hoki catch by number.

## 1.4 Catch history

The total annual catches of hoki within the EEZ from 1969 to 2015–16 are given in Tables 1 and 2. The hoki fishery was developed by Japanese and Soviet vessels in the early 1970s (Table 1). Catches increased to 100 000 t in 1977, but dropped to less than 10 000 t in 1978 when the 200 n. mile Exclusive Economic Zone (EEZ) was declared and a quota limit of 60 000 t was introduced (Figure 1). Hoki remained a relatively small fishery of up to 50 000 t a year until 1986, when the TACC was increased. The fishery expanded to an estimated catch of about 255 000 t in 1987–88 (Table 2). Reported annual catches ranged between 175 000 and 215 000 t from 1988–89 to 1995–96, increased to 246 000 t in 1996–97, and peaked at 269 000 t in 1997–98, when the TACC was over-caught by 19 000 t. The TACC was reduced to 90 000 t by 2007–08 and catches declined accordingly (Table 2). The TACC was increased in five steps since 2009–10 and catches have increased. The TACC for 2015–16 was 150 000 t, reduced from the 160 000 t TACC in 2014–15.

Catches by area since 1988–89 are given in Table 3 and Figure 2. The pattern of fishing has changed markedly since 1988–89 when over 90% of the total catch was taken in the WCSI spawning fishery. This was due to a combination of TAC changes and redistribution of fishing effort. The catch from the WCSI declined steadily from 1988–89 to 1995–96, increased again to between 90 000 and 107 000 t from 1996–97 until 2001–02, then dropped sharply to a low of 20 500 t in 2008–09. The WCSI catch decreased to 69 900 t in 2015–16, after increasing over the previous six years. This was about 50% of the total hoki catch in 2015–16, making the WCSI the largest fishery in New Zealand for the six most recent years (Table 3). In Cook Strait, catches peaked at 67 000 t in 1995–96, declined to 14 900 in 2010–11, but then increased to 20 100 t by 2014–15, and decreased to 18 400 t in 2015–16. Non-spawning catches on the Chatham Rise peaked at about 75 000 t in 1997–98 and 1998–99, decreased to a low of 30 700 t in 2004–05, increased again to 40 100 in 2014–15, and decreased slightly to 36 700 t in 2015–16. The Chatham Rise was the largest hoki fishery from 2006–07 to 2009–10, but contributed only about 27% of the total catch in 2015–16. Catches from the Sub-Antarctic peaked at over 30 000 t from 1999–2000 to 2001–02, declined to a low of 6200 t in 2004–05, increased slowly to 19 900 t in 2013–14, and decreased to 6600 t in 2015–16. Catches from other areas remained at relatively low levels (Table 3).

From 1999–2000 to 2001–02, there was a redistribution in catch from eastern stock areas (Chatham Rise, ECSI, ECNI, and Cook Strait) to western stock areas (WCSI, Puysegur, and Sub-Antarctic) (Figure 2). This was initially due to industry initiatives to reduce the catch of small fish in the area of the Mernoo Bank, but from 1 October 2001 was part of an informal agreement with the Minister of Fisheries that 65% of the catch should be taken from the western fisheries to reduce pressure on the eastern stock. This agreement was removed in 2002–03 following the 2003 hoki assessment, which indicated that the eastern hoki stock was less depleted than the western stock, and effort was shifted back into eastern areas, particularly Cook Strait. From 2004–05 to 2006–07 there was a further agreement with the Minister that only 40% of the catch should be taken from western fisheries. From 1 October 2007 the target catch from the western fishing grounds was further reduced to 25 000 t within the overall TACC of 90 000 t. This target was exceeded in both 2007–08 and 2008–09, with about 30 000 t taken from western areas. In 2009–10, the target catch from the western fishing grounds was increased to 50 000 t within the overall TACC of 110 000 t, and



catches were at about the industry-agreed catch split. The target western catch was further increased to 60 000 t in 2010–11 (within the overall TACC of 120 000 t), to 70 000 t in 2011–12 and 2012–13 (overall TACC of 130 000 t), to 90 000 t in 2013–14 (overall TACC 150 000 t), to 100 000 t in 2014–15 (overall TACC 160 000 t), and then decreased to 90 000 t in 2014–15 (overall TACC 150 000 t). The split between eastern and western catches was within 2000 t of the management targets since 2011–12, except in 2014–15 where the eastern catch was 4600 t over the target, and in 2015–16 where the western catch fell short of the target total by 13 400 t.

## **1.5 Recent hoki research**

McKenzie (2017) reported the stock assessment carried out in 2016, using the Bayesian model developed in 2002 (Francis et al. 2003) and implemented in the general-purpose stock-assessment program CASAL (Bull et al. 2012). The Ministry for Primary Industries Deepwater Fisheries Assessment Working Group agreed on a single base model run. In this base model, the problem of the lack of old fish in both fishery-based and survey-based observations was dealt with by allowing natural mortality to be age dependent. For the previous assessment, the fits to the Sub-Antarctic trawl series were improved by using two catchabilities instead of one but for the 2017 assessment a single catchability was used, but with a higher estimated process error for the trawl survey. The western stock was estimated to be 40–79%  $B_0$  and the eastern stock 44–75%  $B_0$  (values are 95% CIs for the base case). The western stock experienced an extended period of poor recruitment from 1995 to 2001 inclusive. Western recruitment was near or just below average from 2002 to 2009; below average in 2010, 2012 and 2013; and well above average in 2011 and 2014 (McKenzie 2017).

Castillo-Jordan et al. (2016) reviewed recruitment patterns in 30 southern hemisphere stocks, including New Zealand hoki, using fisheries data from 1980 to 2010. Cluster and dynamic factor analysis showed a detectable degree of synchrony between recruitment indices for the hakes and lings from Australia, New Zealand, Chile, and South Africa. They also showed that stock recruitment patterns for hoki were strongly correlated with the Interdecadal Pacific Oscillation (IPO) and Southern Oscillation Index (SOI) (Pearson correlation coefficients,  $r = 0.50$  and  $r = -0.44$  respectively).

Simmons et al. (2016) published a contentious report that suggested widespread unreported catch and discarding in New Zealand fisheries, including the hoki fishery, from 1950–2010.

The only new fisheries-independent estimates of hoki abundance since the 2016 hoki assessment was the Sub-Antarctic trawl survey carried out in November–December 2016. Results from this survey are summarised in Section 3.1. A trawl survey was also carried out on the WCSI in August 2016 (O’Driscoll & Ballara submitted), but trawl estimates from this survey are not considered to provide reliable indices of hoki abundance. Results from previous hoki surveys published in the past year were the 2015 Cook Strait acoustic survey (O’Driscoll et al. 2016), and the 2016 Chatham Rise trawl (Stevens et al. 2017).

## **2. HOKI FISHERY 2015–16**

### **2.1 Catch and effort information**

#### **2.1.1 Total Allowable Commercial Catch (TACC) and other management controls**

In the 2015–16 fishing year the TACC for HOK1 was 150 000 t. This TACC applied to all areas of the EEZ except the Kermadec FMA which had a TACC of 10 t. There was an agreement with the Minister for Primary Industries that no more than 90 000 t of the TACC should be taken from western stock areas. The TACC for the 2016–17 fishing year remains at 150 000 t.

Chartered vessels may not fish inside the 12-mile Territorial Sea and there are various vessel size restrictions around some parts of the coast. On the WCSI, a 25-mile line closes much of the hoki spawning area in the Hokitika Canyon and most of the area south to the Cook Canyon to vessels over 46 m overall length. In Cook Strait, the whole spawning area is closed to vessels over 46 m overall length.

## 2.1.2 Catch

The overall catch of 136 719 t was about 25 000 t lower than the catch in 2014–15, and about 13 000 t lower than the TACC (see Table 2). The total estimated catch from catch-effort-and-landing-return (CELRL), lining-catch-effort-return (LCER), net-catch-effort-and-landing-return (NCELRL), trawl-catch-effort-return (TCER), lining-trip-catch-effort-return (LTCER), tuna-long-lining-catch-effort-return (TLCER), and trawl-catch-effort-and-processing-return (TCEPR) data was 136 087 t. A small amount of data may not have been entered into the database as the data extraction was done in mid-December 2016. Estimated catches were scaled up to the total monthly harvest return MHR catch of 136 719 t because estimated catches did not match the (MHR) catch.

Relative to 2014–15, catches in 2015–16 decreased in all areas except for ECSI and ECNI (Figure 2a, Table 3). The WCSI was the largest fishery for the sixth consecutive year, with the catch decreasing by 10 000 t to 68 900 t in 2015–16. Catches inside the 25 n. mile line were 15 600 t and made up 23% of the total WCSI catch in 2015–16, an increase from 2014–15, but down from a peak of 41% in 2003–04 (Table A1a). The Chatham Rise was the second largest hoki fishery, with 36 700 t taken from this area in 2015–16, down 3000 t from the 2014–15 level. The catch from Cook Strait of 18 400 t was down by about 1900 t from that in 2014–15, and at a level similar to that in 2013–14. The 6600 t catch from the Sub-Antarctic in 2015–16 was about 10 000 t lower than in 2014–15 (see Table 3). Catches from Puysegur decreased to 800 t, whereas catches from ECNI and ECSI increased by 100 and 500 t respectively. Overall, about 76 600 t of the total catch in 2015–16 was taken from western stock areas, with 60 100 t from the eastern stock areas (Figure 2a). The western catch was well below the industry-agreed western catch limit of 90 000 t.

### Catch recording

Over 92% of the hoki catch was recorded on the TCEPR form (126 300 t), with the WCSI and Cook Strait the only areas where a substantial amount of catch was recorded on the TCER form (Table A1, Figure 2b). Most hoki catch on the WCSI and in Cook Strait was taken by midwater trawling, whereas most catch on the Chatham Rise and Sub-Antarctic was taken by bottom trawling (Figure 2b).

### Target fishing

Up until 2003–04 almost all of the hoki catch was from target hoki tows. Hoki targeting then decreased on the Sub-Antarctic, WCSI and Chatham Rise until 2008–09, when only 86% of the overall hoki catch was from tows targeting hoki (Figure 3). With the increases in TACC from 2009–10, hoki targeting has again increased, and in 2015–16, 95% of the overall catch was taken from hoki target tows (98% of the hoki catch on the WCSI, 86% on the Sub-Antarctic, and 93% on the Chatham Rise). Cook Strait has remained almost exclusively a hoki target fishery.

### Season

A high proportion of the hoki catch in 2015–16 was taken during the spawning season from June to September (Figure 4). Peak catches on the WCSI spawning grounds were in July and August, as in previous years (Figure 5). In the 2016 season there was some fishing on the WCSI in May (particularly inside the 25 n. mile line), with high catches through to mid-August, but the season ended early i.e., in September (as it did in 2015) compared to the previous three seasons (Figure 4b, Figure 5). In Cook Strait, peak catches were from mid-July to mid-September, with about 2900 t caught outside the spawning season (Figure 5). The seasonal pattern of fishing in Cook Strait was similar to that in previous years, but catches were lower in early August 2016 and higher in early-September 2016 compared to those in the 2015 season (Figure 4b, Figure 5). Fishing during the spawning season on the ECSI occurred mainly in September. Fishing at Puysegur was mainly in June, August and September (Figure 5). Outside the spawning fisheries most of

the catch was taken from October 2015 to June 2016 on the Chatham Rise and in the Sub-Antarctic, with small amounts of catch taken over the rest of the year in these areas (see Figures 4 and 5). Small catches were taken year-round from the ECNI (Figures 4 and 5).

### 2.1.3 CPUE analysis

Unstandardised catch and effort from TCEPR data for the six largest hoki fisheries (WCSI, Cook Strait, Chatham Rise, ECSI, Sub-Antarctic, and Puysegur) are summarised in Appendix 1. Standardised CPUE analyses on tow-by-tow target hoki catches reported on TCEPR for the WCSI, Cook Strait, Chatham Rise, and Sub-Antarctic were also carried out (Appendix 1 and Figure 6). Catch rate analysis did not include data from CELR forms (which account for up to a third of the catch in some years in Cook Strait and some catch from the WCSI, but do not provide tow-by-tow effort data), from TCER forms (which have been in use for only seven years), or from the LCER, LTCER, TLCER or NCELR forms. Standardised CPUE analyses using observer tow-by-tow target hoki catches for the WCSI, Cook Strait, Chatham Rise, and Sub-Antarctic were also carried out.

Standardised analyses were carried out only to explore trends in catch rate. CPUE indices are not believed to provide reliable estimates of hoki abundance and were not used in the 2016 hoki stock assessment (McKenzie 2017). Changes in fishing methods (e.g., use of twin trawls), fishing practices (e.g., target fishing, use of escapement panels on smaller boats, incorrect recording of tow duration as some vessels leave the catch in the water until ready to process, changes in target bag size to reflect the processing capacity of the vessel and improve the quality of fish product), and the reliability of gear parameters recorded on the fishing returns are problems for CPUE analyses. There are also other effects on catching ability that cannot be quantified, such as improvements or changes in net and bottom rig design, and electronic equipment. Precision seafood harvesting (PSH) trawls were identified in TCEPR and observer data based on information from the Ministry for Primary Industries and were excluded from unstandardised and standardised analyses.

#### Model

A lognormal linear model was used for all standardised analysis models, following Dunn (2002). A forward stepwise Generalised Linear Model (Chambers & Hastie 1991) implemented in R code (R Development Core Team 2016) was used to select variables in the model. *Fishing year* was forced into the model as the first term, and the algorithm added variables based on changes in residual deviance. The explanatory power of a particular model is described by the reduction in residual deviance relative to the null deviance defined by a simple intercept model. Variables were added to the model until an improvement of less than 1% of residual deviance explained was seen following inclusion of an additional variable. Variables were either categorical or continuous. Categorical variables offered to the model included *vessel key*, *primary method*, *month*, *vessel experience* (number of years vessel participated in the fishery), *twin vessel* (true/false variable for a vessel that has used a twin trawl), and *statistical area*. Continuous variables included *fishing duration*, *fishing distance* (calculated from positions at start and end of tow), *distance 2* (calculated as *fishing duration* × *speed*), *start latitude*, *start longitude*, *start time*, *mid time* (mid time of tow), *depth of bottom*, *effort depth* (depth of net), *depth above bottom* (*depth of bottom* minus *effort depth*), *effort width* (wing spread), *day of season*, and *effort height* (headline height). Model fits to continuous variables were made as third-order polynomials, though a fourth-order polynomial was also offered to the models for duration. As the WCSI dataset included both midwater and bottom tows, nested effects between *method* and *effort duration*, *effort depth*, *effort height*, *effort speed*, *depth above bottom* and *effort width* were investigated. The dependent variable was the log-transformed estimated catch per tow, with positive catches retained and zero hoki catches excluded.

A vessel variable was incorporated into the CPUE standardisation to allow for differences in fishing power between vessels. A subset of “core” vessels was chosen for each analysis, with vessels not involved in the fishery for a certain number of years (varied by analysis) and with a minimum level of annual effort excluded because they provided little information for the standardisations, which could result in model over-fitting (Francis 2001b). TCEPR data were investigated for level of catch and effort for different years of vessel participation in the fishery, and “core” vessels were defined as those which reported approximately 80% of hoki catches in the defined fishery. For observer data (where there were fewer tows), core vessels were defined as those that were in the fishery for at least two years, and contributed more than 35 observed tows.

The standardised indices were calculated using GLM, with associated standard errors. Indices were presented using the canonical form (Francis 1999) so that the year effects for an area were standardised to have a geometric mean of 1. The CVs represent the ratio of the standard error to the index. The 95% confidence intervals were also calculated for each index.

The influence of each variable accepted into the lognormal models was described by influence plots (Bentley et al. 2012). They show the combined effect of (a) the expected log catch for each level of the variable (model coefficients) and (b) the distribution of the levels of the variable in each year, and therefore describe the influence that the variable has on the unstandardised CPUE and which is accounted for by the standardisation.

Fits to the model were investigated using standard residual diagnostics. For each model, a plot of residuals against fitted values and a plot of residuals against quantiles of the standard normal distribution were produced to check for departures from the regression assumptions of homoscedasticity and normality of errors in log-space (i.e., log-normal errors).

On the WCSI, CPUE models were run for midwater and bottom tows that targeted hoki, with sensitivity runs using other target species (“main target species”), for midwater and bottom trawls separately, and included TCER tows that targeted hoki. For Cook Strait CPUE models were for midwater tows that targeted hoki, with a sensitivity run using both TCEPR and TCER data. For the Chatham Rise and ECSI, and Sub-Antarctic, CPUE models were run for bottom tows that targeted hoki, with a sensitivity run using main target species. An analysis of hoki target tows in Statistical Areas 602 and 603 was also run for the Sub-Antarctic as this was a consistently fished area. Selected explanatory variables for target hoki runs are listed in Table 4.

### WCSI

Unstandardised catch rates are presented for both midwater and bottom trawls (Table A2). Midwater trawl catches accounted for 65% of the total spawning season catch in 2015–16. The unstandardised catch rate from all non-zero midwater tows in 2015–16 decreased from 2014–15, and dropped to the seventh highest in the series, with a median catch of 5.0 t per hour, and a median tow duration of 2.2 hours. As most of the catch and tows were from target hoki tows, catch rates and median tow duration were the same for target hoki tows. Catch rates in bottom trawls were lower than those for midwater trawls, with median catch rates of 3.4 t per hour for all non-zero hoki catches, although catch rates were higher for midwater tows at 5.1 t per hour for target hoki tows. Median tow duration of bottom trawls decreased slightly in 2015–16, to 3.7 hours for all target species, and 3.2 hours for target hoki tows. From 1999–2000 to 2003–04, standardised catch rates from non-zero tows showed a similar decline to non-standardised catch rates. However, standardised indices increased at a much higher rate than unstandardised indices since 2003–04 (Table A3–A4, Figure 6a), although they were similar in 2015–16. All datasets showed similar overall trends, although target hoki bottom tows showed an increase in 2015–16, whereas all other datasets showed a decrease.

### Cook Strait

Midwater trawl catches accounted for 96.7% of the spawning season catch of 11 351 t reported on TCEPR forms in 2015–16. A further 4108 t of catch was reported on TCER forms (see Figure 2b). PSH catch accounted for nearly 3000 t. Non-standardised catch rates continued to be high, with an increase in median catch rate from 20.7 to 26.2 t per hour in non-zero mid-water tows from 2014–15 to 2015–

16 (equivalent to a median catch of 16.8 t per tow), and with the same median tow duration to 0.7 hour for both years. Overall, the non-standardised catch rates showed a slight increase from 1989–90 to 2001–02 and have been flat since then, and standardised catch rates showed no trend (Table A3–A4, Figure 6). Catch rates appear to reflect a fishing strategy where vessels limit the size of catches to maintain fish quality.

#### Chatham Rise

Over 92% of the catch in 2015–16 was taken in bottom trawls, with most of the catch reported on TCEPR forms (see Figure 2b), and PSH tows accounted for nearly 6% of the catch. There was a general increase in tow duration since the 1990s, with a median tow duration of 4.6 hour in 2015–16. The median non-standardised catch rate in bottom trawls in 2015–16 of 1.2 t per hour was similar to that in the previous seven years. The catch rate in hoki target trawls increased from 0.6 t per hour in 2002–03 to 1.7 t per hour in 2008–09, and levelled off to 1.4–1.6 t per hour from 2009–10 to 2015–16. Standardised catch rates generally decreased from 1991–92 to 2003–04, increased to 2008–09 and levelled off, but decreased in 2015–16 (Table A3–A4, Figure 6a).

#### Sub-Antarctic

Bottom trawl catches reported on TCEPR forms accounted for 99% of the catch taken in 2015–16, with midwater trawling accounting for the balance (see Figure 2b). Median tow duration in 2015–16 remained at 5.1 hours, but non-standardised catch rates in bottom trawls decreased to 0.3 t per hour. Catch rates for hoki target bottom trawls (1.1 t per hour in 2015–16) were higher than those for all target trawls, and only slightly lower than those on the Chatham Rise. Standardised catch rates generally decreased from 1996–97 to 2003–04, increased to much higher levels since, but decreased in 2015–16 (Figure 6a). Observed vessels had lower CPUE indices from 2010–11 to 2012–13 (Figure 6a).

#### Other fisheries

Spawning season catches from the ECSI were mainly reported on TCEPR forms (see Figure 2b). Midwater tow target hoki catch rates in 2015–16 decreased to 5.8 t per hour, and bottom tow catch rates decreased to 1.2 t per hour. Spawning season catches from Puysegur were also mainly reported on TCEPR (see Figure 2b), and midwater and bottom tow target hoki catch rates in 2015–16 decreased to 1.6 t and 2.9 t per hour respectively.

#### CPUE trends

Standardised CPUE indices for WCSI, Chatham Rise, and Sub-Antarctic all showed similar trends: decreased from 1991–92 to 2003–04 and increased since then, although Chatham Rise indices were flatter than those in the other areas since 2008–09 (Figure 6b). Observer CPUE indices for WCSI, Chatham Rise, and Sub-Antarctic also showed overall similar trends, but were spiky due to fewer data.

### **2.1.4 Bycatch**

Estimates of bycatch in the hoki fishery were determined from data collected by Ministry for Primary Industries observers. For target hoki trawls, the observer data in 2015–16 represented about 46% of vessels, 9% of tows, and 13% of the total catch (Table 5). The bycatch rate (defined as the percentage of the hoki catch) was estimated for the main bycatch species by fishery in Table 6. Other bycatch species were also taken, particularly in the non-spawning fisheries, but bycatch rates for these species were usually less than 1%. Some of the apparent changes in bycatch rates may have been related to changes in observer coverage between years (e.g., Livingston et al. 2002), so the data should be treated with caution. There were changes in the proportion of hoki target catches (see Figure 3, Section 2.1.2), so caution also needs to be exercised when interpreting the definition of the hoki target fishery. A more comprehensive analysis of catch and discards in the hoki, hake, and ling fishery from 1990–91 to 2012–13 was provided by Ballara & O'Driscoll (2015b).

Overall bycatch rates in the spawning areas in 2015–16 were generally low (less than 2%) for most species.

On the WCSI, bottom trawl fishery bycatch rates in 2015–16 were higher than those in 2014–15 for hake (2.7%), ling (3.9%), and silver warehou (2.9%), whereas midwater trawl bycatch rates were similar for hake at 2.2%, and much lower for ling at 2.0%, and silver warehou at 0.2% (Figure 7). In midwater trawls on the WCSI, frofish, hake, and ling increased slightly, and barracouta and jack mackerel decreased, from 2014–15 to 2015–16. As in the past, there was very little bycatch in the midwater Cook Strait fishery (Figure 7), and spiny dogfish and ling had the largest observed bycatch rates (both 0.4% respectively).

In the non-spawning fisheries, bycatch rates in 2015–16 were generally higher than those for spawning fisheries (Figure 7). In the Chatham Rise bottom trawl fishery, bycatch rates for silver warehou (3.0%), hake (1.2%), ling (4.0%), javelinfish (8.9%), rattails (7.5%), spiny dogfish (1.8%) and pale ghost shark (1.0%) all decreased from 2014–15. Of the main Sub-Antarctic bottom trawl bycatch species, bycatch rates increased for hake (5.0%), ling (27%), javelinfish (6.0%), and white warehou (0.9%), and decreased for silver warehou (0.9%), spiny dogfish (3.0%), and rattails (2.5%).

## 2.2 Size and age composition of commercial catches

Data to estimate length frequency distributions in 2015–16 were available from the at-sea Ministry for Primary Industries Observer Programme (OP). Land-based ('market') sampling of landed hoki in Cook Strait and WCSI was also carried out in Cook Strait and WCSI in 2015–16 for Ministry for Primary Industries research project DEE2016/19. The fishing industry observer programme formerly run by the Hoki Fishery Management Company (HMC) was discontinued and no data were provided since 2004–05.

Density plots of all commercial TCEPR and TCER trawls in which hoki was caught in 2015–16 are shown in Figure 8. Observed positions of all tows sampled for hoki length frequency by the OP are provided in the TCEPR plot. Hoki were measured by OP observers from 1251 tows (fewer than in 2012–13 to 2014–15), of which 646 came from the WCSI, 35 from Cook Strait, 332 from the Chatham Rise, 209 from the Sub-Antarctic, 9 from the ECSI (June–September), 10 from Puysegur (June–September), and 10 from ECNI. In Cook Strait and WCSI, 34 and 14 land-based samples were collected respectively by NIWA scientists. Tables 7 and 8 describe timing of sampling in the main areas.

Length frequency distributions were estimated for each of the major fisheries as the weighted (by the catch weight) average of individual length samples. Length frequency data from each area were post-stratified. Data from the WCSI were stratified by area (inside or outside 25 n. miles) and time. Length frequency data from outside and inside the line from May to September were split into weekly or fortnightly time periods and scaled, respectively, to the outside or inside the line catch in each period (Table 8). Adjacent strata were combined if there were few length samples available, e.g., strata 1, 7, 8 and 22 (Table 8). Length frequency data from Cook Strait are normally stratified by month and vessel size. However, because of patchy OP coverage in 2016, Cook Strait samples from June and July were combined for vessels less than 40 m. Stratification for vessels greater than 40 m was by month, although June was further subdivided as the early part of the June catch and samples came from PSH tows (Table 8). A regression tree method (described below) was used to stratify the two non-spawning fisheries.

Catch-at-age from spawning fisheries was estimated using age-length keys derived from otolith ageing. Otoliths were available from the OP and land-based samples. A subsample of 767 otoliths from WCSI (586 from OP samples, and 181 from land-based samples), and 754 otoliths from Cook Strait (25 from OP samples, and 729 from land-based samples) were selected, prepared, and read using the validated technique of Horn & Sullivan (1996) as modified by Cordue et al. (2000). The sub-sample was derived by randomly selecting a set number of otoliths from each of a series of 1 cm length bins covering the bulk of the catch and then systematically selecting additional otoliths to ensure that the tails of the length distribution were represented. The chosen sample sizes approximated those necessary to produce mean weighted CVs of less than 20% across all age classes, in each of the spawning fisheries.

Age-length keys were constructed for each spawning fishery and applied to the total length frequency distribution to produce an age frequency distribution for the catch of each sex separately. Catch-at-age estimates were determined using the 'catch.at.age' software (Bull & Dunn 2002). This software also incorporates data from otolith zone measurements using the consistency scoring method of Francis (2001a) in the age-length key.

Catch-at-age in both the Chatham Rise and Sub-Antarctic fisheries was estimated by sampling directly for age. This continued the approach used since 1998–99 for the Chatham Rise (Francis 2002) and since 2000–01 for the Sub-Antarctic (Ballara et al. 2003). Sampling directly for age is necessary because a single age-length key is not appropriate in non-spawning fisheries. The fisheries are spread over much of the year and there will be substantial fish growth. This means that for any given length the proportions-at-age will change through the fishery. To sample directly for age, observer coverage must be sufficient to provide a random sample of otoliths from the fishery. Francis (2002) suggested that even a sample size of 1200 otoliths may not be sufficient to achieve a target CV of 0.20 in some years.

On the Chatham Rise in 2015–16, 1268 otoliths out of the available 3102 collected from 316 tows that caught over 150 kg of hoki were selected at random for age estimation as follows:

1. Rejected all otoliths from tows that caught less than 1 t of hoki.
2. For tows that caught 1–4 t of hoki, 1 otolith from each tow.
3. For tows that caught 4–6 t of hoki, 3 otoliths from each tow.
4. For tows that caught 6–9 t of hoki, 5 otoliths from each tow.
5. For tows that caught more than 9 t of hoki, 6 otoliths from each tow.

On the Sub-Antarctic in 2015–16, 1217 otoliths out of the available 1896 collected from 202 tows that caught over 150 kg of hoki were selected at random for age estimation as follows:

1. Rejected all otoliths from tows that caught less than 1 t of hoki.
2. For tows that caught 1–3 t of hoki, 5 otoliths from each tow.
3. For tows that caught 3–6 t of hoki, 8 otoliths from each tow.
4. For tows that caught 6–12 t of hoki, 9 otoliths from each tow.
5. For tows that caught more than 12 t of hoki, 11 otoliths from each tow.

Observers also collected 72 Chatham Rise 'non-random' otoliths, but these were not considered for the selected otolith samples. However for the Sub-Antarctic 759 out of 1743 were classified as non-random and these were considered for the selection of otolith samples, otherwise there were not enough otoliths for age estimation after rejecting otoliths from tows catching less than 1 t of hoki.

The method to estimate catch-at-age for the Chatham Rise and Sub-Antarctic followed that of Francis (2002) as modified by Smith (2005). First, the regression tree method (Breiman et al. 1984) was used to stratify the two fishing areas by minimising the weighted least squares of the mean lengths (as a proxy for age) of fish in the observed tows (see Smith (2005) for details). Next, the estimated age frequencies by sex for the observed tows within each stratum were obtained by scaling the otolith ages and sexes up by the estimated numbers of hoki of each sex caught in the tow and averaging over all tows in the stratum. Finally, the number of fish caught in each stratum was estimated from the TCEPR data, and catch-at-age frequencies were calculated as the weighted average, over the strata, of the estimated age frequencies by sex. Numbers of fish were estimated from catch weights using the length-weight relationship of Francis (2003).

Estimates of catch-at-age before 1999–2000 in the Sub-Antarctic and up to 1997–98 on the Chatham Rise were based on an optimised length frequency model (OLF) described in detail by Hicks et al. (2002).

## **2.2.1 Size and age composition in spawning fisheries**

### **West coast South Island**

The length distribution of female hoki in 2016 had two broad modes, while males were strongly unimodal

(Figure 9). The male mode was 75 cm and the smaller of the two female modes at 78 cm were fish from the 2011 year class (age 5). The larger female mode corresponded to older fish from the 2005–09 year classes (Figures 9 and 10). Few hoki less than 65 cm (2012–14 year-classes at ages 2–4) were caught (Figures 9 and 10).

From 2000 to 2004, the sex ratio of the sampled catch was highly skewed (Figure 11a), with many more females caught than males. In 2005–11, as the catch of younger fish increased, the sex ratio reversed with more males than females caught, and in 2012, the catch contained about 50% males and females (Figure 11a). In 2014 and 2015 the percentage of males decreased slightly to 47%, and in 2016 further decreased to 43%. The percentage of hoki aged 7 and older in the catch declined steeply from 68% in 2003–04 to 16% in 2005–06, increased again to 47–49% in 2013 and 2014, but decreased to 43% in 2015 owing to the abundance of the 2011 year-class (Figure 11a). In 2016 the percentage of fish 7 and older increased substantially to 62%. Conversely, the percentage of small fish (less than 65 cm, which is approximately equivalent to ages 3 years and younger) by number in the catch increased from 20% in 2006–07 to 31% in 2008–09, then decreased to 7–14% in 2013–16 (Figure 11b). Some of these small fish were spawning: 32% of the female fish less than 55 cm (i.e., mostly 2 year-olds) were in spawning condition (ripe and running ripe) in 2016, compared to 47% of all fish (Table 9). The spawning state of male hoki is not recorded by observers, but observations from research tows in other areas suggest that a higher proportion of small males than females would be mature.

Previous comparisons showed that in most years there were differences in the length frequency distributions from land-based samples of hoki caught inside the 25 n. mile line and at-sea samples of fish outside this area, with a higher proportion of larger fish (greater than 70 cm) from samples taken inside the line (Ballara & O’Driscoll 2014, 2015a, 2016). In 2016, the observer and land-based sampling data from inside the line in May–September had very few fish less than 70 cm (Figure 12).

The overall mean length of both female and male hoki decreased as the 2016 spawning season progressed (Figure 13). This pattern of declining mean length over the spawning season used to be a common feature of the fishery, but was not observed between 1999 and 2006. The mean length at age for hoki aged from 3–8 years increased from 1988 to about 2005–06, but then decreased (Figure 14).

The OP data used to estimate catch-at-age was representative of the overall spatial, depth, and temporal distribution of the catch in 2015–16, although small vessels and vessels from 65–70 m were not well sampled (Figure 15).

### **Cook Strait**

The length distribution of female hoki in 2016 was broad, while the male length distribution was strongly unimodal with a peak at 71 cm (Figure 16). There was a broad age distribution of females 2–12 years, while most males were aged 5 (2011 year-class) with a group at 2–10 years (Figure 17). Few fish from the 2012 or 2010 year-classes (ages 4 and 6 respectively) were caught in 2016, and only 10% of the catch was fish less than 65 cm (see Figure 11b).

In 2016, the OP data used to estimate catch-at-age was poorly representative of the overall spatial and depth and temporal distribution of the catch (Figures 18 and 19, see Table 8). However, land-based samples were well spread throughout the spawning season (Table 8, Figure 19).

Length frequency distributions by strata showed that the size distribution of the catch was broadly similar in each month, by each sampling method, and by vessel size category (Figure 20). The sex ratio of the catch fluctuated over time, but was dominated by females from 2001–05, and was mostly male-dominated since then (see Figure 11a). The apparent change in sex ratio may be related to a sampling bias, as there is some evidence that larger vessels catch a higher proportion of female hoki in Cook Strait (O’Driscoll et al. 2015b). There was a slight decreasing trend in the mean length of hoki in June and July (Figure 21). As on the WCSI, the mean length-at-age increased from 1988 to the mid-2000s and there is now a slight decreasing trend, especially at ages 4–10 (Figure 22).



The catch-at-age data for 2011–2013 were not used in the 2015 hoki stock assessment model as they were not considered representative of the commercial catch due to poor observer coverage and the rapidly changing sex ratio. The catch-at-age data for 2014–2016 were included in the 2017 assessment model because of improved coverage due to reinstatement of the land-based sampling programme for Cook Strait hoki.

### **Puysegur**

In 2015–16, 10 OP samples were collected during the spawning season, and these were mainly fish of 45–90 cm (Figure 23).

### **East coast South Island**

Nine samples were collected during the 2016 spawning season (Figure 24). Fish from this area were larger than those observed in the non-spawning fishery on the Chatham Rise, especially the females.

## **2.2.2 Size and age composition in non-spawning fisheries**

### **Chatham Rise**

About 93 % of the commercial catch, 83% of length frequency data, and 90% of the available otoliths came from the hoki target fishery in 2015–16 (Figure 25). The tree-based regression split the OP data into three strata based on depth of net, and date (Table 10). The mean length of hoki was smaller in shallower water, and later in the season. The length distribution of hoki in 2015–16 was bimodal and similar for males and females (Figure 26). The catch was dominated by hoki of 45–90 cm, with the small left-hand mode from 45–55 cm representing the 2014 year-class (age 2+), and the stronger right-hand mode from the 2011 year-class (age 4+) peaking at 70 and 72 cm for males and females respectively. The modal age was 5+ (2010 year-class) for females and 4+ (2011 year class) for males (Figure 27). More females than males were caught in 2015–16, and males comprised 39% of the catch (see Figure 11a). There was a lower proportion of large old fish (males and females) than in other areas, with only 20% of the catch aged 7 years or older, and only 30% of these were male (see Figure 11a). About 27% of the catch by number in 2015–16 was less than 65 cm, a decrease from 2012–13 (45%), due to the high numbers of 4+ and 5+ hoki caught (see Figure 11b).

The OP data used to estimate catch-at-age was representative of the overall spatial and temporal distribution of the catch in 2015–16 (Figure 28), although coverage was lower than ideal in January, March and April (Figure 28).

### **Sub-Antarctic**

About 86% of the commercial catch, but only 48% of length frequency data and 75% of the available otoliths came from the hoki target fishery in 2015–16 (Figure 29). The remainder of the otoliths were from tows that targeted jack mackerel, hake, ling, squid, silver warehou, or white warehou. The tree-based regression split the OP data into three strata based on date and latitude (Table 10). As for 2013–14 and 2014–15 data, stratum 1 was further stratified by depth of net at 400 m, as squid and silver warehou target trawls that are in shallower depths and tend to catch smaller hoki could overwhelm catches in the overall length frequency distribution for this stratum – although this did not greatly change the overall length frequency distribution. Smaller fish tended to occur on the Snares Shelf, especially in shallower water, and early on in the season.

The length distribution of hoki in 2015–16 was bimodal and similar for males and females (Figure 30). The catch was dominated by hoki of 40–55 cm from the 2014 year-class (age 1+), and fish from 58–90 cm from ages 2–10 (Figure 31). The modal age of females was 5+ (2010 year-class) and for the males was 4+ (2011 year-class). There was a higher proportion of old fish caught than on the Chatham Rise (see Figure 11a), but the proportion of the catch of fish less than 65 cm was similar to that on the Chatham Rise at 28% (see Figure 11b). About 40% of the fish caught in 2015–16 were males (see Figure 11a).

The OP sampling was not representative of the overall spatial and temporal distribution of the catch (Figure 32). Deeper tows were under-sampled and there was little coverage in Statistical Area 603.

### **Problems with estimation of catch-at-age in non-spawning fisheries**

In addition to the problems associated with whether OP coverage was representative of the catch, there was an ongoing problem with selection of otoliths. Observers collected otoliths from 10 fish out of the 50–150 sampled per tow for length measurement (and three otoliths per tow in the spawning fisheries). As in previous years, a rank sums test showed that the observers tended to select larger fish for extraction of otoliths from the Sub-Antarctic in 2015–16, although the Chatham Rise otolith selection appeared unbiased in 2015–16 (Figure 33). This introduces a bias into the age estimates which is difficult to correct. Electronic aids introduced to help observers take random samples for otoliths should have solved this problem, but this does not seem to have worked.

## **2.2.3 Comparison of size and age composition between the main fisheries**

Length distributions from the main fisheries in 2015–16 are compared in Figure 34. The catch in all areas was mainly fish 50–90 cm. The 2011 year class (68–75 cm) was important in all areas and comprised age 4+ fish on the Chatham Rise and Sub-Antarctic, and age 5 fish in Cook Strait (both sexes) and on the WCSI (mainly as males). In all areas there were few hoki from the 2012 year class. The 2010 year class (age 5+) also appeared strong on the Chatham Rise and in the Sub-Antarctic. The 2014 year class appeared in all the main fishery areas. Most fish on the Chatham Rise were less than 90 cm. Large female fish (over 90 cm) were proportionately more abundant in Cook Strait, ECSI, and WCSI.

## **3. HOKI RESEARCH**

### **3.1 Resource surveys**

#### **3.1.1 Trawl surveys**

##### **Chatham Rise**

No Chatham Rise trawl survey was carried out in January 2017.

##### **Sub-Antarctic**

The seventeenth *Tangaroa* trawl survey of the Sub-Antarctic summer series was conducted from 22 November to 22 December 2016, with previous surveys in 1991–1993, 2000–2009, 2011, 2012 and 2014. An unprecedented amount of survey time was lost due to bad weather (8 days) which meant that survey tows in the core trawl survey strata (strata 1–25) were unable to be completed for the first time in the survey time-series. Only 56 of the planned 82 phase 1 stations were successfully completed, with fewer than 2 stations carried out in some core strata: 1 (1 station), 3A (1 station), and 2 (no stations), and no stations in deep strata 25 and 27.

Abundance index estimates and CVs for core strata with more than two stations surveyed (i.e., strata 2, 3B, and 4–15) were 33 390 t (CV 14.1%) for hoki, up by 7% from the 2014 estimate for the equivalent strata. Estimates were scaled up using a factor based on the proportion of hoki abundance in missing strata (1, 2, 3A) in previous surveys from 2000–14. The scaling factor for hoki was 1.137 with CV = 9.5%. The scaled biomass index of 37 992 t (CV 17.0%) was up 21% from 31 329 t in 2014 (Table 11).

Several modes were present in the hoki scaled length frequency including some 1+ fish at 32–42 cm and relatively few 2+ fish at 50–60 cm (2015 and 2014 year-classes) (Figures 35 and 36). There was a broad mode of older age classes of hoki aged 5–12. Age 1+ and 2+ hoki were concentrated in stratum 3B, south of the Snares Islands. The distribution of these younger year classes was consistent with previous surveys.

## WCSI

The fourth *Tangaroa* trawl survey of the WCSI was carried out 2–20 August 2016, with previous surveys in 2000, 2012, and 2013. Hoki was not a target species for the 2016 survey (O’Driscoll & Ballara in press). The trawl survey was extended deeper in 2016 (from 800 to 1000 m) to improve coverage of deepwater bycatch species. A total of 58 successful random tows were completed in 11 strata north of Hokitika Canyon. Trawl abundance estimates and sampling CVs were estimated for three different areas: ‘core’ (2000 survey area from 300–650 m); ‘all’ (2012–13 survey area from 200–800 m); and ‘deep’ (2016 survey area from 200–1000 m)

Hoki estimates were highly variable between the 2000, 2012, 2013, and 2016 trawl surveys (Table 12). There was a six-fold increase in estimated hoki abundance in the core trawl strata between 2000 and 2012, a halving in 2013, and a further large reduction (by 45%) in 2016. The variability in trawl indices may be explained by changes in vertical availability (O’Driscoll et al. 2015a), and trawl estimates from the northern area do not provide reliable indices of hoki abundance (O’Driscoll et al. 2015b).

Several modes were present in the hoki scaled length frequency distribution (Figure 37), including small hoki at 27–38 cm (age 1 year from the 2015 year-class) and relatively high numbers at 40–58 cm (age 2 years from the 2014 year-class). The main length mode of larger hoki was between 60 and 100 cm. Hoki were also actively spawning throughout the survey period, with 38% of female hoki maturing (stage 3), 16% spawning (stages 4–6) and 18% spent (stage 7) in research catches. A high proportion of hoki caught in deeper strata (greater than 500 m) were spent females.

### 3.1.2 Acoustic surveys

No hoki acoustic surveys were carried out in 2016.

## 4. CONCLUSIONS

The total reported hoki catch in 2015–16 was 136 719 t, 13 000 t lower than the TACC of 150 000 t, and 25 000 t lower than the catch in 2014–15. Catches in 2015–16 decreased in all areas except for the ECSI and ECNI. The WCSI was the largest hoki fishery for the sixth consecutive season, followed by the non-spawning fishery on the Chatham Rise.

Length and age frequency distributions from the commercial fishery show that most of the catch in 2015–16 was fish of 50–90 cm. The 2011 year class (68–75 cm) was important in all areas as age 4+ on the Chatham Rise and Sub-Antarctic, and as age 5 in Cook Strait (both sexes) and on the WCSI (mainly as males). The 2010 year class (age 5+) also appeared strong on the Chatham Rise and in the Sub-Antarctic. In all fisheries there were few hoki from the 2012 year classes. The 2014 year class appeared in all main fishery areas. The percentage of small fish in the catch decreased in the non-spawning fisheries, mainly due to the presence of the strong 2010 and 2011 year class. Most fish caught on the Chatham Rise in 2015–16 were less than 90 cm, and the largest hoki came from the Cook Strait, ECSI, and WCSI fisheries.

The relative biomass index for hoki from the core strata in the 2016 Sub-Antarctic trawl survey increased by 21% from 2014. Several modes were present in the hoki scaled length frequency including some 1+ fish (2015 year-class), relatively few 2+ fish (2014 year-class), and a broad mode of older age classes of hoki ages 5–12. A trawl survey was also carried out on the WCSI in August 2016, but trawl estimates from this survey are not considered to provide reliable indices of hoki abundance.

## 5. ACKNOWLEDGMENTS

This work was funded by Ministry for Primary Industries Research Projects DEE2016-08HOK, DEE2016-19, and DEE2016-20. It also incorporated trawl survey results from Ministry for Primary Industries Research Projects DEE2016-01 and DEE2016-03. Thanks to the many scientific and industry staff who contributed to the collection and analysis of data used in this report. We especially thank Colin Sutton for otolith preparation, and Peter Horn and Debbie Hulston for their ageing work. We also thank Peter McMillan for reviewing this report.

## 6. REFERENCES

- Annala, J.H. (comp.) (1990). Report from the Fishery Assessment Plenary, April–May 1990: stock assessments and yield estimates. 165 p. (Unpublished report held in NIWA library, Wellington.)
- Ballara, S.L.; O’Driscoll, R.L. (2014). Catches, size, and age structure of the 2011–12 hoki fishery, and a summary of input data used for the 2013 stock assessment. *New Zealand Fisheries Assessment Report 2014/05*. 117 p.
- Ballara, S.L.; O’Driscoll, R.L. (2015a). Catches, size, and age structure of the 2013–14 hoki fishery, and a summary of input data used for the 2015 stock assessment. *New Zealand Fisheries Assessment Report 2015/57*. 122 p.
- Ballara, S.L.; O’Driscoll, R.L. (2015b). Fish and invertebrate bycatch and discards in New Zealand hoki, hake, and ling fisheries from 1990–91 until 2012–13. *New Zealand Aquatic Environment and Biodiversity Report No. 163*. 65 p.
- Ballara, S.L.; O’Driscoll, R.L. (2016). Catches, size, and age structure of the 2014–15 hoki fishery, and a summary of input data used for the 2016 stock assessment. *New Zealand Fisheries Assessment Report 2016/40*. 122 p.
- Ballara, S.L.; O’Driscoll, R.L.; Phillips, N.L.; Livingston, M.E.; Smith, M.H.; Kim, S.W. (2003). Catches, size, and age structure of the 2001–02 hoki fishery, and a summary of input data used for the 2003 stock assessment. *New Zealand Fisheries Assessment Report 2003/42*. 77 p.
- Bentley, N.; Kendrick, T.H.; Starr, P.J.; Breen, P.A. (2012). Influence plots and metrics: tools for better understanding fisheries catch-per-unit-effort standardizations. *ICES Journal of Marine Science* 69(1): 84–88.
- Breiman, L.; Friedman, J.H.; Olshen, R.A.; Stone, C.J. (1984). Classification and regression trees. Wadsworth, Belmont, California. 358 p.
- Bull, B.; Dunn, A. (2002). Catch-at-age user manual v1.06.2002/09/12. NIWA Internal Report 114. 23 p. (Unpublished report held in NIWA library, Wellington.)
- Bull, B.; Francis, R.I.C.C.; Dunn, A.; McKenzie, A.; Gilbert, D.J.; Smith, M.H.; Bian, R. (2012). CASAL (C++ algorithmic stock assessment laboratory): CASAL User Manual v2.30-2012/03/21. *NIWA Technical Report 135*. 280 p.
- Butterworth, D.; Hilary, R.; Ianelli, J. (2014). Report on the review of the New Zealand hoki stock assessment model. Report to the Ministry for Primary Industries, 4 March 2014. 17 p. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Castillo-Jordán, C.; Klaer, N.L.; Tuck, G.N.; Frusher, S.D.; Cubillos, L.A.; Tracey, S.R.; Salinger, M.J. (2016). Coincident recruitment patterns of Southern Hemisphere fishes. *Canadian Journal of Fisheries & Aquatic Sciences* 73: 270–278.
- Chambers, J.M.; Hastie, T.J. (1991). Statistical models in S. Wadsworth & Brooks-Cole, Pacific Grove, CA. 608 p.
- Cordue, P.L.; Ballara, S.L.; Horn P.L. (2000). Hoki ageing: recommendation of which data to routinely record for hoki otoliths. Final Research Report for Ministry of Fisheries Research Project MOF1999/01. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Dunn, A. (2002). Updated catch-per-unit-effort indices for hoki (*Macruronus novaezelandiae*) on the west coast South Island, Cook Strait, Chatham Rise, and sub-Antarctic for the years 1990 to 2001. *New Zealand Fisheries Assessment Report 2002/47*. 51 p.
- Francis, R.I.C.C. (1999). The impact of correlations in standardised CPUE indices. New Zealand Fisheries Assessment Research Document 99/42. 30 p. (Unpublished report held in NIWA library,

- Wellington.)
- Francis, R.I.C.C. (2001a). Improving the consistency of hoki age estimation. *New Zealand Fisheries Assessment Report 2001/12*. 18 p.
- Francis, R.I.C.C. (2001b). Orange roughy CPUE on the South and East Chatham Rise. *New Zealand Fisheries Assessment Report 2001/26*. 30 p.
- Francis, R.I.C.C. (2002). Estimating catch at age in the Chatham Rise hoki fishery. *New Zealand Fisheries Assessment Report 2002/9*. 22 p.
- Francis, R.I.C.C. (2003). Analyses supporting the 2002 stock assessment of hoki. *New Zealand Fisheries Assessment Report 2003/5*. 34 p.
- Francis, R.I.C.C. (2007). Assessment of hoki (*Macruronus novaezelandiae*) in 2006. *New Zealand Fisheries Assessment Report 2007/15*. 99 p.
- Francis, R.I.C.C. (2008). Assessment of hoki (*Macruronus novaezelandiae*) in 2007. *New Zealand Fisheries Assessment Report 2008/4*. 109 p.
- Francis, R.I.C.C. (2009). Assessment of hoki (*Macruronus novaezelandiae*) in 2008. *New Zealand Fisheries Assessment Report 2009/7*. 80 p.
- Francis, R.I.C.C.; Haist, V.; Bull, B. (2003). Assessment of hoki (*Macruronus novaezelandiae*) in 2002 using a new model. *New Zealand Fisheries Assessment Report 2003/6*. 69 p.
- Francis, R.I.C.C.; Neil, H.L.; Horn, P.L.; Gillanders, B.; Marriott, P.; Vorster, J. (2011). A pilot study to evaluate the utility of otolith microchemistry for determining natal fidelity in New Zealand hoki. Final Research Report for Ministry of Fisheries Research Project HOK2006/05 Objective 1. 24 p. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Hicks, A.C.; Cordue, P.L.; Bull, B. (2002). Estimating proportion at age and sex in the commercial catch of hoki (*Macruronus novaezelandiae*) using length frequency data. *New Zealand Fisheries Assessment Report 2002/43*. 51 p.
- Hicks, A.C.; Gilbert, D.J. (2002). Stock discrimination of hoki (*Macruronus novaezelandiae*) based on otolith ring measurements. *New Zealand Fisheries Assessment Report 2002/2*. 31 p.
- Hicks, A.C.; Smith, P.J.; Horn, P.L.; Gilbert, D.J. (2003). Differences in otolith measurements and gill raker counts between the two major spawning stocks of hoki (*Macruronus novaezelandiae*) in New Zealand. *New Zealand Fisheries Assessment Report 2003/7*. 23 p.
- Horn, P.L. (2011). Natal fidelity: a literature review in relation to the management of the New Zealand hoki (*Macruronus novaezelandiae*) stocks. *New Zealand Fisheries Assessment Report 2011/34*. 18 p.
- Horn, P.L.; Sullivan, K.J. (1996). Validated ageing methodology using otoliths, and growth parameters for hoki (*Macruronus novaezelandiae*) in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* 30: 161–174.
- Kalish, J.M.; Livingston, M.E.; Schofield, K.A. (1996). Trace elements in the otoliths of New Zealand blue grenadier (*Macruronus novaezelandiae*) as an aid to stock discrimination. *Marine and Freshwater Research* 47: 537–542.
- Livingston, M.E. (1990). Stock structure of New Zealand hoki (*Macruronus novaezelandiae*) New Zealand Fisheries Assessment Research Document 90/8. 21 p. (Unpublished report held in NIWA library, Wellington.)
- Livingston, M.E. (1997). The stock structure of hoki: hypotheses and assumptions revised. (Unpublished report presented to the Hoki Working Group 1997, held by Ministry for Primary Industries, Wellington.)
- Livingston, M.E.; Clark, M.R.; Baird, S.-J. (2002). Trends in bycatch of major fisheries in depths over 200 m on the Chatham Rise, for fishing years 1989/90 to 1998/99. Final Research Report for Ministry of Fisheries Research Project ENV1999/05. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Livingston, M.E.; Schofield, K.A. (1996). Stock discrimination of hoki (*Macruronus novaezelandiae*, Merluccidae) in New Zealand waters using morphometrics. *New Zealand Journal of Marine and Freshwater Research* 30: 197–208.
- McKenzie, A. (2013). Assessment of hoki (*Macruronus novaezelandiae*) in 2012. *New Zealand Fisheries Assessment Report 2013/27*. 65 p.
- McKenzie, A. (2015a). Assessment of hoki (*Macruronus novaezelandiae*) in 2013. *New Zealand Fisheries Assessment Report 2015/08*. 73 p.

- McKenzie, A. (2015b). Assessment of hoki (*Macruronus novaezelandiae*) in 2014. *New Zealand Fisheries Assessment Report 2015/09*. 68 p.
- McKenzie, A. (2016). Assessment of hoki (*Macruronus novaezelandiae*) in 2015. *New Zealand Fisheries Assessment Report 2016/01*. 88 p.
- McKenzie, A. (2017). Assessment of hoki (*Macruronus novaezelandiae*) in 2016. *New Zealand Fisheries Assessment Report 2017/11*. 80 p.
- O'Driscoll, R.L.; Bagley, N.W.; Ballara, S.L.; Lacroit, Y. (2015a). Trawl and acoustic survey of hoki and middle depth fish abundance on the west coast South Island, July–August 2013 (TAN1308). *New Zealand Fisheries Assessment Report 2015/20*. 104 p.
- O'Driscoll, R.L.; Ballara, S.L. (2017). Trawl survey of middle depth fish abundance on the west coast South Island, August 2016 (TAN1609). Draft New Zealand Fisheries Assessment Report held by Ministry for Primary Industries. 76 p.
- O'Driscoll, R.L.; Ballara, S.L.; Bagley, N.W. (2015b). Review of performance of west coast South Island trawl/acoustic survey of deepwater stocks Final Research Report for Ministry for Primary Industries Research Project SEA2014/10. 86 p. (Unpublished report held by Ministry for Primary Industries.)
- O'Driscoll, R.L.; Lacroit, Y.; Dunford, A.J.; MacGibbon, D.J. (2016). Acoustic survey of spawning hoki in Cook Strait during winter 2015 and update of acoustic q priors for hoki stock assessment modelling. *New Zealand Fisheries Assessment Report 2016/44*. 55 p.
- R Development Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <http://www.R-project.org>.
- Simmons, G.; Bremner, G.; Whittaker, H.; Clarke, P.; Teh, L.; Zylich, K.; Zeller, D.; Pauly, D.; Stringer, C.; Torkington, B.; Haworth, N. (2016). Reconstruction of marine fisheries catches for New Zealand (1950–2010). Institute for the Oceans and Fisheries, The University of British Columbia, Working Paper Series, Working Paper #2015 – 87. 60 p.
- Smith, M.H. (2005). Direct estimation of year class frequencies for the non-spawning hoki fisheries with estimates of the coefficients of variation. *New Zealand Fisheries Assessment Report 2005/14*. 26 p.
- Smith, P.J.; Bull, B.; McVeagh, S.M. (2001). Evaluation of meristics characters for determining hoki stock relationships. Final Research Report for Ministry of Fisheries Research Project HOK1999/05 Objective 1. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Smith, P.J.; McVeagh, S.M.; Ede, A. (1996). Genetically isolated stocks of orange roughy (*Hoplostethus atlanticus*), but not of hoki (*Macruronus novaezelandiae*), in the Tasman Sea and southwest Pacific Ocean around New Zealand. *Marine Biology* 125: 783–793.
- Smith, P.J.; Patchell, G.; Benson, P.G. (1981). Genetic tags in the New Zealand hoki *Macruronus novaezelandiae*. *Animal Blood Groups and Biochemical Genetics* 12: 37–45.
- Stevens, D.W.; O'Driscoll, R.L.; Ballara, S.L.; Lacroit, Y. (2017). Trawl survey of hoki and middle depth species on the Chatham Rise, January 2016 (TAN1601). *New Zealand Fisheries Assessment Report 2017/08*. 131 p.

## TABLES

**Table 1: Reported trawl catches (t) from 1969 to 1987–88; 1969–83 by calendar year, 1983–84 to 1987–88 by fishing year (1 October to 30 September). Source, FSU data.**

Year	U.S.S.R.	Japan	South Korea	New Zealand		Total
				Domestic	Chartered	
1969		95				95
1970		414				414
1971		411				411
1972	7 300	1 636				8 936
1973	3 900	4 758				8 658
1974	13 700	2 160		125		15 985
1975	36 300	4 748		62		41 110
1976	41 800	24 830		142		66 772
1977	33 500	54 168	9 865	217		97 750
1978*	2 028 +	1 296	4 580	678		8 581
1979	4 007	8 550	1 178	2 395	7 970	24 100
1980	2 516	6 554		2 658	16 042	27 770
1981	2 718	9 141	2	5 284	15 657	32 802
1982	2 251	7 591		6 982	15 192	32 018
1983	3 853	7 748	137	7 706	20 697	40 141
1983–84	4 520	7 897	93	9 229	28 668	50 407
1984–85	1 547	6 807	35	7 213	28 068	43 670
1985–86	4 056	6 413	499	8 280	80 375	99 623
1986–87	1 845	4 107	6	8 091	153 222	167 271
1987–88	2 412	4 159	10	7 078	216 680	230 339

\* Catches for foreign licensed and New Zealand chartered vessels from 1978 to 1984 were based on estimated catches from vessel logbooks. Few data were available for the first 3 months of 1978 because these vessels did not begin completing these logbooks until 1 April 1978.

+ Soviet hoki catches were taken from the estimated catch records and differ from official Ministry for Primary Industries statistics. Estimated catches were used because of the large amount of hoki converted to meal and not recorded as processed fish.

**Table 2: Reported catch (t) from QMS<sup>1</sup>, estimated catch (t) from Monthly Harvest Return (MHR) data, and TACC (t) for HOK 1 from 1986–87 to 2015–16. Estimated catches included TCEPR and CELR data (from 1989–90), LCER data (from 2003–04), NCELR data (from 2006–07), and TCER and LTCER data (from 2007–08).**

Year	Reported catch	Estimated catch (MHR)		TACC
		Exclude HOKET	Include HOKET	
1986–87	158 171		175 000	250 000
1987–88	216 206		255 000	250 000
1988–89	208 500		210 000	250 000
1989–90	210 000		210 000	251 884
1990–91	215 000		210 000	201 897
1991–92	215 000		215 000	201 897
1992–93	195 000		215 000	202 155
1993–94	191 000		195 000	202 155
1994–95	174 000		190 000	220 350
1995–96	210 000		168 000	240 000
1996–97	246 000		194 000	250 000
1997–98	269 000		230 000	250 000
1998–99	244 500		234 000	250 000
1999–00	242 000		237 000	250 000
2000–01	230 625	229 858	229 862	250 000
2001–02	200 054	195 492	195 506	200 000
2002–03	182 560	184 659	184 668	200 000
2003–04	133 764	135 784	135 787	180 000
2004–05	102 885	104 364	106 189	100 000
2005–06	101 984	104 385	105 965	100 000
2006–07	97 790	101 009	102 861	100 000
2007–08	87 815	89 318	91 045	90 000
2008–09	87 598	88 805	89 475	90 000
2009–10	105 105	107 209	107 209	110 000
2010–11	115 782	118 805	118 805	120 000
2011–12	126 184	130 108	130 108	130 000
2012–13	127 962	131 575	132 618	130 000
2013–14	143 705	146 344	146 344	150 000
2014–15	156 471	161 528	161 528	160 000
2015–16	136 087	136 719	136 722	150 000

1. Discrepancies between QMS data and estimated catches from 1986 to 1990 arose from incorrect surimi conversion factors. The estimated catch in those years was corrected from conversion factors measured each year by Ministry observers on the WCSI fishery. Since 1990 the current conversion factor of 5.8 was used, and the total catch reported to the QMS is considered to be more representative of the true level of catch. From 2000–01 MHR catches were shown including and excluding HOK ET catches (catches outside the EEZ).



**Table 3: Estimated total catch (t) of hoki by area<sup>1</sup>, 1988–89 to 2015–16. Estimated (TCEPR and CELR) catches were scaled to reported (QMR or MHR) catch totals. Data also included LCER (from 2003–04), and NCELR estimated data (from 2006–07), and TCER and LTCER data (from 2007–08).**

Fishing Year	Spawning fisheries					Non-spawning fisheries				Total catch
	WCSI	Puysegur	Cook Strait	ECSI	Sub- Antarctic	Chatham Rise	ECNI	WCNI	Other <sup>2</sup>	
1988–89	188 000	3 500	7 000	-	5 000	5 000	-	-	-	208 500
1989–90	165 000	8 000	14 000	-	10 000	13 000	-	-	-	210 000
1990–91	154 000	4 000	26 500	1 000	18 000	11 500	-	-	-	215 000
1991–92	105 000	5 000	25 000	500	34 000	45 500	-	-	-	215 000
1992–93	98 000	2 000	21 000	-	26 000	43 000	2 000	-	3 000	195 000
1993–94	113 000	2 000	37 000	-	12 000	24 000	2 000	-	1 000	191 000
1994–95	80 000	1 000	40 000	-	13 000	39 000	1 000	-	-	174 000
1995–96	73 000	3 000	67 000	1 000	12 000	49 000	3 000	-	2 000	210 000
1996–97	91 000	5 000	61 000	1 500	25 000	56 500	5 000	-	1 000	246 000
1997–98	107 000	2 000	53 000	1 000	24 000	75 000	4 000	-	3 000	269 000
1998–99	94 562	2 883	45 252	1 977	23 753	73 594	2 315	94	97	244 527
1999–00	102 721	2 880	43 192	2 351	33 772	56 014	1 387	98	4	242 419
2000–01	102 235	6 798	36 298	2 411	30 076	49 847	2 035	147	-	229 847
2001–02	92 719	5 322	23 976	2 971	30 175	39 151	1 147	39	-	195 500
2002–03	73 856	5 948	36 713	7 382	20 199	39 092	929	532	8	184 659
2003–04	45 112	1 158	41 034	2 140	11 635	33 650	880	126	-	135 735
2004–05	32 647	5 501	24 485	4 259	6 337	30 434	516	36	-	104 215
2005–06	38 281	1 457	21 405	653	6 961	34 944	673	8	-	104 382
2006–07	33 328	408	20 113	1 006	7 661	37 813	667	8	-	101 004
2007–08	20 928	308	18 470	2 323	8 708	37 920	640	19	-	89 316
2008–09	20 548	233	17 535	1 054	9 807	39 011	588	25	-	88 801
2009–10	36 349	272	17 880	669	12 275	39 138	618	7	-	107 208
2010–11	48 373	1 176	14 937	1 625	12 655	38 447	1 588	2	-	118 803
2011–12	54 532	1 308	15 859	2 531	15 743	39 246	858	31	-	130 108
2012–13	56 208	955	19 415	3 311	14 095	36 530	1 051	9	-	131 574
2013–14	69 400	778	18 400	2 750	19 927	33 752	1 326	9	-	146 342
2014–15	78 701	1 875	20 125	3 598	16 378	40 071	765	11	4	161 528
2015–16	68 869	1 056	18 378	4 126	6 639	36 714	888	20	-	136 690

<sup>1</sup> Estimated catches by area from TCEPR, CELR, LCER, NCELR, and TCER adjusted pro rata to the total reported (QMS or MHR) catches (excluding HOK ET catches) in Table 2.

<sup>2</sup> Area undefined because of missing positions or statistical areas.

- No catches

**Table 4: Variables retained in order of decreasing explanatory value by each lognormal CPUE model for each fishery area and the corresponding total R<sup>2</sup> value.**

**WCSI: TCEPR tow-by-tow, target hoki**

Variable	R-squared
Year	6.88
Day of year	16.76
Vessel	23.90
Mid time of tow	27.18

**WCSI: Observer catch, target hoki**

Variable	R-squared
Year	6.83
Vessel	15.37
Day of year	24.61
Start time of tow	27.43

**Cook Strait: TCEPR tow-by-tow, target hoki**

Variable	R-squared
Year	1.95
Day of year	16.33
Vessel	22.00

**Cook Strait: Observer catch, target hoki**

Variable	R-squared
Year	4.15
Vessel	11.00
Day of year	16.79

**Chatham Rise: TCEPR tow-by-tow, target hoki**

Variable	R-squared
Year	11.47
Vessel	16.03
Start time of tow	19.32
Duration	22.32
Month	23.57

**Chatham Rise: Observer catch, target hoki**

Variable	R-squared
Year	12.35
Vessel	15.64
Start time of tow	18.27
Duration	21.30
Month	22.73

**Sub-Antarctic: TCEPR tow-by-tow, target hoki**

Variable	R-squared
Year	7.11
Start time of tow	12.79
Vessel	17.04
Day of year	20.65
Duration	23.24
Statistical area	24.91
Depth of net	25.94

**Sub-Antarctic: Observer catch, target hoki**

Variable	R-squared
Year	8.55
Start time of tow	14.22
Vessel	18.04
Duration	20.54
Depth of net	22.84
Statistical area	24.20
Month	26.10

**Table 5: Observer coverage 2015–16 by area, for combined trawl methods. WCSI, Cook Strait and ECSI are for June to September only.**

**(a) All target species tows**

Area	Number of vessels			Number of tows			Catch (t)		
	All	Observed	Percent	All	Observed	Percent	All	Observed	Percent
Chatham Rise	48	15	31.2	5 518	335	6.1	34 559	2 686	7.8
Cook Strait	26	4	15.4	999	35	3.5	14 698	585	4.0
ECNI	47	3	6.4	1 837	10	0.5	672	9	1.3
ECSI	18	4	22.2	314	9	2.9	4 120	160	3.9
Puysegur	16	7	43.8	236	22	9.3	1 056	117	11.0
Sub-Antarctic	33	16	48.5	1 832	209	11.4	6 639	1 092	16.5
WCNI	15	1	6.7	176	1	0.6	20	0	-
WCSI	44	17	38.6	5 329	629	11.8	68 291	12 303	18.0
All areas combined	104	33	31.7	16 241	1 249	7.7	130 055	16 951	13.0

**(b) Target hoki tows**

Area	Number of vessels			Number of tows			Catch (t)		
	All	Observed	Percent	All	Observed	Percent	All	Observed	Percent
Chatham Rise	16	12	75.0	4 005	279	7.0	32 103	2 279	7.1
Cook Strait	20	4	20.0	945	35	3.7	14 680	585	4.0
ECNI	17	2	11.8	212	5	2.4	378	6	1.7
ECSI	14	4	28.6	282	9	3.2	3 497	160	4.6
Puysegur	4	1	25.0	64	6	9.4	707	78	11.1
Sub-Antarctic	10	5	50.0	833	100	12.0	5 730	852	14.9
WCNI	1	1	100.0	1	1	100.0	1	0	-
WCSI	34	16	47.1	4 868	561	11.5	67 181	11 834	17.6
All areas combined	59	27	45.8	11 210	995	8.9	124 277	15 794	12.7

**Table 6: Bycatch rates (in parentheses) on vessels with Observer Programme observers in the hoki fishery for tows that targeted hoki from 1990–91 to 2015–16. The WCSI (bottom and midwater trawls), Cook Strait, and ECSI data covered the spawning season (June–September) only. -, less than 0.1 t (except for Cook Strait 1994–95 and 1996–97, Puysegur 1997–98 to 2008–09, and ECSI 2006–07, for which there were no observer data). Bycatch rates were not calculated where observed hoki catch was less than 100 t. Species chosen were the top eight by observed catch in an area. Species include: BAR, barracouta; CSQ, leafscale gulper shark; FRO, frostfish; GSP, pale ghost shark; HAK, hake; HOK, hoki; JAV, javefinfish; JMA, jack mackerels; LIN, ling; RAT, rattails; RCO, red cod; SND, shovelnose dogfish; SPD, spiny dogfish; SPO, rig; SQU, arrow squid; SWA, silver warehou; and WWA, white warehou. – zero or less than 1 t.**

**(a) WCSI (bottom trawl)**

	Catch in t (% of hoki catch)									
Year	HOK	HAK	JAV	LIN	RAT	SPD	SQU	SWA	Other	
1990–91	1 046	25 (2.4)	1 (0.1)	56 (5.3)	2 (0.2)	8 (0.8)	11 (1.1)	67 (6.4)	67 (3.7)	
1991–92	516	7 (1.4)	1 (0.2)	44 (8.5)	2 (0.4)	5 (1)	13 (2.5)	9 (1.7)	9 (8.5)	
1992–93	3 375	82 (2.4)	1 (0)	79 (2.3)	6 (0.2)	30 (0.9)	4 (0.1)	78 (2.3)	78 (1.8)	
1993–94	1 503	52 (3.5)	4 (0.3)	56 (3.7)	4 (0.3)	28 (1.9)	17 (1.1)	23 (1.5)	23 (3.9)	
1994–95	179	24 (13.4)	1 (0.6)	30 (16.8)	3 (1.7)	7 (3.9)	8 (4.5)	9 (5)	9 (14)	
1995–96	360	48 (13.3)	- (-)	31 (8.6)	2 (0.6)	43 (11.9)	1 (0.3)	26 (7.2)	26 (7.8)	
1996–97	1	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	
1997–98	673	69 (10.2)	3 (0.4)	45 (6.7)	5 (0.7)	15 (2.2)	1 (0.2)	19 (2.8)	19 (4.5)	
1998–99	2 660	244 (9.2)	19 (0.7)	159 (6)	24 (0.9)	67 (2.5)	3 (0.1)	85 (3.2)	85 (4.1)	
1999–00	3 033	438 (14.4)	17 (0.6)	122 (4)	26 (0.9)	35 (1.1)	4 (0.1)	84 (2.8)	84 (5.8)	
2000–01	1 462	54 (3.7)	4 (0.3)	66 (4.5)	6 (0.4)	13 (0.9)	6 (0.4)	57 (3.9)	57 (2.5)	
2001–02	7 493	592 (7.9)	33 (0.4)	306 (4.1)	51 (0.7)	80 (1.1)	39 (0.5)	60 (0.8)	60 (3.7)	
2002–03	2 609	213 (8.2)	17 (0.6)	139 (5.3)	19 (0.7)	28 (1.1)	21 (0.8)	49 (1.9)	49 (5.4)	
2003–04	2 034	335 (16.5)	32 (1.6)	270 (13.3)	20 (1)	28 (1.4)	37 (1.8)	182 (8.9)	182 (14.3)	
2004–05	1 507	74 (4.9)	5 (0.3)	126 (8.4)	5 (0.3)	23 (1.5)	13 (0.9)	74 (4.9)	74 (7.7)	
2005–06	2 242	102 (4.5)	26 (1.2)	141 (6.3)	17 (0.8)	50 (2.2)	18 (0.8)	70 (3.1)	70 (6.4)	
2006–07	1 375	71 (5.2)	12 (0.9)	38 (2.8)	11 (0.8)	7 (0.5)	11 (0.8)	42 (3)	42 (4.5)	
2007–08	1 297	23 (1.8)	8 (0.6)	43 (3.3)	6 (0.5)	28 (2.2)	5 (0.4)	36 (2.8)	36 (2.5)	
2008–09	61	31 (50.8)	2 (3.3)	4 (6.6)	- (-)	4 (6.6)	- (-)	1 (1.6)	1 (16.4)	
2009–10	3 888	67 (1.7)	14 (0.4)	132 (3.4)	9 (0.2)	73 (1.9)	6 (0.2)	41 (1.1)	41 (1.6)	
2010–11	2 961	194 (6.5)	18 (0.6)	154 (5.2)	21 (0.7)	49 (1.6)	6 (0.2)	75 (2.5)	75 (4.3)	
2011–12	5 284	169 (3.2)	21 (0.4)	217 (4.1)	16 (0.3)	136 (2.6)	14 (0.3)	61 (1.1)	61 (1.6)	
2012–13	6 874	865 (12.6)	98 (1.4)	449 (6.5)	82 (1.2)	158 (2.3)	43 (0.6)	102 (1.5)	102 (5.3)	
2013–14	4 996	554 (11.1)	67 (1.3)	278 (5.6)	50 (1.0)	55 (1.1)	34 (0.7)	96 (1.9)	96 (6.5)	
2014–15	4 761	389 (8.2)	72 (1.5)	281 (5.9)	65 (1.4)	48 (1)	25 (0.5)	58 (1.2)	58 (5.9)	
2015–16	6 751	184 (2.7)	58 (0.9)	266 (3.9)	64 (1)	56 (0.8)	19 (0.3)	198 (2.9)	198 (3.9)	

**(b) WCSI (midwater trawls)**

Year	Catch in t (% of hoki catch)								
	HOK	BAR	FRO	HAK	JMA	LIN	SPD	SWA	Other
1990–91	27 606	36 (0.1)	285 (1)	1 548 (5.6)	197 (0.7)	187 (0.7)	35 (0.1)	397 (1.4)	397 (0.6)
1991–92	18 157	57 (0.3)	174 (1)	145 (0.8)	83 (0.5)	97 (0.5)	94 (0.5)	147 (0.8)	147 (0.8)
1992–93	15 720	67 (0.4)	82 (0.5)	289 (1.8)	16 (0.1)	104 (0.7)	27 (0.2)	61 (0.4)	61 (0.4)
1993–94	31 065	29 (0.1)	253 (0.8)	166 (0.5)	128 (0.4)	111 (0.4)	187 (0.6)	591 (1.9)	591 (0.8)
1994–95	25 541	30 (0.1)	188 (0.7)	817 (3.2)	261 (1)	191 (0.8)	186 (0.7)	152 (0.6)	152 (0.7)
1995–96	17 346	85 (0.5)	111 (0.6)	1 361 (7.8)	157 (0.9)	247 (1.4)	272 (1.6)	446 (2.6)	446 (1.8)
1996–97	14 270	10 (0.1)	192 (1.4)	647 (4.5)	89 (0.6)	131 (0.9)	59 (0.4)	422 (3)	422 (0.6)
1997–98	17 981	15 (0.1)	84 (0.5)	1 008 (5.6)	20 (0.1)	281 (1.6)	230 (1.3)	427 (2.4)	427 (1.4)
1998–99	14 768	12 (0.1)	203 (1.4)	781 (5.3)	21 (0.1)	130 (0.9)	151 (1)	136 (0.9)	136 (0.9)
1999–00	15 729	2 (0)	155 (1)	643 (4.1)	6 (0)	169 (1.1)	76 (0.5)	299 (1.9)	299 (1)
2000–01	14 971	90 (0.6)	270 (1.8)	460 (3.1)	3 (0)	196 (1.3)	69 (0.5)	238 (1.6)	238 (1.1)
2001–02	9 175	38 (0.4)	215 (2.3)	868 (9.5)	1 (0)	206 (2.2)	39 (0.4)	65 (0.7)	65 (2.8)
2002–03	7 582	73 (1)	166 (2.2)	315 (4.2)	2 (0)	52 (0.7)	13 (0.2)	47 (0.6)	47 (1.1)
2003–04	6 396	63 (1)	213 (3.3)	482 (7.5)	63 (1)	237 (3.7)	22 (0.3)	87 (1.4)	87 (2.9)
2004–05	5 671	221 (3.9)	262 (4.6)	271 (4.8)	8 (0.1)	155 (2.7)	15 (0.3)	25 (0.4)	25 (3.2)
2005–06	7 283	35 (0.5)	152 (2.1)	302 (4.2)	3 (0)	91 (1.2)	12 (0.2)	26 (0.4)	26 (0.9)
2006–07	8 410	26 (0.3)	175 (2.1)	41 (0.5)	1 (0)	41 (0.5)	23 (0.3)	38 (0.4)	38 (0.8)
2007–08	6 478	6 (0.1)	158 (2.4)	24 (0.4)	1 (0)	29 (0.4)	20 (0.3)	17 (0.3)	17 (1.2)
2008–09	8 733	3 (0)	106 (1.2)	37 (0.4)	24 (0.3)	64 (0.7)	21 (0.2)	58 (0.7)	58 (1.8)
2009–10	7 731	- (-)	17 (0.2)	18 (0.2)	- (-)	30 (0.4)	6 (0.1)	24 (0.3)	24 (0.6)
2010–11	6 595	25 (0.4)	15 (0.2)	37 (0.6)	- (-)	35 (0.5)	13 (0.2)	23 (0.4)	23 (0.7)
2011–12	13 150	2 (0)	43 (0.3)	132 (1.0)	1 (0)	117 (0.9)	108 (0.8)	29 (0.2)	29 (0.6)
2012–13	24 970	3 (0)	162 (0.6)	647 (2.6)	18 (0.1)	372 (1.5)	91 (0.4)	44 (0.2)	44 (0.9)
2013–14	26 989	7 (0)	295 (1.1)	476 (1.8)	1 (-)	331 (1.2)	89 (0.3)	50 (0.2)	50 (1)
2014–15	33 042	164 (0.5)	312 (0.9)	435 (1.3)	62 (0.2)	388 (1.2)	98 (0.3)	75 (0.2)	75 (0.9)
2015–16	21 974	10 (0)	292 (1.3)	486 (2.2)	20 (0.1)	435 (2)	58 (0.3)	51 (0.2)	51 (1.1)

**Table 6: continued.**

**(c) Cook Strait (midwater trawls)**

	Catch in t (% of hoki catch)											
Year	HOK	CSQ	LIN	RAT	RCO	SPD	SPO	SWA	Other			
1992-93	107	- (-)	- (-)	- (-)	- (-)	1 (0.9)	- (-)	- (-)	- (-)	- (-)		
1993-94	495	- (-)	6 (1.2)	- (-)	- (-)	1 (0.2)	- (-)	- (-)	- (-)	- (0.2)		
1995-96	734	- (-)	2 (0.3)	- (-)	- (-)	13 (1.8)	- (-)	- (-)	- (-)	- (-)		
1997-98	3 435	- (-)	7 (0.2)	- (-)	- (-)	55 (1.6)	7 (0.2)	- (-)	- (-)	- (0.2)		
1998-99	3 513	- (-)	16 (0.5)	- (-)	- (-)	76 (2.2)	- (-)	- (-)	- (-)	- (0.2)		
1999-00	3 017	- (-)	9 (0.3)	- (-)	- (-)	103 (3.4)	- (-)	- (-)	- (-)	- (0.1)		
2000-01	4 089	- (-)	15 (0.4)	- (-)	- (-)	84 (2)	- (-)	1 (0)	(0)	1 (0.2)		
2001-02	1 991	- (-)	6 (0.3)	- (-)	- (-)	44 (2.2)	- (-)	- (-)	- (-)	- (0.1)		
2002-03	2 416	- (-)	5 (0.2)	- (-)	- (-)	104 (4.3)	- (-)	- (-)	- (-)	- (0.1)		
2003-04	2 482	- (-)	4 (0.2)	- (-)	- (-)	39 (1.6)	- (-)	- (-)	- (-)	- (0.2)		
2004-05	2 176	- (-)	4 (0.2)	- (-)	- (-)	38 (1.8)	- (-)	2 (0.1)	(0.1)	2 (0.4)		
2005-06	1 080	- (-)	2 (0.2)	- (-)	- (-)	15 (1.4)	- (-)	- (-)	- (-)	- (0.2)		
2006-07	2 102	- (-)	10 (0.5)	- (-)	2 (0.1)	84 (4.0)	- (-)	2 (0.1)	(0.1)	2 (0.2)		
2007-08	3 437	3 (0.1)	8 (0.2)	2 (0.1)	1 (0)	63 (1.8)	- (-)	1 (0)	(0)	1 (0.1)		
2008-09	2 290	- (-)	3 (0.1)	1 (0)	- (-)	27 (1.2)	- (-)	- (-)	- (-)	- (0.1)		
2009-10	3 353	- (-)	4 (0.1)	3 (0.1)	- (-)	27 (0.8)	- (-)	- (-)	- (-)	- (0.2)		
2010-11	1 590	1 (0.1)	- (-)	- (-)	- (-)	13 (0.8)	- (-)	2 (0.1)	(0.1)	2 (0.1)		
2011-12	1 551	1 (0.1)	4 (0.3)	3 (0.2)	1 (0.1)	27 (1.7)	- (-)	7 (0.4)	(0.4)	7 (0.4)		
2012-13	956	- (-)	3 (0.3)	- (-)	- (-)	6 (0.6)	- (-)	- (-)	- (-)	- (0.2)		
2013-14	2 537	6 (0.2)	7 (0.3)	3 (0.1)	1 (0)	24 (1)	- (-)	6 (0.2)	(0.2)	6 (0.6)		
2014-15	320	- (-)	1 (0.3)	- (-)	- (-)	2 (0.6)	- (-)	- (-)	- (-)	- (-)		
2015-16	2 139	- (-)	9 (0.4)	1 (0)	- (-)	9 (0.4)	- (-)	- (-)	- (-)	- (0.1)		

**(d) Puysegur (bottom and midwater trawls)**

	Catch in t (% of hoki catch)									
Year	HOK	CSQ	HAK	LIN	RCO	SND	SPD	SWA	Other	
1990-91	986	- (-)	3 (0.3)	25 (2.5)	2 (0.2)	2 (0.2)	1 (0.1)	1 (0.1)	1 (1.4)	
1991-92	1 028	1 (0.1)	27 (2.6)	431 (41.9)	16 (1.6)	4 (0.4)	4 (0.4)	2 (0.2)	2 (5.3)	
1992-93	530	3 (0.6)	3 (0.6)	80 (15.1)	2 (0.4)	1 (0.2)	- (-)	1 (0.2)	1 (3)	
1993-94	959	- (-)	- (-)	8 (0.8)	5 (0.5)	- (-)	6 (0.6)	7 (0.7)	7 (1)	
1994-95	226	- (-)	- (-)	8 (3.5)	- (-)	- (-)	- (-)	- (-)	- (0.4)	
1995-96	719	- (-)	2 (0.3)	33 (4.6)	3 (0.4)	- (-)	2 (0.3)	3 (0.4)	3 (0.7)	
1996-97	455	1 (0.2)	- (-)	6 (1.3)	- (-)	- (-)	3 (0.7)	3 (0.7)	3 (2)	
1998-99	226	- (-)	4 (1.8)	25 (11.1)	6 (2.6)	1 (0.4)	9 (4)	6 (2.6)	6 (6.2)	
1999-00	370	- (-)	- (-)	25 (6.8)	- (-)	2 (0.5)	7 (1.9)	17 (4.6)	17 (4)	
2000-01	823	- (-)	6 (0.7)	30 (3.6)	4 (0.5)	5 (0.6)	16 (1.9)	221 (26.9)	221 (6.3)	
2001-02	561	- (-)	- (-)	20 (3.6)	1 (0.2)	- (-)	1 (0.2)	34 (6.1)	34 (1.2)	
2002-03	678	2 (0.3)	2 (0.3)	52 (7.7)	- (-)	1 (0.2)	2 (0.3)	25 (3.7)	25 (3.8)	
2003-04	549	- (-)	- (-)	32 (5.8)	1 (0.2)	2 (0.4)	2 (0.4)	14 (2.5)	14 (3.1)	
2004-05	1 237	- (-)	1 (0.1)	20 (1.6)	1 (0.1)	- (-)	11 (0.9)	1 (0.1)	1 (1.4)	
2005-06	478	5 (1.1)	3 (0.6)	105 (22)	1 (0.2)	10 (2.1)	1 (0.2)	26 (5.4)	26 (7.7)	
2006-07	10	- (-)	- (-)	4 (40.0)	- (-)	- (-)	- (-)	- (-)	- (20.0)	
2009-10	31	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	1 (3.2)	1 (-)	
2010-11	1	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	
2011-12	381	6 (1.6)	6 (1.6)	19 (5)	- (-)	2 (0.5)	- (-)	5 (1.3)	5 (2.6)	
2012-13	444	13 (2.9)	12 (2.7)	22 (5)	- (-)	12 (2.7)	1 (0.2)	30 (6.8)	30 (7.7)	
2013-14	69	3 (4.3)	1 (1.4)	6 (8.7)	- (-)	1 (1.4)	- (-)	- (-)	- (5.8)	
2014-15	8	- (-)	- (-)	5 (62.5)	- (-)	- (-)	- (-)	1 (12.5)	1 (-)	
2015-16	163	7 (4.3)	1 (0.6)	7 (4.3)	- (-)	15 (9.2)	- (-)	1 (0.6)	1 (4.9)	

Table 6: continued.

## (e) Sub-Antarctic (bottom trawls)

	Catch in t (% of hoki catch)										
Year	HOK	HAK	JAV	LIN	RAT	SPD	SWA	WWA	Other		
1990–91	1 974	204 (10.3)	17 (0.9)	91 (4.6)	14 (0.7)	3 (0.2)	- (-)	3 (0.2)	3 (10.9)		
1991–92	3 452	332 (9.6)	47 (1.4)	248 (7.2)	39 (1.1)	15 (0.4)	9 (0.3)	35 (1)	35 (5.9)		
1992–93	2 566	509 (19.8)	30 (1.2)	224 (8.7)	21 (0.8)	8 (0.3)	5 (0.2)	22 (0.9)	22 (4.4)		
1993–94	1 118	31 (2.8)	11 (1)	98 (8.8)	10 (0.9)	12 (1.1)	11 (1)	5 (0.4)	5 (5.5)		
1994–95	877	22 (2.5)	14 (1.6)	57 (6.5)	12 (1.4)	15 (1.7)	- (-)	8 (0.9)	8 (6.2)		
1995–96	742	27 (3.6)	9 (1.2)	95 (12.8)	15 (2)	5 (0.7)	8 (1.1)	22 (3)	22 (6.7)		
1996–97	66	8 (12.1)	4 (6.1)	3 (4.5)	3 (4.5)	- (-)	- (-)	- (-)	-(30.3)		
1997–98	1 893	127 (6.7)	66 (3.5)	190 (10)	59 (3.1)	20 (1.1)	3 (0.2)	28 (1.5)	28 (6)		
1998–99	4 727	133 (2.8)	74 (1.6)	256 (5.4)	77 (1.6)	20 (0.4)	26 (0.6)	18 (0.4)	18 (4.8)		
1999–00	5 020	212 (4.2)	186 (3.7)	336 (6.7)	65 (1.3)	47 (0.9)	158 (3.1)	25 (0.5)	25 (6.7)		
2000–01	2 739	87 (3.2)	76 (2.8)	369 (13.5)	50 (1.8)	58 (2.1)	159 (5.8)	26 (1)	26 (7.7)		
2001–02	3 889	154 (4)	308 (7.9)	193 (5)	94 (2.4)	97 (2.5)	35 (0.9)	27 (0.7)	27 (7.7)		
2002–03	2 003	81 (4)	99 (4.9)	363 (18.1)	47 (2.4)	80 (4)	21 (1.1)	20 (1.0)	20 (10.3)		
2003–04	548	37 (6.8)	36 (6.6)	309 (56.4)	16 (2.9)	171 (31.2)	54 (9.8)	13 (2.4)	13 (15.7)		
2004–05	391	24 (6.1)	71 (18.2)	189 (48.3)	15 (3.8)	6 (1.5)	5 (1.3)	10 (2.6)	10 (11.5)		
2005–06	1 170	14 (1.2)	29 (2.5)	118 (10.1)	14 (1.2)	63 (5.4)	68 (5.8)	70 (6)	70 (4)		
2006–07	1 225	16 (1.3)	50 (4.1)	225 (18.4)	18 (1.5)	85 (6.9)	82 (6.7)	85 (6.9)	85 (7.3)		
2007–08	2 670	100 (3.8)	176 (6.6)	1 002 (37.5)	28 (1.1)	30 (1.1)	9 (0.3)	76 (2.8)	76 (10.8)		
2008–09	2 890	93 (3.2)	127 (4.4)	359 (12.4)	40 (1.4)	83 (2.9)	52 (1.8)	39 (1.4)	39 (6.6)		
2009–10	2 905	64 (2.2)	147 (5.1)	232 (8)	85 (2.9)	68 (2.3)	26 (0.9)	30 (1)	30 (7.2)		
2010–11	2 014	34 (1.7)	61 (3)	208 (10.3)	58 (2.9)	105 (5.2)	58 (2.9)	56 (2.8)	56 (8.3)		
2011–12	2 141	46 (2.1)	64 (3)	404 (18.9)	48 (2.2)	46 (2.1)	1 (0)	30 (1.4)	30 (5.4)		
2012–13	6 059	58 (1)	197 (3.2)	647 (10.7)	129 (2.1)	132 (2.2)	226 (3.7)	39 (0.6)	39 (4.7)		
2013–14	5 327	81 (1.5)	225 (4.2)	593 (11.1)	149 (2.8)	240 (4.5)	58 (1.1)	74 (1.4)	74 (7.3)		
2014–15	2 463	27 (1.1)	131 (5.3)	299 (12.1)	120 (4.9)	185 (7.5)	115 (4.7)	6 (0.2)	6 (9.5)		
2015–16	1 739	87 (5.0)	105 (6)	470 (27)	41 (2.4)	53 (3)	16 (0.9)	16 (0.9)	16 (13.3)		

## (f) Chatham Rise and ECSI (excluded ECSI from June–September) (bottom trawl).

	Catch in t (% of hoki catch)									
Year	HOK	GSP	HAK	JAV	LIN	RAT	SPD	SWA	Other	
1990–91	3 323	33 (1)	132 (4)	142 (4.3)	157 (4.7)	100 (3)	24 (0.7)	210 (6.3)	210(15.9)	
1991–92	4 853	24 (0.5)	59 (1.2)	70 (1.4)	144 (3)	129 (2.7)	5 (0.1)	28 (0.6)	28 (7.1)	
1992–93	455	- (-)	46 (10.1)	13 (2.9)	8 (1.8)	7 (1.5)	- (-)	9 (2)	9 (11)	
1993–94	3 526	6 (0.2)	50 (1.4)	76 (2.2)	112 (3.2)	108 (3.1)	16 (0.4)	15 (0.4)	15 (7.5)	
1994–95	1 463	- (-)	23 (1.6)	58 (4)	51 (3.5)	38 (2.6)	7 (0.5)	6 (0.4)	6 (6)	
1995–96	3 477	- (-)	112 (3.2)	103 (3)	131 (3.8)	190 (5.5)	47 (1.4)	128 (3.7)	128 (5.6)	
1996–97	1 027	- (-)	86 (8.4)	38 (3.7)	49 (4.8)	90 (8.8)	6 (0.6)	116 (11.3)	116(10.4)	
1997–98	5 388	- (-)	107 (2)	291 (5.4)	184 (3.4)	297 (5.5)	67 (1.2)	90 (1.7)	90(10.5)	
1998–99	7 745	61 (0.8)	90 (1.2)	339 (4.4)	267 (3.5)	313 (4)	131 (1.7)	84 (1.1)	84 (7)	
1999–00	3 457	68 (2)	62 (1.8)	217 (6.3)	103 (3)	156 (4.5)	135 (3.9)	126 (3.6)	126(11.7)	
2000–01	4 262	117 (2.8)	125 (2.9)	344 (8.1)	214 (5)	291 (6.8)	91 (2.1)	214 (5)	214(14.2)	
2001–02	4 643	117 (2.5)	95 (2)	385 (8.3)	223 (4.8)	385 (8.3)	121 (2.6)	50 (1.1)	50(13.1)	
2002–03	2 260	104 (4.6)	67 (3)	429 (19)	199 (8.8)	330 (14.6)	48 (2.1)	139 (6.2)	139(21.1)	
2003–04	2 384	69 (2.9)	52 (2.2)	250 (10.5)	157 (6.6)	265 (11.1)	58 (2.4)	245 (10.3)	245(16.4)	
2004–05	4 768	104 (2.2)	52 (1.1)	528 (11.1)	177 (3.7)	338 (7.1)	106 (2.2)	134 (2.8)	134(11.6)	
2005–06	5 182	99 (1.9)	49 (1)	396 (7.6)	132 (2.5)	316 (6.1)	93 (1.8)	260 (5)	260 (9.3)	
2006–07	5 533	69 (1.2)	80 (1.4)	500 (9)	155 (2.8)	165 (3)	39 (0.7)	195 (3.5)	195 (7)	
2007–08	5 695	62 (1.1)	77 (1.4)	408 (7.2)	121 (2.1)	323 (5.7)	73 (1.3)	153 (2.7)	153 (9.1)	
2008–09	4 427	28 (0.6)	50 (1.1)	355 (8)	96 (2.2)	289 (6.5)	45 (1)	71 (1.6)	71 (4.9)	
2009–10	5 743	39 (0.7)	73 (1.3)	546 (9.5)	136 (2.4)	437 (7.6)	48 (0.8)	244 (4.2)	244 (6.3)	
2010–11	6 012	51 (0.8)	52 (0.9)	385 (6.4)	143 (2.4)	317 (5.3)	46 (0.8)	222 (3.7)	222 (6)	
2011–12	7 868	69 (0.9)	42 (0.5)	329 (4.2)	185 (2.4)	350 (4.5)	107 (1.4)	237 (3)	237 (5.8)	
2012–13	10 254	116 (1.1)	106 (1)	754 (7.3)	390 (3.8)	822 (8)	96 (0.9)	321 (3.1)	321 (9.8)	
2013–14	8 808	83 (0.9)	88 (1.0)	444 (5)	350 (4)	462 (5.2)	229 (2.6)	389 (4.4)	389 (9.8)	
2014–15	5 025	75 (1.5)	116 (2.3)	581 (11.6)	262 (5.2)	415 (8.3)	94 (1.9)	242 (4.8)	242(11.2)	
2015–16	7 303	74 (1)	85 (1.2)	650 (8.9)	290 (4)	550 (7.5)	129 (1.8)	217 (3)	217 (8.1)	

**Table 6: continued.**

**(g) ECSI, June–September (bottom and midwater trawls).**

Year	Catch in t (% of hoki catch)											
	HOK	HAK	JAV	LIN	RAT	SPD	SWA	WWA	Other			
2000–01	5	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)
2001–02	97	- (-)	- (-)	1 (1)	1 (1)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (1)
2002–03	914	22 (2.4)	6 (0.7)	8 (0.9)	18 (2)	5 (0.6)	20 (2.2)	2 (0.2)	2 (2.7)			
2003–04	939	2 (0.2)	4 (0.4)	4 (0.4)	6 (0.6)	1 (0.1)	1 (0.1)	2 (0.2)	2 (0.8)			
2004–05	280	- (-)	1 (0.4)	1 (0.4)	2 (0.7)	- (-)	- (-)	- (-)	- (0.4)			
2005–06	505	5 (1)	1 (0.2)	- (-)	3 (0.6)	1 (0.2)	35 (6.9)	- (-)	- (0.6)			
2007–08	72	2 (2.8)	2 (2.8)	1 (1.4)	9 (12.5)	- (-)	2 (2.8)	2 (2.8)	2 (4.2)			
2008–09	311	- (-)	- (-)	- (-)	1 (0.3)	- (-)	- (-)	- (-)	- (-)			
2009–10	41	- (-)	1 (2.4)	1 (2.4)	18 (43.9)	- (-)	- (-)	2 (4.9)	2 (4.9)			
2010–11	413	2 (0.5)	- (-)	1 (0.2)	4 (1)	- (-)	- (-)	2 (0.5)	2 (0.2)			
2011–12	355	1 (0.3)	2 (0.6)	1 (0.3)	15 (4.2)	- (-)	10 (2.8)	3 (0.8)	3 (0.6)			
2012–13	1 451	7 (0.5)	3 (0.2)	4 (0.3)	17 (1.2)	4 (0.3)	99 (6.8)	- (-)	- (0.2)			
2013–14	43	3 (7)	1 (2.3)	1 (2.3)	2 (4.7)	- (-)	- (-)	- (-)	- (4.7)			
2014–15	627	7 (1.1)	1 (0.2)	2 (0.3)	- (-)	2 (0.3)	5 (0.8)	- (-)	- (0.5)			
2015–16	249	2 (0.8)	8 (3.2)	5 (2)	17 (6.8)	7 (2.8)	6 (2.4)	5 (2)	5 (6.8)			

**Table 7: Number of 2015–16 hoki length frequency samples and otoliths by observer trips and land-based sampling programme, target species, and monthly timing. Length frequency samples with errors, missing data or outside the sample period (e.g. non-spawning in a spawning area) were removed. Four observer programme length frequency samples were excluded as these were from large vessels inside the 25 n. mile line, which may have position errors. – no data.**

**(a) WCSI observer samples**

Trip	Month	Target species	Number of samples	
			Length	Otoliths
1	May	HOK	19	27
2	Jun/Jul	HOK	39	57
3	Jun	HOK	3	4
4	May/Jul	HOK	3	-
5	Jun/Jul	HAK	30	12
6	Jun/Jul	HAK/HOK	26	8
7	Jun/Jul	HOK	24	40
8	Jun/Jul	HOK	27	43
9	Jun/Jul	HOK	5	1
10	Jul	HAK	2	2
11	Jun/Jul	HOK	21	27
12	Jul	HAK	2	2
13	Jul/Aug	HOK	39	42
14	Jul	HOK	35	47
15	Jul/Aug	HOK	34	30
16	Jul/Aug	HAK/HOK	33	30
17	Jul/Aug	HAK/HOK	14	28
18	Jul/Aug	HOK	19	17
19	Jul/Aug	HOK	15	20
20	Jul/Aug	HOK	26	8
21	Jul/Aug	HOK	16	8
22	Jul/Aug	HOK	16	23
23	Jul/Aug	HOK	20	18
24	Jul/Aug	HOK	11	3
25	Aug	HOK	22	2
26	Aug	HAK/HOK	16	7
27	Aug/Sep	HOK	22	21
28	Aug	HOK	4	-
29	Aug/Sep	HOK	31	24
30	Aug/Sep	HOK	21	17
31	Aug	HOK	14	7
32	Aug/Sep	HOK	4	-
33	Aug	HOK	8	4
34	Aug	HOK	7	5
35	Aug/Sep	HAK/HOK	15	2
36	Sep	HAK/HOK	2	-
37	Sep	HAK	1	-
Total			646	586
		HAK (68), HOK (578) (602 outside, 44 inside)		
Land-based Jun	Jun	HOK	4	51
Land-based Jul	Jul	HOK	4	49
Land-based Aug	Aug	HOK	6	81
Land-based Sep			-	-
Total land-based			14	181
Total			660	767



**Table 7: continued.**

**(b) Cook Strait observer and land-based samples.**

Trip	Month	Target species	Number of samples	
			Length	Otoliths
1			2	-
2	Jun/Aug	HOK	9	11
3	Jun/Aug	HOK	15	-
4	Aug/Sep	HOK	9	14
Observer total	Sep	HOK	35	25
Land-based				
	Jun	HOK	3	55
Land-based	Jul	HOK	10	213
Land-based	Aug	HOK	13	279
Land-based	Sep	HOK	8	182
Land-based total			34	729
Total			69	754

**Table 7: continued.**

**(c) Chatham Rise and ECSI observer data; Chatham Rise includes ECSI non-spawning data. – no data.**

Trip	Month	Target species	Number of length samples		Number of otoliths
			Chatham Rise	ECSI spawning	
1	Oct	SWA	2	-	-
2	Oct	SWA	1	-	-
3	Oct/Nov	HOK	32	-	99
4	Oct	LIN/SWA	2	-	2
5	Nov/Dec	BAR/HOK/SQU	9	-	25
6	Nov	HOK	4	-	14
7	Nov	SSO	1	-	1
8	Nov/Dec	HOK	33	-	79
9	Nov/Dec	HOK	28	-	126
10	Dec	HOK/SWA	30	-	109
11	Jan	BOE	3	-	-
12	Dec	HOK	3	-	17
13	Dec/Jan	HOK	31	-	152
14	Jan	HOK	3	-	-
15	Jan	BOE/ORH/SSO	7	-	-
16	Feb/Mar	HOK/SWA	35	-	145
17	May	HOK	9	-	31
18	Apr/May/Jun	HOK	32	1	181
19	Jun	ORH	1	-	-
20	May/Jun	HOK/SWA/WWA	33	3	157
21	Jun	SQU	2	-	-
22	Aug	ORH	3	-	-
23	Sep	HOK	3	1	12
24	Aug/Sep	BYS/BYX	5	-	-
25	Sep	HOK	-	4	-
26	Sep	HOK	1	-	5
27	Sep	HOK/SWA	22	-	101
Total	-	-	335	9	1256

**Table 7: continued.**

**(d) Sub-Antarctic observer data**

Trip	Month	Target species	Number of samples	
			Length	Otoliths
1	Oct	LIN	6	37
2	Oct	HOK	3	30
3	Oct	HOK	5	49
4	Nov	HOK/SWA	2	10
5	Oct	HAK/WWA	3	10
6	Nov	HAK/SQU/WWA	3	14
7	Dec	HAK	8	44
8	Jan/Feb	JMA/SQU	8	5
9	Jan	HOK	1	9
10	Jan	SQU/SWA	7	19
11	Jan	HAK/SQU/SWA	12	79
12	Jan/Feb	BOE/SSO	4	-
13	Mar	SQU	3	-
14	Feb	HAK/WWA	3	15
15	Feb	SQU	1	-
16	Mar	SQU	1	-
17	Mar	SQU	2	-
18	Mar	SQU	2	-
19	Feb	SWA	1	10
20	Mar	SQU	1	-
21	Mar/Apr	SQU/SWA	10	30
22	Mar/Apr	HOK	10	93
23	Mar	SQU	6	-
24	Mar	SQU	2	-
25	Mar	SQU	2	-
26	Mar/Apr	SQU	5	-
27	Apr	SQU	2	-
28	May	HOK	1	10
29	May/Jun	HOK	42	367
30	Jun	SWA	1	9
31	Jul	SCI	1	-
32	Jul	LIN/WWA	7	33
33	Sep	SBW	2	-
34	Sep	SBW	3	-
35	Sep	HOK	37	342
36	Sep	SBW	2	-
Total	-	-	209	1215

**Table 8: Stratification for the 2016 WCSI and Cook Strait length samples.**

**(a) Stratification of WCSI hoki 2016 fishery length frequency data.**

Stratum	25 n. mile line	Number of length samples			Date	Catch Tonnes
		Date	Observer	Land-based		
1	Inside	12–20 May	10	-	1–19 May	141.6
2	Inside	21–27 May	7	-	20 May – 9 Jul	1 274.5
3	Inside	13–30 Jun	4	4	10–30 Jun	3 611.0
4	Inside	3–12 Jul	1	2	1–14 Jul	2 633.9
5	Inside	20–27 Jul	2	2	15–28 Jul	2 102.1
6	Inside	30 Jul – 11 Aug	11	2	29 Jul – 11 Aug	2 654.7
7	Inside	14–29 Aug	9	4	12 Aug – 30 Sep	3 747.1
8	Outside	8 May – 2 Jun	5	-	8 May – 2 Jun	120.3
9	Outside	3–9 Jun	10	-	3–9 Jun	113.0
10	Outside	10–16 Jun	9	-	10–16 Jun	501.5
11	Outside	17–23 Jun	9	-	17–23 Jun	1 533.4
12	Outside	24–30 Jun	43	-	24–30 Jun	3 274.2
13	Outside	1–7 Jul	63	-	1–7 Jul	4 633.4
14	Outside	8–14 Jul	67	-	8–14 Jul	6 202.4
15	Outside	15–21 Jul	41	-	15–21 Jul	4 800.2
16	Outside	22–28 Jul	47	-	22–28 Jul	5 466.0
17	Outside	29 Jul – 4 Aug	54	-	29 Jul – 4 Aug	5 843.6
18	Outside	5–11 Aug	55	-	5–11 Aug	6 742.4
19	Outside	12–18 Aug	67	-	12–18 Aug	5 847.7
20	Outside	19–25 Aug	77	-	19–25 Aug	5 019.4
21	Outside	26 Aug – 1 Sep	40	-	26 Aug – 1 Sep	2 242.8
22	Outside	2–13 Sep	15	-	2–13 Sep	355.8

**(b) Cook Strait 2016 stratification**

Stratum	Vessel size	Date range	Catch (t)	Number of length samples	
				Land-based	Observer
1	< 40 m	Jun–Jul	2 796.4	7	4
2	< 40 m	Aug	3 108.4	9	21
3	< 40 m	Sep	1 035.9	4	1
4	≥ 40m	Jun 1–12	462.9	-	8
5	≥ 40m	Jun 13–30	380.2	2	-
6	≥ 40m	Jul	1 762.6	4	1
7	≥ 40m	Aug	3 266.0	4	-
8	≥ 40m	Sep	2 619.3	4	-

**Table 9: Percentage of female hoki by observer gonad stages on the WCSI for female fish less than or equal to 55 cm (n = 1398) and female fish greater than 55 cm (n = 33 137) for the 2016 spawning season.**

Stage		Females ≤ 55 cm		Females > 55 cm	
		Number	Percent	Number	Percent
1	Immature and resting	377	27	1 155	3
2	Maturing	522	37	13 560	41
3	Mature/Ripe	350	25	10 765	33
4	Running ripe	101	7	4 589	14
5	Spent	48	3	3 068	9

**Table 10: Strata for the 2015–16 non spawning fisheries based on the tree regression of all data (Observer Programme only), with comparison of the TCEPR, Observer Programme (OP), tows and otolith data sampled by stratum. The catch for OP is the total catch for the observed tows.**

**(a) Chatham Rise**

Stratum	Splitting variables		Mean length (cm)	Hoki catch (t)		No. of tows		No. of otoliths	No. of fish measured
	Depth of net (m)	Date		TCEPR	OP	TCEPR	OP		
1	< 584	31 Mar–30 Sep 2016	66.2	8 974.6	1 168.1	1 348	101	343	10 568
2	< 584	1 Oct 2015–30 Mar 2016	71.7	22 653.0	1 204.9	3 270	172	571	16 855
3	≥ 584	-	79.0	5 079.6	291.1	1 258	59	154	4 600

**(b) Sub-Antarctic**

Stratum	Splitting variables			Mean length (cm)	Hoki catch (t)		No. of tows		No. of otoliths	No. of fish measured
	Latitude	Date	Depth of net (m)		TCEPR	OP	TCEPR	OP		
1A	≤ 48.9° S	-	< 400 m	63.8	183.3	97.1	113	27	76	991
1B	≤ 48.9° S	-	≥ 400 m	71.5	2 175.6	344.6	568	49	302	3 974
2	> 48.9° S	Oct 2015 – Jun 2016	-	78.8	3 178.9	333.9	941	92	432	5 169
3	> 48.9° S	Jul – Sep 2016	-	89.1	1 101.0	316.7	220	41	313	2 943


**Table 11: Relative abundance estimates (t in thousands) of hoki for the core strata (300–800 m), and all strata (200–1000 m) from the surveys in the Sub-Antarctic summer *Tangaroa* time series. 2016 core abundance is corrected for strata with fewer than two tows. The CV is the coefficient of variation as % (in parentheses).**

Survey	Abundance (t)	
	Core	All
1991	81 631 (6.8)	81 816 (6.8)
1992	88 053 (6.1)	88 384 (6.1)
1993	100 629 (9.2)	101 112 (9.2)
2000	55 663 (12.6)	56 407 (12.4)
2001	38 145 (15.5)	39 396 (15.0)
2002	39 890 (13.7)	40 502 (13.5)
2003	14 318 (12.9)	14 723 (12.6)
2004	17 593 (11.8)	18 114 (11.6)
2005	20 440 (12.8)	20 680 (12.7)
2006	14 336 (10.7)	14 747 (10.5)
2007	45 876 (15.8)	46 003 (15.7)
2008	46 981 (13.9)	48 341 (13.6)
2009	65 017 (16.2)	66 157 (16.0)
2011	46 070 (14.7)	46 757 (14.5)
2012	55 739 (15.2)	56 131 (15.1)
2014	31 329 (12.9)	31 727 (12.8)
2016	37 992 (17.0)	-



**Table 12: Relative abundance estimates (t in thousands) of hoki for the core strata (300–650 m), all strata (200–800 m), and deep strata (200–1000 m) from the 2000, 2012, 2013, and 2016 WCSI *Tangaroa* trawl surveys. The 2000 survey abundance estimates were re-calculated using 2012–13 stratum areas. The CV is the coefficient of variation as % (in parentheses).**

Survey	Abundance (t)		
	Core	All	Deep
2000	5 385 (20.6)	-	-
2012	32 495 (24.2)	32 602 (24.1)	-
2013	14 184 (26.9)	14 356 (26.5)	-
2016	7 734 (35.7)	7 797 (35.4)	7 830 (35.3)

## FIGURES




**Figure 1: Total New Zealand hoki catch (tonnes, × 1000) estimated from reported landings for calendar years 1972 to 1983 and fishing years 1983–84 (1984) to 2015–16.**




**Figure 2a: Estimated total catch (t) of hoki by 'stock' area (upper panel) and fishing area (lower panel) from 1988-89 (1989) to 2015-16 (2016). "Eastern" areas include Chatham Rise, east coast South Island (ECSI), Cook Strait, and east coast North Island (ECNI). "Western" areas include west coast South Island (WCSI), Sub-Antarctic, and Puysegur.**







**Figure 2b: Total catches and catches by form type by hoki fishing area and fishing year. All areas (except Cook Strait) also show TCEPR data split by MW (midwater trawl) and BT (Bottom trawl). Sub-Antarctic and Puysegur had very little CELR or TCER data.**



**Figure 3: Hoki catch by target species and area for the 1989–90 to 2015–16 fishing years. Hoki catches by target species include HOK, hoki; HAK, hake; LIN, ling; SWA, silver warehou; SQU, arrow squid; SPE, sea perch; WWA, white warehou; and SCI, scampi.**



**Figure 4a: Hoki catch by month and area for the 2015–16 fishing year (maximum circle size is 30 000 t).**



**Figure 4b: Daily distribution of hoki catch by area (in 5-day bins) by main area for 2011–12 to 2015–16 fishing years.**




Figure 5: Distribution of hoki catch by month and area for the 1989–90 to 2015–16 fishing years.









Figure 5 continued.





**Figure 6a: Model catch, unstandardised geometric, and standardised CPUE indices for core data TCEPR and observed (OP) hoki tows from the WCSI for 1990–2016.**





**Figure 6a continued. Model catch, and unstandardised geometric and standardised CPUE indices for core data TCEPR and observed hoki tows from Cook Strait for 1990–2016. Cook Strait included only midwater tows.**







**Figure 6a continued. Model catch, and unstandardised geometric and standardised CPUE indices for core data hoki tows from the Chatham Rise for 1990–2015. Dataset for Chatham Rise included only bottom tows.**



**Figure 6a continued. Model catch, and unstandardised geometric and standardised CPUE indices for core data hoki tows from the Sub-Antarctic for 1990–2015. Datasets for Sub-Antarctic included only bottom tows.**



**Figure 6b: Comparison of relative standardised CPUE indices from model runs for each area.**



**Figure 7: Bycatch rates on vessels with Observer Programme observers in the hoki fishery for tows targeting hoki from 1990–91 to 2015–16. WCSI (bottom and midwater trawls), Cook Strait, and ECSI data cover the spawning season (June–September) only. No observer data for Cook Strait 1994–95 and 1996–97, Puysegur 1997–98 to 2008–09, and ECSI 2006–07. Bycatch rates not calculated where observed hoki catch was less than 100 t. Species chosen are the top eight by observed catch in an area. Species include: BAR, barracouta; BOE, black oreo; BNS, bluenose; BYS, alfonsino; CSQ, leafscale gulper shark; FRO, frostfish; GSP, pale ghost shark; HAK, hake; HOK, hoki; JAV, javelinfish; JMA, jack mackerels; LDO, lookdown dory; LIN, ling; MDO, mirror dory; ORH, orange roughy; RAT, rattails; RBM, rays bream; RCO, red cod; RIB, ribaldo; RBT, redbait; SBW, southern blue whiting; SCH, school shark; SKI, gemfish; SND, shovelnose dogfish; SPD, spiny dogfish; SPE, sea perch; SPO, rig; SQU, arrow squid; SWA, silver warehou; and WWA, white warehou.**







Figure 7: continued.






**Figure 8: Density plots of all commercial TCEPR and TCER trawls where hoki was caught in the 2015–16 fishing year. TCEPR plot also shows observed positions as black dots.**




**Figure 9: Length frequency distributions of hoki in commercial catches from the west coast South Island spawning fishery from 1989 to 1997 sampled at sea by the Observer Programme. n, number of tows sampled; no., number of fish sampled. Numbers above the histograms mark estimated year-class modes, e.g., 91 = 1991 year-class.**




**Figure 9 continued.** Length frequency distributions of hoki in commercial catches from the west coast South Island spawning fishery from 1998 to 2009 sampled at sea by the Observer Programme. In 2003–05 and 2007–09, Observer Programme data were combined with samples of landings from inside the 25 n. mile line sampled by NIWA. n, number of tows sampled; N, number of landings sampled by NIWA; no., number of fish sampled. Numbers above the histograms mark estimated year-class modes, e.g., 2004 = 2004 year-class.







**Figure 9 continued.** Length frequency distributions of hoki in commercial catches from the west coast South Island spawning fishery from 2010 to 2016. In 2010, 2015, and 2016, Observer Programme data were combined with land-based samples from inside the 25 n. mile line sampled by NIWA. n, number of tows sampled; no., number of fish sampled; N, number of landings sampled by NIWA. Numbers above the histograms mark estimated year-class modes, e.g., 2007 = 2007 year-class.




**Figure 10: Catch-at-age of hoki in commercial catches from the west coast South Island spawning fishery from 1988 to 2016. n, number of fish aged. Black bars show 1987 and 1988 year-classes and dark grey bars show 1991–94 year-classes.**




**Figure 10 continued. Black bars show 1987 and 1988 year-classes and dark grey bars show 1991–94 year-classes, and light grey bars (from 2004 on) represent the 2002 and 2003 year classes.**



**Figure 10 continued.** Light grey bars in the represent the 2003 and 2002 year classes, and dark grey bars represent the 2007–2009 year classes.




**Figure 11a: Percentage of males in the catch, percentage of all fish aged 7 and older in the catch, and percentage of male fish (older than seven) in the catch, by area and fishing year. Legend for middle panel also applies to upper and lower panels.**




**Figure 11b: Percentage of small fish in the catch by area and fishing year. Legend for top panel also applies to other panels.**





**Figure 12: Comparison of length frequency distributions from inside and outside the 25 n. mile line in 2016. Inside the line length frequencies came from fish sampled at sea by the Observer Programme and from fish sampled in processing sheds by the land-based sampling programme, and outside the line length frequencies came from fish sampled at sea by the Observer Programme.**



**Figure 13: Mean length of female (black) and male (blue) hoki taken in commercial catches from the west coast South Island spawning fishery 1987–2016 sampled at sea by the Observer Programme. Dashed lines are a loess fit.**









Figure 13 continued.



**Figure 14: Mean length-at-age of female and male hoki taken in commercial catches from the west coast South Island spawning fishery 1988–2016 sampled at sea by the Observer Programme. Lines are a loess fit. Points with fewer than ten records were excluded.**



**Figure 15: Comparison of WCSI 2015–16 Observer Programme catch coverage with TCEPR catches by day of year, depth, latitude, longitude, and vessel length. If sampling is representative of the fishery, then the blue lines (observed catches) should overlay the black lines (TCEPR catch).**



**Figure 16: Length frequency distributions of hoki in commercial catches from the Cook Strait spawning fishery from 1991 to 2016 sampled by the land-based sampling programme, and at sea by the Observer Programme. n, number of landings sampled; no., number of fish sampled. Numbers above the histograms mark year-class modes, e.g., 91 = 1991 year-class.**






Figure 16 continued:



**Figure 16 continued:**





**Figure 17: Catch-at-age of hoki in commercial catches from the Cook Strait spawning fishery from 1988 to 2016 sampled by the land-based sampling programme, and at sea by observers. 2006 data excluded Nelson land-based samples from vessels of at least 40 m length which sorted their catch at sea. Black bars show 1987 and 1988 year-classes in the 1990–2003 seasons; dark grey bars show 1991–94 year-classes, light grey bars show the 2000 year-class, and black bars show the 2002–2003 year-classes from the 2004 season onwards.**






Figure 17 continued.







**Figure 17 continued.**




**Figure 18: Comparison of Cook Strait 2015–16 Observer Programme catch coverage for TCEPR and TCER catches by day of year, depth, latitude, longitude, and vessel length. If sampling is representative of the fishery, then the blue lines (sampled catches) should overlay the black lines (catches).**



**Figure 19: Cook Strait 2015–16 catch by day for vessels less than 40 m and 40 m or longer during the spawning season, showing timing of Observer Programme samples (black dots), and land-based samples (hollow dots).**



**Figure 20: Comparison of length frequency strata of hoki taken in commercial catches from Cook Strait during 2016. Data from Observer Programme and land-based sampling data. n, number of tows or landings sampled; no., number of fish sampled.**



**Figure 21: Mean length of female (black) and male (blue) hoki taken in commercial catches from the Cook Strait spawning fishery 1989–2016 from landings sampled by the Observer Programme. Lines are a loess fit.**






Figure 21 continued.



**Figure 22: Mean length-at-age of female and male hoki taken in commercial catches from the Cook Strait spawning fishery 1988–2016 sampled at sea by the Observer Programme and NIWA land-based sampling programme. Lines are a loess fit. Points with fewer than ten records were excluded.**



**Figure 23: Length frequency distributions of hoki in commercial catches from the Puysegur spawning fishery from 1989 to 1997, and 1999 to 2016 sampled at sea by the Observer Programme. n, number of tows sampled; no., number of fish sampled.**









Figure 23 continued.




**Figure 23 continued.**



**Figure 24: Length frequency distributions of hoki taken in commercial catches from the ECSI spawning fishery from 2001 to 2016 sampled by the Scientific Observer Programme (2001–2006, 2008–2013, 2015, 2016), combined with Hoki Management Company data (2001 to 2005).. n, number of tows sampled; no., number of fish sampled.**



**Figure 25: Percentage of hoki TCEPR, CELR and TCER catch, hoki length frequency samples, and hoki otoliths collected by the Observer Programme, by target species for the Chatham Rise fishery from 2000–01 to 2015–16. Three-letter codes denote target species: HOK, hoki; ORH, orange roughy; OEO, oreos; SQU, squid; SWA, silver warehou; HAK, hake; SCI, scampi; LIN, ling; BAR, barracouta; SPE, sea perch; Other, all other target species combined.**



**Figure 26: Length frequency distributions of hoki taken in commercial catches from the Chatham Rise fishery from 1990-91 to 2015-16 sampled by the Observer Programme (and combined with Hoki Management Company data in 2000-01 to 2003-04). 2006-07 data included target hoki and hake tows. n, number of tows sampled; no., number of fish sampled.**






Figure 26 continued.




**Figure 26 continued.**




**Figure 27: Proportions at age and sex in the catch from the Chatham Rise fishery estimated by direct ageing of otoliths from 2000–01 to 2015–16. Dark grey bars show 1997–99 year-classes; black bars show 2000–02 year-classes; light grey bars show 2003–05 year-classes.**






**Figure 28: Comparison of Chatham Rise 2015–16 Observer Programme catch coverage with TCEPR catches by day of year, depth, latitude, longitude, and vessel length. If sampling is representative of the fishery, then the blue lines (observed catches) should overlay the black lines (TCEPR catch).**



**Figure 29: Percentages of hoki TCEPR, TCER and CELR catch, hoki length frequency samples, and hoki otoliths collected by the Observer Programme, by target species for the Sub-Antarctic fishery from 2000–01 to 2015–16. Three-letter codes denote target species: HOK, hoki; HAK, hake; SQU, squid; ORH, orange roughy, SSO, smooth oreo; OEO, oreo; SWA, silver warehou; SBW, southern blue whiting; SCI, scampi; LIN, ling; WWA, white warehou; Other, other target species combined.**



**Figure 30: Length frequency distributions of hoki taken in commercial catches from the Sub-Antarctic fishery from 1990-91 to 2015-16 sampled by the Observer Programme (and combined with Hoki Management Company data in 2000-01 to 2004-05). n, number of tows sampled; no., number of fish sampled.**







Figure 30 continued.





**Figure 30 continued.**



**Figure 31: Proportions-at-age and sex in the catch from the Sub-Antarctic fishery as estimated by direct ageing of otoliths from 2000–01 to 2015–16. Dark grey bars show 1997–99 year-classes; black bars show 2000–02 year-classes; light grey bars show 2003–05 year-classes.**




**Figure 32: Comparison of Sub-Antarctic 2015–16 Observer Programme catch coverage with TCEPR catches by day of year, depth, latitude, longitude and vessel length (m). If sampling is representative of the fishery, then the blue lines (observed catches) should overlay the black lines (TCEPR catch).**




**Figure 33: Histograms of ranks of the lengths that yielded 2015–16 Chatham Rise and Sub-Antarctic otoliths relative to the lengths of hoki measured for each tow. If sampling was random then the expected counts are given by the dotted line. The p-value was calculated using the rank-sum test.**





**Figure 34: Length frequency distributions of female and male hoki taken in commercial catches from different areas during the 2015–16 fishing year.**



**Figure 35: Scaled length frequency for hoki from Sub-Antarctic November–December Tangaroa trawl surveys. n, population numbers of fish; n.a., population numbers of fish for all strata; n.c., population numbers of fish for core strata; CV, coefficients of variation in parentheses; no., number of fish measured.**





Figure 35 continued.



**Figure 36: Scaled age frequency (years) for hoki from Sub-Antarctic November–December *Tangaroa* trawl surveys. nf, number of female ages; nm, number of male ages; mean weighted CV in parentheses.**





Figure 36 continued.



**Figure 37: Length frequency distributions by sex of hoki for core (grey), all (white), and deep (black) strata from the 2000, 2012, 2013, and 2016 WCSI August *Tangaroa* trawl surveys. n.d, estimated scaled total number of fish for deep strata; n.a, estimated scaled total number of fish for all strata; n.c, estimated scaled total number of fish for core strata; and CV, the coefficient of variation (in parentheses).**

## APPENDICES

**Table A1a: Number of vessels, tows, and total catch inside and outside the 25 n. mile line off WCSI, by year. Data source was un-groomed non-zero TCEPR, TCER, and CELR data. Year defined as June to October. There were no October data available for 2016. It was assumed that CELR data all came from inside the 25 nautical mile line, and included mid-water and bottom trawl tows reported on the CELR form only.**

Fishing year	Number of vessels					Number of tows				
	TCEPR Outside	TCER Outside	TCEPR Inside	TCER Inside	CELR	TCEPR Outside	TCER Outside	TCEPR Inside	TCER Inside	CELR
1990	79	-	37	-	13	7 989	-	83	-	196
1991	75	-	41	-	17	8 135	-	68	-	302
1992	71	-	25	-	17	6 171	-	47	-	358
1993	64	-	22	-	18	6 886	-	108	-	511
1994	69	-	30	-	18	8 463	-	137	-	425
1995	65	-	36	-	21	8 521	-	189	-	319
1996	59	-	27	-	23	6 631	-	157	-	583
1997	73	-	45	-	23	7 597	-	440	-	747
1998	67	-	35	-	23	7 609	-	365	-	449
1999	53	-	34	-	18	6 835	-	280	-	624
2000	47	-	28	-	15	6 624	-	725	-	855
2001	52	-	45	-	16	6 960	-	1 380	-	819
2002	47	-	37	-	13	6 401	-	1 253	-	563
2003	44	-	29	-	8	6 619	-	829	-	680
2004	42	-	31	-	10	5 133	-	1 271	-	748
2005	37	-	15	-	10	3 623	-	530	-	464
2006	35	-	20	-	5	3 993	-	210	-	348
2007	30	-	9	-	6	2 620	-	146	-	253
2008	24	5	8	9	-	2 335	18	45	155	-
2009	25	6	3	11	-	1 961	15	3	253	-
2010	28	5	8	12	-	2 318	13	56	313	-
2011	29	6	9	16	-	2 802	40	298	474	-
2012	29	9	12	14	-	2 848	54	379	488	-
2013	24	10	11	13	-	3 057	91	181	588	-
2014	25	9	9	13	-	3 469	127	383	666	-
2015	29	10	16	15	-	3 927	103	559	698	-
2016	27	8	9	15	-	3 608	78	818	844	-

Fishing year	Catches (kg)								
	TCEPR Outside	TCER Outside	Total Outside	TCEPR Inside	TCER Inside	CELR	Total Inside	Total Overall	Percent Inside
1990	158 447	-	158 447	1 585	-	339	1 924	160 371	1
1991	128 259	-	128 259	1 015	-	222	1 237	129 496	1
1992	100 507	-	100 507	849	-	184	1 033	101 540	1
1993	95 402	-	95 402	737	-	522	1 259	96 661	1
1994	113 833	-	113 833	1 110	-	693	1 803	115 636	1
1995	79 083	-	79 083	1 851	-	747	2 598	81 681	3
1996	67 247	-	67 247	2 492	-	1 908	4 400	71 647	3
1997	82 141	-	82 141	5 637	-	2 360	7 997	90 138	6
1998	96 144	-	96 144	5 522	-	2 610	8 132	104 276	5
1999	85 486	-	85 486	4 295	-	3 846	8 141	93 627	5
2000	87 547	-	87 547	9 443	-	4 719	14 162	101 709	9
2001	80 508	-	80 508	16 627	-	4 979	21 606	102 114	16
2002	70 674	-	70 674	17 846	-	4 180	22 026	92 700	19
2003	57 211	-	57 211	11 583	-	4 944	16 527	73 738	16
2004	26 287	-	26 287	13 922	-	4 885	18 807	45 094	31
2005	24 820	-	24 820	5 574	-	2 223	7 797	32 617	17
2006	33 131	-	33 131	2 681	-	2 438	5 119	38 250	7
2007	30 192	-	30 192	1 128	-	1 962	3 090	33 282	3
2008	19 926	32	19 958	327	567	-	894	20 852	4
2009	19 285	23	19 308	36	1 102	-	1 138	20 446	6
2010	33 178	36	33 214	951	1 983	-	2 934	36 148	8
2011	40 653	168	40 821	4 047	3 441	-	7 488	48 309	16
2012	45 837	148	45 985	4 641	3 598	-	8 239	54 224	15
2013	49 028	97	49 125	2 596	3 589	-	6 185	55 310	11
2014	58 654	124	58 778	4 593	4 451	-	9 044	67 822	13
2015	64 491	268	64 759	7 481	5 107	-	12 588	77 347	16
2016	52 424	203	52 627	9 964	5 701	-	15 665	68 292	23

**Table A1b: Number of TCEPR, TCER and CELR Cook Strait number of vessels, tows, and total catch by year. Data source was un-groomed non-zero TCEPR, TCER, and CELR tows that caught hoki. ‘CELR trawl’ includes mid-water and bottom trawl tows reported on the CELR form only. Year defined as June to October. There were no October data available for 2016.**

Fishing year	Number of vessels				Number of tows			
	TCEPR	TCER	CELR	Total	TCEPR	TCER	CELR	Total
1990	18	-	30	48	1 071	-	568	1 639
1991	22	-	41	63	2 097	-	1 510	3 607
1992	24	-	31	55	1 684	-	845	2 529
1993	20	-	30	50	1 532	-	934	2 466
1994	31	-	39	70	1 957	-	1 377	3 334
1995	26	-	33	59	2 291	-	1 266	3 557
1996	42	-	37	79	4 700	-	1 485	6 185
1997	40	-	28	68	4 921	-	1 061	5 982
1998	31	-	28	59	3 022	-	1 317	4 339
1999	21	-	28	49	2 656	-	942	3 598
2000	22	-	32	54	2 372	-	1 157	3 529
2001	25	-	23	48	2 042	-	981	3 023
2002	19	-	22	41	1 127	-	531	1 658
2003	21	-	25	46	1 933	-	998	2 931
2004	20	-	31	51	1 863	-	1 134	2 997
2005	15	-	15	30	1 454	-	476	1 930
2006	13	-	13	26	1 067	-	328	1 395
2007	8	-	14	22	980	-	491	1 471
2008	7	20	-	27	668	581	-	1 249
2009	10	21	1	32	878	551	1	1 430
2010	8	18	-	26	841	523	-	1 364
2011	7	20	-	27	519	571	-	1 090
2012	9	20	-	29	779	401	-	1 180
2013	10	20	-	30	973	359	-	1 332
2014	10	20	-	30	965	385	-	1 350
2015	10	19	-	29	910	372	-	1 282
2016	9	19	-	28	757	395	-	1 152

Fishing year	Catches (kg)			
	TCEPR	TCER	CELR	Total
1990	12 109	-	2 596	14 705
1991	22 153	-	7 013	29 166
1992	19 583	-	4 973	24 556
1993	17 533	-	4 199	21 732
1994	26 785	-	9 071	35 856
1995	26 600	-	7 361	33 962
1996	50 986	-	8 018	59 005
1997	49 946	-	6 562	56 508
1998	36 308	-	9 408	45 716
1999	34 040	-	6 222	40 262
2000	30 603	-	8 986	39 588
2001	24 630	-	8 188	32 818
2002	17 628	-	4 104	21 732
2003	27 341	-	7 271	34 613
2004	28 509	-	10 520	39 030
2005	18 482	-	4 369	22 851
2006	16 670	-	3 035	19 704
2007	12 594	-	5 403	17 997
2008	9 215	6 661	-	15 876
2009	10 044	5 112	-	15 156
2010	10 916	4 875	-	15 791
2011	7 315	4 519	-	11 834
2012	9 998	3 154	-	13 152
2013	12 330	3 119	-	15 449
2014	11 691	3 163	-	14 854
2015	12 906	3 523	-	16 428
2016	11 628	4 081	-	15 709



**Table A1c: Number of Chatham Rise and ECSI vessels, tows and catch for all vessels by year for the non-spawning season. Data source was un-groomed non-zero TCEPR, TCER, and CELR tows that caught hoki. ‘CELR’ includes all fishing methods reported on the CELR form, and ‘CELR trawl’ includes mid-water and bottom trawl tows only. Chatham Rise data included data from October to September, and ECSI data included data from October to May.**

Fishing year	Number of vessels				Number of tows			
	TCEPR	TCER	CELR	Total	TCEPR	TCER	CELR	Total
1990	47	-	23	70	3 325	-	529	3 854
1991	69	-	38	107	5 724	-	900	6 624
1992	76	-	30	106	8 601	-	539	9 140
1993	75	-	29	104	8 575	-	512	9 087
1994	78	-	26	104	6 447	-	525	6 972
1995	87	-	31	118	10 028	-	675	10 703
1996	102	-	26	128	11 651	-	405	12 056
1997	105	-	18	123	12 609	-	303	12 912
1998	97	-	18	115	16 176	-	212	16 388
1999	87	-	24	111	14 984	-	421	15 405
2000	70	-	16	86	13 432	-	330	13 762
2001	68	-	11	79	12 360	-	373	12 733
2002	60	-	14	74	10 343	-	280	10 623
2003	63	-	15	78	11 400	-	255	11 655
2004	59	-	11	70	9 511	-	211	9 722
2005	51	-	12	63	7 418	-	132	7 550
2006	52	-	14	66	7 314	-	134	7 448
2007	47	-	11	58	7 324	-	153	7 477
2008	42	11	-	53	7 012	65	-	7 077
2009	37	12	1	50	6 227	79	2	6 308
2010	39	16	-	55	6 003	278	-	6 281
2011	39	14	-	53	5 446	140	-	5 586
2012	37	13	-	50	5 647	190	-	5 837
2013	39	14	-	53	5 458	416	-	5 874
2014	39	17	-	56	5 421	414	-	5 835
2015	41	13	-	54	5 813	203	-	6 016
2016	35	13	-	48	5 638	234	-	5 872

Fishing year	Catches (kg)			
	TCEPR	TCER	CELR	Total
1990	13 091	-	71	13 161
1991	29 965	-	162	30 126
1992	48 036	-	99	48 134
1993	44 169	-	63	44 231
1994	22 662	-	63	22 725
1995	38 991	-	182	39 173
1996	50 283	-	86	50 368
1997	55 726	-	93	55 819
1998	77 105	-	93	77 197
1999	72 656	-	929	73 585
2000	55 912	-	98	56 010
2001	49 307	-	532	49 840
2002	39 105	-	38	39 144
2003	39 071	-	17	39 088
2004	33 608	-	39	33 647
2005	30 423	-	8	30 432
2006	34 934	-	6	34 941
2007	37 797	-	10	37 806
2008	37 855	60	-	37 915
2009	38 997	8	-	39 005
2010	39 086	47	-	39 133
2011	38 402	40	-	38 442
2012	39 169	72	-	39 241
2013	36 433	89	-	36 523
2014	33 700	45	-	33 746
2015	40 020	45	-	40 065
2016	36 679	29	-	36 708

**Table A1d: Number of ECSI vessels, tows and catch for all vessels by year for the spawning season. Data source was un-groomed non-zero TCEPR, TCER, and CELR tows that caught hoki. Year defined as June to October. ‘CELR trawl’ included mid-water and bottom trawl tows reported on the CELR form only. There were no data available for October 2016.**

Fishing year	Number of vessels				Number of tows			
	TCEPR	TCER	CELR	Total	TCEPR	TCER	CELR	Total
1990	8	-	17	25	45	-	123	168
1991	12	-	20	32	134	-	234	368
1992	10	-	12	22	106	-	242	348
1993	9	-	13	22	32	-	274	306
1994	9	-	12	21	44	-	215	259
1995	12	-	10	22	48	-	72	120
1996	26	-	10	36	192	-	77	269
1997	21	-	6	27	194	-	154	348
1998	20	-	6	26	213	-	81	294
1999	19	-	9	28	141	-	151	292
2000	16	-	9	25	126	-	229	355
2001	16	-	8	24	197	-	251	448
2002	17	-	10	27	257	-	146	403
2003	21	-	11	32	555	-	219	774
2004	14	-	10	24	114	-	248	362
2005	12	-	3	15	284	-	69	353
2006	6	-	5	11	141	-	76	217
2007	12	-	4	16	108	-	27	135
2008	10	4	-	14	239	47	-	286
2009	11	3	-	14	103	37	-	140
2010	10	4	-	14	78	97	-	175
2011	8	5	-	13	129	74	-	203
2012	11	6	-	17	183	88	-	271
2013	12	6	-	18	245	55	-	300
2014	13	6	-	19	188	18	-	206
2015	9	4	-	13	238	10	-	248
2016	11	7	-	18	313	15	-	328

Fishing year	Catches (kg)			
	TCEPR	TCER	CELR	Total
1990	51	-	229	280
1991	841	-	503	1 345
1992	547	-	396	943
1993	137	-	172	309
1994	164	-	353	517
1995	52	-	103	155
1996	1 199	-	103	1 301
1997	817	-	973	1 790
1998	1 300	-	371	1 671
1999	765	-	1 329	2 094
2000	599	-	1 822	2 421
2001	1 658	-	760	2 418
2002	2 806	-	225	3 031
2003	6 460	-	1 006	7 466
2004	1 370	-	927	2 297
2005	4 683	-	50	4 733
2006	1 137	-	57	1 194
2007	1 001	-	63	1 064
2008	2 302	40	-	2 342
2009	1 117	29	-	1 146
2010	600	138	-	738
2011	1 504	152	-	1 657
2012	2 355	175	-	2 530
2013	3 283	110	-	3 393
2014	2 758	38	-	2 797
2015	3 660	40	-	3 700
2016	4 150	29	-	4 178

**Table A1e: Number of Sub-Antarctic vessels, tows and catch for all vessels by fishing year. Data source was un-groomed non-zero TCEPR, TCER, and CELR tows that caught hoki. ‘CELR trawl’ included mid-water and bottom trawl tows reported on the CELR form only.**

Fishing year	Number of vessels				Number of tows			
	TCEPR	TCER	CELR	Total	TCEPR	TCER	CELR	Total
1990	64	-	-	64	2 787	-	-	2 787
1991	66	-	-	66	4 617	-	-	4 617
1992	76	-	-	76	7 025	-	-	7 025
1993	63	-	2	65	6 143	-	4	6 147
1994	65	-	-	65	3 718	-	-	3 718
1995	62	-	-	62	3 585	-	-	3 585
1996	68	-	1	69	4 170	-	2	4 172
1997	74	-	-	74	5 003	-	-	5 003
1998	68	-	1	69	5 419	-	4	5 423
1999	68	-	-	68	5 145	-	-	5 145
2000	56	-	1	57	7 677	-	3	7 680
2001	56	-	-	56	7 401	-	-	7 401
2002	55	-	1	56	8 443	-	25	8 468
2003	50	-	3	53	5 689	-	10	5 699
2004	46	-	-	46	3 850	-	-	3 850
2005	43	-	-	43	2 638	-	-	2 638
2006	41	-	-	41	2 507	-	-	2 507
2007	36	-	-	36	3 004	-	-	3 004
2008	35	-	-	35	2 731	-	-	2 731
2009	32	1	-	33	2 914	1	-	2 915
2010	34	2	-	36	3 171	2	-	3 173
2011	35	1	-	36	2 931	1	-	2 932
2012	34	3	-	37	2 731	3	-	2 734
2013	35	1	-	36	2 844	1	-	2 845
2014	30	1	-	31	3 385	2	-	3 387
2015	31	3	-	34	3 115	16	-	3 131
2016	30	3	-	33	1 819	13	-	1 832

Fishing year	Catches (kg)			
	TCEPR	TCER	CELR	Total
1990	11 748	-	-	11 748
1991	16 669	-	-	16 669
1992	30 688	-	-	30 688
1993	24 836	-	-	24 836
1994	11 636	-	-	11 636
1995	13 128	-	-	13 128
1996	14 269	-	1	14 270
1997	21 771	-	-	21 771
1998	25 129	-	1	25 129
1999	23 753	-	-	23 753
2000	33 772	-	-	33 772
2001	30 076	-	-	30 076
2002	30 175	-	-	30 175
2003	20 194	-	5	20 199
2004	11 635	-	-	11 635
2005	6 337	-	-	6 337
2006	6 961	-	-	6 961
2007	7 661	-	-	7 661
2008	8 708	-	-	8 708
2009	9 807	-	-	9 807
2010	12 275	-	-	12 275
2011	12 655	-	-	12 655
2012	15 743	-	-	15 743
2013	14 095	-	-	14 095
2014	19 926	1	-	19 927
2015	16 376	1	-	16 378
2016	6 638	-	-	6 639

**Table A1f: Number of Puysegur vessels, tows and catch for all vessels by year for the spawning season. Data source was un-groomed non-zero TCEPR, TCER, and CELR tows that caught hoki. Year defined as June to December. ‘CELR trawl’ included mid-water and bottom trawl tows reported on the CELR form only. There were no October to December data available for 2016.**

Fishing year	Number of vessels				Number of tows			
	TCEPR	TCER	CELR	Total	TCEPR	TCER	CELR	Total
1990	64	-	-	64	2 787	-	-	2 787
1991	66	-	-	66	4 617	-	-	4 617
1992	76	-	-	76	7 025	-	-	7 025
1993	63	-	2	65	6 143	-	4	6 147
1994	65	-	-	65	3 718	-	-	3 718
1995	62	-	-	62	3 585	-	-	3 585
1996	68	-	1	69	4 170	-	2	4 172
1997	74	-	-	74	5 003	-	-	5 003
1998	68	-	1	69	5 419	-	4	5 423
1999	68	-	-	68	5 145	-	-	5 145
2000	56	-	1	57	7 677	-	3	7 680
2001	56	-	-	56	7 401	-	-	7 401
2002	55	-	1	56	8 443	-	25	8 468
2003	50	-	3	53	5 689	-	10	5 699
2004	46	-	-	46	3 850	-	-	3 850
2005	43	-	-	43	2 638	-	-	2 638
2006	41	-	-	41	2 507	-	-	2 507
2007	36	-	-	36	3 004	-	-	3 004
2008	35	-	-	35	2 731	-	-	2 731
2009	32	1	-	33	2 914	1	-	2 915
2010	34	2	-	36	3 171	2	-	3 173
2011	35	1	-	36	2 931	1	-	2 932
2012	34	3	-	37	2 731	3	-	2 734
2013	35	1	-	36	2 844	1	-	2 845
2014	30	1	-	31	3 385	2	-	3 387
2015	31	3	-	34	3 115	16	-	3 131
2016	30	3	-	33	1 819	13	-	1 832

Fishing year	Catches (kg)			
	TCEPR	TCER	CELR	Total
1990	11 748	-	-	11 748
1991	16 669	-	-	16 669
1992	30 688	-	-	30 688
1993	24 836	-	-	24 836
1994	11 636	-	-	11 636
1995	13 128	-	-	13 128
1996	14 269	-	1	14 270
1997	21 771	-	-	21 771
1998	25 129	-	1	25 129
1999	23 753	-	-	23 753
2000	33 772	-	-	33 772
2001	30 076	-	-	30 076
2002	30 175	-	-	30 175
2003	20 194	-	5	20 199
2004	11 635	-	-	11 635
2005	6 337	-	-	6 337
2006	6 961	-	-	6 961
2007	7 661	-	-	7 661
2008	8 708	-	-	8 708
2009	9 807	-	-	9 807
2010	12 275	-	-	12 275
2011	12 655	-	-	12 655
2012	15 743	-	-	15 743
2013	14 095	-	-	14 095
2014	19 926	1	-	19 927
2015	16 376	1	-	16 378
2016	6 638	-	-	6 639

**Table A2a: Number of vessels, total hoki catch, number of tows, median tow duration, median catch per tow, and median catch per hour for all WCSI vessels by year. Year defined as June to October. There were no October data available for 2016. Data are non-zero catches for TCEPR midwater tows.**

**MW tows (all target species):**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	69	149 295	6 780	4.2	10.3	2.6
1991	66	118 323	6 744	4.0	10.2	2.6
1992	61	92 024	5 193	3.6	12.4	3.5
1993	57	82 529	5 263	3.2	10.3	3.7
1994	63	105 195	7 139	3.0	8.9	3.2
1995	59	75 148	7 408	3.5	4.9	1.4
1996	59	64 802	5 171	3.5	6.8	1.9
1997	76	82 639	6 611	3.8	7.4	2.0
1998	66	95 864	6 695	3.5	10.4	2.8
1999	56	76 767	5 256	3.1	10.3	3.3
2000	52	79 535	5 316	2.8	12.0	4.3
2001	62	78 853	5 879	2.6	9.0	3.4
2002	56	61 528	4 654	2.3	9.8	4.1
2003	51	51 751	4 312	3.0	8.1	2.4
2004	51	32 049	4 230	2.4	4.6	1.5
2005	37	19 682	2 365	2.5	5.1	1.8
2006	36	21 067	2 015	3.0	6.7	2.5
2007	31	21 093	1 432	3.5	9.3	3.5
2008	15	12 047	886	1.8	6.4	3.8
2009	23	12 590	887	3.2	8.9	3.1
2010	26	23 033	1 216	2.6	15.3	5.2
2011	24	29 603	1 514	2.0	17.2	8.4
2012	27	30 122	1 567	2.1	16.3	7.9
2013	24	33 851	1 811	2.6	15.4	6.2
2014	26	43 805	2 317	2.8	15.1	5.9
2015	27	50 970	2 681	2.7	15.4	6.0
2016	24	38 311	2 641	2.2	11.7	5.0
All years	241	1 582 479	107 983	3.2	9.5	3.0

**MW tows (Target hoki tows):**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	69	149 263	6 736	4.2	10.3	2.6
1991	66	118 202	6 727	4.0	10.2	2.6
1992	60	91 904	5 141	3.6	12.4	3.6
1993	56	82 133	5 030	3.1	10.5	4.1
1994	62	105 007	6 978	3.0	9.5	3.3
1995	59	74 715	7 145	3.5	4.9	1.4
1996	59	64 735	5 115	3.5	6.8	1.9
1997	76	82 222	6 505	3.8	7.9	2.1
1998	66	95 670	6 630	3.5	10.4	2.8
1999	56	76 532	5 142	3.1	10.3	3.4
2000	51	79 269	5 194	2.7	12.0	4.5
2001	62	78 512	5 726	2.6	9.3	3.6
2002	56	61 336	4 579	2.3	9.8	4.3
2003	51	51 466	4 208	3.0	8.1	2.5
2004	51	31 874	4 152	2.3	4.9	1.6
2005	37	19 620	2 266	2.4	5.7	2.0
2006	34	20 729	1 734	2.6	8.5	3.2
2007	31	20 786	1 136	2.8	15.0	5.5
2008	13	11 841	806	1.7	7.3	4.7
2009	15	12 367	685	2.7	14.2	5.0
2010	23	22 884	1 172	2.5	17.1	5.5
2011	24	29 468	1 495	2.0	17.4	8.5
2012	27	30 071	1 559	2.1	16.3	7.9
2013	24	33 697	1 793	2.6	15.4	6.2
2014	26	43 770	2 298	2.8	15.2	6.0
2015	27	50 906	2 649	2.7	15.5	6.1
2016	24	38 311	2 639	2.2	11.7	5.0
All years	241	1 577 292	105 240	3.2	9.9	3.1

**Table A2b: Number of vessels, total hoki catch, number of tows, median tow duration, median catch per tow, and median catch per hour for all WCSI vessels by year. Year defined as June to October. There were no October data available for 2016. Data are non-zero catches for TCEPR bottom tows.**

**All target species BT tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	41	10 737	1 292	4.0	3.2	0.8
1991	36	10 951	1 458	4.0	3.6	0.9
1992	38	9 334	1 036	4.1	4.1	1.0
1993	33	13 656	1 727	3.8	5.2	1.4
1994	32	9 703	1 468	4.2	3.7	0.8
1995	27	5 809	1 331	4.5	2.5	0.5
1996	38	4 914	1 586	4.7	2.1	0.4
1997	47	5 145	1 442	5.0	2.2	0.5
1998	40	5 925	1 308	5.2	2.9	0.5
1999	39	12 894	1 835	4.7	4.1	0.8
2000	34	17 487	2 064	4.5	6.0	1.2
2001	40	18 238	2 399	4.5	5.0	0.9
2002	35	26 993	3 005	5.0	5.2	1.0
2003	39	17 057	3 197	5.3	2.3	0.4
2004	35	8 174	2 154	6.0	1.5	0.3
2005	30	10 708	1 801	6.6	2.5	0.4
2006	26	14 723	2 145	8.3	2.8	0.4
2007	22	10 252	1 344	7.1	3.1	0.4
2008	17	8 179	1 472	9.0	2.4	0.3
2009	18	6 735	1 083	9.2	3.0	0.3
2010	21	11 116	1 171	7.2	4.9	0.8
2011	21	15 075	1 565	6.1	6.2	1.0
2012	23	20 353	1 656	5.2	9.9	1.9
2013	18	17 773	1 427	5.1	10.8	2.3
2014	18	19 321	1 533	5.2	10.2	1.8
2015	19	20 997	1 795	5.2	8.4	1.5
2016	21	24 076	1 785	3.7	12.6	3.4
All years	146	356 324	46 079	5.0	4.0	0.7

**Target hoki BT tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	34	10 597	1 129	4.2	4.1	1.1
1991	31	10 877	1 321	4.0	4.1	1.1
1992	28	9 152	791	4.0	7.0	1.7
1993	29	13 611	1 588	3.8	5.9	1.6
1994	29	9 679	1 369	4.3	4.2	0.9
1995	24	5 794	1 290	4.5	2.5	0.5
1996	37	4 885	1 544	4.7	2.1	0.4
1997	42	5 115	1 354	5.0	2.5	0.5
1998	34	5 888	1 217	5.3	3.1	0.5
1999	35	12 856	1 689	4.7	5.1	1.0
2000	32	17 417	1 903	4.4	6.3	1.4
2001	37	18 216	2 314	4.6	5.0	1.0
2002	34	26 724	2 839	5.0	5.9	1.1
2003	39	16 793	2 791	5.1	3.0	0.6
2004	34	7 911	1 799	5.7	2.0	0.4
2005	27	9 732	1 240	5.6	4.5	0.8
2006	24	13 087	1 405	7.0	5.0	0.8
2007	20	8 874	731	4.8	9.3	1.7
2008	13	5 246	480	5.0	8.6	1.7
2009	13	4 460	350	4.5	11.2	2.6
2010	19	9 214	611	3.2	13.5	4.7
2011	17	11 707	908	4.1	11.4	2.9
2012	20	18 853	1 184	3.8	15.0	4.1
2013	16	16 060	996	3.5	15.6	4.8
2014	15	17 203	1 075	3.9	15.3	3.8
2015	17	18 209	1 216	3.6	14.3	4.0
2016	17	23 071	1 494	3.2	15.1	5.1
All years	131	331 232	36 628	4.5	5.2	1.1

**Table A2c: Number of vessels, total hoki catch, number of tows, median tow duration, median catch per tow, and median catch per hour for all Cook Strait vessels by year. Year defined as June to October. There were no October data available for 2016. Data are non-zero catches for TCEPR midwater tows.**

**All target species tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	17	11 894	1 048	1.2	9.1	7.4
1991	22	21 976	2 069	1.5	8.2	5.0
1992	22	19 345	1 642	1.2	8.3	6.5
1993	20	16 977	1 499	1.0	8.3	7.0
1994	29	25 106	1 810	1.0	11.8	11.8
1995	24	24 376	2 162	1.0	8.3	9.9
1996	36	41 820	3 087	0.8	11.1	16.7
1997	34	43 248	3 592	1.0	10.6	11.2
1998	28	30 711	2 373	1.0	11.4	11.7
1999	21	28 084	2 037	1.0	12.7	14.8
2000	21	27 935	1 989	0.7	12.0	19.3
2001	25	23 581	1 842	0.8	11.0	14.0
2002	15	17 147	1 068	1.0	14.9	17.2
2003	20	26 979	1 816	1.0	12.6	16.2
2004	19	27 712	1 793	1.0	12.2	14.2
2005	13	18 166	1 344	1.0	13.0	16.9
2006	11	16 330	1 015	0.8	15.1	20.5
2007	7	12 444	952	1.0	11.0	13.8
2008	6	7 558	404	0.8	18.4	23.3
2009	8	9 095	740	0.6	10.1	18.2
2010	8	10 839	820	0.8	11.2	14.9
2011	6	7 346	527	0.8	11.3	16.7
2012	9	9 778	759	0.9	10.7	13.7
2013	9	11 659	858	0.7	11.9	17.2
2014	10	11 388	944	1.0	10.2	11.1
2015	9	12 122	805	0.7	14.4	20.7
2016	9	10 585	621	0.7	16.8	26.2
All years	71	524 203	39 616	1.0	10.9	12.7

**Target hoki tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	17	11 894	1 048	1.2	9.1	7.4
1991	22	21 976	2 069	1.5	8.2	5.0
1992	22	19 345	1 642	1.2	8.3	6.5
1993	18	16 957	1 493	1.0	8.3	7.0
1994	29	25 065	1 804	1.0	11.8	11.9
1995	24	24 320	2 158	1.0	8.3	9.9
1996	36	41 744	3 076	0.8	11.2	16.7
1997	34	43 179	3 585	1.0	10.6	11.2
1998	28	30 674	2 371	1.0	11.4	11.7
1999	21	28 081	2 036	1.0	12.7	14.8
2000	21	27 935	1 989	0.7	12.0	19.3
2001	25	23 553	1 839	0.8	11.0	14.0
2002	15	17 147	1 068	1.0	14.9	17.2
2003	20	26 979	1 814	1.0	12.6	16.3
2004	19	27 712	1 791	1.0	12.2	14.2
2005	13	18 162	1 343	1.0	13.0	16.9
2006	11	16 330	1 014	0.8	15.1	20.5
2007	7	12 396	949	1.0	10.9	13.8
2008	5	7 555	397	0.8	18.8	24.3
2009	8	9 083	739	0.6	10.1	18.2
2010	8	10 783	818	0.8	11.2	14.8
2011	6	7 346	527	0.8	11.3	16.7
2012	9	9 778	759	0.9	10.7	13.7
2013	9	11 659	858	0.7	11.9	17.2
2014	10	11 388	944	1.0	10.2	11.1
2015	9	12 119	804	0.7	14.4	20.9
2016	9	10 585	621	0.7	16.8	26.2
All years	71	523 746	39 556	1.0	10.9	12.7

**Table A2d: Number of vessels, total hoki catch, number of tows, median tow duration, median catch per**

tow, and median catch per hour for Chatham Rise and ECSI by fishing year. Data source was un-groomed bottom trawl non-zero TCEPR tows that caught hoki. Chatham Rise data includes data from October to September, and ECSI data includes data from October to May.

**All target species tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	47	13 001	3 297	4.0	1.5	0.5
1991	59	18 080	4 787	4.0	2.0	0.5
1992	72	43 456	8 169	4.0	3.1	0.8
1993	61	39 238	7 523	3.9	3.4	1.0
1994	64	18 125	5 305	3.5	2.1	0.7
1995	70	30 585	7 914	3.8	3.0	0.9
1996	84	37 620	9 294	3.6	2.9	0.9
1997	96	42 898	10 330	3.7	3.2	0.9
1998	82	55 824	12 479	4.0	3.3	0.9
1999	77	61 528	12 620	4.0	4.1	1.0
2000	60	44 753	10 746	4.1	3.0	0.8
2001	60	46 150	11 429	4.5	3.0	0.7
2002	55	36 271	9 491	4.5	2.9	0.7
2003	62	37 415	10 912	4.7	2.5	0.5
2004	58	31 656	9 131	5.0	2.3	0.5
2005	50	28 914	7 048	5.0	2.8	0.6
2006	50	34 077	7 145	4.8	3.5	0.8
2007	46	37 640	7 267	4.6	3.5	0.8
2008	38	37 375	6 890	4.8	3.6	0.8
2009	37	38 956	6 186	4.3	4.6	1.1
2010	38	38 454	5 833	4.5	5.3	1.2
2011	38	38 136	5 286	4.7	5.9	1.2
2012	35	38 818	5 428	4.8	5.7	1.3
2013	35	35 085	5 179	4.8	5.6	1.2
2014	32	32 511	5 147	4.8	5.1	1.1
2015	35	38 385	5 458	5.0	5.7	1.2
2016	32	33 967	5 089	4.6	5.4	1.2
All years	200	988 918	205 383	4.2	3.2	0.8

**Target hoki tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	31	11 788	1 902	4.0	3.8	1.0
1991	41	16 761	3 285	4.0	3.5	0.9
1992	47	42 305	5 408	3.8	5.7	1.6
1993	40	38 354	5 169	3.5	5.7	1.6
1994	36	17 525	3 372	3.2	4.2	1.3
1995	42	30 097	6 485	3.5	3.9	1.1
1996	58	37 177	7 969	3.5	3.2	1.0
1997	73	42 380	8 988	3.5	3.7	1.1
1998	63	55 315	11 159	4.0	4.2	1.0
1999	46	60 838	11 244	4.0	4.4	1.1
2000	34	44 113	9 413	4.1	3.7	0.9
2001	40	44 928	9 762	4.5	3.5	0.8
2002	31	35 087	7 773	4.4	3.4	0.8
2003	32	36 051	9 196	4.8	3.0	0.6
2004	28	30 207	7 142	4.9	3.0	0.6
2005	21	27 472	4 973	5.0	4.1	0.8
2006	20	32 329	4 997	4.8	5.0	1.0
2007	21	34 746	4 733	4.5	5.8	1.2
2008	22	33 527	4 187	4.8	6.6	1.4
2009	21	33 645	3 896	4.2	7.3	1.7
2010	21	35 151	4 349	4.6	6.9	1.5
2011	23	34 811	4 056	4.8	7.2	1.5
2012	24	37 639	4 380	4.8	7.2	1.6
2013	22	33 919	4 160	4.8	6.7	1.4
2014	18	31 788	3 935	4.9	6.6	1.4
2015	21	37 437	4 309	5.0	7.2	1.4
2016	14	31 580	3 899	4.8	6.8	1.5
All years	165	946 970	160 141	4.2	4.4	1.1



**Table A2e: Number of vessels, total hoki catch, number of tows, median tow duration, median catch per tow, and median catch per hour for ECSI. Data source was un-groomed midwater or bottom non-zero TCEPR tows that caught hoki. Year defined as June to October. There were no October data available for 2016. Data are not shown for MW vessels in 2009 or 2010 as there was only one vessel.**

**All target species mid-water tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
2000	7	289	24	2.7	7.5	2.4
2001	15	1 264	123	2.4	6.0	2.2
2002	10	2 003	145	2.2	10.9	4.2
2003	18	4 453	301	2.1	13.1	5.2
2004	5	1 438	85	2.2	10.4	6.0
2005	6	4 037	221	2.0	15.0	8.4
2006	4	485	41	1.5	10.0	5.7
2007	4	299	26	1.1	8.7	8.8
2008	3	263	28	3.0	8.1	2.7
2009	1	462	-	-	-	-
2010	1	28	-	-	-	-
2011	4	879	57	1.0	14.7	10.8
2012	8	1 686	117	1.6	10.3	5.2
2013	10	2 945	191	2.0	14.2	6.6
2014	12	2 586	134	2.2	16.5	7.9
2015	8	3 345	189	2.3	17.3	6.9
2016	8	3 536	201	2.2	15.1	5.8
All years	34	30 000	1 917	2.0	13.4	5.9

**Target hoki mid-water tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
2000	7	289	24	2.7	7.5	2.4
2001	15	1 264	123	2.4	6.0	2.2
2002	10	2 003	145	2.2	10.9	4.2
2003	18	4 453	301	2.1	13.1	5.2
2004	5	1 438	85	2.2	10.4	6.0
2005	6	4 037	221	2.0	15.0	8.4
2006	4	485	41	1.5	10.0	5.7
2007	4	299	26	1.1	8.7	8.8
2008	3	263	28	3.0	8.1	2.7
2009	1	462	-	-	-	-
2010	1	28	-	-	-	-
2011	4	879	57	1.0	14.7	10.8
2012	8	1 686	117	1.6	10.3	5.2
2013	10	2 945	191	2.0	14.2	6.6
2014	12	2 586	134	2.2	16.5	7.9
2015	8	3 345	189	2.3	17.3	6.9
2016	8	3 536	201	2.2	15.1	5.8
All years	34	29 094	1 884	2.0	13.3	5.7

**Table A2e ECSI continued.**

**All target bottom tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
2000	10	250	69	2.5	2.5	1.0
2001	13	441	85	2.7	3.5	1.2
2002	16	828	126	2.6	3.9	1.5
2003	16	2 081	255	2.9	5.4	1.9
2004	7	250	44	2.4	3.1	1.0
2005	8	717	98	3.0	4.1	1.8
2006	7	163	31	2.1	2.1	1.6
2007	11	666	81	2.0	6.2	2.9
2008	12	2 112	215	2.8	7.5	2.5
2009	8	635	76	2.8	6.2	2.4
2010	8	533	70	2.8	7.3	2.1
2011	6	592	56	3.5	10.8	3.0
2012	9	753	89	2.5	7.2	2.4
2013	10	302	48	2.9	3.6	1.4
2014	12	227	55	2.5	2.5	1.0
2015	7	254	41	2.8	4.1	1.4
2016	10	555	98	3.0	3.5	1.2
All years	39	11 430	1 579	2.8	4.7	1.8

**Target hoki bottom tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
2000	10	250	69	2.5	2.5	1.0
2001	13	441	85	2.7	3.5	1.2
2002	16	828	126	2.6	3.9	1.5
2003	16	2 081	255	2.9	5.4	1.9
2004	7	250	44	2.4	3.1	1.0
2005	8	717	98	3.0	4.1	1.8
2006	7	163	31	2.1	2.1	1.6
2007	11	666	81	2.0	6.2	2.9
2008	12	2 112	215	2.8	7.5	2.5
2009	8	635	76	2.8	6.2	2.4
2010	8	533	70	2.8	7.3	2.1
2011	6	592	56	3.5	10.8	3.0
2012	9	753	89	2.5	7.2	2.4
2013	10	302	48	2.9	3.6	1.4
2014	12	227	55	2.5	2.5	1.0
2015	7	254	41	2.8	4.1	1.4
2016	10	555	98	3.0	3.5	1.2
All years	28	10 856	1 433	2.8	5.0	1.9

**Table A2f: Number of vessels, total hoki catch, number of tows, median tow duration, median catch per tow, and median catch per hour for Sub-Antarctic by fishing year. Data source was un-groomed non-zero TCEPR bottom tows that caught hoki.**

**All target species tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	36	11 542	2 589	4.0	2.6	0.6
1991	43	16 177	4 420	4.3	2.6	0.6
1992	58	29 688	6 877	4.2	3.1	0.8
1993	39	22 304	5 647	4.0	3.1	0.8
1994	45	9 051	3 163	4.2	1.6	0.4
1995	42	11 716	3 223	4.3	2.2	0.6
1996	46	10 889	3 483	4.2	1.9	0.5
1997	58	19 288	4 522	4.5	3.2	0.7
1998	49	24 217	5 192	4.3	3.3	0.8
1999	49	20 966	4 673	4.5	2.9	0.7
2000	43	31 576	7 155	4.2	3.0	0.8
2001	46	26 222	6 669	4.5	2.7	0.6
2002	47	29 568	8 093	4.4	2.1	0.6
2003	44	19 870	5 556	4.9	2.4	0.5
2004	41	11 168	3 728	5.0	2.0	0.4
2005	40	6 148	2 542	5.3	1.0	0.2
2006	34	6 491	2 360	5.3	0.7	0.1
2007	31	7 420	2 878	5.2	0.8	0.2
2008	29	8 015	2 625	5.5	1.0	0.2
2009	25	9 195	2 807	5.0	1.0	0.2
2010	29	11 551	3 023	5.4	1.0	0.2
2011	28	10 973	2 689	5.0	1.5	0.3
2012	28	13 665	2 580	5.0	2.2	0.5
2013	29	14 050	2 773	4.8	2.6	0.5
2014	22	19 786	3 211	5.0	3.6	0.7
2015	25	15 606	2 835	5.1	3.3	0.7
2016	25	6 550	1 787	5.1	1.5	0.3
All years	167	423 691	107 100	4.5	2.3	0.5

**Hoki target tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	20	10 922	2 048	4.0	3.6	0.9
1991	30	15 229	3 862	4.4	2.8	0.6
1992	33	28 278	5 314	4.1	4.1	1.0
1993	24	21 359	4 817	3.8	3.6	0.9
1994	22	8 748	1 977	4.0	3.2	0.9
1995	25	11 453	2 297	4.0	3.9	1.0
1996	25	10 628	2 437	4.0	3.1	0.9
1997	42	18 919	3 293	4.2	4.6	1.1
1998	34	23 669	4 267	4.2	4.2	1.0
1999	33	20 391	3 563	4.2	4.1	1.1
2000	30	30 884	5 806	4.0	3.9	1.0
2001	31	25 397	5 324	4.2	3.5	0.8
2002	33	28 612	6 253	4.2	2.9	0.8
2003	33	19 101	4 322	4.8	3.0	0.7
2004	26	10 815	2 864	4.9	3.0	0.6
2005	25	5 151	1 351	5.1	2.5	0.5
2006	16	4 636	720	5.0	4.0	0.8
2007	20	5 143	1 136	4.5	2.2	0.5
2008	13	5 828	909	4.8	4.5	0.9
2009	12	6 883	918	4.4	5.1	1.2
2010	12	9 687	1 231	4.5	6.1	1.3
2011	15	9 210	1 237	4.5	5.5	1.2
2012	17	11 538	1 193	4.6	7.6	1.6
2013	16	11 703	1 363	4.3	6.2	1.5
2014	13	17 217	1 864	4.5	7.1	1.6
2015	15	13 052	1 610	4.9	6.2	1.3
2016	9	5 651	824	4.9	5.2	1.1
All years	110	390 103	72 800	4.2	3.9	0.9

**Table A2g: Number of vessels, total hoki catch, number of tows, median tow duration, median catch per tow, and median catch per hour for Puysegur by year. Data source was un-groomed midwater or bottom non-zero TCEPR tows that caught hoki. Year defined as June to December. There were no October to December data available for 2015. Data were removed where there was one or two vessels only.**

**All target species midwater tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	25	7 154	759	2.5	7.9	3.2
1991	16	3 188	269	2.4	10.2	4.1
1992	13	1 058	141	3.0	5.2	2.0
1993	8	660	71	2.0	6.2	2.8
1994	17	2 209	266	3.0	3.9	1.1
1995	15	1 015	163	2.3	3.0	1.4
1996	12	1 447	155	2.7	7.1	2.9
1997	20	4 742	410	3.5	8.5	2.5
1998	7	884	95	3.0	8.2	2.4
1999	16	1 416	141	3.4	4.8	1.3
2000	13	2 054	161	4.2	8.0	2.0
2001	22	5 212	372	4.3	10.0	2.2
2002	19	3 128	260	3.6	6.8	1.6
2003	20	5 137	309	2.8	12.1	3.6
2004	4	574	33	3.7	12.2	3.0
2005	9	4 953	220	2.1	22.0	9.6
2006	4	236	16	2.8	14.8	5.0
2007	1	-	-	-	-	-
2008	1	-	-	-	-	-
2009	1	-	-	-	-	-
2010	1	-	-	-	-	-
2011	2	-	-	-	-	-
2012	2	-	-	-	-	-
2013	2	-	-	-	-	-
2014	3	9	4	3.5	2.2	0.8
2015	3	864	50	2.5	16.4	3.7
2016	3	411	36	3.5	6.6	1.6
All years	105	48 563	4 114	3.0	8.0	2.6

**Hoki target mid-water tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	25	7 149	758	2.5	7.9	3.2
1991	16	3 173	268	2.4	10.2	4.1
1992	12	1 027	129	3.0	5.2	2.0
1993	8	660	71	2.0	6.2	2.8
1994	17	2 189	264	3.0	3.9	1.1
1995	15	1 015	163	2.3	3.0	1.4
1996	12	1 447	155	2.7	7.1	2.9
1997	20	4 742	410	3.5	8.5	2.5
1998	7	884	95	3.0	8.2	2.4
1999	16	1 416	141	3.4	4.8	1.3
2000	13	2 054	161	4.2	8.0	2.0
2001	22	5 206	371	4.3	10.0	2.2
2002	19	3 128	260	3.6	6.8	1.6
2003	20	5 137	309	2.8	12.1	3.6
2004	3	571	29	3.5	13.2	5.1
2005	8	4 942	216	2.1	22.0	10.0
2006	4	236	16	2.8	14.8	5.0
2007	1	-	-	-	-	-
2008	1	-	-	-	-	-
2009	1	-	-	-	-	-
2010	1	-	-	-	-	-
2011	2	-	-	-	-	-
2012	2	-	-	-	-	-
2013	2	-	-	-	-	-
2014	3	9	4	3.5	2.2	0.8
2015	3	864	50	2.5	16.4	3.7
2016	3	411	36	3.5	6.6	1.6
All years	103	48 471	4 089	3.0	8.1	2.6

**Table A2g continued.: Puysegur.**

**All target species bottom tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	15	104	207	3.3	0.2	0.1
1991	24	1 663	372	4.3	3.1	0.8
1992	30	4 012	842	4.3	3.0	0.6
1993	12	1 044	220	4.2	3.4	0.8
1994	20	394	175	4.2	1.1	0.3
1995	12	252	200	5.8	0.5	0.1
1996	16	955	354	4.3	1.0	0.2
1997	25	1 162	336	5.5	0.8	0.2
1998	19	1 295	252	4.8	2.8	0.6
1999	22	966	265	5.2	1.1	0.2
2000	20	849	273	5.4	1.0	0.2
2001	24	919	221	4.2	2.0	0.5
2002	18	1 852	193	3.8	6.5	1.5
2003	20	796	181	4.5	1.8	0.4
2004	14	198	81	4.8	0.6	0.1
2005	21	582	291	5.8	0.9	0.1
2006	16	1 002	256	4.0	1.0	0.3
2007	13	253	118	5.0	0.7	0.1
2008	6	134	56	4.8	1.4	0.3
2009	7	126	57	3.1	1.0	0.3
2010	7	121	110	4.8	0.5	0.1
2011	11	208	108	4.5	1.0	0.2
2012	7	163	59	3.7	1.0	0.4
2013	9	395	79	4.2	3.1	0.6
2014	11	796	134	3.9	4.1	1.0
2015	12	500	117	4.6	1.1	0.2
2016	8	517	68	4.5	5.5	1.3
All years	96	21 258	5 625	4.5	1.5	0.3

**Hoki target bottom tows:**

<b>Fishing year</b>	<b>Number of vessels</b>	<b>Total catch (t)</b>	<b>Number of tows</b>	<b>Median tow duration (h)</b>	<b>Median catch per tow (t)</b>	<b>Median catch per hour (t/h)</b>
1990	8	22	20	3.5	0.7	0.2
1991	20	1 541	310	4.1	4.1	0.9
1992	26	3 778	701	4.2	3.1	0.8
1993	11	1 019	201	4.0	4.0	0.9
1994	16	356	138	4.4	1.1	0.3
1995	9	217	144	5.8	0.6	0.1
1996	16	892	272	4.1	1.5	0.3
1997	22	983	295	5.3	0.9	0.2
1998	18	1 262	237	4.8	3.0	0.7
1999	21	931	238	5.1	1.2	0.2
2000	18	817	224	5.0	1.6	0.3
2001	22	910	198	4.2	2.5	0.6
2002	16	1 836	184	3.8	7.0	1.7
2003	14	774	135	4.5	3.0	0.7
2004	5	152	24	3.3	4.2	1.2
2005	8	240	51	3.2	2.2	0.9
2006	6	707	79	3.5	6.0	2.1
2007	2	-	-	-	-	-
2008	1	-	-	-	-	-
2009	1	-	-	-	-	-
2010	1	-	-	-	-	-
2011	1	-	-	-	-	-
2012	1	-	-	-	-	-
2013	4	221	21	3.9	11.5	2.7
2014	3	241	33	3.5	4.1	1.1
2015	4	371	26	3.2	16.0	4.5
2016	4	297	28	3.3	10.3	2.9
All years	78	17 721	3 586	4.3	2.6	0.6

**Table A3: CPUE datasets for all vessels and for core (TCEPR) or final (observer) vessels for each year (1990–2016) for the main hoki fishing areas. Prop. zeros: proportion of tows with zero hoki catch.**

**WCSI: TCEPR tow-by-tow, target hoki**

Fishing year	All vessels					Core vessels				
	No. vessels	Catch	Effort	Prop. zeros	CPUE	No. vessels	Catch	Effort	Prop. zeros	CPUE
1990	76	114 865.5	7 424	0.04	15.47	20	45 703.3	2 342	0.02	19.51
1991	73	104 310.3	7 709	0.04	13.53	28	56 589.3	2 813	0.02	20.12
1992	66	88 465.8	5 787	0.06	15.29	29	46 377.0	2 568	0.03	18.06
1993	60	86 525.0	6 448	0.06	13.42	35	54 625.5	3 799	0.05	14.38
1994	66	99 164.0	8 162	0.09	12.15	41	72 321.7	5 449	0.08	13.27
1995	62	67 746.1	8 241	0.10	8.22	43	51 623.4	6 057	0.09	8.52
1996	61	59 507.6	6 536	0.07	9.10	42	53 129.2	5 300	0.07	10.02
1997	76	78 705.7	7 654	0.07	10.28	50	67 412.1	6 195	0.07	10.88
1998	68	91 522.6	7 680	0.04	11.92	55	85 700.4	7 137	0.04	12.01
1999	59	82 908.7	6 711	0.03	12.35	51	82 530.6	6 658	0.03	12.40
2000	51	93 697.0	6 999	0.02	13.39	44	92 875.7	6 908	0.02	13.44
2001	63	93 775.8	7 972	0.02	11.76	48	89 975.2	7 494	0.02	12.01
2002	56	85 316.3	7 277	0.02	11.72	47	82 110.1	6 963	0.02	11.79
2003	51	68 096.1	6 958	0.02	9.79	45	67 823.0	6 901	0.02	9.83
2004	51	39 675.1	5 922	0.02	6.70	41	35 211.4	5 408	0.02	6.51
2005	37	29 243.2	3 491	0.02	8.38	36	28 714.9	3 439	0.02	8.35
2006	36	33 620.2	3 113	0.01	10.80	31	32 916.3	3 018	0.01	10.91
2007	32	29 194.3	1 818	0.01	16.06	28	28 401.8	1 776	0.01	15.99
2008	22	16 956.9	1 277	0.01	13.28	15	16 126.4	1 213	0.01	13.29
2009	20	16 786.1	1 030	0.01	16.30	13	15 785.3	966	0.01	16.34
2010	27	31 982.6	1 774	0.01	18.03	24	31 674.0	1 760	0.01	18
2011	27	41 002.0	2 390	0.01	17.16	24	40 564.6	2 356	0.01	17.22
2012	30	48 759.8	2 731	0.01	17.85	29	48 701.2	2 723	0.01	17.89
2013	26	49 485.3	2 770	0.01	17.86	23	49 015.9	2 741	0.01	17.88
2014	26	60 570.6	3 343	-	18.12	23	55 701.7	3 116	-	17.88
2015	28	68 779.9	3 835	-	17.93	23	63 468.6	3 593	-	17.66
2016	25	60 946.5	4 121	0.01	14.79	21	55 096.7	3 743	0.01	14.72

**WCSI: Observer catch for target hoki**

Fishing year	All vessels					Final vessels				
	No. vessels	Catch	Effort	Prop. zeros	CPUE	No. vessels	Catch	Effort	Prop. zeros	CPUE
1990	14	47 607.1	1 512	0.02	31.49	5	9 290.8	482	0.02	19.28
1991	14	28 132.4	1 228	0.02	22.91	5	11 583.2	493	0.02	23.50
1992	10	18 562.5	754	0.03	24.62	6	6 486.6	252	0.01	25.74
1993	15	17 298.0	1 020	0.02	16.96	12	8 401.6	694	0.01	12.11
1994	15	32 398.4	1 549	0.02	20.92	11	16 566.4	985	0.01	16.82
1995	9	25 689.7	797	0.01	32.23	6	6 411.4	401	0.01	15.99
1996	15	17 676.5	1 030	0.03	17.16	10	9 999.4	768	0.02	13.02
1997	12	14 180.4	674	0.02	21.04	11	9 651.2	601	0.02	16.06
1998	16	18 622.0	896	0.01	20.78	14	13 560.6	803	0.01	16.89
1999	14	17 313.0	1 073	0.02	16.14	14	14 234.6	1 050	0.02	13.56
2000	17	18 525.3	1 154	-	16.05	16	17 662.3	1 141	-	15.48
2001	21	15 344.9	1 007	0.01	15.24	21	14 648.9	998	0.01	14.68
2002	16	16 461.8	1 303	0.01	12.63	15	16 090.9	1 275	0.01	12.62
2003	13	10 106.2	912	0.01	11.08	13	10 094.2	909	0.01	11.10
2004	16	8 397.7	1 299	0.01	6.46	14	7 564.0	1 186	0.01	6.38
2005	13	7 178.2	974	-	7.37	12	7 125.9	961	-	7.42
2006	13	9 522.3	780	-	12.21	13	9 519.6	778	-	12.24
2007	16	9 782.3	514	-	19.03	16	9 284.9	461	-	20.14
2008	11	7 054.5	426	-	16.56	11	7 020.8	422	-	16.64
2009	12	7 325.8	367	-	19.96	12	7 273.1	364	-	19.98
2010	14	9 163.6	543	0.02	16.88	13	8 102.0	468	0.02	17.31
2011	11	9 122.6	536	0.01	17.02	11	8 637.5	492	0.01	17.56
2012	15	16 707.3	881	0.01	18.96	14	15 975.7	840	0.01	19.02
2013	17	30 077.1	1 669	-	18.02	17	29 295.6	1 613	-	18.16
2014	17	28 194.2	1 557	0.01	18.11	16	26 589.6	1 449	0.01	18.35
2015	20	35 209.8	1 716	0.01	20.52	20	34 159.8	1 651	0.01	20.69
2016	16	27 510.2	1 678	-	16.39	16	25 955.1	1 578	-	16.45

Table A3 continued.

## Cook Strait: TCEPR MW tow-by-tow, target hoki

Fishing year	All vessels					Final vessels				
	No. vessels	Catch	Effort	Prop. zeros	CPUE	No. vessels	Catch	Effort	Prop. zeros	CPUE
1990	17	11 751.7	1 040	0.03	11.30	11	8 595.4	772	0.03	11.13
1991	22	21 708.2	2 040	0.02	10.64	17	19 123.3	1 793	0.02	10.67
1992	22	17 839.8	1 567	0.03	11.38	16	15 069.1	1 307	0.03	11.53
1993	18	16 317.6	1 430	0.04	11.41	14	15 547.5	1 336	0.04	11.64
1994	28	23 871.0	1 722	0.04	13.86	15	19 143.1	1 308	0.03	14.64
1995	24	21 556.8	1 922	0.02	11.22	17	18 117.2	1 353	0.02	13.39
1996	36	34 736.3	2 587	0.03	13.43	22	27 531.7	1 927	0.03	14.29
1997	34	37 320.1	3 036	0.04	12.29	22	29 626.0	2 413	0.03	12.28
1998	28	26 455.7	2 034	0.03	13.01	19	23 767.8	1 815	0.02	13.10
1999	20	25 780.9	1 821	0.02	14.16	19	25 529.3	1 809	0.02	14.11
2000	21	22 193.8	1 584	0.02	14.01	19	21 895.7	1 553	0.02	14.10
2001	25	20 249.3	1 593	0.03	12.71	20	19 232.3	1 504	0.02	12.79
2002	15	15 274.8	917	0.01	16.66	9	13 892.6	822	0.01	16.90
2003	19	22 949.7	1 532	0.03	14.98	12	21 651.6	1 408	0.03	15.38
2004	19	25 280.6	1 626	0.01	15.55	14	23 476.4	1 533	0.01	15.31
2005	12	15 571.9	1 141	0.01	13.65	9	15 199.7	1 118	0.01	13.60
2006	11	13 928.3	883	0.01	15.77	9	13 709.5	865	0.01	15.85
2007	7	10 773.4	843	0.01	12.78	6	10 611.5	832	0.01	12.75
2008	5	6 239.5	324	0.01	19.26	3	6 147.6	315	0.01	19.52
2009	8	6 856.9	589	0.01	11.64	6	6 491.2	555	0.01	11.70
2010	8	9 423.8	729	0.01	12.93	5	9 301.3	719	0.01	12.94
2011	6	6 202.5	453	0.01	13.69	5	6 184.9	450	0.01	13.74
2012	9	8 879.8	705	0.01	12.60	6	8 711.9	693	0.01	12.57
2013	9	9 150.1	730	-	12.53	5	8 899.8	695	-	12.81
2014	10	9 537.9	838	0.01	11.38	8	9 138.8	805	0.01	11.35
2015	9	9 855.2	693	-	14.22	8	9 790.7	690	-	14.19
2016	9	8 928.4	534	0.01	16.72	6	8 611.5	508	-	16.95

## Cook Strait: Observer catch for target hoki

Fishing year	All vessels					Final vessels				
	No. vessels	Catch	Effort	Prop. zeros	CPUE	No. vessels	Catch	Effort	Prop. zeros	CPUE
1998	11	3 355.4	197	0.05	17.03	9	2 429.5	141	0.03	17.23
1999	10	3 458.1	212	0.01	16.31	8	2 518.0	152	0.01	16.57
2000	7	3 016.9	151	0.01	19.98	7	2 236.5	111	-	20.15
2001	9	4 020.5	228	0.01	17.63	6	2 350.2	143	-	16.43
2002	9	1 989.8	135	0.04	14.74	5	1 052.5	75	-	14.03
2003	5	2 415.9	131	0.02	18.44	4	1 368.4	81	-	16.89
2004	7	2 482.5	126	0.03	19.70	6	1 726.0	90	0.02	19.18
2005	9	2 151.2	122	-	17.63	4	1 384.3	65	-	21.30
2006	5	1 079.8	65	-	16.61	4	831.3	47	-	17.69
2007	7	2 013.0	164	0.02	12.27	7	1 553.3	133	0.01	11.68
2008	6	3 409.2	198	0.02	17.22	5	2 611.1	145	0.01	18.01
2009	4	1 956.4	149	0.03	13.13	4	1 409.5	108	-	13.05
2010	9	3 181.6	229	0.01	13.89	7	2 378.2	183	0.01	13
2011	5	1 588.3	86	-	18.47	5	1 281.8	70	-	18.31
2012	7	1 516.1	141	0.03	10.75	7	1 298.1	122	0.04	10.64
2013	4	955.6	85	-	11.24	4	827.3	74	-	11.18
2014	4	2 288.5	187	0.01	12.24	4	1 984.7	175	0.01	11.34
2015	2	319.4	20	-	15.97	2	303.4	19	-	15.97
2016	4	2 104.5	133	0.01	15.82	4	1 864.0	119	0.01	15.66

Table A3 continued.

## Chatham Rise: TCEPR tow-by-tow, target hoki

Fishing year	All vessels					Final vessels				
	No. vessels	Catch	Effort	Prop. zeros	CPUE	No. vessels	Catch	Effort	Prop. zeros	CPUE
1992	39	31 885.7	4 408	0.02	7.23	4	6 544.3	1 028	0.01	6.37
1993	37	27 904.8	4 054	0.01	6.88	3	9 829.4	1 842	0.01	5.34
1994	29	13 998.7	2 872	0.02	4.87	4	9 022.5	1 874	0.01	4.81
1995	35	20 268.9	4 404	0.03	4.60	8	15 174.6	3 272	0.02	4.64
1996	45	28 474.7	6 105	0.03	4.66	9	23 465.7	4 081	0.02	5.75
1997	63	34 656.4	7 253	0.03	4.78	10	27 635.0	4 773	0.02	5.79
1998	61	43 226.9	8 732	0.02	4.95	16	40 580.3	7 615	0.02	5.33
1999	44	54 455.4	9 735	0.01	5.59	15	53 190.8	9 216	0.01	5.77
2000	33	38 013.7	8 126	0.01	4.68	15	37 579.1	7 873	0.01	4.77
2001	38	37 590.9	8 361	0.01	4.50	15	35 758.1	7 849	0.01	4.56
2002	29	28 745.8	6 584	0.01	4.37	14	27 412.3	6 134	-	4.47
2003	29	28 990.5	7 699	0.01	3.77	15	26 922.9	7 081	0.01	3.80
2004	27	19 357.0	5 280	0.01	3.67	12	18 161.4	4 887	0.01	3.72
2005	23	21 196.3	3 982	0.01	5.32	10	20 061.3	3 621	0.01	5.54
2006	17	26 831.9	4 217	-	6.36	11	25 218.4	3 904	-	6.46
2007	17	27 518.0	3 805	0.01	7.23	9	25 735.8	3 508	0.01	7.34
2008	22	25 856.1	3 252	-	7.95	8	23 421.4	2 866	-	8.17
2009	19	27 248.2	3 195	-	8.53	7	23 632.9	2 714	-	8.71
2010	20	29 668.8	3 718	-	7.98	8	27 726.2	3 428	-	8.09
2011	21	30 551.1	3 619	-	8.44	8	27 263.6	3 208	-	8.50
2012	24	31 571.8	3 699	-	8.54	8	28 685.5	3 238	-	8.86
2013	20	28 407.6	3 619	-	7.85	7	25 930.7	3 238	-	8.01
2014	17	28 515.8	3 487	-	8.18	8	27 032.3	3 104	-	8.71
2015	20	34 618.3	3 910	-	8.85	8	31 717.6	3 536	-	8.97
2016	14	29 269.4	3 540	-	8.27	8	28 707.3	3 421	-	8.39

## Chatham Rise: Observer catch for target hoki

Fishing year	All vessels					Final vessels				
	No. vessels	Catch	Effort	No. zeros	CPUE	No. vessels	Catch	Effort	No. zeros	CPUE
1994	5	2 915.8	568	0.01	5.13	2	1 246.8	252	-	4.95
1995	3	1 098.9	268	0.01	4.10	2	1 095.9	264	0.01	4.15
1996	4	1 560.7	274	0.01	5.70	2	1 376.0	225	-	6.12
1997	6	756.3	186	0.01	4.07	2	278.3	78	0.03	3.57
1998	13	4 334.7	864	0.01	5.02	12	4 248.8	837	0.01	5.08
1999	12	7 291.4	1 125	0.01	6.48	12	7 258.6	1 118	0.01	6.49
2000	8	2 948.8	536	0.02	5.50	6	2 934.6	522	0.02	5.62
2001	13	3 135.0	765	0.09	4.10	11	3 104.7	752	0.09	4.13
2002	10	4 224.9	921	-	4.59	10	4 224.8	920	-	4.59
2003	9	1 677.1	615	-	2.73	8	1 676.6	613	-	2.74
2004	6	1 306.8	298	0.07	4.39	4	1 298.7	289	0.01	4.49
2005	7	3 902.6	629	-	6.20	5	3 839.8	614	-	6.25
2006	8	3 393.3	461	-	7.36	7	3 381.9	452	-	7.48
2007	8	5 121.8	736	-	6.96	7	5 114.5	732	-	6.99
2008	7	4 832.6	642	-	7.53	7	4 662.4	600	-	7.77
2009	10	3 548.5	449	0.02	7.90	9	3 315.9	394	0.02	8.42
2010	9	5 379.0	577	-	9.32	9	2 804.5	344	0.01	8.15
2011	10	5 497.9	668	0.01	8.23	8	4 275.6	514	-	8.32
2012	8	6 622.9	791	0.01	8.37	6	5 234.2	649	0.01	8.07
2013	15	8 889.0	1 295	0.01	6.86	12	7 431.7	1 089	0.01	6.82
2014	11	8 657.4	1 096	0.01	7.90	10	7 544.0	942	0.01	8.01
2015	12	4 186.7	448	-	9.35	10	3 210.2	341	-	9.41
2016	12	6 168.0	923	0.01	6.68	11	5 225.3	744	-	7.02



Table A3 continued.

## Sub-Antarctic: TCEPR tow-by-tow, target hoki

Fishing year	All vessels					Final vessels				
	No. vessels	Catch	Effort	Prop. zeros	CPUE	No. vessels	Catch	Effort	Prop. zeros	CPUE
1992	25	21 855.6	4 085	0.02	5.35	4	7 422.2	1 355	0.01	5.48
1993	23	18 731.0	4 187	0.02	4.47	4	8 119.1	1 739	0.01	4.67
1994	15	6 974.4	1 481	0.02	4.71	3	4 288.2	925	0.01	4.64
1995	24	10 308.1	2 012	0.03	5.12	6	6 406.1	1 391	0.03	4.61
1996	22	9 155.4	2 039	0.06	4.49	6	5 996.6	1 304	0.02	4.60
1997	37	12 362.3	2 173	0.03	5.69	13	11 411.6	1 829	0.03	6.24
1998	35	18 571.1	3 391	0.02	5.48	14	17 785.1	3 126	0.02	5.69
1999	30	15 982.1	2 775	0.02	5.76	12	13 590.3	2 268	0.02	5.99
2000	26	23 440.4	4 821	0.02	4.86	16	22 683.2	4 480	0.02	5.06
2001	31	17 463.0	4 033	0.02	4.33	15	16 589.9	3 715	0.02	4.47
2002	32	20 901.3	5 021	0.03	4.16	16	20 007.0	4 633	0.02	4.32
2003	29	15 929.1	3 727	0.02	4.27	15	15 311.0	3 474	0.01	4.41
2004	23	8 302.4	2 393	0.06	3.47	10	7 946.9	2 229	0.05	3.57
2005	25	4 524.9	1 216	0.06	3.72	7	4 150.1	1 026	0.05	4.04
2006	15	3 336.7	525	0.03	6.36	6	3 062.3	450	0.02	6.81
2007	20	3 431.3	802	0.05	4.28	9	3 078.8	721	0.03	4.27
2008	13	4 742.6	751	0.01	6.32	5	4 675.3	720	0.01	6.49
2009	12	5 696.0	743	0.01	7.67	4	5 616.0	723	0.01	7.77
2010	12	7 912.1	961	0.02	8.23	5	7 562.7	911	0.01	8.30
2011	15	8 032.7	1 112	0.01	7.22	5	7 329.4	1 021	0.01	7.18
2012	17	10 275.7	1 032	-	9.96	7	9 306.7	954	-	9.76
2013	16	9 796.2	1 134	0.01	8.64	6	9 503.0	1 082	0.01	8.78
2014	13	15 056.0	1 564	-	9.63	6	14 723.6	1 492	-	9.87
2015	15	11 430.4	1 349	-	8.47	8	10 608.1	1 205	-	8.80
2016	10	4 102.8	626	0.02	6.55	6	4 066.8	609	0.01	6.68

## Sub-Antarctic: Observer catch for target hoki

Fishing year	All vessels					Final vessels				
	No. vessels	Catch	Effort	Prop. zeros	CPUE	No. vessels	Catch	Effort	Prop. zeros	CPUE
1998	6	1 304.4	219	-	5.96	4	1 246.6	201	-	6.20
1999	9	4 546.2	686	0.02	6.63	7	4 436.0	658	0.01	6.74
2000	8	3 374.0	809	0.01	4.17	7	3 370.9	803	0.01	4.20
2001	13	1 546.4	355	0.01	4.36	7	1 463.1	315	-	4.64
2002	7	3 035.5	628	-	4.83	5	2 998.5	616	-	4.87
2003	9	1 311.6	349	0.01	3.76	6	1 307.0	344	0.01	3.80
2004	7	443.4	156	0.17	2.84	3	413.7	144	0.15	2.87
2005	3	354.0	88	0.03	4.02	3	354.0	88	0.03	4.02
2006	4	797.2	92	-	8.67	3	762.1	80	-	9.53
2007	6	875.2	154	-	5.68	4	849.6	152	-	5.59
2008	5	1 898.6	317	0.01	5.99	3	1 612.9	213	-	7.57
2009	4	2 644.1	364	0.01	7.26	3	2 621.6	360	0.01	7.28
2010	6	2 311.8	280	0.07	8.26	4	2 023.8	238	0.03	8.50
2011	8	2 006.8	274	0.02	7.32	3	1 466.5	197	0.02	7.44
2012	5	1 631.3	192	-	8.50	4	1 316.7	147	-	8.96
2013	14	4 264.5	587	0.01	7.26	9	3 926.4	530	-	7.41
2014	12	4 257.2	539	0.01	7.90	7	3 375.4	388	-	8.70
2015	11	2 258.5	313	-	7.22	6	1 827.2	240	-	7.61
2016	6	1 026.2	163	0.02	6.30	2	734.0	110	0.02	6.67

**Table A4: Lognormal CPUE standardised indices, and binomial, and combined CPUE indices (with 95% confidence intervals).**

**WCSI: TCEPR tow-by-tow, target hoki**

Year	Index	CI
1990	1.09	1.04–1.14
1991	1.16	1.11–1.21
1992	1.16	1.11–1.21
1993	1.00	0.96–1.04
1994	0.95	0.93–0.98
1995	0.62	0.60–0.63
1996	0.72	0.70–0.74
1997	0.74	0.72–0.75
1998	0.90	0.87–0.92
1999	0.94	0.91–0.96
2000	1.06	1.03–1.09
2001	0.78	0.76–0.80
2002	0.76	0.75–0.78
2003	0.59	0.58–0.61
2004	0.38	0.37–0.39
2005	0.46	0.45–0.48
2006	0.69	0.67–0.72
2007	1.17	1.11–1.22
2008	1.10	1.04–1.16
2009	1.52	1.43–1.62
2010	1.47	1.40–1.54
2011	1.48	1.42–1.55
2012	1.72	1.65–1.79
2013	1.88	1.80–1.95
2014	1.67	1.61–1.73
2015	1.79	1.73–1.85
2016	1.39	1.35–1.44

**WCSI: Observer catch, target hoki**

Year	Index	CI
1990	0.98	0.85–1.13
1991	1.04	0.93–1.17
1992	1.13	0.97–1.33
1993	0.92	0.82–1.02
1994	0.96	0.89–1.04
1995	0.70	0.62–0.79
1996	0.75	0.69–0.82
1997	0.86	0.78–0.94
1998	0.93	0.86–1.01
1999	0.83	0.77–0.90
2000	1.06	0.99–1.13
2001	0.71	0.66–0.77
2002	0.80	0.75–0.86
2003	0.50	0.46–0.54
2004	0.30	0.28–0.32
2005	0.45	0.42–0.48
2006	0.78	0.72–0.84
2007	1.29	1.18–1.42
2008	1.08	0.97–1.19
2009	1.59	1.43–1.76
2010	1.38	1.25–1.53
2011	1.77	1.61–1.95
2012	1.84	1.70–1.99
2013	1.95	1.84–2.07
2014	1.79	1.69–1.90
2015	1.88	1.77–1.99
2016	1.33	1.26–1.41

**Cook Strait: TCEPR MW tow-by-tow, target hoki**

Year	Index	CI
1990	1.17	1.08–1.26
1991	0.98	0.93–1.04
1992	1.11	1.04–1.18
1993	1.02	0.96–1.09
1994	1.26	1.18–1.33
1995	1.25	1.18–1.33
1996	1.11	1.05–1.16
1997	0.95	0.91–0.99
1998	1.02	0.97–1.07
1999	1.00	0.95–1.05
2000	1.00	0.95–1.06
2001	0.83	0.79–0.88
2002	1.25	1.17–1.35
2003	1.00	0.95–1.06
2004	0.91	0.86–0.96
2005	0.85	0.80–0.90
2006	1.00	0.94–1.07
2007	0.77	0.72–0.83
2008	1.16	1.04–1.30
2009	0.77	0.71–0.84
2010	0.96	0.89–1.03
2011	1.14	1.04–1.26
2012	0.84	0.78–0.90
2013	0.92	0.85–0.99
2014	0.90	0.84–0.97
2015	0.96	0.89–1.04
2016	1.11	1.02–1.22

**Cook Strait: Observer catch, target hoki**

Year	Index	CI
1998	0.97	0.80–1.18
1999	1.03	0.85–1.25
2000	1.01	0.82–1.25
2001	0.83	0.67–1.02
2002	0.79	0.61–1.02
2003	0.62	0.47–0.81
2004	1.23	0.98–1.54
2005	1.54	1.17–2.02
2006	1.18	0.88–1.58
2007	0.61	0.50–0.74
2008	1.14	0.95–1.38
2009	0.69	0.55–0.87
2010	0.94	0.78–1.13
2011	1.40	1.09–1.79
2012	0.93	0.76–1.15
2013	1.08	0.84–1.40
2014	1.14	0.95–1.37
2015	0.94	0.56–1.57
2016	1.61	1.30–2.00

**Table A4: continued.**

**Chatham Rise: TCEPR tow-by-tow, target hoki**

Year	Index	CI
1992	1.07	1.00–1.14
1993	1.00	0.95–1.05
1994	0.97	0.93–1.02
1995	0.84	0.81–0.88
1996	1.02	0.98–1.05
1997	0.94	0.91–0.97
1998	0.86	0.84–0.88
1999	0.97	0.95–1.00
2000	0.78	0.77–0.80
2001	0.74	0.72–0.75
2002	0.73	0.72–0.75
2003	0.56	0.55–0.58
2004	0.54	0.52–0.55
2005	0.76	0.74–0.79
2006	1.00	0.97–1.04
2007	1.04	1.01–1.08
2008	1.29	1.24–1.34
2009	1.43	1.38–1.49
2010	1.24	1.20–1.29
2011	1.32	1.28–1.37
2012	1.41	1.36–1.46
2013	1.29	1.24–1.34
2014	1.39	1.34–1.44
2015	1.41	1.36–1.46
2016	1.29	1.24–1.33

**Chatham Rise: Observer catch, target hoki**

Year	Index	CI
1994	1.19	1.04–1.37
1995	0.75	0.64–0.87
1996	0.99	0.85–1.15
1997	0.46	0.35–0.59
1998	0.75	0.69–0.81
1999	0.92	0.85–0.99
2000	0.87	0.79–0.96
2001	0.63	0.57–0.68
2002	0.73	0.67–0.79
2003	0.52	0.48–0.58
2004	0.79	0.66–0.95
2005	0.87	0.79–0.96
2006	1.18	1.06–1.32
2007	1.16	1.07–1.25
2008	1.37	1.25–1.50
2009	1.45	1.30–1.62
2010	1.50	1.33–1.69
2011	1.30	1.18–1.43
2012	1.36	1.24–1.49
2013	1.23	1.14–1.33
2014	1.50	1.38–1.62
2015	1.58	1.39–1.78
2016	1.20	1.11–1.31

**Sub-Antarctic: TCEPR tow-by-tow, target hoki**

Year	Index	CI
1992	0.98	0.92–1.05
1993	0.92	0.87–0.98
1994	1.08	1.00–1.17
1995	0.89	0.83–0.94
1996	0.93	0.88–0.99
1997	1.21	1.15–1.28
1998	1.03	0.99–1.07
1999	0.94	0.90–0.98
2000	0.87	0.84–0.90
2001	0.78	0.75–0.81
2002	0.76	0.73–0.78
2003	0.75	0.72–0.78
2004	0.53	0.50–0.55
2005	0.56	0.53–0.60
2006	0.78	0.71–0.86
2007	0.66	0.61–0.71
2008	1.09	1.01–1.18
2009	1.35	1.26–1.46
2010	1.41	1.32–1.51
2011	1.30	1.22–1.39
2012	1.67	1.56–1.78
2013	1.47	1.38–1.56
2014	1.61	1.53–1.70
2015	1.47	1.39–1.56
2016	1.14	1.05–1.24

**Sub-Antarctic: Observer catch, target hoki**

Year	Index	CI
1998	1.09	0.88–1.35
1999	1.00	0.89–1.12
2000	0.66	0.59–0.74
2001	0.99	0.85–1.16
2002	0.68	0.61–0.77
2003	0.55	0.48–0.64
2004	0.84	0.68–1.02
2005	0.58	0.46–0.73
2006	1.12	0.86–1.47
2007	1.08	0.87–1.35
2008	1.39	1.20–1.62
2009	1.14	1.00–1.31
2010	1.33	1.14–1.55
2011	1.38	1.16–1.64
2012	1.07	0.89–1.29
2013	1.12	1.00–1.25
2014	1.51	1.35–1.70
2015	1.17	1.01–1.35
2016	1.03	0.85–1.26