



Settlement indices for 2016/17 fishing year for the red rock lobster (*Jasus edwardsii*)

New Zealand Fisheries Assessment Report 2018/16

J. Forman,
A. McKenzie,
D. Stotter

ISSN 1179-5352 (online)
ISBN 978-1-77665-849-7 (online)

May 2018



Requests for further copies should be directed to:

Publications Logistics Officer
Ministry for Primary Industries
PO Box 2526
WELLINGTON 6140

Email: brand@mpi.govt.nz
Telephone: 0800 00 83 33
Facsimile: 04-894 0300

This publication is also available on the Ministry for Primary Industries websites at:
<http://www.mpi.govt.nz/news-and-resources/publications>
<http://fs.fish.govt.nz> go to Document library/Research reports

© Crown Copyright - Ministry for Primary Industries

Table of Contents

Executive Summary	1
1. INTRODUCTION	2
2. METHODS	3
2.1 Recording settlement on collectors	3
2.2 Calculating indices of settlement	5
3. RESULTS	8
3.1 Standardised indices	8
4. SUMMARY AND DISCUSSION	10
5. CONCLUSIONS	19
6. ACKNOWLEDGMENTS	19
7. REFERENCES	19
Appendix 1: Data characteristics and diagnostics	21

EXECUTIVE SUMMARY

Forman, J.S.; McKenzie, A.; Stotter, D.R. (2018). Settlement indices for 2016/17 fishing year for the red rock lobster (*Jasus edwardsii*).

New Zealand Fisheries Assessment Report 2018/16. 43 p.

This report addresses objective one of the Ministry for Primary Industries project CRA201502C (Estimating settlement).

We update the information on annual patterns of settlement for the red rock lobster (*Jasus edwardsii*) on crevice collectors at key sites in CRA 3 (Gisborne), CRA 4 (Napier and Castlepoint), CRA 5 (Kaikoura), CRA 7 (Moeraki), and CRA 8 (Halfmoon Bay and Jackson Bay).

In the 2016/17 fishing year, two groups of collectors in Gisborne, Napier, and Castlepoint, four groups in Kaikoura, and one group in each of Moeraki, Halfmoon Bay and Jackson Bay were monitored. Each group has at least three collectors that are checked monthly when possible and a monthly mean catch per group of collectors is calculated. A raw and standardised index based on the rock lobster fishing year, April to March, is produced from the groups of collectors at each site.

Puerulus settlement in 2016/17 was notable for the high levels of settlement recorded at Castlepoint, Kaikoura, Moeraki, and Jackson Bay. Settlement was just above the long term mean at Gisborne and Halfmoon Bay, and below the long term mean at Napier. This is the fifth consecutive year of very high settlement at Jackson Bay and the third out of the last four years of higher than average settlement at Castlepoint.

Due to the Kaikoura earthquake in November 2016, the collectors at South Bay (KAI001) and North Bay (KAI003) were uplifted out of the water. Fortunately, the collectors could be moved into deeper water close to the original location, however, it remains unclear how much the movement will affect future settlement. The other two sites to the south of the Kaikoura peninsula were seemingly unaffected.

1. INTRODUCTION

Rock lobsters are one of New Zealand's most valuable fisheries. Understanding larval recruitment processes will greatly assist the management of this fishery because it may explain changes in levels of recruitment to the fishery and enable the prediction of trends in catch levels at least four years in advance, allowing management and commercial strategies to be implemented. This report updates the patterns of spatial and temporal settlement of *Jasus edwardsii* on crevice collectors in New Zealand.

Rock lobsters spend several months as phyllosoma larvae in waters tens to hundreds of kilometres offshore. They return to the shore as postlarvae (pueruli) after metamorphosing near the shelf break. The puerulus is the settling stage: it resembles the juvenile in shape and is 9–13 mm in carapace length, but it is transparent. Pueruli settle when they cease extensive forward swimming and take up residence on the substrate. Some older pueruli and young juveniles, however, move after first settling elsewhere. Post-settlement migration (secondary dispersal) such as this is not uncommon among invertebrates (e.g., Reynolds & Eggleston 2004), the young redistributing from high-density settlement habitats is thought to be a strategy to reduce density-dependent mortality. The puerulus moults into the first juvenile instar (sometimes referred to as the first-moult postpuerulus) a few days to three weeks after settlement. Higher water temperatures reduce the time taken to moult. Depending on sex and locality, the rock lobster then takes about 4–11 years to reach minimum legal size.

The development of sampling programmes to estimate levels of postlarval settlement that can be used to predict fishery performance is a goal for both palinurids (e.g., Phillips et al. 2000, Gardner et al. 2001) and homarids (e.g., Wahle et al. 2004), with encouraging or well-demonstrated success for some projects. In New Zealand there are significant correlations between the level of settlement and the fishery catch per unit effort (CPUE) for most fishery areas. The best correlations occur in fisheries with shorter intervals between settlement and recruitment, and in those with large contrasts in the settlement record (Booth & McKenzie 2008).

Monthly occurrence of pueruli and young juveniles on crevice collectors (Booth & Tarring 1986) has been followed at up to nine key sites within the main New Zealand rock lobster fishery since the early 1980s. The indices of settlement are now reported annually. It has become clear from this and other monitoring, that settlement is not uniform in time or space. Settlement occurs mainly at night and at any lunar phase, is seasonal, and levels of settlement can vary by an order of magnitude or more from year to year (Booth & Stewart 1993, Forman et al. 2014).

For detailed further information on the puerulus sampling program in New Zealand see Booth et al. (2006).

OBJECTIVES

1. To determine trends in puerulus settlement at selected key sites around New Zealand.

Specific Objectives

To estimate monthly and annual indices of puerulus settlement at key sites in CRA 3, CRA 4, CRA 5, CRA 7 and CRA 8 (Gisborne, Napier, Castlepoint, Kaikoura, Moeraki, Halfmoon Bay, and Jackson Bay).

2. METHODS

2.1 Recording settlement on collectors

Levels of puerulus settlement are monitored using ‘crevice’ collectors (Booth & Tarring 1986, Booth et al. 1991) at seven key sites that encompass much of the main rock lobster fishing coast of New Zealand. The collector was developed in New Zealand to catch *J. edwardsii* pueruli and is now used throughout much of the range of *J. edwardsii*. They are inexpensive, easily set and checked, and provide (unlike many other types of collector) a standard settlement surface for between-month and between-site comparisons.

Each key site is separated from its neighbour by 150–400 km, and most sites were chosen after trying many locations (Figure 1). Criteria for the establishment of key sites included the distance from the neighbouring site, proximity to the open ocean, accessibility, tractability, and the level of puerulus catch.

At each key site, collectors are set in groups of between 3 and 20, with at least 2–3 m between individual collectors. It is unclear whether or not there is interference in the catch between collectors at these spacings, but because the distances remain unaltered, any interference is likely to have a minimal impact on the overall monthly and annual index. At each site there is a core group of at least three (although usually five) collectors. At most sites there have been up to three additional groups of three or more collectors, set in both directions along the coast as conditions allow. Since 2002, however, fewer of these additional groups of collectors have been monitored; the focus is now on the core group (usually the one first established, and therefore with the longest record of settlement). Where feasible, one other group of collectors is also monitored. See Table 1 for a summary of the collector sites, the number of collectors by site, and the method of collector deployment. Methods of deployment include shore based collectors which are attached to concrete weights in sheltered subtidal locations, suspended collectors which are hung from wharf piles with the collectors suspended just off the bottom, and closing collectors which have a closing mechanism that surrounds the collector as it is hauled up by boat.

Collectors are generally checked monthly as weather and tides allow and are cleaned of heavy growth so that the condition of collectors is consistent. Repairs required are noted at each collector check and these are made in the field where possible. Spare (and conditioned) collectors are maintained at each site or nearby as replacements. If possible, collector replacement is made outside the main settlement season.

At most sites, local people are employed to check the collectors, under NIWA’s direction. Quality control of checks and equipment is maintained with direct contact once or twice a year. A standard result form is filled out and sent to NIWA after each check. At Castlepoint and Moeraki, NIWA staff check the collectors. Monthly checks, especially during the main winter settlement season, are not always possible for all groups of collectors because of logistical issues. Two groups of collectors in Kaikoura (KAI005 and KAI006) are fully managed by CRAMAC 5 and one other group of collectors in Kaikoura (KAI003) is funded by CRAMAC 5 but is maintained by NIWA.




Figure 1: Map of New Zealand showing the location of collectors at the key monitoring sites (although not all groups are now checked). The sites that are checked are in bold and the number of collectors in that set is in brackets. Also shown are the CRA areas; CRA 6 is the Chatham Islands and CRA 10 is the Kermadec Islands (to the northeast of the North Island).

Table 1: Number of collectors, method of collector deployment, and years of operation of all collectors used in the settlement index. For definitions of collector type see Section 2.1, Booth & Tarring (1986), and Phillips & Booth (1994).

Site	Number of collectors	Location	Method of deployment	Years of operation
Gisborne	5	Harbour (GIS001)	Shore	1987–2003
	5	Whangara (GIS002)	Shore	1991–Present
	5	Tatapouri (GIS003)	Shore	1994–2006
	5	Kaiti (GIS004)	Shore	1994–Present
Napier	5	Harbour (NAP001)	Suspended	1979–Present
	3	Westshore (NAP002)	Closing	1991–1999
	5	Cape Kidnappers (NAP003)	Shore	1994–Present
	3	Breakwater (NAP004)	Shore	1991–2002
Castlepoint	9	Castlepoint (CPT001)	Shore	1983–Present
	5	Orui (CPT002)	Shore	1991–Present
	5	Mataikona (CPT003)	Shore	1991–2006
Kaikoura	5	South peninsula (KAI001)	Shore	1981–Present
	3	South peninsula (KAI002)	Shore	1988–2003
	5	North peninsula (KAI003)	Shore	1980–Present
	3	North peninsula (KAI004)	Shore	1992–2003
	4	South Kaikoura (KAI005)	Shore	2008–Present
	3	Haumuri Bluff (KAI006)	Shore	2008–Present
	3	Gooch Bay (KAI008)	Shore	1980–1983
	3	Middle South Coast (KAI009)	Shore	1981–1988
	Moeraki	3	Wharf (MOE002)	Closing
6		Pier (MOE007)	Suspended	1998–Present
Halfmoon Bay	8	Wharf (HMB001)	Suspended	1980–Present
	3	Thompsons (HMB002)	Closing	1988–2002
	3	Old Mill (HMB003)	Closing	1990–2002
	3	The Neck (HMB004)	Closing	1992–2002
	3	Mamaku Point (HMB005)	Closing	1992–2002
Jackson Bay	5	Jackson wharf (JAC001)	Suspended	1999–Present
	3	Jackson Head (JAC002)	Closing	1999–2006
Chalky Inlet	4	Chalky Inlet (CHI001)	Closing	1986–2012

2.2 Calculating indices of settlement

All standardisations before 2014 used settlement data based on the Calendar year. In contrast all standardisations presented here use the fishing year from 1 April to 31 March, with the year label being that which April is in. For example, 20 April 2004 is in the 2004 fishing year, while 11 Feb 2004 is in the 2003 fishing year. An alternative labelling also used is to call the 2004 fishing year the 2004/05 fishing year. The change from a calendar to a fishing year was requested by the Rock Lobster Working Group (RLWG), and aligns with the year used in stock assessments, for which the standardised puerulus indices are offered as data inputs.

The standardised index of annual settlement used here incorporates all settlement for the year for each site, irrespective of month. This approach to the standardisation was based on Bentley et al. (2004), but with

the adjustments noted below: assignment of the month for settlement, and the groups of collectors used. The term ‘settlement’ refers to the presence of pueruli and juveniles up to 14.5 mm carapace length (CL, the maximum size for a first-instar juvenile observed in laboratory studies).

Following Bentley et al. (2004) the standardisation used collectors that were sampled at least 36 times (equivalent to three years of monthly sampling). No outliers were removed from any of the data sets after fitting. In Bentley et al. (2004) outliers were removed, but the effect on the standardised indices was minor.

Because a collector check on any one day is thought to be a snapshot of what has been going on for about the last 14 days, then the appropriate month to label as settlement may not be the nominal month. In standardisations prior to 2014, if the check took place up to the seventh of the month its catch was attributed to the previous month. This also avoids the situation where if a collector is checked on the first and last day of a month, there are two records for that month, but none for the previous or subsequent months. Nonetheless, it was decided by the RLWG that the nominal month should be used for standardisations.

At three sites (Gisborne, Jackson Bay, and Moeraki) some pilot groups of collectors were dropped because of logistical reasons or for very low counts. For Jackson Bay and Moeraki even the best groups of collectors, after dropping of pilot groups, recorded very low counts (Forman et al. 2014, Appendix 1). To reduce the proportion of zero counts at some sites, the months used for the standardisation are restricted to those where counts are high (Table 2). In some fishing years the number of samples is low, and that year is dropped from the standardisation if the number of samples to the site is less than 10.

The annual index takes into account changes in collector location and sampling to date. A generalised linear model framework was used, in which the response (dependent) variable is the log of numbers of settlers per collector sample and a negative binomial distribution is used. For Kaikoura and Moeraki alternative distributions were investigated (quasi-Poisson, zero-inflated Poisson, zero-inflated negative binomial) and the negative binomial was chosen as the best. In a previous standardisation for Gisborne alternative distributions were investigated and the negative binomial was chosen as the best (Forman et al. 2015). For the other sites the RLWG decided that the negative binomial distribution should be used.

The predictor variables available to the standardisations were year, group, and month. The year variable was included in all models; the other independent variables group and month were added to the model in a stepwise process. At each step the variable that most improved the fit of the model measured by the Akaike Information Criterion (AIC) was included (Akaike 1974).

In summary, the standardisation method common to all sites was to:

1. use the fishing year from 1st April to 31st March
2. use the actual month in which a sample was taken (instead of samples taken up to the 7th of the month being assigned to the previous month)
3. drop collectors with fewer than 36 samples
4. restrict where necessary the months used (to reduce the proportion of zero counts in the data)
5. drop fishing years with fewer than 10 samples
6. use a negative binomial model for the data
7. use year, month, and group (collector is not offered as an alternative to group) as the predictor variables in the standardisation.

Each set of annual indices is presented as the annual value divided by the geometric mean of the annual values, or where the annual values are close to zero (Moeraki and Halfmoon Bay) by dividing by the arithmetic mean of the annual values. In either case, a value for the index above 1 represents above average settlement for that year, and a value below 1 indicates below average settlement. For comparison, a raw form of these indices is also given (arithmetic mean for each year), which is also scaled to have an average value of 1 over all years.

The data set used for all sites is an extract from the *rocklob* database and is complete for the 2016/17 fishing year (i.e. data is complete up to 31 March 2017).

3. RESULTS

3.1 Standardised indices

Standardised indices were produced for the sites Gisborne (CRA 3), Napier (CRA 4), Castlepoint (CRA 4), Kaikoura (CRA 5), Moeraki (CRA 7), Halfmoon Bay (CRA 8), and Jackson Bay (CRA 8). There were no new data for Chalky Inlet (CRA 8). To reduce the number of zeros a subset of months was used for many of the standardisations (Table 2).

For each site plots are given in the following sections for puerulus data characteristics, standardised index, and standardisation diagnostics. Diagnostics look reasonable for all sites (Appendix 1). The standardised indices for all sites are summarised in Table 3.

Table 2: Months for which data were used in standardisation.

Site	Months
Gisborne	May–September
Napier	All
Castlepoint	All
Kaikoura	January–September
Moeraki	May–October
Halfmoon Bay	May–December
Jackson Bay	All

Table 3: Standardised annual indices for each site. Year is fishing year 1 April–31 March. ‘-’: no usable sampling was done; 0.00: no observed settlement.

Fishing year	Gisborne CRA 3	Napier CRA 4	Castlepoint CRA 4	Kaikoura CRA 5	Moeraki CRA 7	Halfmoon Bay CRA 8	Chalky Inlet CRA 8	Jackson Bay CRA 8
1979	–	0.78	–	–	–	–	–	–
1980	–	1.25	–	–	–	–	–	–
1981	–	2.05	–	0.53	–	8.14	–	–
1982	–	1.14	2.44	0.72	–	0.39	–	–
1983	–	1.33	1.19	0.16	–	3.92	–	–
1984	–	0.41	0.72	0.37	–	0.30	–	–
1985	–	0.22	0.57	0.23	–	0.00	0.36	–
1986	–	–	0.84	0.08	–	0.12	0.21	–
1987	3.24	–	1.64	1.03	–	1.59	1.42	–
1988	2.76	1.36	0.93	0.39	–	0.22	1.31	–
1989	0.97	1.18	1.14	0.78	–	0.60	1.64	–
1990	0.43	1.04	1.09	1.54	–	0.43	1.84	–
1991	1.05	2.45	2.12	6.58	0.00	0.93	1.03	–
1992	2.80	2.09	2.10	5.13	0.09	0.54	0.52	–
1993	1.75	2.21	1.05	2.01	0.00	0.00	0.14	–
1994	3.00	1.53	0.87	1.06	0.00	1.19	1.64	–
1995	1.07	1.06	0.91	0.59	0.07	0.40	0.40	–
1996	1.64	1.54	1.26	0.62	0.61	0.33	1.76	–
1997	0.98	1.08	1.68	1.94	0.26	0.56	1.41	–
1998	1.77	0.97	1.05	1.88	0.35	0.30	0.50	–
1999	0.28	0.43	0.34	1.25	0.06	0.23	1.70	0.24
2000	0.90	0.73	0.52	1.27	2.67	1.22	1.26	0.50
2001	1.12	1.23	0.70	0.53	1.11	1.75	0.60	0.20
2002	0.94	1.45	0.76	3.25	0.58	1.47	1.42	1.28
2003	2.71	1.31	0.93	3.31	4.82	3.94	1.56	0.48
2004	0.71	1.06	0.49	1.00	0.24	0.16	0.30	0.36
2005	2.46	1.28	1.26	2.20	0.05	0.00	–	1.20
2006	0.27	0.65	0.47	1.07	0.04	0.13	–	0.23
2007	0.36	0.92	1.03	1.66	0.04	0.48	–	0.21
2008	0.63	0.64	1.04	1.59	0.07	0.09	–	0.08
2009	1.69	0.89	1.07	0.52	0.44	1.03	–	0.14
2010	0.61	0.94	1.16	1.25	0.97	1.66	7.03	1.80
2011	0.18	0.49	0.89	0.56	0.69	0.14	1.44	1.97
2012	0.66	0.70	0.58	1.11	0.80	0.18	4.37	6.83
2013	0.92	0.95	1.69	0.71	1.17	0.76	–	11.95
2014	0.39	1.03	0.69	1.28	0.34	0.87	–	19.06
2015	1.48	1.05	1.65	0.86	7.73	0.56	–	4.92
2016	1.15	0.68	1.85	2.78	2.81	1.38	–	11.64

4. SUMMARY AND DISCUSSION

In this section trends over time for each site are discussed, and monthly puerulus settlement for currently operating collectors is plotted.

Gisborne

Settlement at Gisborne in 2016/17 was just above the long-term mean (Figure 2). The monthly settlement pattern between Whangara and Kaiti were similar except for two high settlement peaks recorded in Whangara during May and July (Figure 3).

Napier

Settlement at Napier was below the long-term mean (Figure 4). Over the last 11 years, settlement has been close to or below the long-term mean, and no significantly high settlement has occurred since the 1990s. The monthly settlement pattern between Napier Port and Kaiti were similar, unfortunately there was no check in July, to compare with the peak settlement month at Cape Kidnappers (Figure 5).

Castlepoint

For the third time in four years settlement at Castlepoint was above the long-term mean (Figure 6). It was also the highest annual settlement since 1992 and the fourth highest overall. Monthly levels of settlement between Castlepoint and Orui were similar with exceptionally high settlement from April to July then dropping down to low levels of settlement from September to November and rising to a summer peak in February (Figure 7).

Kaikoura

Kaikoura was well above the long-term mean and the fifth highest on record (Figure 8). North Bay, South Bay, and Haumuri Bluff had similar levels of settlement with South Kaikoura recording much higher levels in April and May (Figure 9). The South Bay and North Bay collectors were uplifted out of the water by the November 2016 Kaikoura earthquake and the affect this will have on the settlement to these collectors is unclear. To the south, the two sites South Kaikoura and Haumuri Bluff, were seemingly unaffected and comparisons in settlement patterns between the affected and unaffected areas will be made once more data is collected. The uplifted collectors were dragged a short distance into deeper water and at this stage all collectors seem to be catching well.

Moeraki

Although much lower than last year's record, settlement was still well above the long term mean and was the third highest on record (Figure 10). Most of the settlement occurred in June and July (Figure 11). No checks were done after November due to safety concerns as the old pier the collectors are suspended from is now too unstable and dangerous to use. Another location close to the pier has been established.

Halfmoon Bay

Settlement was above the long-term mean in Halfmoon Bay (Figure 12). The settlement peak, like Moeraki, occurred in July (Figure 13).

Jackson Bay

Settlement was the third highest on record and continues a five year series of very high settlement in this area (Figure 14). Settlement was high in all months of the year, peaking in August, and, like previous years, showed no clear pattern of settlement (Figure 15).

Chalky Inlet

No new data were received in 2016/17.

Monthly settlement comparisons for all key sites are shown in Figure 16.

Gisborne (001,002,003,004)




Figure 2: Standardised and raw indices of annual settlement for Gisborne with 95% confidence intervals.




Figure 3: Whangara and Kaiti monthly settlement, 2016/17 fishing year. Mean number of *Jasus edwardsii* pueruli plus juveniles less than 14.5 mm carapace length per collector.

Napier (001,002,003,004)




Figure 4: Standardised and raw indices of annual settlement for Napier with 95% confidence intervals. Note that there were no checks in 1986–87.

Napier




Figure 5: Napier Port and Cape Kidnappers monthly settlement, 2016/17 fishing year. Mean number of *Jasus edwardsii* pueruli plus juveniles less than 14.5 mm carapace length per collector.

Castlepoint (001,002,003)




Figure 6: Standardised and raw indices of annual settlement for Castlepoint with 95% confidence intervals.

Castlepoint




Figure 7: Castlepoint and Orui monthly settlement, 2016/17 fishing year. Mean number of *Jasus edwardsii* pueruli plus juveniles less than 14.5 mm carapace length per collector.

Kaikoura (001,002,003,004,005,006,008,009)




Figure 8: Standardised and raw indices of annual settlement for Kaikoura with 95% confidence interval.

Kaikoura




Figure 9: South Bay, North Bay, South Kaikoura, and Haumuri Bluffs monthly settlement, 2016/17 fishing year. Mean number of *Jasus edwardsii* pueruli plus juveniles less than 14.5 mm carapace length per collector.

Moeraki (002,007)




Figure 10: Standardised and raw indices of annual settlement for Moeraki with 95% confidence intervals.

Moeraki (MOE007)




Figure 11: Moeraki monthly settlement, 2016/17 fishing year. Mean number of *Jasus edwardsii pueruli* plus juveniles less than 14.5 mm carapace length per collector.

Halfmoon Bay (001,002,003,004,005)




Figure 12: Standardised and raw indices of annual settlement for Halfmoon Bay with 95% confidence intervals.

Halfmoon Bay (HMB001)




Figure 13: Halfmoon Bay monthly settlement, 2016/17 fishing year. Mean number of *Jasus edwardsii* pueruli plus juveniles less than 14.5 mm carapace length per collector.

Jackson Bay (001,002)




Figure 14: Standardised and raw indices of annual settlement for Jackson Bay with 95% confidence intervals.

Jackson Bay (JKB001)




Figure 15: Jackson Bay monthly settlement, 2016/17 fishing year. Mean number of *Jasus edwardsii* pueruli plus juveniles less than 14.5 mm carapace length per collector.




Figure 16: The mean settlement by month, over all years, for each key collector site. See Table 1 for the collector groups.

5. CONCLUSIONS

All sites except Napier recorded settlement above the long term-mean. At Jackson Bay, it was the seventh year of above average settlement with the last five years being exceptionally high. High levels of settlement also continued at Castlepoint, with three of the last four years well above the long-term mean, and at Moeraki, with very high settlement occurring over the last two years. Kaikoura recorded its highest level of settlement in 13 years and Gisborne was just above the long-term mean. For the last 11 years Napier has had average to below average settlement.

For Gisborne, Napier, and Castlepoint the puerulus index is potentially a signal for recruited abundance 4–6 years into the future. For Moeraki, the estimated interval is 4–5 years and for Halfmoon Bay it is 6–8 years (Booth & McKenzie 2008).

6. ACKNOWLEDGMENTS

Thank you to Evan Baddock, Andy and Glenys Bassett, Neil and Murray Burden, Phred Dobbins, Derek Kater, Ian Maze, Shane Metcalfe, Craig and Helen Petherick, Port of Napier, Paul Reinke, Eric Stevens, and CRAMAC 5 for collector checks and field assistance. Thank you to the Rock Lobster Working Group for suggestions made regarding the work, and Reyn Naylor for reviewing the document. This project was funded by the Ministry for Primary Industries under project CRA201502C.

7. REFERENCES

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19 (6): 716–723.
- Bentley, N.; Booth, J.D.; Breen, P.A. (2004). Calculating standardised indices of annual rock lobster settlement. *New Zealand Fisheries Assessment Report 2004/32*. 45 p.
- Booth, J.D.; Carruthers, A.D.; Bolt, C.D.; Stewart, R.A. (1991). Measuring depth of settlement in the red rock lobster, *Jasus edwardsii*. *New Zealand Journal of Marine and Freshwater Research* 25: 123–132.
- Booth, J.D.; McKenzie, A. (2008). Strong relationships between levels of puerulus settlement and recruited stock abundance in the red rock lobster (*Jasus edwardsii*) in New Zealand. *Fisheries Research* 95: 161–168.
- Booth, J.D.; McKenzie, A.; Forman, J.S.; Stotter, D.R. (2006). Monitoring puerulus settlement in the red rock lobster (*Jasus edwardsii*), 1974–2005, with analyses of correlation between settlement and subsequent stock abundance. Final Research Report for Ministry of Fisheries Research Project CRA2004-02. 76 p. (Unpublished report held by Ministry for Primary Industries, Wellington).
- Booth, J.D.; Stewart, R.A. (1993). Puerulus settlement in the red rock lobster, *Jasus edwardsii*. New Zealand Fisheries Assessment Research Document 93/5. 39 p. (Unpublished report held in NIWA Greta Point library, Wellington.)
- Booth, J.D.; Tarring, S.C. (1986). Settlement of the red rock lobster, *Jasus edwardsii*, near Gisborne, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 20: 291–297.
- Forman, J.; McKenzie, A.; Stotter, D. (2014). Settlement indices for 2012 for the red rock lobster (*Jasus edwardsii*). *New Zealand Fisheries Assessment Report 2014/47*. 56 p.

- Forman, J.; McKenzie, A.; Stotter, D (2015). Settlement indices for 2013 for the red rock lobster (*Jasus edwardsii*). *New Zealand Fisheries Assessment Report 2015/13*. 61 p.
- Gardner, C.; Frusher, S.D.; Kennedy, R.B.; Cawthorn, A. (2001). Relationship between settlement of southern rock lobster pueruli, *Jasus edwardsii*, and recruitment to the fishery in Tasmania, Australia. *Marine and Freshwater Research 52*: 1067–1075.
- Phillips, B.F.; Booth, J.D. (1994). Design, use, and effectiveness of collectors for catching the puerulus stage of spiny lobsters. *Reviews in Fisheries Science 2*: 255–289.
- Phillips, B.F.; Cruz, R.; Caputi, N.; Brown, R.S. (2000). Predicting the catch of spiny lobster fisheries. *In: Spiny lobsters. Fisheries and culture*. Phillips, B.F.; Kittaka, J. (eds) pp. 357–375. Blackwell Science, Oxford.
- Reyns, N.B.; Eggleston, D.B. (2004). Environmentally-controlled, density-dependent secondary dispersal in a local estuarine crab population. *Oecologia 140*: 280–288.
- Wahle, R.A.; Incze, L.S.; Fogarty, M.J. (2004). First projections of American lobster fishery recruitment using a settlement index and variable growth. *Bulletin of Marine Science 74*: 101–114.

APPENDIX 1: DATA CHARACTERISTICS AND DIAGNOSTICS

For each site plots are given for puerulus data characteristics, and standardisation diagnostics (Figures 17–44). To reduce the number of zeros a subset of months was used for many of the standardisations (see Table 2). The number of puerulus samples by group and fishing year at each site are given in Tables 4–10. Diagnostics look reasonable for all sites.

Gisborne




Figure 17: Characteristics of the Gisborne puerulus standardisation data. The top-left figure shows the distribution of puerulus counts, and the bottom-left the proportion of these that are non-zero. The top right figure shows the percentage of samples by month, and the bottom right shows the arithmetic mean of the number of pueruli measured in a sample.




Figure 18: Deviance residuals for the negative binomial model for Gisborne. Predicted values are in natural space.




Figure 19: Quantile-quantile plot for the negative binomial standardisation model for Gisborne.




Figure 20: Data distribution and that from the fitted negative binomial model for Gisborne.

Table 4: Gisborne standardisation data set. Number of puerulus samples by group and fishing year.

Year	GIS001	GIS002	GIS003	GIS004	Total
1987	15	0	0	0	15
1988	23	0	0	0	23
1989	25	0	0	0	25
1990	25	0	0	0	25
1991	25	5	0	0	30
1992	24	17	0	0	41
1993	25	20	0	0	45
1994	25	20	25	23	93
1995	25	24	25	25	99
1996	25	20	0	25	70
1997	25	20	23	25	93
1998	25	25	25	25	100
1999	20	25	21	18	84
2000	23	25	25	25	98
2001	24	25	25	25	99
2002	20	25	19	25	89
2003	0	18	19	30	67
2004	0	20	20	25	65
2005	0	25	19	25	69
2006	0	23	24	30	77
2007	0	24	0	0	24
2008	0	20	0	25	45
2009	0	18	0	25	43
2010	0	15	0	25	40
2011	0	20	0	20	40
2012	0	20	0	25	45
2013	0	20	0	15	35
2014	0	20	0	18	38
2015	0	20	0	19	39
2016	0	20	0	19	39

Napier




Figure 21: Characteristics of the Napier puerulus standardisation data. The top-left figure shows the distribution of puerulus counts, and the bottom-left the proportion of these that are non-zero. The top right figure shows the percentage of samples by month, and the bottom right shows the arithmetic mean of the number of pueruli measured in a sample.

Negative binomial




Figure 22: Deviance residuals for the negative binomial model for Napier. Predicted values are in natural space.




Figure 23: Quantile-quantile plot for the negative binomial standardisation model for Napier.




Figure 24: Data distribution and that from the fitted negative binomial model for Napier.

Table 5: Napier standardisation data set. Number of puerulus samples by group and fishing year.

	NAP001	NAP002	NAP003	NAP004	Total
1979	52	0	0	0	52
1980	65	0	0	0	65
1981	66	0	0	0	66
1982	60	0	0	0	60
1983	48	0	0	0	48
1984	60	0	0	0	60
1985	36	0	0	0	36
1988	18	0	0	0	18
1989	36	0	0	0	36
1990	36	0	0	3	39
1991	60	21	0	26	107
1992	69	21	0	32	122
1993	69	17	0	33	119
1994	65	27	25	33	150
1995	59	29	41	30	159
1996	72	33	50	33	188
1997	71	24	65	36	196
1998	66	18	58	27	169
1999	72	6	55	27	160
2000	47	0	48	27	122
2001	65	0	61	21	147
2002	57	0	52	18	127
2003	66	0	54	0	120
2004	71	0	59	0	130
2005	72	0	53	0	125
2006	72	0	47	0	119
2007	53	0	40	0	93
2008	56	0	59	0	115
2009	60	0	59	0	119
2010	60	0	52	0	112
2011	60	0	53	0	113
2012	50	0	36	0	86
2013	50	0	50	0	100
2014	50	0	59	0	109
2015	55	0	59	0	114
2016	40	0	53	0	93

Castlepoint




Figure 25: Characteristics of the Castlepoint puerulus standardisation data. The top-left figure shows the distribution of puerulus counts, and the bottom-left the proportion of these that are non-zero. The top right figure shows the percentage of samples by month, and the bottom right shows the arithmetic mean of the number of pueruli measured in a sample.

Negative binomial




Figure 26: Deviance residuals for the negative binomial model for Castlepoint. Predicted values are in natural space.




Figure 27: Quantile-quantile plot for the negative binomial standardisation model for Castlepoint.




Figure 28: Data distribution and that from the fitted negative binomial model for Castlepoint.

Table 6: Castlepoint standardisation data set. Number of puerulus samples by group and fishing year.

	CPT001	CPT002	CPT003	Total
1982	18	0	0	18
1983	68	0	0	68
1984	57	0	0	57
1985	41	0	0	41
1986	70	0	0	70
1987	66	0	0	66
1988	66	0	0	66
1989	67	0	0	67
1990	72	0	0	72
1991	72	17	16	105
1992	71	46	38	155
1993	70	63	61	194
1994	102	60	50	212
1995	97	48	37	182
1996	108	60	60	228
1997	108	60	55	223
1998	98	36	35	169
1999	116	18	65	199
2000	105	21	60	186
2001	99	36	53	188
2002	104	52	62	218
2003	99	51	55	205
2004	114	53	65	232
2005	107	60	60	227
2006	108	58	45	211
2007	106	50	0	156
2008	107	55	0	162
2009	99	55	0	154
2010	117	65	0	182
2011	108	60	0	168
2012	108	46	0	154
2013	117	70	0	187
2014	99	59	0	158
2015	105	64	0	169
2016	103	52	0	155

Kaikoura




Figure 29: Characteristics of the Kaikoura puerulus standardisation data. The top-left figure shows the distribution of puerulus counts, and the bottom-left the proportion of these that are non-zero. The top right figure shows the percentage of samples by month, and the bottom right shows the arithmetic mean of the number of pueruli measured in a sample.

Negative binomial




Figure 30: Deviance residuals for the negative binomial model for Kaikoura. Predicted values are in natural space.




Figure 31: Quantile-quantile plot for the negative binomial standardisation model for Kaikoura.




Figure 32: Data distribution and that from the fitted negative binomial for Kaikoura.

Table 7: Kaikoura standardisation data set. Number of puerulus samples by group and fishing year.

	KAI001	KAI002	KAI003	KAI004	KAI005	KAI006	KAI008	KAI009	Total
1981	15	0	15	0	0	0	15	0	45
1982	18	0	18	0	0	0	18	0	54
1983	15	0	15	0	0	0	18	0	48
1984	21	0	21	0	0	0	24	0	66
1985	18	0	17	0	0	0	24	0	59
1986	21	0	20	0	0	0	24	0	65
1987	24	0	24	0	0	0	27	0	75
1988	24	6	24	0	0	0	18	9	81
1989	24	24	27	0	0	0	0	27	102
1990	27	27	27	0	0	0	0	27	108
1991	27	24	27	0	0	0	0	24	102
1992	21	21	21	21	0	0	0	21	105
1993	24	24	24	24	0	0	0	15	111
1994	26	27	27	27	0	0	0	0	107
1995	27	27	27	27	0	0	0	0	108
1996	15	15	15	15	0	0	0	0	60
1997	12	12	12	9	0	0	0	0	45
1998	12	12	15	15	0	0	0	0	54
1999	15	15	18	18	0	0	0	0	66
2000	26	26	27	27	0	0	0	0	106
2001	27	27	27	27	0	0	0	0	108
2002	27	24	27	26	0	0	0	0	104
2003	45	0	45	0	0	0	0	0	90
2004	44	0	45	0	0	0	0	0	89
2005	45	0	45	0	0	0	0	0	90
2006	45	0	45	0	0	0	0	0	90
2007	45	0	50	0	12	9	0	0	116
2008	45	0	44	0	20	18	0	0	127
2009	44	0	45	0	24	21	0	0	134
2010	45	0	45	0	21	15	0	0	126
2011	45	0	45	0	15	15	0	0	120
2012	45	0	40	0	11	6	0	0	102
2013	39	0	40	0	3	3	0	0	85
2014	40	0	40	0	24	24	0	0	128
2015	45	0	45	0	12	9	0	0	111
2016	45	0	45	0	9	9	0	0	108

Moeraki




Figure 33: Characteristics of the Moeraki puerulus standardisation data. The top-left figure shows the distribution of puerulus counts, and the bottom-left the proportion of these that are non-zero. The top right figure shows the percentage of samples by month, and the bottom right shows the arithmetic mean of the number of pueruli measured in a sample.

Negative binomial




Figure 34: Deviance residuals for the negative binomial model for Moeraki. Predicted values are in natural space.




Figure 35: Quantile-quantile plot for the negative binomial standardisation model for Moeraki.




Figure 36: Data distribution and that from the fitted negative binomial for Moeraki.

Table 8: Moeraki standardisation data set. Number of puerulus samples by group and fishing year.

	MOE002	MOE007	Total
1991	16	0	16
1992	14	0	14
1993	12	0	12
1994	15	0	15
1995	15	0	15
1996	18	0	18
1997	15	0	15
1998	18	0	18
1999	15	0	15
2000	15	0	15
2001	17	4	21
2002	18	9	27
2003	15	67	82
2004	6	71	77
2005	15	71	86
2006	6	73	79
2007	0	52	52
2008	0	73	73
2009	0	39	39
2010	0	46	46
2011	0	50	50
2012	0	36	36
2013	0	36	36
2014	0	24	24
2015	0	36	36
2016	0	30	30

Halfmoon Bay




Figure 37: Characteristics of the Halfmoon Bay puerulus standardisation data. The top-left figure shows the distribution of puerulus counts, and the bottom-left the proportion of these that are non-zero. The top right figure shows the percentage of samples by month, and the bottom right shows the arithmetic mean of the number of pueruli measured in a sample.




Figure 38: Deviance residuals for the negative binomial model at Halfmoon Bay. Predicted values are in natural space.




Figure 39: Quantile-quantile plot for the negative binomial standardisation model for Halfmoon Bay.




Figure 40: Data distribution and that from the fitted negative binomial model for Halfmoon Bay.

Table 9: Halfmoon Bay standardisation data set. Number of puerulus samples by group and fishing year.

	HMB001	HMB002	HMB003	HMB004	HMB005	Total
1981	12	0	0	0	0	12
1982	23	0	0	0	0	23
1983	18	0	0	0	0	18
1984	18	0	0	0	0	18
1985	18	0	0	0	0	18
1986	18	21	0	0	0	39
1987	24	15	0	0	0	39
1988	21	15	0	0	0	36
1989	18	15	0	0	0	33
1990	22	15	15	0	0	52
1991	21	18	18	0	0	57
1992	18	15	11	15	15	74
1993	21	21	21	21	17	101
1994	18	21	21	21	21	102
1995	21	18	18	18	18	93
1996	18	21	21	21	21	102
1997	21	21	21	21	21	105
1998	15	24	21	21	21	102
1999	9	21	21	21	21	93
2000	18	18	18	21	18	93
2001	24	18	18	18	18	96
2002	18	21	21	21	21	102
2003	30	0	0	0	0	30
2004	16	0	0	0	0	16
2005	40	0	0	0	0	40
2006	72	0	0	0	0	72
2007	59	0	0	0	0	59
2008	48	0	0	0	0	48
2009	53	0	0	0	0	53
2010	64	0	0	0	0	64
2011	64	0	0	0	0	64
2012	56	0	0	0	0	56
2013	48	0	0	0	0	48
2014	48	0	0	0	0	48
2015	64	0	0	0	0	64
2016	56	0	0	0	0	56

Jackson Bay




Figure 41: Characteristics of the Jackson Bay puerulus standardisation data. The top-left figure shows the distribution of puerulus counts, and the bottom-left the proportion of these that are non-zero. The top right figure shows the percentage of samples by month, and the bottom right shows the arithmetic mean of the number of pueruli measured in a sample.

Negative binomial




Figure 42: Deviance residuals for the negative binomial model for the Jackson Bay site. Predicted values are in natural space.




Figure 43: Quantile-quantile plot for the negative binomial standardisation model for Jackson Bay.




Figure 44: Data distribution and that from the fitted negative binomial for Jackson Bay.

Table 10: Jackson Bay standardisation data set. Number of puerulus samples by group and fishing year.

	JKB001	JKB002	Total
1999	18	20	38
2000	48	34	82
2001	50	36	86
2002	48	30	78
2003	40	21	61
2004	38	24	62
2005	35	16	51
2006	19	4	23
2007	40	0	40
2008	30	0	30
2009	20	0	20
2010	22	0	22
2011	39	0	39
2012	52	0	52
2013	49	0	49
2014	48	0	48
2015	34	0	34
2016	46	0	46