



# **Aquatic Environment and Biodiversity Annual Review 2012**

**A summary of environmental interactions between  
fisheries and the aquatic environment**

**MINISTRY FOR PRIMARY  
INDUSTRIES**

**AQUATIC ENVIRONMENT AND  
BIODIVERSITY ANNUAL REVIEW  
(2012)**

**Fisheries Management Science Team**

### Acknowledgements

In addition to the thanks due to members of AEWG and BRAG working groups, special acknowledgement is due to Owen Anderson (NIWA) for his contribution to the new chapter on Non-protected Bycatch; Igor Debski (DOC) on the new seabirds chapter, and Ian Tuck (NIWA) on the benthic chapter. Notwithstanding these contributions, any errors or omissions are the Ministry's.

### Image on cover

Chatham-Challenger Ocean Survey 20/20

### Disclaimer

This document is published by the Ministry Primary Industries which was formed from the merger of the Ministry of Fisheries, the Ministry of Agriculture and Forestry and the New Zealand Food Safety Authority in 2010 and 2011. All references to the Ministry of Fisheries in this document should, therefore, be taken to refer also to the legal entity, the Ministry for Primary Industries. The information in this publication is not government policy. While every effort has been made to ensure the information is accurate, the Ministry for Primary Industries does not accept any responsibility or liability for error of fact, omission, interpretation or opinion that may be present, nor for the consequences of any decisions based on this information. Any view or opinion expressed does not necessarily represent the view of the Ministry for Primary Industries.

### Publisher

Fisheries Management Science Team  
Ministry for Primary Industries  
Pastoral House, 25 The Terrace  
PO Box 2526, Wellington 6140  
New Zealand

[www.mpi.govt.nz](http://www.mpi.govt.nz)

Telephone: 0800 00 83 33

Facsimile: +64 4 894 0300

ISBN 978-0-478-40503-3 (print)

ISBN 978-0-478-40504-0 (online)

© Crown Copyright March 2013 – Ministry for Primary Industries

### Preferred citation

Ministry for Primary Industries (2012). Aquatic Environment and Biodiversity Annual Review 2012. Compiled by the Fisheries Management Science Team, Ministry for Primary Industries, Wellington, New Zealand. 387 p.

## PREFACE

This, the 2012 edition of the Aquatic Environment and Biodiversity Review, expands and updates the first edition published in 2011. It summarises information on a range of issues related to the environmental effects of fishing and aspects of marine biodiversity and productivity relevant to fish and fisheries. This review is a conceptual analogue of the Ministry's annual Reports from the Fisheries Assessment Plenary. It summarises the most recent data and analyses on particular aquatic environment issues and, where appropriate, assesses current status against any specified targets or limits. Whereas the Reports from the Fisheries Assessment Plenary are organised by fishstock, however, the Aquatic Environment and Biodiversity Review is organised by issue (for example, protected species bycatch, benthic impacts), and almost all issues involve more than one fishstock or fishery.

Several Fisheries Assessment Working Groups (FAWGs) contribute to the Fisheries Assessment Plenary, but only two generally contribute to the Aquatic Environment and Biodiversity Review. These are the Aquatic Environment Working Group (AEWG) and the Biodiversity Research Advisory Group (BRAG). However, a wider variety of research is summarised in the Aquatic Environment and Biodiversity Review than in the Reports from the Fisheries Assessment Plenary, and some of this is peer-reviewed through processes other than the Ministry's science working groups. In particular, the Department of Conservation funds and reviews research on protected species, and the Ministry of Business Innovation and Employment funds a wide variety of research, some of which is relevant to fisheries. Where such research is relevant to fisheries it will be considered for inclusion in the review.

As has happened with the Reports from the Fisheries Assessment Plenary, continual future expansion and improvement of the Aquatic Environment and Biodiversity Review is anticipated and additional chapters will be developed to provide increasingly comprehensive coverage of the issues. New chapters are included this year for seabirds (Chapter 5) and the bycatch and discards of fish and invertebrates (Chapter 6), and a new appendix summarising aquatic environment and marine biodiversity research since 1998 has now been developed (Appendix 12.9). A chapter on Hector's/Maui's dolphins has been identified as a priority for development in 2013. Data acquisition, modelling, and assessment techniques will also progressively improve, and it is expected that reference points to guide fisheries management decisions will be developed. Both will lead to changes to the current chapters. We hope the condensation in this review of the information from previously scattered reports will assist fisheries managers, stakeholders and other interested parties to understand the issues, locate relevant documents, track research progress and make informed decisions.

This revision has been led by the Science Team within the Directorate of Fisheries Management of the Ministry for Primary Industries (primarily Martin Cryer, Rohan Currey, Rich Ford, and Mary Livingston) but has relied critically on the input of members of the Ministry's Aquatic Environment Working Group (AEWG) and Biodiversity Research Advisory Group (BRAG) and the Department of Conservation's Conservation Services Technical Working Group (DOC-CSTWG). I would especially like to recognise and thank the large number of research providers and scientists from research organisations, academia, the seafood industry, environmental NGOs, Māori customary, DOC, and MPI, along with all other technical and non-technical participants in present and past AEWG and BRAG meetings for their substantial contributions to this review. My sincere thanks to each and all who have contributed.

I am pleased to endorse this document as representing the best available scientific information relevant to the aspects of the environmental effects of fishing and marine biodiversity covered as at December 2012.



Pamela Mace  
Principal Adviser Fisheries Science  
Ministry for Primary Industries

## Contents

PREFACE .....	3
1. INTRODUCTION .....	6
1.1. Context and purpose .....	6
1.2. Legislation .....	6
1.3. Policy Setting .....	9
1.4. Science processes .....	11
1.5. References .....	12
2. Research themes covered in this document .....	13
THEME 1: PROTECTED SPECIES .....	16
3. New Zealand sea lions ( <i>Phocarcos hookeri</i> ) .....	17
3.1. Context .....	17
3.2. Biology .....	19
3.3. Global understanding of fisheries interactions .....	25
3.4. State of knowledge in New Zealand .....	25
3.5. Indicators and trends .....	39
3.6. References .....	41
4. New Zealand fur seal ( <i>Arctocephalus forsteri</i> ) .....	44
4.1. Context .....	44
4.2. Biology .....	45
4.3. Global understanding of fisheries interactions .....	50
4.4. State of knowledge in New Zealand .....	50
4.5. Indicators and trends .....	59
4.6. References .....	60
5. New Zealand seabirds .....	63
5.1. Context .....	64
5.2. Biology .....	68
5.3. Global understanding of fisheries interactions .....	68
5.4. State of knowledge in New Zealand .....	70
5.5. Indicators and trends .....	113
5.6. References .....	115
THEME 2: NON-PROTECTED BYCATCH .....	120
6. Non-protected species (fish and invertebrates) bycatch .....	121
6.1. Context .....	122
6.2. Global understanding .....	123
6.3. State of knowledge in New Zealand .....	124
6.4. Indicators and trends .....	156
6.5. References .....	157
THEME 3: BENTHIC IMPACTS .....	159
7. Benthic (seabed) impacts .....	160
7.1. Context .....	161
7.2. Global understanding .....	164
7.3. State of knowledge in New Zealand .....	168
7.4. Indicators and trends .....	183
7.5. References .....	184
THEME 4: ECOSYSTEM EFFECTS .....	187
8. New Zealand Climate and Oceanic Setting .....	188
8.1. Context .....	189
8.2. Indicators and trends .....	193
8.3. Ocean climate trends and New Zealand fisheries .....	201
8.4. References .....	203
9. Habitats of particular significance for fisheries management .....	205

9.1.	Context .....	205
9.2.	Global understanding .....	207
9.3.	State of knowledge in New Zealand.....	210
9.4.	Indicators and trends .....	213
9.5.	References .....	213
10.	Land-based effects on fisheries, aquaculture and supporting biodiversity.....	216
10.1.	Context .....	216
10.2.	Global understanding .....	218
10.3.	State of knowledge in New Zealand.....	220
10.4.	Indicators and trends .....	224
10.5.	References .....	225
THEME 5: MARINE BIODIVERSITY .....		228
11.	Biodiversity .....	229
11.1.	Introduction .....	231
11.2.	Global understanding and developments.....	235
11.3.	State of knowledge in New Zealand.....	243
11.4.	Progress and re-alignment .....	281
11.5.	References .....	286
11.6.	Appendix .....	295
12.	Appendices .....	298
12.1.	Terms of Reference for the Aquatic Environment Working Group in 2012.....	298
12.2.	AEWG Membership 2012.....	303
12.3.	Terms of Reference for the Biodiversity Research Advisory Group (BRAG) 2012.....	304
12.4.	BRAG attendance 2011-2012.....	310
12.5.	Generic Terms of Reference for Research Advisory Groups (Sept 2010) .....	310
12.6.	Fisheries 2030.....	314
12.7.	OUR STRATEGY 2030: Growing and protecting New Zealand .....	316
12.8.	Other strategic policy documents .....	317
12.9.	Appendix of Aquatic Environment and Biodiversity funded and related projects.....	323

# **1. INTRODUCTION**

## **1.1. Context and purpose**

This document contains a summary of information and research on aquatic environment issues relevant to the management of New Zealand fisheries and expands and updates the first version published in 2011 (MAF 2011). It is designed to complement the Ministry's annual Reports from Fisheries Assessment Plenaries (e.g., MPI 2012a & b) and emulate those documents' dual role in providing an authoritative summary of current understanding and an assessment of status relative to any overall targets and limits. However, whereas the Reports from Fisheries Assessment Plenaries have a focus on individual fishstocks, this report has a focus on aquatic environment fisheries management issues and biodiversity responsibilities that often cut across many fishstocks, fisheries, or activities, and sometimes across the responsibilities of multiple agencies.

This update has been developed by the Science Team within the Fisheries Management Directorate of the Resource Management and Programmes branch, Ministry for Primary Industries (MPI). It does not cover all issues but, as anticipated, includes more chapters than the first edition in 2011. As with the Reports from Fisheries Assessment Plenaries, it is expected to change and grow as new information becomes available, more issues are considered, and as feedback and ideas are received. This synopsis has a broad, national focus on each issue and the general approach has been to avoid too much detail at a fishery or fishstock level. For instance, the benthic (seabed) effects of mobile bottom-fishing methods are dealt with at the level of all bottom trawl and dredge fisheries combined rather than at the level of a target fishery that might contribute only a small proportion of the total impact. The details of benthic impacts by individual fisheries will be documented in the respective chapters in the May or November Report from the Fisheries Assessment Plenary, and linked there to the fine detail and analysis in Aquatic Environment and Biodiversity Reports (AEBRs), Fisheries Assessment Reports (FARs), and Final Research Reports (FRRs). Such sections have already been developed for several species in both 2012 Fishery Assessment Plenary Reports, and others will follow.

The first part of this document describes the legislative and broad policy context for aquatic environment and biodiversity research commissioned by MPI, and the science processes used to generate and review that research. The second, and main, part of the document contains chapters focused on various aquatic environment issues for fisheries management. Those chapters are divided into five broad themes: protected species; non-QMS fish bycatch; benthic effects; ecosystem issues (including New Zealand's oceanic setting); and marine biodiversity. A third part of the review includes a number of appendices for reference. This review is not comprehensive in its coverage of all issues or of all research within each issue, but attempts to summarise the best available information on the issues covered. Each chapter has been considered by the appropriate working group at least once.

## **1.2. Legislation**

The primary legislation for the management of fisheries, including effects on the aquatic environment, is the Fisheries Act 1996. The main sections setting out the obligation to avoid, remedy, or mitigate any adverse effect of fishing on the aquatic environment are sections 8, 9, and 15, although sections 10, 11, and 13 are also relevant to decision-making under this Act (Table 1.1). The Ministry also administers the residual parts of the Fisheries Act 1983, the Treaty of Waitangi (Fisheries Claims) Settlement Act 1992, the Fisheries (Quota Operations Validation) Act 1997, the Maori Fisheries Act 2004, the Maori Commercial Aquaculture Claims Settlement Act 2004, the Aquaculture Reform (Repeals and Transitional Provisions) Act 2004, the Driftnet Prohibition Act 1991, and the Antarctic Marine Living Resources Act 1981. Other Acts are relevant in specific circumstances: the Wildlife Act 1953 and the Marine Mammals Protection Act 1978 for protected species; the Marine Reserves Act 1971 for "no take" marine reserves; the Conservation Act 1987; the Hauraki Gulf Marine Park Act

2000; the Resource Management Act 1991 for issues in coastal marine areas that could affect fisheries interests or be the subject of sustainability measures under section 11 of the Fisheries Act; and the Exclusive Economic Zone and Continental Shelf (Environmental Effects) Act 2012 for issues outside the Territorial Sea. These Acts are administered by other agencies and this leads to a requirement for the Ministry for Primary Industries to work with other government departments (especially the Department of Conservation and through the Natural Resource Sector<sup>1</sup>) and with various territorial authorities (especially Regional Councils) to a greater extent than is required for most fisheries stock assessment issues.

**Table 1.1: Sections of the Fisheries Act 1996 relevant to the management of the effects of fishing on the aquatic environment.**

<b>Fisheries Act 1996</b>
<p><b>s8 Purpose –</b>            (1) The purpose of this Act is to provide for the utilisation of fisheries resources while ensuring sustainability, where            (2) “Ensuring sustainability” means –                (a) Maintaining the potential of fisheries resources to meet the reasonably foreseeable needs of future generations; and                (b) Avoiding, remedying, or mitigating any adverse effects of fishing on the aquatic environment:            “Utilisation” means conserving, using, enhancing, and developing fisheries resources to enable people to provide for their social, economic, and cultural well-being.</p> <p><b>s9 Environmental Principles.</b>            associated or dependent species should be maintained above a level that ensures their long-term viability;            biological diversity of the aquatic environment should be maintained;            habitat of particular significance for fisheries management should be protected.</p> <p><b>s11 Sustainability Measures.</b> The Minister may take into account, in setting any sustainability measure, (a) any effects of fishing on any stock and the aquatic environment;</p> <p><b>s15 Fishing-related mortality of marine mammals or other wildlife.</b> A range of management considerations are set out in the Fisheries Act 1996, which empower the Minister to take measures to avoid, remedy or mitigate any adverse effects of fishing on associated or dependent species and any effect of fishing-related mortality on any protected species. These measures include the setting of catch limits or the prohibition of fishing methods or all fishing in an area, to ensure that such catch limits are not exceeded.</p>

Under the primary legislation lie various layers of Regulations and Orders in Council (see <http://www.legislation.govt.nz/>). It is beyond the scope of this document to summarise these.

In addition to its domestic legislation, the New Zealand government is a signatory to a wide variety of International Instruments and Agreements that bring with them various International Obligations (Table 1.2). Section 5 of the Fisheries Act requires that the Act be interpreted in a manner that is consistent with international obligations and with the Treaty of Waitangi (Fisheries Claims) Settlement Act 1992.

<sup>1</sup> The Natural Resources Sector is a network of government agencies established to enhance collaboration. Its main purpose is to ensure a strategic, integrated and aligned approach is taken to natural resources development and management across government agencies. The network is chaired by MfE’s Chief Executive. The Sector aims to provide high-quality advice to government and provide effective implementation and execution of major government policies through coordination and integration across agencies, management of relationships, and alignment of the policies and practices of individual agencies.

**Table 1.2: International agreements and regional agreements to which New Zealand is a signatory, that are relevant to the management of the effects of fishing on the aquatic environment.**

International Instruments	Regional Fisheries Agreements
<ul style="list-style-type: none"> <li>• <b>Convention on the Conservation of Migratory Species of Wild Animals (CMS).</b> Aims to conserve terrestrial, marine and avian migratory species throughout their range.</li> <li>• <b>Agreement on the Conservation of Albatrosses and Petrels (ACAP).</b> Aims to introduce a number of conservation measures to reduce the threat of extinction to the Albatross and Petrel species.</li> <li>• <b>Convention on Biological Diversity (CBD)</b> Provides for conservation of biological diversity and sustainable use of components. States accorded the right to exploit resources pursuant to environmental policies.</li> <li>• <b>United Nations Convention on the Law of the Sea (UNCLOS)</b> Acknowledges the right to explore and exploit, conserve and manage natural resources in the State's EEZ...with regard to the protection and preservation of the marine environment including associated and dependent species, pursuant to the State's environmental policies.</li> <li>• <b>Convention on the International Trade in Endangered Species of Wild Fauna and Flora (CITES).</b> Aims to ensure that international trade in wild animals and plants does not threaten their survival.</li> <li>• <b>United Nations Fishstocks Agreements.</b> Aims to lay down a comprehensive regime for the conservation and management of straddling and highly migratory fish stocks.</li> <li>• <b>International Whaling Commission (IWC)</b> Aims to provide for the proper conservation of whale stocks and thus make possible the orderly development of the whaling industry.</li> <li>• <b>Wellington Convention</b> Aims to prohibit drift net fishing activity in the convention area.</li> <li>• <b>Food and Agriculture Organisation – International Plan of Action for Seabirds (FAO-IPOA Seabirds)</b> Voluntary framework for reducing the incidental catch of seabirds in longline fisheries.</li> <li>• <b>Food and Agriculture Organisation – International Plan of Action for Sharks (FAO –IPOA Sharks)</b> Voluntary framework for the conservation and management of sharks.</li> <li>• <b>Noumea Convention.</b> Promotes protection and management of natural resources. Parties to regulate or prohibit activity likely to have adverse effects on species, ecosystems and biological processes.</li> <li>• <b>Food and Agriculture Organisation - Code of Conduct for Responsible Fisheries</b> Provides principles and standards applicable to the conservation, management and development of all fisheries, to be interpreted and applied to conform to the rights, jurisdiction and duties of States contained in UNCLOS.</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Convention for the Conservation of Southern Bluefin Tuna (CCSBT)</b> Aims to ensure, through appropriate management, the conservation and optimum utilisation of the global Southern Bluefin Tuna fishery. The Convention specifically provides for the exchange of data on ecologically related species to aid in the conservation of these species when fishing for southern bluefin tuna.</li> <li>• <b>Convention for the Conservation of Antarctic Marine Living Resources (CCAMLR).</b> Aims to conserve, including rational use of Antarctic marine living resources. This includes supporting research to understand the effects of CCAMLR fishing on associated and dependent species, and monitoring levels of incidental take of these species on New Zealand vessels fishing in CCAMLR waters.</li> <li>• <b>Convention on the Conservation and Management of Highly Migratory Fish Stocks in the Western and Central Pacific Ocean (WCPFC).</b> The objective is to ensure, through effective management, the long-term conservation and sustainable use of highly migratory fish stocks in accordance with UNCLOS.</li> <li>• <b>South Tasman Rise Orange Roughy Arrangement.</b> The arrangement puts in place the requirement for New Zealand and Australian fishers to have approval from the appropriate authorities to trawl or carry out other demersal fishing for any species in the STR area</li> <li>• <b>Convention on the Conservation and Management of High Seas Fishery Resources in the South Pacific Ocean</b> (a Regional Fisheries Management Organisation, colloquially <b>SPRFMO</b>) has recently been negotiated to facilitate management of non-highly migratory species in the South Pacific.</li> </ul>

## **1.3. Policy Setting**

### **1.3.1. Our Strategy 2030 and MPI's Statement of Intent 2012/15**

The Ministry for Primary Industries' Statement of Intent, SOI, is an important guiding document for the short to medium term. That for 2012–15 is available on the Ministry's website at:

<http://www.mpi.govt.nz/Default.aspx?TabId=126&id=1341>

The SOI sets out the Ministry's strategic direction for the coming three years, primarily through implementation of Our Strategy 2030 (Appendix 12.7). This strategy was agreed by Cabinet in August 2011 and sets out MPI's vision of "growing and protecting New Zealand" and defines the focus and approach of the organisation. The strategy includes four focus areas and outcomes: maximising export opportunities; improving sector productivity; increasing sustainable resource use; and protecting from biological risk.

MPI is the single key adviser to the Government across all aspects of the primary industries, food production and related trade issues. MPI is the principal adviser to the Government on agriculture, horticulture, aquaculture, fisheries, forestry, and food industries, animal welfare, and the protection of New Zealand's primary industries from biological risk. Aspects of the role specific to fisheries itemized in the SOI include supporting the development of sustainable limits to natural resource use. To that end, MPI contracts the following types of research (relevant to this document):

- aquatic environment research to assess the effects of fishing on marine habitats, protected species, trophic linkages, and to understand habitats of special significance for fisheries;
- biodiversity research to increase our understanding of the systems that support resilient ecosystems and productive fisheries.

### **1.3.2. Fisheries 2030**

New Zealand's Quota Management System (QMS) forms the overall framework for management of domestic fisheries (see <http://www.fish.govt.nz/en-nz/Commercial/Quota+Management+System/default.htm>). Within that framework, Fisheries 2030 provides a long-term goal for the New Zealand fisheries sector. After endorsement by Cabinet, it was released by the Minister of Fisheries in September 2009. It can be found on the MPI website at:

[http://www.fish.govt.nz/en-nz/Fisheries+2030/default.htm?wbc\\_purpose=bas](http://www.fish.govt.nz/en-nz/Fisheries+2030/default.htm?wbc_purpose=bas)

*(noting that the Ministry of Fisheries merged with the Ministry of Agriculture and Forestry on 1 July 2011 and became the Ministry for Primary Industries on 30 April 2012. This URL and subsequent links in this document will eventually change as the new Ministry's systems are progressively merged).*

Fisheries 2030 sets out a goal to have *New Zealanders maximising benefits from the use of fisheries within environmental limits*. To support this goal, major outcomes for Use (of fisheries) and Environment are specified. The Environment outcome is the main driver for aquatic environment research: *The capacity and integrity of the aquatic environment, habitats and species are sustained at levels that provide for current and future use*. Fisheries 2030 states that this means:

- Biodiversity and the function of ecological systems, including trophic linkages, are conserved
- Habitats of special significance to fisheries are protected
- Adverse effects on protected species are reduced or avoided

- Impacts, including cumulative impacts, of activities on land, air or water on aquatic ecosystems are addressed.

### **1.3.3. Fisheries Plans**

Fisheries planning processes for deepwater, highly migratory species, inshore finfish, inshore shellfish and freshwater fisheries use objective-based management to drive the delivery of services, as described in Fisheries 2030 and affirmed in the 2012/15 SOI and Our Strategy 2030. The planning processes are guided by five National Fisheries Plans, which recognise the distinctive characteristics of these fisheries. Plans for Deepwater and Highly Migratory species have been approved by the Minister and a suite of three plans for inshore species has been released in prototype form. These plans establish management objectives for each fishery, including those related to the environmental effects of fishing. All are available on the Ministry's websites.

Deepwater and middle depth fisheries:

<http://www.fish.govt.nz/en-nz/Consultations/Archive/2010/National+Fisheries+Plan+for+Deepwater+and+Middle-Depth+Fisheries/default.htm>

Highly migratory species (HMS) fisheries:

<http://www.fish.govt.nz/en-nz/Consultations/Archive/2010/National+Fisheries+Plan+for+Highly+Migratory+Species/default.htm>

Inshore fisheries (comprising finfish, shellfish, and freshwater fisheries):

<http://www.fish.govt.nz/en-nz/Fisheries+Planning/default.htm>

Certain research areas (aquatic environment, recreational and biodiversity) are not entirely covered by fish plans, as many of these issues span multiple fisheries and plans. Antarctic research is also excluded from fish plans as it is beyond their spatial scope. These areas are administered by the science team and subject to the drivers in Tables 1.1, 1.2 and Fisheries 2030.

### **1.3.4. Other strategic documents**

A number of strategies or reviews have been published that potentially affect fisheries values and research. These include: the New Zealand Biodiversity Strategy (2000); the Biosecurity Strategy (2003, followed by its science strategy 2007); the MPA Policy and Implementation Plan (2005); MfE's discussion paper on Management of Activities in the EEZ (2007); MRST's Roadmap for Environment Research (2007); the Revised Coastal Policy Statement (2010); the National Plan of Action to Reduce the Incidental Catch of Seabirds in New Zealand Fisheries (2004, soon to be revised); and the New Zealand National Plan of Action for the Conservation and Management of Sharks (2008). Links to these documents are provided in Appendix 12.8 because they provide some of the broad policy setting for aquatic environment issues and research across multiple organisations and agencies.

In 2012, the Natural Resource Sector cluster formed a Marine Director's Group to improve data sharing and information exchange across key agencies with marine environmental responsibilities, particularly MPI, DOC, MfE, EPA, LINZ, MBIE. The Marine Director's Group is chaired by MPI and DOC and a substantial amount of cross-agency work has been initiated to: summarise relevant marine information held by different agencies and current marine research investment; identify knowledge and funding gaps; and to develop a long-term Marine Research Strategy for New Zealand.

## **1.4. Science processes**

### **1.4.1. Research Planning**

Until 2010 the Ministry of Fisheries ran an iterative planning process to determine, in conjunction with stakeholders and subject to government policy, the future directions and priorities for fisheries research. Subsequently, the Ministry has adopted an overall approach of specifying objectives for fisheries in Fisheries Plans and using these plans to develop associated implementation strategies and required services, including research. These services are identified in Annual Operational Plans that are updated each year.

For deepwater fisheries and highly migratory stocks (HMS), the transition to the new research planning approach is well advanced because fisheries plans for these areas have been approved by the Minister. Research for these fisheries are already being developed using Fisheries Plan and Annual Operating Plan processes as primary drivers, and, as necessary, Research Advisory Groups (RAGs) to develop the technical detail of particular projects. The Ministry's website contains more information on this approach, developed during the Research Services Strategy Review, at: [http://www.fish.govt.nz/NR/rdonlyres/04D579E5-6DCC-42A6-BF68-9CAB800D6392/0/Research\\_Services\\_Strategy\\_Review\\_Report.pdf](http://www.fish.govt.nz/NR/rdonlyres/04D579E5-6DCC-42A6-BF68-9CAB800D6392/0/Research_Services_Strategy_Review_Report.pdf) (see Section 5.2, pages 14 to 21) and in summary at: [http://www.fish.govt.nz/NR/rdonlyres/432EA3A0-AEA7-41DD-8E5C-D0DCA9A3B96B/0/RSS\\_letter.pdf](http://www.fish.govt.nz/NR/rdonlyres/432EA3A0-AEA7-41DD-8E5C-D0DCA9A3B96B/0/RSS_letter.pdf). Generic terms of reference for Research Advisory Groups are in Appendix 12.5. For inshore fisheries, the three Fisheries Plans (inshore finfish, shellfish, and freshwater) are still under development, so a transitional research planning process was established for 2010 and developed slightly in 2011. This included the following steps:

- Identification of the main management information needs using:
  - Fisheries Plans or Fisheries Operational Plans where available
  - Any relevant Medium Term Research Plan
  - Fishery managers' understanding of decisions likely to require research information in the next 1–3 years.
- Technical discussions as required (i.e., tailored to the needs of the different research areas) to consider:
  - The feasibility and utility of each project
  - The likely cost of each project
  - Any synergies or overlaps with work being conducted by other providers (including industry, CRIs, MBIE, Universities, etc.)
- Stakeholder meetings as required to discuss relative priorities for particular projects

The process for aquatic environment research for 2011/12 and 2012/13 (other than aspects driven by deepwater and HMS plans or the specific needs of inshore fishery managers) followed essentially these same steps.

The Ministry runs a separate planning group to design and prioritise its research programme on marine biodiversity. Given its much broader and more strategic focus, the Biodiversity Research Advisory Group (BRAG) has both peer review and planning roles and therefore differs slightly in constitution from the Ministry's other working and planning groups.

### **1.4.2. Contributing Working Groups**

The main contributing working groups for this document are the Ministry's Aquatic Environment Working Group (AEWG) and Biodiversity Research Advisory Group (BRAG). The Department of Conservation's Conservation Services Programme and National Plan of Action Seabirds Technical Working Group (CSP/NPOA-TWG, see <http://www.doc.govt.nz/conservation/marine-and->

[coastal/commercial-fishing/marine-conservation-services/meetings-and-project-updates/](#)) also considers a wide range of DOC-funded projects related to protected species, sometimes in joint meetings with the AEWG. The Ministry's Fishery Assessment Working Groups occasionally consider research relevant to this synopsis. Terms of reference for AEWG and BRAG are periodically revised and updated (see Appendix 12.1 and 12.3 for the 2012 Terms of Reference for AEWG and BRAG, respectively).

AEWG is convened for the Ministry's peer review purposes with an overall purpose of assessing, based on scientific information, the effects of fishing, aquaculture, and enhancement on the aquatic environment for all New Zealand fisheries. The purview of AEWG includes: bycatch and unobserved mortality of protected species, fish, and other marine life; effects of bottom fisheries on benthic biodiversity, species, and habitat; effects of fishing on biodiversity, including genetic diversity; changes to ecosystem structure and function as a result of fishing, including trophic effects; and effects of aquaculture and fishery enhancement on the environment and on fishing. Where possible, AEWG may explore the implications of any effects, including with respect to any standards, reference points, and relevant indicators. The AEWG is a technical forum to assess the effects of fishing or environmental status and make projections. It has no mandate to make management recommendations or decisions. Membership of AEWG is open (attendees for 2012 are listed in Appendix 12.2).

The two main responsibilities of BRAG are: to review, discuss, and convey views on the results of marine biodiversity research projects contracted by the Ministry; and to discuss, evaluate, make recommendations and convey views on Medium Term Biodiversity Research Plans and constituent individual projects. Both tasks have hitherto been undertaken in the context the strategic goals in the New Zealand Biodiversity Strategy (2000) and the Strategy for New Zealand Science in Antarctica and the Southern Ocean (2010), but the focus of the programme is currently being reviewed to align it with more recent strategic documents. BRAG also administers some large cross-government projects such as NORFANZ, BIOROSS, Fisheries and Biodiversity Ocean Survey 20/20; and International Polar Year (IPY) Census of Antarctic Marine Life (IPY-CAML). Membership of BRAG is also open (attendees for 2011 and 2012 are listed in Appendix 12.4).

Following consideration at one or more meetings of appropriate working groups, reports from individual projects are also technically reviewed by the Ministry before they are finalised for use in management and/or for public release. Fisheries Assessment Reports, FARs, and Aquatic Environment and Biodiversity reports, AEBRs, are also subject to editorial review whereas Final Research Reports, FRRs, and Research Progress Reports, RPRs, are not. Finalised FARs, AEBRs, historical FARDs (Fisheries Assessment Research Documents) and MMBRs (Marine Biodiversity and Biosecurity Reports), and some FRRs can be found at:

<http://fs.fish.govt.nz/Page.aspx?pk=61&tk=209>.

Increasingly, reports will be available from the MPI website at: <http://www.mpi.govt.nz/news-resources/publications>.

## 1.5. References

- Ministry of Agriculture and Forestry (2011). Aquatic Environment and Biodiversity Annual Review 2011: a summary of environmental interactions between fisheries and the aquatic environment. Compiled by the Fisheries Science Group, Ministry of Agriculture and Forestry, Wellington, New Zealand. 199 p.
- Ministry for Primary Industries (2012a). Report from the Fisheries Assessment Plenary, May 2012: stock assessments and yield estimates. Compiled by the Fisheries Science Group, Ministry for Primary Industries, Wellington, New Zealand. 1194 p.
- Ministry for Primary Industries (2012b). Fisheries Assessment Plenary, November 2012: stock assessments and yield estimates. Compiled by the Fisheries Science Group, Ministry for Primary Industries, Wellington, New Zealand. 531 p.

## **2. Research themes covered in this document**

The Ministry has identified four broad categories of research on the environmental effects of fishing (Figure 2.1): bycatch and fishing-related mortality of protected species; bycatch of non-protected species, primarily non-QMS fish; modification of benthic habitats (including seamounts); and various ecosystem effects (including fishing and non-fishing effects on habitats of particular significance for fisheries management and trophic relationships). Other emerging issues (such as the genetic consequences of selective fishing and the impacts of aquaculture) are not dealt with in detail in this synopsis but it is anticipated that those that turn out to be important will be dealt with in future iterations. A fifth theme for this document is MPI research on marine biodiversity. The research has been driven largely by the Biodiversity Strategy but has strategic importance for fisheries in that it provides for better understanding of the ecosystems that support fisheries productivity.

Our understanding is not uniform across these themes and, for example, our knowledge of the quantum and consequences of fishing-related mortality of protected species is much better developed than our knowledge of the consequences of mortalities of non-target fish, bottom trawl impacts, or land management choices for ecosystem processes or fisheries productivity. Ultimately, the goal of research described in this synopsis is to complement information on fishstocks to ensure that the Ministry has the information required to underpin the ecosystem approach to fisheries management envisaged in Fisheries 2030. Stock assessment results have been published for many years in Fisheries Assessment Reports, and Final Research Reports, and the Annual Report from the Fishery Assessment Plenary. Collectively, these provide a rich and well-understood resource for fisheries managers and stakeholders. In 2005, an environmental section was included in the hoki plenary report as part of the characterisation of that fishery and to highlight any particular environmental issues associated with the fishery. Similar, fishery-specific sections have since been developed for other working group reports and the plenary, including many fisheries for highly migratory species and the trawl fisheries for scampi and squid, but work on environmental issues has otherwise been more difficult to access for fisheries managers and stakeholders. The Ministry is, therefore, looking at improving ways to document, review, publicise, and integrate information from environmental assessments with traditional fishery assessments. This will rely heavily on studies that are published in Aquatic Environment and Biodiversity Reports and Final Research Reports but, given the overlapping mandates and broader scope of work in this area, also on results published by other organisations. The integration of all this work into a single source document analogous to the Report from the Fishery Assessment Plenary will take time and not all issues will be covered for some years.

THEME	RESEARCH QUESTIONS	CURRENTWORK
 <b>1.PROTECTED SPECIES</b> <ul style="list-style-type: none"> <li>• Marine mammals</li> <li>• Seabirds</li> <li>• Turtles</li> <li>• Protected fish</li> <li>• Corals</li> </ul>	<ul style="list-style-type: none"> <li>• How many of each NZ-breeding protected species are caught and killed in our fisheries (and out of zone)?</li> <li>• How many unobserved deaths are caused?</li> <li>• What is the likely effect of fishing-related mortality on protected species populations?</li> <li>• Which species or populations are most at risk?</li> <li>• Which fisheries cause the most risk and where are the most cost-effective gains to be made?</li> <li>• What mitigation approaches are most successful and in what circumstances?</li> <li>• What levels of bycatch would lead to different population outcomes?</li> </ul>	<ul style="list-style-type: none"> <li>• Estimation of annual bycatch of protected species by fishery</li> <li>• Abundance and productivity of key seabird populations</li> <li>• Abundance and productivity of Hector's &amp; Maui's dolphins</li> <li>• Semi-quantitative risk assessment for all seabirds</li> <li>• Semi-quantitative risk assessment for other protected species</li> <li>• Full quantitative risk assessment for selected seabird populations</li> <li>• Modelling to assess robust links between observed bycatch and population outcomes</li> </ul>
 <b>2. OTHER BYCATCH</b> <ul style="list-style-type: none"> <li>• Non-QMS fish &amp; invertebrates</li> </ul>	<ul style="list-style-type: none"> <li>• How much non-target fish is caught and discarded in our fisheries?</li> <li>• What is the effect of that bycatch?</li> <li>• What do trends in bycatch show?</li> </ul>	<ul style="list-style-type: none"> <li>• Continued monitoring cycle for deepwater and highly migratory</li> <li>• Risk assessment for tier 3 deepwater bycatch species</li> </ul>
 <b>3.BENTHIC EFFECTS</b> <ul style="list-style-type: none"> <li>• Distribution of habitats &amp; trawling</li> <li>• Effects of trawling on each</li> </ul>	<ul style="list-style-type: none"> <li>• What seabed habitats occur where in our TS/EEZ and how much of each is affected by trawling or shellfish dredging?</li> <li>• How sensitive is each habitat to disturbance and what do we lose when each is disturbed?</li> <li>• What are the consequences of different management approaches?</li> </ul>	<ul style="list-style-type: none"> <li>• Testing of habitat classifications</li> <li>• Assessment of recovery rate of some key inshore habitats</li> <li>• Assessment of relative sensitivity of habitats</li> <li>• Mapping of sensitive biogenic habitats, and deepwater and inshore trawl footprints</li> </ul>
 <b>4. ECOSYSTEM EFFECTS</b> <ul style="list-style-type: none"> <li>• Trophic studies</li> <li>• Habitats of significance</li> <li>• Ecosystem indicators</li> <li>• Land-use effects</li> <li>• Climate variability</li> <li>• Climate Change</li> <li>• System productivity</li> </ul>	<ul style="list-style-type: none"> <li>• How do the ecosystems that support our fisheries function?</li> <li>• What are the key predator-prey or synergistic relationships in these systems?</li> <li>• Are our fisheries affecting food webs or ecosystem services?</li> <li>• What changes are occurring in the ecosystems that support our fisheries?</li> <li>• What is "habitat of particular significance for fisheries management"?</li> <li>• How do fisheries and/or land management affect fish habitat and fisheries production?</li> <li>• What are the major risks and opportunities from ocean-climate variability and trends?</li> </ul>	<ul style="list-style-type: none"> <li>• Habitat of significance: Kaipara Harbour fish habitats (SNA)</li> <li>• Habitat of significance: review of information for inshore finfish</li> <li>• Habitat of significance: coastal shark nursery areas (starting with rig)</li> <li>• Multi-impact risk assessment</li> <li>• Monitoring and indicators of environmental change for deepwater fisheries</li> <li>• Ecotrophic factors affecting highly migratory species</li> </ul>
 <b>5. MARINE BIODIVERSITY</b> <ul style="list-style-type: none"> <li>• Characterising NZ biodiversity</li> <li>• Functional ecology</li> <li>• Genetic diversity</li> <li>• Ocean climate</li> <li>• Metrics &amp; indicators</li> <li>• Threats &amp; impacts</li> <li>• Ross Sea &amp; IPY</li> </ul>	<ul style="list-style-type: none"> <li>• What are the key drivers of pattern in New Zealand's marine biodiversity?</li> <li>• How does biodiversity contribute to the resilience of ecosystems to perturbation and climate change?</li> <li>• What drives genetic connectivity within species?</li> <li>• What do we need to measure and monitor to assess risks and change?</li> <li>• How are biota adapted to polar conditions and what is their sensitivity to perturbation?</li> </ul>	<ul style="list-style-type: none"> <li>• Mapping key biogenic habitats</li> <li>• SPRFMO benthic habitats</li> <li>• Modelling seabed response and recovery from disturbance</li> <li>• Ocean acidification in fish habitat</li> <li>• Experimental response of shellfish to warming and acidification</li> <li>• Monitoring surface plankton</li> <li>• Implications of ocean acidification for plankton productivity</li> <li>• Marine environmental monitoring</li> </ul>

Figure 2.1: Summary of themes in the Aquatic Environment and Biodiversity Annual Review 2011.

CURRENT STATE OF KNOWLEDGE
<ul style="list-style-type: none"> <li>• Aggregate “on deck” bycatch of seabirds (and approximate species composition), marine mammals, and large sharks known reasonably well for offshore trawl and longline fisheries, but less well for inshore fisheries (where observer coverage has historically been low).</li> <li>• Incidental, cryptic, or unobserved mortality very poorly known (and difficult to assess).</li> <li>• Factors affecting fishing related mortality are well known for most seabirds and marine mammals.</li> <li>• Knowledge of population abundance is increasing for some key seabird species and well known for sea lions, but poorly known or dated for other seabirds, some species of dolphins, fur seals, and most sharks.</li> <li>• Qualitative or semi-quantitative risk assessments have been completed for almost all seabirds and marine mammals.</li> <li>• Fully quantitative risk assessments have been completed for two seabird populations, Hector’s / Maui’s dolphins, and sea lions.</li> <li>• Impact of fishing-related mortality on most protected species remains uncertain because of some key knowledge gaps.</li> <li>• Some methods of mitigating bycatch have been formally tested.</li> </ul>
<ul style="list-style-type: none"> <li>• Bycatch and discards are monitored and reported using observer records for the main deepwater and highly migratory fisheries.</li> <li>• Bycatch and discards for inshore vessels remain poorly known.</li> <li>• Some mitigation approaches have been assessed (e.g., for scampi trawl).</li> </ul>
<ul style="list-style-type: none"> <li>• Modelled predictions (that have been tested in deepwater) are available of the distribution of seabed habitats at a broad scale using classifications (BOMECS) and at finer scale for seamounts and some biogenic habitats.</li> <li>• Excellent understanding of the distribution of bottom trawling in offshore waters (but not in coastal waters, especially for most shellfish dredge fisheries).</li> <li>• Good understanding of the effects of trawling on some nearshore habitats.</li> <li>• General understanding of the effects of trawling on biogeochemical processes.</li> <li>• General understanding of the relative sensitivity of different habitats.</li> </ul>
<ul style="list-style-type: none"> <li>• Variability in the diets of key commercial species in the Chatham Rise ecosystem have been described as part of a wider biodiversity and MSI programme.</li> <li>• A preliminary trophic model of the Subantarctic ecosystem suggests a low productivity system supporting a simple food chain with high transfer efficiencies.</li> <li>• Atlases have been developed showing the distribution of spawning, pupping, egg-laying, and juveniles of key species (this needs finalising for inshore species).</li> <li>• A review of land-based effects on fish habitat and coastal biodiversity has been completed.</li> <li>• A start has been made on assessing ecosystem change over time (through fish-based indicators calculated from trawl survey data and acoustic time series of mesopelagic biomass)</li> <li>• A summary of ocean climate variability and change has been produced.</li> <li>• Broad reviews have been completed of the impacts of climate variability on fisheries (especially recruitment), but the likely impacts of ocean climate change or acidification remain poorly known.</li> <li>• This theme has links and synergies with MBIE, DOC, universities and the MPI biodiversity programme.s</li> </ul>
<ul style="list-style-type: none"> <li>• Taxonomy and ID Guides have been produced and specimens recorded in National Collections.</li> <li>• Biodiversity surveys completed on local scale (Fiordland, Spirits Bay, seamounts) and larger fishery scale (Norfolk ridge, Chatham Rise, Challenger Plateau, BOI).</li> <li>• Measures and indicators for marine biodiversity measures and ecosystem have been developed.</li> <li>• Predictive modelling techniques have been applied and habitat classification methods improved</li> <li>• Productivity in benthic communities has been measured.</li> <li>• Specimens from New Zealand have been genetically assessed and entered into the barcode of life.</li> <li>• Seamount connectivity, land-sea connectivity, and endemism have been studied.</li> <li>• A plan for monitoring the marine environment for long-term change is under development.</li> <li>• Demersal fish trophic studies on the Chatham Rise have been completed.</li> <li>• A review of NZ data from deep-sea and abyssal habitats has been completed.</li> <li>• A multidisciplinary study of longterm (1000 years) changes to NZ marine ecosystem is ongoing.</li> <li>• Latitudinal gradient project, ICECUBE and 2 large scale surveys in the Ross Sea have been conducted.</li> <li>• This theme has links and synergies with MBIE, DOC, universities and the MPI AEWG programmes</li> </ul>




Figure 2.1 continued: Summary of Themes in the Aquatic Environment & Biodiversity Review 2011

## **THEME 1: PROTECTED SPECIES**

### 3. New Zealand sea lions (*Phocarctos hookeri*)

<i>Scope of chapter</i>	This chapter outlines the biology of New Zealand (or Hooker's) sea lions ( <i>Phocarctos hookeri</i> ), the nature of fishing interactions, the management approach, trends in key indicators of fishing effects and major sources of uncertainty.
<i>Area</i>	Southern parts of the New Zealand EEZ and Territorial Sea.
<i>Focal localities</i>	Areas with significant fisheries interactions include the Auckland Islands Shelf, the Stewart/Snares Shelf and Campbell Plateau.
<i>Key issues</i>	Improving estimates of incidental bycatch in some trawl fisheries (e.g. scampi), improving estimates of SLED post-exit survival, improving understanding of interaction rate and improving understanding of the demographic processes underlying recent population trends.
<i>Emerging issues</i>	Assessing potential impacts of resource competition and/or resource limitation through ecosystem effects on NZ sea lion population viability. The role of fisheries impacts in light of ongoing declines in population size. Estimation of interactions given low numbers of observed captures.
<i>MPI Research (current)</i>	PRO2010-01 <i>Estimating the nature &amp; extent of incidental captures of seabirds, marine mammals &amp; turtles in New Zealand commercial fisheries</i> ; PRO2012-02 <i>Assess the risk posed to marine mammal populations from New Zealand fisheries</i> ; External review of the Breen-Fu-Gilbert model (SRP2011-04).
<i>Other Govt Research (current)</i>	<u>DOC Marine Conservation Services Programme (CSP)</u> : INT2012-01 <i>To understand the nature and extent of protected species interactions with New Zealand commercial fishing activities</i> ; POP2012-01 <i>To provide information on the population level and dynamics of the New Zealand sea lion at the Auckland Islands relevant to assessing the impacts of commercial fishing impacts on this population</i> ; POP2012-02 <i>To determine the key demographic factors driving the observed population decline of New Zealand sea lions at the Auckland Islands</i> . <u>NIWA Research</u> : SA123098 <i>Multispecies modelling to evaluate the potential drivers of decline in New Zealand sea lions</i> ; TMMA103 <i>Conservation of New Zealand's threatened iconic marine megafauna</i> .
<i>Links to 2030 objectives</i>	Objective 6: Manage impacts of fishing and aquaculture. Strategic Action 6.2: Set and monitor environmental standards, including for threatened and protected species and seabed impacts.
<i>Related issues/chapters</i>	See the New Zealand fur seal chapter.

#### 3.1. Context

Management of fisheries impacts on New Zealand (NZ) sea lions is legislated under the Marine Mammals Protection Act (MMPA) 1978 and the Fisheries Act (FA) 1996. Under s.3E of the MMPA, the Minister of Conservation, with the concurrence of the Minister for Primary Industries (formerly the Minister of Fisheries), may approve a population management plan (PMP). Although a NZ sea lion PMP was proposed by the Department of Conservation (DOC) in 2007 (DOC 2007), following consultation DOC decided not to proceed with the PMP.

All marine mammal species are designated as protected species under s.2(1) of the FA. In 2005, the Minister of Conservation approved the Conservation General Policy, which specifies in Policy 4.4 (f) that “*Protected marine species should be managed for their long-term viability and recovery*

*throughout their natural range.*” DOC’s Regional Conservation Management Strategies outline specific policies and objectives for protected marine species at a regional level. New Zealand’s sub-Antarctic islands, including Auckland and Campbell islands, were inscribed as a World Heritage area in 1998.

The Minister of Conservation gazetted the NZ sea lion as a threatened species in 1997. In 2009, DOC approved the *New Zealand sea lion species management plan<sup>2</sup>: 2009–2014* (DOC 2009). It aims: *“To make significant progress in facilitating an increase in the New Zealand sea lion population size and distribution.”* The plan specifies a number of goals, of which the following are most relevant for fisheries interactions:

*“To avoid or minimise adverse human interactions on the population and individuals.*

*To ensure comprehensive protection provisions are in place and enforced.*

*To ensure widespread stakeholder understanding, support and involvement in management measures.”*

In the absence of a PMP, the Ministry for Primary Industries (MPI, formerly the Ministry of Fisheries, MFish) manages fishing-related mortality of NZ sea lions under s.15(2) of the FA. Under that section, the Minister *“may take such measures as he or she considers are necessary to avoid, remedy, or mitigate the effect of fishing-related mortality on any protected species, and such measures may include setting a limit on fishing-related mortality.”*

Management of NZ sea lion bycatch aligns with Fisheries 2030 Objective 6: *Manage impacts of fishing and aquaculture.* Further, the management actions follow Strategic Action 6.2: *Set and monitor environmental standards, including for threatened and protected species and seabed impacts.*

The relevant National Fisheries Plan for the management of NZ sea lion bycatch is the National Fisheries Plan for Deepwater and Middle-depth Fisheries (the National Deepwater Plan). Under the National Deepwater Plan, the objective most relevant for management of NZ sea lions is Management Objective 2.5: *Manage deepwater and middle-depth fisheries to avoid or minimise adverse effects on the long-term viability of endangered, threatened and protected species.*

Specific objectives for the management of NZ sea lion bycatch will be outlined in the fishery-specific chapters of the National Deepwater Plan for the fisheries with which NZ sea lions are most likely to interact. These fisheries include trawl fisheries for arrow squid (SQU1T and SQU6T), southern blue whiting (SBW) and scampi (SCI). The SBW chapter of the National Deepwater Plan is complete and includes Operational Objective 2.2: *Ensure that incidental New Zealand sea lion mortalities, in the southern blue whiting fishery at the Campbell Islands (SBW6I), do not impact the long term viability of the sea lion population and captures are minimised through good operational practices.* Chapters in the National Deepwater Plan for arrow squid and scampi are under development.

Currently, MPI limits the actual or estimated bycatch of sea lions in the SQU6T trawl fishery based on tests of the likely performance of candidate bycatch control rules (and, hence, bycatch limits) using an integrated population and fishery model (Breen *et al.* 2010). Candidate rules are assessed against the following two criteria:

- a. A rule should provide for an increase in the sea lion population to more than 90% of carrying capacity<sup>3</sup>, or to within 10% of the population size that would have been attained in the

---

<sup>2</sup> The species management plan differs from the draft Population Management Plan in that it is quite broad in scope; providing a framework to guide the Department of Conservation in its management of the NZ sea lion over the next 5 years. The draft population management plan focused on options for managing the extent of incidental mortality of NZ sea lions from fishing through establishing a maximum allowable level of fishing-related mortality (MALFiRM) for all New Zealand fisheries waters.

<sup>3</sup> Carrying capacity in this instance applies to the current range. For managing the SQU6T fishery, carrying capacity refers to the maximum number of NZ sea lions that could be sustained on the Auckland Islands.

absence of fishing, and that these levels must be attained with 90% certainty, over 20-year and 100-year projections.

- b. A rule should attain a mean number of mature mammals that exceeded 90% of carrying capacity in the second 50 years of 100-year projection runs.

These management criteria were developed and approved in 2003 by a Technical Working Group comprised of MFish, DOC, squid industry representatives, and environmental groups.

Likely performance is also assessed against two additional criteria proposed by DOC:

- a) A rule should maintain numbers above 90% of the carrying capacity in at least 18 of the first 20 years.
- b) A rule should lead to at least a 50% chance of an increase in the number of mature animals over the first 20 years of the model projections.

## **3.2. Biology**

### **3.2.1. Taxonomy**

The NZ sea lion (*Phocarctos hookeri*, Gray, 1844) is one of only two species of otariid (eared seals, including fur seals and sea lions) native to New Zealand, the other being the NZ fur seal (*Arctocephalus forsteri*, Lesson, 1828). The NZ sea lion is also New Zealand's only endemic pinniped.

### **3.2.2. Distribution**

Before human habitation, NZ sea lions ranged around the North and South Islands of New Zealand. Pre-European remains of NZ sea lions have been identified from at least 47 archaeological sites, ranging from Stewart Island to North Cape, with most occurring in the southern half of the South Island (Smith 1989, 2011, Childerhouse and Gales 1998, Gill 1998). Subsistence hunting on the mainland and subsequent commercial harvest from outlying islands of NZ sea lions for skins and oil resulted in population decline and contraction of the species' range (Gales 1995, Childerhouse and Gales 1998, Nagaoka 2001, 2006). Currently, most NZ sea lions are found in the New Zealand Sub-Antarctic, with individuals ranging to the NZ mainland and Macquarie Island.

NZ sea lion breeding colonies<sup>4</sup> are highly localized, with most pups being born at two main breeding areas, the Auckland Islands and Campbell Island (Wilkinson *et al.* 2003, Chilvers 2008). At the Auckland Islands, there are three breeding colonies: Enderby Island (mainly at Sandy Bay and South East Point); Dundas Island; and Figure of Eight Island. On Campbell Island there is one breeding colony at Davis Point, another colony at Paradise Point, plus a small number of non-colonial breeders (Wilkinson *et al.* 2003, Chilvers 2008, Maloney *et al.* 2009, Maloney *et al.* 2012). Twenty-five sea lion pups were captured and tagged around Stewart Island during a DOC recreational hut and track maintenance trip in March 2012. Breeding on the Auckland Islands represents 71–87% of the pup production for the species, with the remaining 13–29% occurring on Campbell Island (based on concurrent pup counts in 2003, 2008 and 2010; see section 3.2.5).

---

<sup>4</sup> DOC (2009) defines colonies as “haul-out sites where 35 pups or more are born each year for a period of 5 years or more.” Haul-out sites are defined as “terrestrial sites where NZ sea lions occur but where pups are not born, or where less than 35 pups are born per year over 5 consecutive years.”

Although breeding is concentrated on the Auckland Islands and Campbell Island, occasional births have been reported from the Snares and Stewart Islands (Wilkinson *et al.* 2003, Chilvers *et al.* 2007). Breeding is also taking place on the New Zealand mainland at the Otago peninsula, mainly the result of a single female arriving in 1992 and giving birth in 1993 (McConkey *et al.*, 2002).

On land, NZ sea lions are able to travel long distances and climb high hills, and are found in a variety of habitats including sandy beaches, grass fields, bedrock, and dense bush and forest (Gales 1995, Augé *et al.* 2012). Following the end of the females' oestrus cycle in late January, adult and sub-adult males disperse throughout the species' range, whereas dispersal of females (both breeding and non-breeding) appears more restricted (Marlow 1975, Robertson *et al.* 2006, Chilvers and Wilkinson 2008).

### 3.2.3. Foraging ecology

Most foraging studies have been conducted on lactating female NZ sea lions from Enderby Island (Chilvers *et al.* 2005a, 2006, Chilvers and Wilkinson 2009, although work is underway at Campbell Island under NIWA project TMMA103, *Conservation of New Zealand's threatened iconic marine megafauna*). These show that females from this place forage primarily within the Auckland Islands continental shelf and its northern edge, and that individuals show strong foraging site fidelity both within and across years. Satellite tagging data from lactating females showed that the mean return distance travelled per foraging trip is  $423 \pm 43$  km ( $n = 26$ ), which is greater than that recorded for any other sea lion species (Chilvers *et al.* 2005a). While foraging, about half of the time is spent submerged, with a mean dive depth of  $130 \pm 5$  m (max. 597 m) and a mean dive duration of  $4 \pm 1$  minutes (max. 14.5 minutes; Chilvers *et al.* 2006). NZ sea lions, like most pinnipeds, may use their whiskers to help them capture prey at depths where light does not penetrate (Marshall 2008, Hanke *et al.* 2010).

Studies conducted on female NZ sea lions suggest that the foraging behaviour of each individual falls into one of two distinct categories, benthic or meso-pelagic (Chilvers and Wilkinson 2009). Benthic divers have fairly consistent dive profiles, reaching similar depths (120 m on average) on consecutive dives in relatively shallow water to presumably feed on benthic prey. Meso-pelagic divers, by contrast, exhibit more varied dive profiles, undertaking both deep ( $> 200$  m) and shallow ( $< 50$  m) dives over deeper water. Benthic divers tend to forage further from their breeding colonies, making their way to the north-eastern limits of Auckland Islands' shelf, whereas meso-pelagic divers tend to forage along the north-western edge of the shelf over depths of approximately 3000 m (Chilvers and Wilkinson 2009).

The differences in dive profiles have further implications for the animals' estimated aerobic dive limits (ADL; Chilvers *et al.* 2006), defined as the maximum amount of time that can be spent underwater without increasing blood lactate concentrations (a by-product of anaerobic metabolism). If animals exceed their ADL and accumulate lactate, they must surface and go through a recovery period in order to aerobically metabolize the lactate before they can undertake subsequent dives. Chilvers *et al.* (2006) estimated that lactating female NZ sea lions exceed their ADL on 69% of all dives, a much higher proportion than most other otariids (which exceed their ADL for only 4–10% of dives; Chilvers *et al.* 2006). NZ sea lions that exhibit benthic diving profiles are estimated to exceed their ADL on 82% of dives, compared with 51% for meso-pelagic divers (Chilvers 2008).

Chilvers *et al.* (2006) and Chilvers and Wilkinson (2009) suggested that the long, deep diving behaviour, the propensity to exceed their estimated ADL, and differences in physical condition and age at first reproduction from animals at Otago together indicate that females from the Auckland Islands may be foraging at or near their physiological limits. However, Bowen (2012) suggested a lack of relationship between surface time and anaerobic diving would seem to indicate that ADL has been underestimated. Further, given a number of studies of diving behaviour were conducted during

early lactation when the demands of offspring are less than they would be later in lactation, Bowen (2012) considered it unlikely that females are operating at or near a physiological limit.

Adult females at Otago are generally heavier for a given age, breed earlier, undertake shorter foraging trips, and have shallower dive profiles compared with females from the Auckland Islands (Table 3.1). Any observed differences may reflect differences in environment between the Auckland Islands and the Otago peninsula, a founder effect, or a combination of these or other factors.

**Table 3.1: Comparison of select characteristics between adult female NZ sea lions from the Auckland Islands and those from the Otago peninsula (Chilvers *et al.* 2006, Augé *et al.* 2011a, 2011b, 2011c). Data are means  $\pm$  SE (where available).**

Characteristic	Auckland Islands	Otago
Reproduction at age 4	< 5% of females	> 85% of females
Average mass at 8-13 years of age	112 kg	152 kg
Foraging distance from shore	102.0 $\pm$ 7.7 km (max = 175 km)	4.7 $\pm$ 1.6 km (max = 25 km)
Time spent foraging at sea	66.2 $\pm$ 4.2 hrs	11.8 $\pm$ 1.5 hrs
Dive depth	129.4 $\pm$ 5.3 m (max = 597 m)	20.2 $\pm$ 24.5 m (max = 389 m)
Dives estimated to exceed ADL	68.7 $\pm$ 4.4 percent	7.1 $\pm$ 8.1 percent

NZ sea lions are generalist predators with a varied diet that includes fish (rattail, red cod, opalfish, hoki), cephalopods (octopus, squid), crustaceans (lobster krill, scampi), and salps (Cawthorn *et al.* 1985; Childerhouse *et al.* 2001; Meynier *et al.* 2009). The three main methods used to assess NZ sea lion diets involve analyses of stomach contents, scats and regurgitate, and the fatty acid composition of blubber (Meynier *et al.* 2008). Stomach contents of by-caught animals tend to be biased towards the target species of the fishery concerned (e.g. squid in the SQU6T fishery), whereas scats and regurgitates are biased towards less digestible prey (Meynier *et al.* 2008). Stomach, scat and regurgitate approaches tend to reflect only recent prey (Meynier *et al.* 2008). By contrast, analysis of the fatty acid composition of blubber provides a longer-term perspective on diets ranging from weeks to months (although individual prey species are not identifiable). This approach suggests that the diet of female NZ sea lions tends to include proportionally more arrow squid (*Nototodarus sloanii*) and proportionally less red cod (*Pseudophycis bachus*) and scampi (*Metanephrops challengeri*) than for male NZ sea lions, while lactating and non-lactating females do not differ in their diet (Meynier *et al.* 2008; Meynier 2010).

### 3.2.4. Reproductive biology

NZ sea lions exhibit marked sexual dimorphism, with adult males being larger and darker in colour than adult females (Walker and Ling 1981, Cawthorn *et al.* 1985). Cawthorn *et al.* (1985) and Dickie (1999) estimated the maximum age of males and females to be 21 and 23 years, respectively, but Childerhouse *et al.* (2010a) recently reported a maximum estimated age for females of 28 years (although the AEWG had some concerns about the methods used and this estimate may not be reliable). Although females can become sexually mature as early as age 2 and give birth the following year, most do not breed until they are 6 years old (Childerhouse *et al.* 2010a). Males generally reach sexual maturity at 4 years of age, but because of their polygynous colonial breeding strategy (i.e., males actively defend territories and mate with multiple females within a harem) they are only able to successfully breed at 7–9 years old, once they have attained sufficient physical size (Marlow 1975, Cawthorn *et al.* 1985). Reproductive rate in females increases rapidly between the ages of 3 and 7, reaching a plateau until the age of approximately 15 and declining rapidly thereafter, with the maximum recorded age at reproduction being 26 years (Breen *et al.* 2010, Childerhouse *et al.* 2010b, Chilvers *et al.* 2010). Chilvers *et al.* (2010) estimated from tagged sea lions that the median lifetime

reproductive output of a female NZ sea lion was 4.4 pups, and 27% of all females that survive to age 3 never breed. Analysis of tag-resight data from female New Zealand sea lions on Enderby Island indicates average annual breeding probability is approximately 0.30-0.35 for prime-age females that did not breed in the previous year (ranges reflect variation relating to the definition of breeders) and 0.65-0.68 for prime-age females that did breed in the previous year (MacKenzie 2011).

NZ sea lions are philopatric (i.e., they return to breed at the same location where they were born, although more so for females than males). Breeding is highly synchronised and starts in late November when adult males establish territories for their harems (Robertson *et al.* 2006, Chilvers and Wilkinson 2008). Pregnant and non-pregnant females appear at the breeding colonies in December and early January, with pregnant females giving birth to a single pup in late December before entering oestrus 7–10 days later and mating again (Marlow 1975). Twin births and the fostering of pups in NZ sea lions are rare (Childerhouse and Gales 2001). Shortly after the breeding season ends in mid-January, the harems break up with the males dispersing offshore and females often moving away from the rookeries with their pups (Marlow 1975, Cawthorn *et al.* 1985).

Pups at birth weigh 8–12 kg with parental care restricted to females (Walker and Ling 1981, Cawthorn *et al.* 1985, Chilvers *et al.* 2006). Females remain ashore for about 10 days after giving birth before alternating between foraging trips lasting approximately two days out at sea and returning for about one day to suckle their pups (Gales and Mattlin 1997, Chilvers *et al.* 2005). New Zealand pup growth rates are lower than those reported for other sea lion species, and may be linked to a relatively low concentration of lipids in the females' milk during early lactation (Riet-Sapirza *et al.* 2012, Chilvers 2008). Pups are weaned after about 10–12 months (Marlow 1975, Gales and Mattlin 1997).

### 3.2.5. Population biology

For NZ sea lions, the overall size of the population is indexed using estimates of the number of pups that are born each year (Chilvers *et al.* 2007). Since 1995, the Department of Conservation (DOC) has conducted mark-recapture counts at each of the main breeding colonies at the Auckland Islands to estimate annual pup production (i.e., the total number of pups born each year, including dead and live animals; Robertson and Chilvers 2011). The data show a decline in pup production from a peak of 3021 in 1997/98 to a low of  $1501 \pm 16$  pups in 2008/09 (Chilvers and Wilkinson 2011, Robertson and Chilvers 2011; Table 3.2), with the largest single-year decline (31%) occurring between the 2007/08 and 2008/09 counts. The most recent estimate of pup production for the Auckland Islands population was  $1683 \pm 16$  pups in 2011/12 (Chilvers 2012a) and a project is underway to obtain a comparable estimate for 2012/13 (POP2012-01).

Total NZ sea lion abundance (including pups) at the Auckland Islands has been estimated using Bayesian population models (Breen *et al.* 2003, Breen and Kim 2006a, Breen and Kim 2006b, Breen *et al.* 2010). Although other abundance estimates are available (e.g. Gales and Fletcher 1999), the integrated models are preferred because they take into account a variety of age-specific factors (breeding, survival, maturity, vulnerability to fishing, and the proportion incidentally captured by fishing), as well as data on the re-sighting of tagged animals and pup production estimates, to generate estimates of the overall size of the NZ sea lion population inhabiting the Auckland Islands (Table 3.2). The most recent estimate of NZ sea lion abundance for the Auckland Islands population was 12 065 animals (90% CI: 11 160–13 061) in 2009. The integrated model suggested a net decline at the Auckland Islands of 23% between 1995 and 2009, or 29% between the maximum estimated population size in 1998 and 2009.

**Table 3.2: Pup production and population estimates of NZ sea lions from the Auckland Islands from 1995 to 2010. Pup production data are direct counts or mark-recapture estimates from Chilvers *et al.* (2007), Robertson and Chilvers (2011) and Chilvers (2012a). Standard errors only apply to the portion of pup production estimated using mark-recapture methods. Population estimates from P. Breen, estimated in the model by Breen *et al.* 2010. Year refers to the second year of a breeding season (e.g., 2010 refers to the 2009-10 season).**

Year	Pup production estimate		Population size estimate	
	Mean	Standard error (for mark recapture estimates)	Median	90% confidence interval
1995	2 518	21	15 675	14 732–16 757
1996	2 685	22	16 226	15 238–17 318
1997	2 975	26	16 693	15 656–17 829
1998	3 021	94	16 911	15 786–18 128
1999	2 867	33	15 091	13 932–16 456
2000	2 856	43	15 248	14 078–16 586
2001	2 859	24	15 005	13 870–16 282
2002	2 282	34	13 890	12 856–15 079
2003	2 518	38	14 141	13 107–15 295
2004	2 515	40	14 096	13 057–15 278
2005	2 148	34	13 369	12 383–14 518
2006	2 089	30	13 110	12 150–14 156
2007	2 224	38	13 199	12 231–14 215
2008	2 175	44	12 733	11 786–13 757
2009	1 501	16	12 065	11 160–13 061
2010	1 814	36		
2011	1 550 <sup>5</sup>	41		
2012	1 683	16		

For the Campbell Island population, pup production was estimated at 681–726 pups in 2010 (Robertson and Chilvers 2011, Maloney *et al.* 2012). Pup production estimates at Campbell Island are increasing over time, although there have been changes to the methodology (Maloney *et al.* 2009). Previous estimates of total pup production were: 150 in 1992/93; 385 in 2003; and 583 in 2007-08 (Cawthorn 1993, Childerhouse *et al.* 2005, Maloney *et al.* 2009). There were also minimum pup counts of 51 in 1987/88, 122 in 1991/92 and 78 (from a partial count) in 1997/98 (Moore and Moffat 1990, McNally *et al.* 2001, M. Fraser, unpubl. data cited in Maloney *et al.* 2009).

For the Otago sub-population, annual pup production has ranged from 0 to 7 pups since the 1994/95 breeding season, with five pups recorded in 2010/11 (McConkey *et al.* 2002, Augé 2011). A modelling exercise suggested that this population can expand to 9–22 adult females by 2018 (Lalas and Bradshaw 2003). The sub-population at Otago is of special interest because it highlights the potential for establishing new breeding colonies, in this case from a single pregnant female (McConkey *et al.* 2002).

Established anthropogenic sources of mortality in NZ sea lion include: historic subsistence hunting and commercial harvest (Gales 1995, Childerhouse and Gales 1998); pup entrapment in rabbit burrows prior to rabbit eradication from Enderby Island in 1993 (Gales and Fletcher 1999); human disturbance, including attacks by dogs, vehicle strikes and deliberate shooting on mainland New Zealand (Gales 1995); and fisheries bycatch (see below).

In addition to the established effects, there are a number of other anthropogenic effects that may also influence NZ sea lion mortality. However their role, if any, is presently unclear. These include: possible competition for resources between NZ sea lions and the various fisheries (Robertson and

<sup>5</sup> Due to extreme weather conditions there was some delay in making the 2010/11 pup count which may affect comparability with previous years. However DOC's analysis suggests any such effect is unlikely to be large (Chilvers and Wilkinson 2011).

Chilvers 2011, Bowen 2012); effects of organic and inorganic pollutants, including polychlorinated biphenyls (PCBs), dichlorodiphenyltrichloroethane (DDT) and heavy metals such as mercury and cadmium (Baker 1999, Robertson and Chilvers 2011); and impacts of eco-tourism.

Other sources of mortality include epizootics, particularly *Campylobacter* which killed 1600 pups (53% of pup production) and at least 74 adult females on the Auckland Islands in 1997/98 (Wilkinson *et al.* 2003, Robertson and Chilvers 2011) and *Klebsiella pneumoniae* which killed 33% and 21% of pups on the Auckland Islands in 2001/02 and 2002/03 respectively (Wilkinson *et al.* 2006). The 1998 epizootic event may have affected the fecundity of the surviving pups; reducing their breeding rate relative to other cohorts (Gilbert and Chilvers 2008). There are also occurrences of predation by sharks (Cawthorn *et al.* 1985, Robertson and Chilvers 2011), starvation of pups if they become separated from their mothers (Walker and Ling 1981, Castinel *et al.* 2007), drowning in wallows and male aggression towards females and pups (Wilkinson *et al.* 2000, Chilvers *et al.* 2005b).

Analysis of tag-resight data on Enderby Island yielded estimates of average annual survival for prime-age females of 0.90 for females that did not breed and 0.95 for females that did breed, with no indication of a systematic change in survival during the period 1997/98 to 2010/11 (MacKenzie 2011). Further analysis of tag-resight data is planned under DOC project POP2012-02 to determine the key demographic factors driving the observed population decline of New Zealand sea lions at the Auckland Islands.

Despite a historic reduction in population size as a result of subsistence hunting and commercial harvest, the NZ sea lion population does not display low genetic diversity at microsatellite loci and thus does not appear to have suffered effects of genetic drift and inbreeding depression (Robertson and Chilvers 2011).

### 3.2.6. Conservation biology and threat classification

Threat classification is an established approach for identifying species at risk of extinction (IUCN 2010). The risk of extinction for NZ sea lions has been assessed under two threat classification systems, the International Union for the Conservation of Nature (IUCN) Red List of Threatened Species (IUCN 2010) and the New Zealand Threat Classification System (Townsend *et al.* 2008).

In 2008, the IUCN updated the Red List status of NZ sea lions, listing them as Vulnerable, A3b<sup>6</sup> on the basis of a marked (30%) decline in pup production in the last 10 years, at some of the major rookeries (Gales 2008). The IUCN further recommended that the species should be reviewed within a decade in light of what they considered to be the current status of NZ sea lions (i.e., declining pup production, reducing population size, severe disease outbreaks).

In 2010, DOC updated the New Zealand Threat Classification status of all NZ marine mammals (Baker *et al.* 2010). In the revised list, NZ sea lions had their threat classification increased from At Risk, Range Restricted<sup>7</sup> to Nationally Critical under criterion C<sup>8</sup> with a Range Restricted qualifier based on the recent rate of decline (Baker *et al.* 2010).

---

<sup>6</sup> A taxon is listed as 'Vulnerable' if it is considered to be facing a high risk of extinction in the wild. A3b refers to a reduction in population size (A), based on a reduction of  $\geq 30\%$  over the last 10 years or three generations (whichever is longer up to a maximum of 100 years (3); and when considering an index of abundance that is appropriate to the taxon (b; IUCN 2010).

<sup>7</sup> A taxon is listed as 'Range Restricted' if it is confined to specific substrates, habitats or geographic areas of less than 1000 km<sup>2</sup> (100 000 ha); this is assessed by taking into account the area of occupied habitat of all sub-populations (Townsend *et al.* 2008).

<sup>8</sup> A taxon is listed as 'Nationally Critical' under criterion C if the population (irrespective of size or number of sub-populations) has a very high (rate of) ongoing or predicted decline; greater than 70% over 10 years or three generations, whichever is longer (Townsend *et al.* 2008).

### 3.3. **Global understanding of fisheries interactions**

Reviews of fisheries interactions among pinnipeds globally can be found in Read *et al.* 2006, Woodley and Lavigne (1991), Katsanevakis (2008) and Moore *et al.* (2009). Because NZ sea lions are endemic to New Zealand, the global understanding of fisheries interactions for this species is outlined under state of knowledge in New Zealand. For related information on fishing interactions for NZ fur seals, both within New Zealand and overseas, see the NZ fur seal chapter.

### 3.4. **State of knowledge in New Zealand**

NZ sea lions interact with trawl fisheries resulting in incidental bycatch, specifically from animals being caught and drowned in the trawl nets. These interactions are largely confined to trawl fisheries in Sub-Antarctic waters (Figure 3.1); particularly the Auckland Islands arrow squid fishery (SQU6T), but also the Auckland Islands scampi fishery (SCI6A), other Auckland Islands trawl fisheries, the Campbell Island southern blue whiting (*Micromesistius australis*) fishery (SBW6I) and the Stewart-Snares shelf fisheries targeting mainly arrow squid (SQU1T; Thompson and Abraham 2010, Thompson *et al.* 2011, Thompson *et al.* 2012).<sup>9</sup>

NZ sea lions forage to depths of up to 600 m (Table 3.1), within the habitat where depth ranges for prey species range from 0–500 m for arrow squid, 250–600 m for spawning southern blue whiting and 350–550 m for scampi (Tuck 2009, Ministry of Fisheries 2011). There is seasonal variation in the distribution overlap between NZ sea lions and the target species fisheries (Table 3.3). Breeding male sea lions, breeding ashore between November and January with occasional trips to sea, then migrate away from the Auckland island area (Robertson *et al.* 2006). Breeding females are in the Auckland island area year round, ashore to give birth for up to 10 days during December and January and then dividing their time between foraging at sea (~2days) and suckling their pup ashore (~1.5 days; Chilvers *et al.* 2005a). The SQU6T fishery currently operates between February and July, peaking between February and May, whereas the SQU1T fishery operates between December and May, peaking between January and April, before the squid spawn. The SBW6I fishery operates in August and September, peaking in the latter month, when the fish aggregate to spawn. The SCI6A fishery may operate at any time of the year but does not operate continuously.

#### 3.4.1. **Quantifying fisheries interactions**

Since 1988, the level of NZ sea lion bycatch has been monitored by government observers aboard a proportion of the fishing fleet in the SQU6T fishery (Wilkinson *et al.* 2003), generally amounting to around 20–40% observer coverage between 1995 and 2010 but reaching almost 100% during the 2001/02 season (see Table 3.4). Over the same period, there has also been 1–15% observer coverage for non-squid trawl fisheries operating around the Auckland Islands (primarily targeting scampi, but also jack mackerel, orange roughy and hoki), 20–60% observer coverage in the Campbell Island southern blue whiting fishery, and 8–43% observer coverage for the Stewart-Snares shelf trawl fisheries (primarily targeting squid, but also hoki, jack mackerel and barracouta; Table 3.4). Unobserved trips have tended to report NZ sea lion captures at a lower rate than observed trips across all observed fisheries. Fishers reported 177 NZ sea lion captures between 1998–99 and 2008–09, while observers reported 196 captures over the same period (Abraham and Thompson 2011). Observers observed an overall average of 4.7–11.2% of trawl tows each year over this time period, but fisheries where most sea lions are caught had higher observer coverage.

<sup>9</sup> See the Report from the Fisheries Assessment Plenary, May 2011 (Ministry of Fisheries 2011) for further information regarding the biology and stock assessments for these species.

**Table 3.3: Monthly distribution of NZ sea lion activity and the main trawl fisheries with observed reports of NZ sea lion incidental captures (see text for details).**

NZ sea lions	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
Breeding males	Dispersed at sea or at haulouts		At breeding colony			Dispersed at sea or at haulouts						
Breeding females	At sea			At breeding colony		At breeding colony and at-sea foraging and suckling						
Pups	At sea			At breeding colony								
Non-breeders	Dispersed at sea, at haulouts, or breeding colony periphery											
Major fisheries	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
Squid				Stewart-Snares Shelf		Auckland Islands and Stewart-Snares Shelf				Auckland Islands		
Southern blue whiting	Pukaki Rise and Campbell Rise											Bounty Islands
Scampi	Auckland Islands											

The number of NZ sea lion captures reported by observers has been incorporated in increasingly sophisticated models to estimate the total number of captures across the entire fishing fleet in each fishing year (Smith and Baird 2007b, Thompson and Abraham 2010, Abraham and Thompson 2011). This approach is currently applied using information collected under DOC project INT2012-01 and analysed under MPI project PRO2010-01 (Thompson *et al.* 2011, Thompson *et al.* 2012). Estimates in Table 3.4 for the SQU6T and Campbell Island fisheries were generated using Bayesian models, whereas those for the Stewart-Snares and the Auckland Islands scampi and Auckland Islands other fisheries were generated using ratio estimates (Thompson *et al.* 2012). Captures comprise the number of NZ sea lions brought on deck (both dead and alive), and necessarily exclude the unknown fraction of animals that exit trawls through Sea Lion Exclusion Devices (SLEDs) as well as those that were decomposed upon capture or that climbed aboard vessels (Smith and Baird 2007b, Thompson and Abraham 2010 Thompson *et al.* 2011). Only 8 of the 248 captures from 1995/96 to 2008/09 were released alive (Thompson and Abraham 2010). Interactions are defined as the number of sea lion that would have been caught if no SLEDs were used (Thompson *et al.* 2012).

In the years since SLEDs were introduced in the SQU6T fishery, both the observed and estimated numbers of NZ sea lion captures have declined overall, except for a slight increase in 2009/10 (Table 3.4). Conversely, for those other fisheries where SLEDs are not deployed, observed and estimated numbers of NZ sea lion captures increased in the Campbell Island southern blue whiting fishery to a peak in 2010 (Table 3.4). For the Stewart-Snares and the Auckland Islands non-squid fisheries, the observed and estimated numbers of NZ sea lion captures have fluctuated without trend (Table 3.4).

Capture rate is defined as the number of NZ sea lions caught per 100 tows. Strike rate is defined as the number of NZ sea lions that would be caught per 100 tows if no SLEDs were fitted. Models indicate that the interaction rate of female NZ sea lions (equivalent to the capture rate were no SLEDs fitted) is influenced by a number of factors including year, distance from rookery, tow duration, and change of tow direction (Smith and Baird 2005). Conversely, the interaction rate of male NZ sea lions is

influenced by year, the number of days into the fishery (males leave the rookeries soon after mating whereas females remain with the pups), and time of day (Smith and Baird 2005).




Figure 3.1: Distribution of trawl fishing effort and observed NZ sea lion captures, 2002-03 to 2010-11 (<http://data.dragonfly.co.nz/psc/>). Fishing effort is mapped into 0.2-degree cells, with the colour of each cell being related to the amount of effort. Observed fishing events are indicated by black dots, and observed captures are indicated by red dots. Fishing is only shown if the effort could be assigned a latitude and longitude, and if there were three or more vessels fishing within a cell. In this case, 96.0% of the effort is shown.

**Table 3.4a: Effort, observed and estimated NZ sea lion captures in trawl fisheries by fishing year in the New Zealand EEZ (<http://data.dragonfly.co.nz/pse/>). For each fishing year, the table gives the the total number of tows; the observer coverage (the percentage of tows that were observed); the number of observed captures (both dead and alive); the capture rate (captures per hundred tows); the estimation method used (model, ratio or both combined); the mean number of estimated total captures (with 95% confidence interval); the mean number of estimated total interactions (with 95% confidence interval), and the stike rate (interactions per hundred tows). For more information on the methods used to prepare the data, see Thompson *et al.* (2012).**

Fishing year	Fishing effort		Observed captures		Estimated captures			Estimated interations		Estimated strike rate	
	All effort	% obs	Number	Rate	Method	Mean	95% c.i.	Mean	95% c.i.	Mean	95% c.i.
<i>Trawl fisheries</i>											
1995–96	10 081	10	16	1.5	Both	148	85–242	148	85–243	1.5	0.8–2.4
1996–97	10 941	15	28	1.7	Both	155	104–221	155	102–225	1.4	0.9–2.1
1997–98	9 964	14	14	1.0	Both	76	47–119	76	45–121	0.8	0.5–1.2
1998–99	10 551	16	6	0.4	Both	33	20–49	33	19–50	0.3	0.2–0.5
1999–00	9 043	22	28	1.4	Both	88	63–129	89	59–130	1.0	0.7–1.4
2000–01	8 910	40	46	1.3	Both	61	52–72	83	59–111	0.9	0.7–1.2
2001–02	9 945	19	23	1.2	Both	64	46–88	94	61–139	0.9	0.6–1.4
2002–03	8 308	19	11	0.7	Both	34	22–48	62	37–97	0.7	0.4–1.2
2003–04	10 033	23	21	0.9	Both	61	43–85	214	120–376	2.1	1.2–3.7
2004–05	11 109	23	14	0.5	Both	53	36–77	181	94–325	1.6	0.8–2.9
2005–06	9 316	21	14	0.7	Both	52	35–75	174	86–334	1.9	0.9–3.6
2006–07	6 728	24	15	0.9	Both	47	32–66	118	59–235	1.8	0.9–3.5
2007–08	6 545	33	8	0.4	Both	29	18–42	118	35–418	1.8	0.5–6.4
2008–09	6 677	27	3	0.2	Both	22	12–36	103	25–383	1.5	0.4–5.7
2009–10	5 541	34	15	0.8	Both	46	32–66	141	51–439	2.5	0.9–7.9
2010–11	6 389	31	6	0.3	Both	29	17–43	81	26–259	1.3	0.4–4.1
<i>Auckland Islands squid</i>											
1995–96	4 467	12	13	2.4	Model	131	69–226	131	67–224	2.9	1.6–5.0
1996–97	3 716	19	28	3.9	Model	142	91–208	142	89–210	3.8	2.6–5.5
1997–98	1 441	22	13	4.2	Model	60	33–102	60	31–104	4.2	2.5–6.9
1998–99	402	38	5	3.2	Model	14	5–27	15	5–29	3.6	2.1–5.9
1999–00	1 206	36	25	5.7	Model	69	45–107	69	42–108	5.8	4.0–8.6
2000–01	583	99	39	6.7	Model	39	39–40	61	39–87	10.4	8.6–13.1
2001–02*	1 648	34	21	3.7	Model	43	30–64	73	43–116	4.4	3.0–6.6
2002–03	1 470	29	11	2.6	Model	19	13–29	48	24–81	3.2	2.0–5.1
2003–04	2 594	30	16	2.0	Model	41	26–62	194	100–356	7.5	4.0–13.5
2004–05^	2 706	30	9	1.1	Model	31	17–51	159	73–303	5.9	2.7–11.1
2005–06	2 462	28	9	1.3	Model	28	15–45	149	62–308	6.0	2.7–12.5
2006–07	1 320	41	7	1.3	Model	16	9–27	87	29–201	6.6	2.3–14.8
2007–08	1 265	46	5	0.9	Model	12	6–21	101	19–396	8.0	1.6–30.9
2008–09	1 925	40	2	0.3	Model	8	3–17	89	12–365	4.6	0.7–18.4
2009–10	1 190	25	3	1.0	Model	13	5–27	107	18–402	9.0	1.7–33.6
2010–11	1 586	34	0	–	Model	4	0–11	56	4–233	3.5	0.4–14.9

\* SLEDs introduced. ^ SLEDs standardised and in widespread use.

**Table 3.4b: Effort, observed and estimated NZ sea lion captures in trawl fisheries by fishing year in the New Zealand EEZ (<http://data.dragonfly.co.nz/psc/>). For each fishing year, the table gives the the total number of tows; the observer coverage (the percentage of tows that were observed); the number of observed captures (both dead and alive); the capture rate (captures per hundred tows or per thousand hooks); the estimation method used (model, ratio or both combined); and the mean number of estimated total captures (with 95% confidence interval). For more information on the methods used to prepare the data, see Thompson *et al.* (2012).**

Fishing year	Fishing effort		Observed captures		Estimated captures		
	All effort	% obs	Number	Rate	Method	Mean	95% c.i.
<i>Auckland Islands scampi</i>							
1995-96	1 303	5	2	3.2	Ratio	11	4–19
1996-97	1 222	15	0	-	Ratio	7	2–15
1997-98	1 107	11	0	-	Ratio	7	1–15
1998-99	1 254	2	0	-	Ratio	9	2–18
1999-00	1 383	5	0	-	Ratio	9	3–18
2000-01	1 417	6	4	4.8	Ratio	14	7–23
2001-02	1 604	9	0	-	Ratio	10	3–20
2002-03	1 351	11	0	-	Ratio	9	2–17
2003-04	1 363	12	3	1.8	Ratio	12	5–20
2004-05	1 275	0	NA	NA	Ratio	9	3–18
2005-06	1 331	9	1	0.9	Ratio	10	3–18
2006-07	1 328	7	1	1.1	Ratio	10	4–19
2007-08	1 327	7	0	-	Ratio	9	2–18
2008-09	1 457	4	1	1.6	Ratio	11	4–21
2009-10	940	10	0	-	Ratio	6	1–13
2010-11	1 401	15	0	-	Ratio	9	2–17
<i>Auckland Islands other</i>							
1995-96	405	6	1	4.0	Ratio	3	1–6
1996-97	296	4	0	-	Ratio	1	0–4
1997-98	684	17	1	0.9	Ratio	3	1–8
1998-99	525	10	1	1.8	Ratio	3	1–7
1999-00	750	13	0	-	Ratio	3	0–8
2000-01	577	7	0	-	Ratio	2	0–7
2001-02	589	4	0	-	Ratio	2	0–7
2002-03	543	13	0	-	Ratio	2	0–7
2003-04	289	17	0	-	Ratio	1	0–4
2004-05	170	7	0	-	Ratio	1	0–3
2005-06	39	15	0	-	Ratio	0	0–1
2006-07	38	5	0	-	Ratio	0	0–1
2007-08	147	45	0	-	Ratio	0	0–2
2008-09	121	50	0	-	Ratio	0	0–2
2009-10	77	66	0	-	Ratio	0	0–1
2010-11	131	37	0	-	Ratio	0	0–2

**Table 3.4c: Effort, observed and estimated NZ sea lion captures in trawl fisheries by fishing year (calendar year for SBW) in the New Zealand EEZ (<http://data.dragonfly.co.nz/psc/>). For each fishing year, the table gives the the total number of tows; the observer coverage (the percentage of tows that were observed); the number of observed captures (both dead and alive); the capture rate (captures per hundred tows or per thousand hooks); the estimation method used (model, ratio or both combined); and the mean number of estimated total captures (with 95% confidence interval). For more information on the methods used to prepare the data, see Thompson *et al.* (2012).**


Fishing year	Fishing effort		Observed captures		Estimated captures		
	All effort	% observed	Number	Rate	Type	Mean	95% c.i.
<i>Campbell Island SBW</i>							
1996	474	27	0	-	Model	0	0–4
1997	641	34	0	-	Model	1	0–3
1998	963	28	0	-	Model	1	0–5
1999	788	28	0	-	Model	1	0–5
2000	447	52	0	-	Model	0	0–3
2001	672	60	0	-	Model	0	0–2
2002	980	28	1	0.4	Model	4	1–11
2003	599	43	0	-	Model	1	0–3
2004	690	34	1	0.4	Model	3	1–9
2005	726	37	2	0.7	Model	5	2–12
2006	521	28	3	2.1	Model	10	3–21
2007	544	32	6	3.5	Model	15	6–29
2008	557	41	2	0.9	Model	8	5–14
2009	627	20	0	-	Model	1	0–7
2010	550	43	11	4.7	Model	24	15–36
2011	815	40	6	1.8	Model	15	8–25
<i>Stewart-Snares (mainly squid)</i>							
1995-96	3432	8	0	-	Ratio	3	0–7
1996-97	5066	10	0	-	Ratio	4	0–9
1997-98	5769	10	0	-	Ratio	5	1–10
1998-99	7582	16	0	-	Ratio	6	1–13
1999-00	5257	23	3	0.3	Ratio	7	3–12
2000-01	5661	43	3	0.1	Ratio	6	3–10
2001-02	5124	18	1	0.1	Ratio	5	1–10
2002-03	4345	16	0	-	Ratio	3	0–8
2003-04	5097	21	1	0.1	Ratio	5	1–10
2004-05	6232	24	3	0.2	Ratio	7	4–13
2005-06	4963	19	1	0.1	Ratio	5	1–10
2006-07	3498	24	1	0.1	Ratio	4	1–7
2007-08	3249	36	1	0.1	Ratio	3	1–7
2008-09	2547	31	0	-	Ratio	2	0–5
2009-10	2784	43	1	0.1	Ratio	3	1–6
2010-11	2456	36	0	-	Ratio	1	0–4

### 3.4.2. Managing fisheries interactions

For NZ sea lions, efforts to mitigate fisheries bycatch have focused on the SQU6T fishery. Spatial and/or temporal closures have been put in place, SLEDs were developed by industry, codes of practice were introduced, and mortality limits imposed. In 1982 the Minister of Fisheries established a 12 nautical mile exclusion zone around the Auckland Islands from which all fishing activities were excluded (Wilkinson *et al.* 2003). In 1995, the exclusion zone was replaced with a Marine Mammal Sanctuary with the same controls on fishing (Chilvers 2008). The area was subsequently also designated as a Marine Reserve in 2003. In addition to these area-based measures, mitigation devices in the form of SLEDs were introduced in the SQU6T fishing fleet in 2001/02 (Figure 3.2), with widespread and standardised use by all the fleet since 2004/05. The use of SLEDs is not mandatory, but is required by the current industry body (the Deepwater Group), fleet wide in application and monitored by MPI observers. In 1992, the Ministry adopted a fisheries-related mortality limit (FRML; previously referred to as a maximum allowable level of fisheries-related mortality or MALFiRM) to set an upper limit on the number of NZ sea lions that could be incidentally drowned each year in the

SQU6T trawl fishery (Chilvers 2008). If this limit is reached, the fishery may be mandatorily closed for the remainder of the season. This has happened seven times (1996 to 1998, 2000, and 2002 to 2004) since this plan was first adopted in 1993 (Table 3.5; Robertson and Chilvers 2011).

### Sea Lion Exclusion Device - SLED



**Figure 3.2: Diagram of a NZ sea lion exclusion device (SLED) inside a trawl net. Image courtesy of the Deepwater Group.**

Before the widespread use of SLEDs, NZ sea lions incidentally caught during fishing were usually retained in trawl nets and hauled on board, allowing observers to gain an accurate assessment of the number of NZ sea lions being captured on observed tows in a given fishery. This enabled a relatively simple estimation of the total number of NZ sea lions killed. However, following the introduction of SLEDs, the number of NZ sea lions interacting with SLEDs and the proportion of those surviving are much more difficult to estimate. Since the introduction of SLEDs, therefore, it has become necessary to estimate the number of NZ sea lions interacting with trawls using a predetermined strike rate to monitor performance against any bycatch limits set. Using a predetermined strike rate enables the FRML to be converted into a number of tows for management purposes. The rate of 5.65% assumed by MPI for the SQU6T fishery is based on rates observed on vessels without SLEDs from 2003/04 to 2005/06 and is also assumed as part of the fishery implementation within an integrated management procedure evaluation model (named the BFG model after its authors, see section 3.3.3). A strike rate of 5.89 will be assumed for the 2012-13 season, reflecting a slight increase in the long-term average. The most recent strike rates are given in Table 3.4 (Thompson *et al.* 2012).

The current management regime for the SQU6T fishery provides for a “discounted” strike rate to apply to all tows when an approved SLED is used (because SLEDs allow some NZ sea lions to escape and survive their encounters with trawl nets; Thompson and Abraham 2010, see Table 3.5). The SLED discount rate is a fisheries management setting and should not be confused with the actual survival of NZ sea lions that encounter a trawl equipped with a SLED, but the discount mechanism is

duplicated in the BFG simulations. The current discount rate of 82% means that the strike rate is reduced from 5.89% to 1.06% so that, for every 100 tows using an approved SLED, 1.06 NZ sea lions are presumed killed. Ideally, the discount rate would be equal to the survival rate of NZ sea lions that encounter a trawl in circumstances that would be fatal if no SLED were fitted. This survival rate is the product of the proportion of animals that exit a trawl with a SLED and their post-exit survival.

**Table 3.5: Maximum allowable level of fisheries-related mortality (MALFiRM) or fisheries-related mortality limit (FRML) from 1991 to 2013. Note, however, that direct comparisons among years of the limits in Table 3.5 are not possible because the assumptions underlying the MALFiRM or FRML changed over time.**

Year	MALFiRM or FRML	Discount rate	Management actions
1991/92	16 (female only)		
1992/93	63		
1993/94	63		
1994/95	69		
1995/96	73		Fishery closed by MFish (4 May)
1996/97	79		Fishery closed by MFish (28 March)
1997/98	63		Fishery closed by MFish (27 March)
1998/99	64		
1999/00	65		Fishery closed by MFish (8 March)
2000/01	75		Voluntary withdrawal by industry
2001/02	79		Fishery closed by MFish (13 April)
2002/03	70		Fishery closed by MFish (29 March), overturned by High Court
2003/04	62 (124)	20%	Fishery closed by MFish (22 March), overturned by High Court FRML increased
2004/05	115	20%	Voluntary withdrawal by industry on reaching the FRML
2005/06	97 (150)	20%	FRML increased in mid-March due to abundance of squid
2006/07	93	20%	
2007/08	81	35%	
2008/09	113 (95)	35%	Lower interim limit agreed due to the decrease in pup numbers
2009/10	76	35%	
2010/11	68	35%	
2011/12	68	35%	
2012/13	68	82%	

In 2004, the Minister of Fisheries requested that the squid fishery industry organisation (Squid Fishery Management Company), government agencies and other stakeholders with an interest in sea lion conservation work collaboratively to develop a plan of action to determine SLED efficacy. In response, an independently chaired working group (the SLED Working Group) was established to develop an action plan to determine the efficacy of SLEDs, with a particular focus on the survivability of NZ sea lions that exit the nets via the exit hole in the SLED. The group undertook a number of initiatives, most notably the standardisation of SLED specifications (including grid spacing) across the fleet (Clement and Associates Ltd. 2007) and the establishment of an underwater video monitoring programme to help understand what happens when a NZ sea lion exits a SLED. White light and infra-red illuminators were tested. Sea lions were observed outside the net on a number of occasions, but only one fur seal and one NZ sea lion were observed exiting the net via the SLED (on tows when white light illumination was used). The footage contributed to understanding of SLED performance, but established that video monitoring was only suitable for tows using mid water gear, as the camera view was often obscured on tows where bottom gear was used. The SLED Working Group was disbanded in early 2010.

The original “MALFiRM” was calculated using the potential biological removal approach (PBR; Wade 1998) and was used from 1992/93 to 2003/04 (Smith and Baird 2007a). Since 2003/04 the FRML has been translated into a maximum permitted number of tows after which the SQU6T fishing

season may be halted by the Minister regardless of the observed NZ sea lion mortality. This approach has been taken because NZ sea lion mortality can no longer be monitored directly since the introduction of SLEDs.

### 3.4.3. Modelling population-level impacts of fisheries interactions

The population-level impact of fisheries interactions has been assessed for the Auckland Islands via a management procedure evaluation model for the SQU6T fishery (see below). The impact of fisheries interactions for all NZ sea lion populations (and other marine mammal populations) will be assessed as part of the marine mammal risk assessment project (PRO2012-02). The goal of this project is to assess the risk posed to marine mammal populations from New Zealand fisheries by applying a similar approach to the recent seabird risk assessment (Richard *et al.* 2011). In this approach, risk is defined as the ratio of total estimated annual fatalities due to bycatch in fisheries, to the level of PBR (Wade 1998). The results of this project should be available in 2014.

Since 2000, an integrated Bayesian management procedure evaluation model having both population and fishery components has been used to assess the likely performance of a variety of management control rules, each of which can be used to determine the FRML for a given SQU6T season (Breen *et al.* 2003, Breen and Kim 2006a, Breen and Kim 2006b, and Breen, Fu and Gilbert 2010). The model underwent several iterations. An early version, developed in 2000/01, was a relatively simple deterministic, partially age-structured population model with density-dependence applied to pup production (Breen *et al.* 2003). An updated version called the Breen-Kim model was built in 2003 to render it fully age-structured and to incorporate various datasets supplied by DOC (Breen and Kim 2006a, 2006b). This model was further revised in 2007/08 to incorporate the latest NZ sea lion population data and to address various model uncertainties and called the BFG model (after its authors, Breen, Fu and Gilbert 2010). In 2009, the model was again updated to incorporate the low NZ sea lion pup counts observed in 2008/09 (and thus better reflect the observed variability in pup survival and pupping rates), as well as NZ sea lion bycatch that occurs in fisheries other than SQU6T. The BFG model was re-run in 2011 using the same underlying data and structure as in 2009 to evaluate the effect of different model assumptions about the survival of NZ sea lions that exit trawl nets via SLEDs (see below). Additional details on the NZ sea lion population model can be found in Breen *et al.* (2010).

The BFG model incorporates various population dynamics observations (tag re-sighting observations, pup births and mortality, age at maturity) as well as bycatch counts and catch-at-age data from the SQU6T trawl fishery. The model was projected into the future by applying the observed dynamics and a virtual fishery model that is managed in roughly the same way as the real SQU6T fishery. A large number of projections were run and used to assess the likely performance of a wide range of different management control rules against the four performance criteria described in Context (two MFish criteria and two DOC criteria). For each set of runs the population indicators were summarised and the rules compared in tables. The BFG model is sensitive to several key parameters (see Sources of uncertainty, below) and is scheduled to be reviewed in 2013.

SLEDs are effective in allowing most NZ sea lions to exit a trawl but some are retained and drowned and others may not survive the encounter. An experimental approach to assessing non-retained fatality rate involved intentionally capturing animals as they exited the escape hole of a SLED between 1999/2000 and 2002/03. Cover nets were added over the escape holes of some SLEDs and sea lions were restrained in these nets after they exited the SLED proper. An underwater video camera was deployed in 2001 to assess the behaviour and the likelihood of post-exit survival of those animals that were retained in the cover nets (Wilkinson *et al.* 2003, Mattlin 2004). The low number of captures filmed and the inability to assess longer term survival meant that this approach could not be used to determine likely survival rates (e.g., Roe 2010).

Necropsies were conducted on animals recovered from the cover net trials and on those incidentally caught and recovered from vessels operating in the SQU6T, SQU1T and SBW6I fisheries. Although all of the NZ sea lions returned for necropsy died as a result of drowning rather than physical trauma (from interactions with the trawl gear including the SLED grid; Roe and Meynier 2010, Roe 2010), necropsies were designed to assess the nature and severity of trauma sustained during capture and to infer the survival prognosis had those animals been able to exit the net (Mattlin 2004). However, problems associated with this approach limited the usefulness of the results. For example, NZ sea lions were frozen on vessels and stored for periods of up to several months before being thawed for 3–5 days to allow necropsy. Roe and Meynier (2010) concluded that this freeze-thaw process created artefactual lesions that mimic trauma but, particularly in the case of brain trauma, could also obscure real lesions. Further, two reviews in 2011 concluded that the lesions in retained animals may not be representative of the injuries sustained by animals that exit a trawl via a SLED (Roe and Meynier 2010, Roe 2010). As a result of these reviews, the use of necropsies to infer the survival of sea lions interacting with SLEDs was discontinued.

Notwithstanding the limitations of the necropsy data in assessing trauma for previously frozen animals, it was possible to determine that none of the necropsied animals sustained sufficient injuries to the body (excluding the head) to compromise survival (Roe and Meynier 2010, Roe 2010). Head trauma, most likely due to impacts with the SLED grid, could not be ruled out as a potential contributing factor (Roe and Meynier 2010, Roe 2010). In order to quantify the likelihood of a NZ sea lion experiencing physical trauma sufficient to render the animal insensible (and therefore likely to drown) after a collision with a SLED grid, a number of factors need to be assessed. These include the likelihood of a head-first impact, the speed of impact, the angle of impact relative to individual grid bars and relative to the grid plane, the location of impact on the grid, head mass, and the risk of brain injury for a given impact speed and head mass. The effect of multiple impacts also needs to be considered. Estimates for each of these factors were derived from a number of sources, including necropsies (for head mass), video footage of Australian fur seals interacting with Seal Exclusion Devices (SEDs) (for impact speed, location and body orientation) and biomechanical modelling of impacts on the SLED grid (for the risk of brain injury).

In the absence of sufficient video footage of NZ sea lion interacting with SLEDs, footage of fur seals (thought to be Australian fur seals) interacting with SEDs in the Tasmanian small pelagic mid-water trawl fishery has been used (Lyle 2011). The SEDs are similar, but not identical, to the New Zealand SLEDs in that both have sloping steel grids to separate the catch from pinnipeds and guide the latter toward an escape hole in the trawl. The angle of slope and the number of sections in the steel grids are variable (either two or three sections, depending on the vessel). Lyle and Willcox (2008) conducted a camera trial between January 2006 and February 2007 to assess the efficacy of the SED and documented 457 interactions for about 170 individual fur seals. Lyle (2011) reanalysed the footage to estimate impact speed, impact location across the SED grid and body orientation at the time of impact. The situation faced by NZ sea lions in a squid trawl is not identical to that faced by the fur seals studied by Lyle and co-workers, but these are closely related otariids of similar size and, in the absence of specific data, Australian fur seals are considered a reasonable proxy to estimate impact speed, impact location and body orientation.

The risk of brain injury was assessed by biomechanical testing and modelling. Tests using an artificial “head form” (as used in vehicular “crash test” studies) were used to assess the likelihood of brain injury to NZ sea lions colliding with a SLED grid (Ponte *et al.* 2010, 2011). In an initial trial (Ponte *et al.* 2010), the head form (weighing 4.8 kg) was launched at three locations on the SLED grid at a speed of  $10 \text{ m.s}^{-1}$  (about 20 knots). This was considered a “worst feasible case” collision representing the combined velocities of a sea lion swimming with a burst speed of  $8 \text{ m.s}^{-1}$  (after Ray 1963, Fish 2008) and a net being towed at  $2 \text{ m.s}^{-1}$  (about 4 knots). A head injury criterion (HIC, a predictor of the risk of brain injury) was calculated based on criteria validated against human-vehicle impact studies and translated into the probability of mild traumatic brain injury (MTBI) for a given collision, taking into account differences between human and sea lion head and brain masses. MTBI is assumed to have the potential to lead to insensibility or disorientation and subsequent death through drowning for

a NZ sea lion experiencing such an injury at depth. Ponte *et al.* (2010) calculated that a collision at the stiffest part of the SLED grid at this highest feasible speed had a very high risk of MTBI, especially for smaller sea lions. This provides an upper bound for the assessment of risk but Ponte *et al.* (2010) also imputed risk at speeds below the maximum.

In a follow-up study, after a research advisory group meeting with other experts, Ponte *et al.* (2011) tested a wider variety of impact locations on the grid and various angles of impact relative to the bars and to the plane of the grid and combined these to produce a HIC “map” for a SLED grid. This HIC map can be used to estimate the risk of MTBI for a collision by a sea lion at any given speed, location, and orientation.

The data collected from the footage of Australian fur seal SED interactions (Lyle 2011) and the biomechanical modelling (Ponte *et al.* 2010, 2011) were combined in a simulation-based probabilistic model to estimate the risk of a sea lion suffering a mild traumatic brain injury when striking a SLED grid (Abraham 2011). The simulation involved selecting an impact location on the SLED grid (from the fur seal data), selecting a head mass (from NZ sea lion necropsy data) and an impact speed (from the fur seal data), calculating the head impact criterion (HIC) (from the HIC map), scaling the HIC to the head mass and impact speed and calculating the expected probability of mild traumatic brain injury, MTBI. Both 45° and 90° degree impacts were considered, with the former, reflecting the angle of a grid when deployed, adopted as the base case. The head masses used may be at the lower end of the range of head masses for NZ sea lions. Impact speeds were drawn from the distribution of speeds observed for fur seals colliding with SEDs (2–6 m.s<sup>-1</sup>) and these are broadly consistent with the combined tow speed and observed swimming speeds of NZ sea lions in the wild (Crocker *et al.* 2001). Different scaling of HIC values was assessed to gauge sensitivity.

For the base case, the simulation results indicated there was a 3.3% chance of a single head-first collision resulting in MTBI with a 95 percentile of 15.7% risk of MTBI (Abraham 2011). Sensitivities modulating single parameters resulted in up to 6.2% probability of a single collision resulting in MTBI. One sensitivity trial involving changes in multiple parameters resulted in a 10.9% probability of MTBI. This scenario considered impact speeds 20% above those measured for fur seals, multiple collisions with the grid, and the least favourable values of scaling exponents used in scaling the test HIC values and calculating MTBI from the HIC (Abraham 2011). These results are probabilities of MTBI resulting from a single head first collision but, because each individual can have multiple interactions with the grid while in a trawl, and some of these will not be head-first, some additional assumptions were made based on the Australian observations. Using these data, Abraham (2011) estimated the number of head-first collisions per interaction as 0.74, leading to an estimated probability of MTBI for a NZ sea lion interacting with a trawl of 2.7%. Single parameter sensitivity runs increased this to up to 4.6% and the multiple parameter sensitivity using the scenario described above increased it to 8.2% (Abraham 2011). Assuming synergistic interaction between successive head-first strikes (each collision carrying 5 times more risk than previous ones) did not appreciably increase the overall risk because few fur seals had multiple head-first collisions. These results indicate that the risk of mortality for NZ sea lions interacting with the SLED grid is probably low, although some remaining areas of uncertainty were identified (see below).

#### **3.4.4. Sources of uncertainty**

There are several outstanding sources of uncertainty in modelling the effects of fisheries interactions on NZ sea lions at the Auckland Islands, including uncertainty relating to the Bayesian management procedure evaluation model (the BFG model, Breen *et al.* 2010), uncertainty in the modelling of stike rate (Thompson *et al.* 2011) and uncertainty relating to the biomechanical modelling (Ponte *et al.* 2010, 2011, Abraham 2011, Lyle 2011).

The BFG model is sensitive to several key parameters. Some relate mostly to uncertainty about the productivity of the NZ sea lion population (including maximum population growth rate, abundance relative to carrying capacity, maximum rate of pup production, and density dependence), whereas others relate to how the fishery works and is managed (including strike rates and the survival of NZ sea lions that interact with SLEDs but are not retained in the net). Conclusions drawn from the BFG model results are sensitive to prior assumptions about how fast this NZ sea lion population is able to grow. The maximum population growth rate ( $\lambda$ ) for this population of NZ sea lions is not known. Fitting the model to the observed data with an uninformative prior led to an estimated maximum rate of less than 1% per year, potentially as a consequence of attempting to estimate  $\lambda$  for a declining population. This is a very low maximum growth rate for a pinniped (some suggest a default value of 12% per year, Wade 1998), so a prior of 8% was applied to the base model. In a sensitivity run, the model was fitted using a prior of 5% per year, and the results were more consistent with the observed data than when 8% was used.

The estimated abundance of NZ sea lions relative to the carrying capacity of mature individuals at the Auckland Islands (K) is another source of uncertainty. When the model is run in the absence of fishing, the median numbers of mature animals after 100 years was only 94.4% of K as estimated from the model. Although the population is not presently near K, over this timescale, the population would normally be expected to approach K. This is thought to be an artefact of the parameterisation of survival rates in the model, which renders the model conservative when assessing performance against K (Breen *et al.* 2010).

The density dependent response for this population of NZ sea lions is largely unknown, although there is presently no evidence of a density dependent response in life-history traits such as pup mass, pup survival or female fecundity (Chilvers 2012b). Ecological principles suggest that, as numbers in a population decline, individuals compete less with one another for resources. Less competition may result in NZ sea lions growing faster as well as having lower mortality rates and higher rates of pup production and survival. The effect of this type of response is that populations tend to recover from events that reduce their numbers, and populations with strong density dependence recover more strongly than those with weak density dependence. In the BFG model, the shape of the density dependent response was “hard wired” in the model and assumed to occur entirely in the mortality rate of pups. The strength of this response is unknown, and there was no information to support a strong preference for any of the assumed values used in sensitivity runs. This means the base model results may be either conservative or optimistic.

The maximum rate of pup production for this population is not known but can be estimated in the population model. Other modelling conducted for DOC (albeit using different assumptions, Breen *et al.* 2010) suggests that the maximum rate of pup production is <0.28 pups per mature adult per year (Gilbert and Chilvers 2008), a level thought to be below that required to replace the population (Breen *et al.* 2010). When this value is fixed in the BFG model, the fitting procedure does not converge successfully. The BFG model authors progressively increased the fixed value until overall fitting was successful at 0.315 pups per mature adult per year. Thus, the BFG model estimates, and can accommodate, only maximum rates of pup production that are roughly 15% higher than those estimated by direct modelling.

In addition to sources of uncertainty for inputs in the BFG model, there are other sources of uncertainty relevant to the management of fisheries interactions. For example, the estimated strike rate has varied considerably over time, and the model estimates of strike rates for recent years are very imprecise (Thompson *et al.* 2011, Table 3.4). Although year on year variation in strike rate is unlikely to appreciably affect the conclusions from the simulations, if the long-term average strike rate is higher or lower than that assumed within the fishery component of the simulations, or if the strike rate or catchability has increased since the introduction of SLEDs, then there may be some bias. If NZ sea lion catchability has increased, as a result of the increased average tow duration in the SQU6T fishery since the introduction of SLEDs (Table 3.6), or by some other factor, then this would make the simulations optimistic.

**Table 3.6: Tow duration in the SQU6T fishery (i.e. for trawl fishers targeting SQU in statistical areas 602, 603, 617 and 618). Years are calendar years. Data from MPI databases.**

Year	No. of tows	Mean tow duration (hours)	Percentage of tows		
			Less than 4 hours	Between 4 & 8 hours	More than 8 hours
1995	4 014	3.7	64.2	33.5	2.2
1996	4 474	3.6	64.3	34.2	1.5
1997	3 719	3.8	62.7	33.7	3.7
1998	1 446	3.2	74.4	24.7	0.9
1999	403	3.5	73.0	24.3	2.7
2000	1 213	3.5	70.3	27.0	2.7
2001	583	3.3	72.9	26.6	0.5
2002	1 647	3.8	59.8	38.8	1.4
2003	1 467	4.1	52.4	44.0	3.6
2004	2 598	5.0	36.7	53.6	9.7
2005	2 693	4.7	43.7	48.6	7.7
2006	2 462	6.3	26.0	49.6	24.3
2007	1 317	7.3	18.9	46.3	34.8
2008	1 265	6.2	20.4	58.7	20.9
2009	1 925	6.5	21.1	51.4	27.5
2010	1 190	7.9	16.4	37.4	46.2
2011	1 585	6.8	24.7	42.8	32.4
2012*	1 283	6.6	23.5	49.3	27.3

\* Includes data up to November 30, 2012.



There are a number of possible sources of uncertainty relating to the biomechanical modelling (Ponte *et al.* 2010, 2011, Abraham 2011, Lyle 2011). The use of linear acceleration, as opposed to rotational (angular) acceleration, in the biomechanical modelling may underestimate the risk of MTBI, although this was thought to be accounted for at least in part by sensitivity analysis of the scaling of HIC values. The testing used an artificial “head form” based on human anatomy, so the effect of NZ sea lion scalp thickness and skull morphology is unknown, although differences in head and brain masses are accounted for. Potential effects of differences in the angle of the head on impact (relative to the neck) were not tested. Impact speeds, locations and orientations of NZ sea lions may differ from those of Australian fur seals, although the fur seal data were considered to be a reasonable proxy by a Research Advisory Group. The head mass values used may be lower than average for NZ sea lions; this would mean risk is likely to be overestimated. This approach assesses risk associated with collisions with the grid of a SLED and cannot be used to assess other sources of mortality resulting, for example, from an animal being retained in a net long enough for them to exceed their dive limit before reaching the surface after escaping from either the SLED or the front of the net. Such sources of cryptic mortality have always existed, are presently unquantified and are not reflected in the estimated overall survival rate of encounters with trawls.

### 3.4.5. Potential indirect threats

In addition to sources of uncertainty associated with direct fisheries interactions, there is the possibility that indirect fisheries effects may have population-level consequences for NZ sea lions. Such indirect effects may include competition for food resources between various fisheries and NZ sea lions (Robertson and Chilvers 2011). In order to determine whether resource competition is present and is having a population-level effect on NZ sea lions, research must identify if there are resources in common for NZ sea lions and the various fisheries within the range of NZ sea lions, and if those resources are limiting. Diet studies have demonstrated overlap in the species consumed by NZ sea lions and those caught in fisheries within the range of NZ sea lions, particularly hoki and arrow squid (Cawthorn *et al.* 1985, Childerhouse *et al.* 2001, Meynier *et al.* 2009). A recent study focused on energy and amino acid content of prey determined that the selected prey species contained all

essential amino acids and were of low to medium energy levels (Meynier 2010). This may indicate that the nutritional content of prey species is not limiting the metabolic activity of NZ sea lions, although vitamin and mineral content were not considered. Meynier (2010) also developed a bio-energetic model and used it to estimate the amount of prey consumed by NZ sea lions at 17 871 tonnes (95% CI 17 738–18 000 t) per year. This is equivalent to ~30% of the tonnage of arrow squid, and ~15% of the hoki harvested annually by the fisheries in the Sub-Antarctic between 2000 and 2006 (Meynier 2010). Comparison of the temporal and spatial distributions of sea lion prey, sea lion foraging and of historical fishing extractions may help to identify the mechanisms whereby resource competition might occur (Bowen 2012). The effects of fishing on sea lion prey species are likely to be complicated by food web interactions and multispecies models may help to assess the extent to which resource competition can impact on sea lion populations, such as those currently being developed by NIWA (Project SA123098). In addition, multispecies models may provide a means for simultaneously assessing multiple drivers of sea lion population change (a review of potential causes is given in Robertson & Chilvers 2011) which may be a more effective approach than focussing on single factor explanations for the recent observed decline in NZ sea lions (Bowen 2012).

### 3.5. Indicators and trends


<i>Population size</i>	<ul style="list-style-type: none"> <li>• 12 065 animals (including pups &lt; 1 yr old) at the Auckland Islands (90% CI: 11 160–13 061) in 2009 (most recent model estimate)<sup>10</sup></li> <li>• 1 683 pups at the Auckland Islands (SE = 16) in 2011/12<sup>11</sup></li> <li>• 681–726 pups at Campbell Island in 2010<sup>12</sup></li> <li>• 25 pups tagged at Stewart Island during a DOC recreational hut and track maintenance trip in March 2012</li> <li>• 5 pups at the Otago Peninsula in 2011/12<sup>13</sup></li> </ul>																																																																						
<i>Population trend</i>	<ul style="list-style-type: none"> <li>• Estimated abundance at the Auckland Islands:</li> </ul>  <table border="1"> <caption>Auckland Islands population size (median &amp; 90% c.i.)</caption> <thead> <tr> <th>Year</th> <th>Population Size (approx.)</th> </tr> </thead> <tbody> <tr><td>1995</td><td>15,500</td></tr> <tr><td>1996</td><td>16,000</td></tr> <tr><td>1997</td><td>16,500</td></tr> <tr><td>1998</td><td>17,000</td></tr> <tr><td>1999</td><td>15,500</td></tr> <tr><td>2000</td><td>15,000</td></tr> <tr><td>2001</td><td>14,500</td></tr> <tr><td>2002</td><td>14,000</td></tr> <tr><td>2003</td><td>14,000</td></tr> <tr><td>2004</td><td>13,500</td></tr> <tr><td>2005</td><td>13,000</td></tr> <tr><td>2006</td><td>13,000</td></tr> <tr><td>2007</td><td>13,000</td></tr> <tr><td>2008</td><td>12,500</td></tr> <tr><td>2009</td><td>12,000</td></tr> </tbody> </table> <ul style="list-style-type: none"> <li>• Pup production at the Auckland Islands:</li> </ul>  <table border="1"> <caption>Auckland Islands pup production (mean &amp; std. err.)</caption> <thead> <tr> <th>Year</th> <th>Pup Production (approx.)</th> </tr> </thead> <tbody> <tr><td>1995</td><td>2,500</td></tr> <tr><td>1996</td><td>2,700</td></tr> <tr><td>1997</td><td>2,900</td></tr> <tr><td>1998</td><td>3,000</td></tr> <tr><td>1999</td><td>2,800</td></tr> <tr><td>2000</td><td>2,800</td></tr> <tr><td>2001</td><td>2,800</td></tr> <tr><td>2002</td><td>2,300</td></tr> <tr><td>2003</td><td>2,500</td></tr> <tr><td>2004</td><td>2,500</td></tr> <tr><td>2005</td><td>2,100</td></tr> <tr><td>2006</td><td>2,100</td></tr> <tr><td>2007</td><td>2,200</td></tr> <tr><td>2008</td><td>2,100</td></tr> <tr><td>2009</td><td>1,500</td></tr> <tr><td>2010</td><td>1,800</td></tr> <tr><td>2011</td><td>1,600</td></tr> <tr><td>2012</td><td>1,700</td></tr> </tbody> </table> <ul style="list-style-type: none"> <li>• The population is probably increasing at Campbell Island based on substantial increases in pup counts (although methodology has changed over time).</li> <li>• The population is increasing at the Otago Peninsula through a combination of reproduction and immigration.</li> </ul>	Year	Population Size (approx.)	1995	15,500	1996	16,000	1997	16,500	1998	17,000	1999	15,500	2000	15,000	2001	14,500	2002	14,000	2003	14,000	2004	13,500	2005	13,000	2006	13,000	2007	13,000	2008	12,500	2009	12,000	Year	Pup Production (approx.)	1995	2,500	1996	2,700	1997	2,900	1998	3,000	1999	2,800	2000	2,800	2001	2,800	2002	2,300	2003	2,500	2004	2,500	2005	2,100	2006	2,100	2007	2,200	2008	2,100	2009	1,500	2010	1,800	2011	1,600	2012	1,700
Year	Population Size (approx.)																																																																						
1995	15,500																																																																						
1996	16,000																																																																						
1997	16,500																																																																						
1998	17,000																																																																						
1999	15,500																																																																						
2000	15,000																																																																						
2001	14,500																																																																						
2002	14,000																																																																						
2003	14,000																																																																						
2004	13,500																																																																						
2005	13,000																																																																						
2006	13,000																																																																						
2007	13,000																																																																						
2008	12,500																																																																						
2009	12,000																																																																						
Year	Pup Production (approx.)																																																																						
1995	2,500																																																																						
1996	2,700																																																																						
1997	2,900																																																																						
1998	3,000																																																																						
1999	2,800																																																																						
2000	2,800																																																																						
2001	2,800																																																																						
2002	2,300																																																																						
2003	2,500																																																																						
2004	2,500																																																																						
2005	2,100																																																																						
2006	2,100																																																																						
2007	2,200																																																																						
2008	2,100																																																																						
2009	1,500																																																																						
2010	1,800																																																																						
2011	1,600																																																																						
2012	1,700																																																																						

<sup>10</sup> Breen *et al.* (2010).

<sup>11</sup> Chilvers (2012).

<sup>12</sup> Robertson and Chilvers (2011), Maloney *et al.* (2012).

<sup>13</sup> For more information, see: <http://www.sealiontrust.org.nz/otago-sea-lion-family-tree/>.

Threat status	<ul style="list-style-type: none"><li>NZ: Nationally Critical, Criterion C<sup>14</sup>, Range Restricted<sup>15</sup>, in 2010<sup>16</sup></li><li>IUCN: Vulnerable, A3b<sup>17</sup>, in 2008<sup>18</sup></li></ul>																																																												
Number of interactions <sup>19</sup>	<ul style="list-style-type: none"><li>81 estimated interactions (95% CI: 26-259) in trawl fisheries in 2010-11</li><li>29 estimated captures (95% CI: 17-43) in trawl fisheries in 2010-11</li><li>6 observed captures in trawl fisheries in 2010-11</li></ul>																																																												
Trend in interactions	<p>Trawl fisheries:</p> <div><p>The top chart displays observed captures from 2003 to 2011. The left y-axis represents 'Observed captures' (0 to 25), and the right y-axis represents 'Captures per 100 tows' (0.00 to 0.35). The x-axis shows 'Fishing year' (03 to 11). The legend indicates 'Dead' (dark blue bars), 'Alive' (light blue bars), and 'Rate' (red line). The bottom chart displays estimated captures from 2003 to 2011. The y-axis represents 'Estimated captures' (0 to 80). The x-axis shows 'Fishing year' (03 to 11). The legend indicates 'Estimated captures' (light blue bars with error bars).</p><table><caption>Observed captures (Top Chart)</caption><tr><th>Fishing year</th><th>Dead</th><th>Alive</th><th>Rate</th></tr><tr><td>03</td><td>12</td><td>0</td><td>0.12</td></tr><tr><td>04</td><td>20</td><td>1</td><td>0.21</td></tr><tr><td>05</td><td>14</td><td>0</td><td>0.14</td></tr><tr><td>06</td><td>15</td><td>0</td><td>0.15</td></tr><tr><td>07</td><td>12</td><td>0</td><td>0.12</td></tr><tr><td>08</td><td>10</td><td>0</td><td>0.10</td></tr><tr><td>09</td><td>3</td><td>0</td><td>0.03</td></tr><tr><td>10</td><td>12</td><td>3</td><td>0.12</td></tr><tr><td>11</td><td>5</td><td>0</td><td>0.05</td></tr></table><table><caption>Estimated captures (Bottom Chart)</caption><tr><th>Fishing year</th><th>Estimated captures</th></tr><tr><td>03</td><td>35</td></tr><tr><td>04</td><td>60</td></tr><tr><td>05</td><td>55</td></tr><tr><td>06</td><td>52</td></tr><tr><td>07</td><td>45</td></tr><tr><td>08</td><td>32</td></tr><tr><td>09</td><td>22</td></tr><tr><td>10</td><td>45</td></tr><tr><td>11</td><td>30</td></tr></table></div>	Fishing year	Dead	Alive	Rate	03	12	0	0.12	04	20	1	0.21	05	14	0	0.14	06	15	0	0.15	07	12	0	0.12	08	10	0	0.10	09	3	0	0.03	10	12	3	0.12	11	5	0	0.05	Fishing year	Estimated captures	03	35	04	60	05	55	06	52	07	45	08	32	09	22	10	45	11	30
Fishing year	Dead	Alive	Rate																																																										
03	12	0	0.12																																																										
04	20	1	0.21																																																										
05	14	0	0.14																																																										
06	15	0	0.15																																																										
07	12	0	0.12																																																										
08	10	0	0.10																																																										
09	3	0	0.03																																																										
10	12	3	0.12																																																										
11	5	0	0.05																																																										
Fishing year	Estimated captures																																																												
03	35																																																												
04	60																																																												
05	55																																																												
06	52																																																												
07	45																																																												
08	32																																																												
09	22																																																												
10	45																																																												
11	30																																																												

<sup>14</sup> A taxon is listed as 'Nationally Critical' under criterion C if the population (irrespective of size or number of sub-populations) has a very high (rate of) ongoing or predicted decline; greater than 70% over 10 years or three generations, whichever is longer (Townsend *et al.* 2008).

<sup>15</sup> A taxon is listed as 'Range Restricted' if it is confined to specific substrates, habitats or geographic areas of less than 1000 km<sup>2</sup> (100 000 ha); this is assessed by taking into account the area of occupied habitat of all sub-populations (Townsend *et al.* 2008).

<sup>16</sup> Baker *et al.* (2010).

<sup>17</sup> A taxon is listed as 'Vulnerable' if it is considered to be facing a high risk of extinction in the wild. A3b refers to a reduction in population size (A), based on a reduction of  $\geq 30\%$  over the last 10 years or three generations (whichever is longer up to a maximum of 100 years (3); and when considering an index of abundance that is appropriate to the taxon (b; IUCN 2010).

<sup>18</sup> Gales (2008).

<sup>19</sup> For more information, see: <http://data.dragonfly.co.nz/psc/>.

### 3.6. References

- Abraham ER (2011). Probability of Mild Traumatic Brain Injury for sea lions interacting with SLEDs. Final Research Report for Ministry of Fisheries project SRP2011-03 (Unpublished report held by the Ministry of Fisheries, Wellington). 21 pages.
- Abraham ER; Thompson FN (2011). Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2008–09. Final Research Report prepared for Ministry of Fisheries project PRO2007/01. (Unpublished report held by the Ministry of Fisheries, Wellington.) 170 p.
- Anonymous (2009). DWG and MFish SLED Specification for SQU 6T 2010 Operation Plan (MK 3/13). Report presented to SLED WG September 2009, Wellington, NZ.
- Augé AA (2006). Terrestrial spatial ecology of female New Zealand sea lions. Unpublished MSc thesis, University of Otago, Dunedin.
- Augé AA; Lallas C; Chilvers BL; Davis LS (2011a). Autumn diet of recolonising female New Zealand sea lions based at Otago Peninsula, South Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 46: 97–110.
- Augé AA; Chilvers BL; Moore AB; Davis LS (2011b). Foraging behaviour indicates marginal habitat for New Zealand sea lions: remnant vs recolonising. *Marine Ecology Progress Series* 432:247–256.
- Augé AA; Chilvers BL; Davis LS; Moore AB (2011c). In the shallow end: diving behaviour of recolonising female New Zealand sea lions (*Phocarctos hookeri*) around the Otago Peninsula. *Canadian Journal of Zoology* 89:1195–1205.
- Baker CS; Chilvers BL; Constantine R; DuFresne S; Mattlin RH; van Helden A; Hitchmough R (2010). Conservation status of New Zealand Marine Mammals (suborders Cetacea and Pinnipedia), 2009. *New Zealand Journal of Marine and Freshwater Research* 44: 101–115.
- Baker AJ (1999). Unusual mortality of the New Zealand sea lion, *Phocarctos hookeri*, Auckland Islands, January–February 1998: A report of a workshop held 8–9 June 1998, Wellington, and a contingency plan for future events. pp. 29–33, Department of Conservation, Te Papa Atawhai, Wellington, New Zealand.
- Breen PA (2008). Sea lion data for use in the population model for Project IPA200609. Unpublished Final Research Report for Ministry of Fisheries, 30 March 2008. 18 pp. plus Confidential Appendix of 6p.
- Breen PA; Hilborn R; Maunders MN; Kim SW (2003). Effects of alternative control rules on the conflict between a fishery and a threatened sea lion (*Phocarctos hookeri*). *Canadian Journal of Fisheries and Aquatic Sciences* 60: 527–541.
- Breen PA; Kim SW (2006a). Exploring alternative management procedures for controlling bycatch of Hooker's sea lions in the SQU 6T squid fishery. Unpublished Final Research Report for Ministry of Fisheries Project M0F2002/03L, Objective 3. Revision 5, 3 February 2006. 88 pp.
- Breen PA; Kim SW (2006b). An integrated Bayesian evaluation of Hooker's sea lion bycatch limits. pp. 471–493 In: A. Trites, S. Atkinson, D. DeMaster, L. Fritz, T. Gelatt, L. Rea, and K. Wynne. (eds.). Sea lions of the world. Alaska Sea Grant College Program, University of Alaska Fairbanks.
- Breen PA; Fu D; Gilbert DJ (2010). Sea lion population modelling and management procedure evaluations: Report for Project SAP2008/14, Objective 2. Presented to AEWG March 22 2010, Wellington, New Zealand.
- Bowen WD (2012). A review of evidence for indirect effects of commercial fishing on New Zealand sea lions (*Phocarctos hookeri*) breeding on the Auckland Islands. Final Report May 2012, Contract Number: POP 2010/01 (Objectives 4 and 5). Department of Conservation, Wellington. 41 p.
- Castinel A; Duignan PJ; Pomroy WE; López-Villalobos N; Gibbs NJ; Chilvers BL; Wilkinson IS (2007). Neonatal mortality in New Zealand sea lions (*Phocarctos hookeri*) at Sandy Bay, Enderby Island, Auckland Islands from 1998 to 2005. *Journal of Wildlife Diseases* 43: 461–474.
- Cawthorn MW (1993). Census and population estimation of Hooker's sea lion at the Auckland Islands, December 1992–February 1993. *DOC Technical Series 2*. Wellington, Department of Conservation. 34 p
- Cawthorn MW; Crawley MC; Mattlin RH; Wilson GJ (1985). Research on pinnipeds in New Zealand. *Wildlife Research Liaison Group Report No. 7*, Wellington, New Zealand.
- Childerhouse, S., and N. Gales. (1998). The historic distribution and abundance of the New Zealand sea lion *Phocarctos hookeri*. *New Zealand Journal of Zoology* 25: 1–16.
- Childerhouse SJ; Gales NJ (2001). Fostering behavior in New Zealand sea lions *Phocarctos hookeri*. *New Zealand Journal of Zoology* 28:189–195.
- Childerhouse SJ; Dix B; Gales NJ (2001). Diet of New Zealand sea lions (*Phocarctos hookeri*) at the Auckland Islands. *Wildlife Research* 28:291–298.
- Childerhouse SJ; Gibbs N; McAlister G; McConkey S; McConnell H; McNally N; Sutherland D (2005). Distribution, abundance and growth of New Zealand sea lion *Phocarctos hookeri* pups on Campbell Island. *New Zealand Journal of Marine and Freshwater Research* 39: 889–898.
- Childerhouse SJ; Dawson SM; Slooten E; Fletcher DJ; Wilkinson IS (2010a). Age distribution of lactating New Zealand sea lions: interannual and intersite variation. *Marine Mammal Science* 26: 123–139.
- Childerhouse SJ; Dawson SM; Fletcher DJ; Slooten E; Chilvers BL (2010b). Growth and reproduction of female New Zealand sea lions. *Journal of Mammalogy* 91:165–176.
- Chilvers BL (2008). New Zealand sea lions *Phocarctos hookeri* and squid trawl fisheries: bycatch problems and management options. *Endangered Species Research* 5: 193–204.
- Chilvers BL (2009). Foraging locations of a decreasing colony of New Zealand sea lions (*Phocarctos hookeri*). *New Zealand Journal of Ecology* 33:106–113.
- Chilvers BL (2012a). Research to assess the demographic parameters of New Zealand sea lions, Auckland Islands: Final Research Report November 2012, Contract Number: POP 2011/01. Department of Conservation, Wellington. 11 p.
- Chilvers BL (2012b). Using life-history traits of New Zealand sea lions, Auckland Islands to clarify potential causes of decline. *Journal of Zoology* 287: 240–249.
- Chilvers BL; Wilkinson IS (2008). Philopatry and site fidelity of New Zealand sea lions (*Phocarctos hookeri*). *Wildlife Research* 35: 463–470.
- Chilvers BL; Wilkinson IS (2009). Diverse foraging strategies in lactating New Zealand sea lions. *Marine Ecology Progress Series* 378: 299–308.
- Chilvers BL; Wilkinson IS (2011). Research to assess the demographic parameters of New Zealand sea lions, Auckland Islands: Draft Final Report November 2011, Contract Number: POP 2010/01. Department of Conservation, Wellington. 20 p.
- Chilvers BL; Wilkinson IS; Duignan PJ; Gemmell NJ (2005a). Summer foraging areas for lactating New Zealand sea lions *Phocarctos hookeri*. *Marine Ecology Progress Series* 304:235–247.

- Chilvers BL; Robertson BC; Wilkinson IS; Duignan PJ; Gemmell NJ (2005b). Male harassment of female New Zealand sea lions: mortality, injury and harassment avoidance. *Canadian Journal of Zoology* 83:642-648.
- Chilvers BL; Wilkinson IS; Duignan PJ; Gemmell NJ (2006). Diving to extremes: are New Zealand sea lions pushing their limits in a marginal habitat? *Journal of Zoology* 269:233-240.
- Chilvers BL; Wilkinson IS; Childerhouse SJ (2007). New Zealand sea lion, *Phocarctos hookeri*, pup production-1995 to 2006. *New Zealand Journal of Marine and Freshwater Research* 41: 205-213.
- Chilvers BL; Wilkinson IS; Mackenzie DI (2010). Predicting life-history traits for female New Zealand sea lions, *Phocarctos hookeri*: integrating short-term mark-recapture data and population modelling. *Journal of Agricultural, Biological, and Environmental Statistics* 15: 259-278.
- Chilvers BL; Amey JM; Huckstadt LA; Costa DP (2011). Partitioning of breeding area foraging locations in New Zealand sea lions. *Polar Biology* 34:565-574.
- Clement & Associates Ltd. (2007). Squid trawl fleet sea lion escape device audit. A report commissioned by the Department of Conservation for Project CSP MIT2004/05. Department of Conservation, Wellington. 19 p.
- Costa DP; Gales NJ (2000). Foraging energetic and diving behaviour of lactating New Zealand sea lions, *Phocarctos hookeri*. *Journal of Experimental Biology* 203:3655-3665.
- Crocker DE; Gales NJ; Costa DP (2001). Swimming speed and foraging strategies of New Zealand sea lions, *Phocarctos hookeri*. *Journal of Zoology* 254: 267-277.
- Department of Conservation. (2007). Draft Population Management Plan for New Zealand Sea Lion. Draft Document for Public Consultation, August 2007. Department of Conservation, Wellington. <http://www.doc.govt.nz/getting-involved/consultations/closed/nz-sea-lion-management/>.
- Department of Conservation. (2009). New Zealand sea lion species management plan: 2009-2014. Department of Conservation, Wellington. 31 p.
- Dickie G (1999). Population dynamics of New Zealand fur seals (*Arctocephalus forsteri*) and New Zealand sea lions (*Phocarctos hookeri*). M.Sc. thesis. University of Otago, Dunedin, New Zealand. 117p.
- Fish F (2008). Streamlining. pp. 1123-26. In W. F. Perrin, B. Wursig, and J. G. M. Thewissen (eds.). *Encyclopedia of Marine Mammals*. Academic Press.
- Gales NJ (1995). New Zealand (Hooker's) sea lion recovery plan. Threatened Species Recovery Plan Series 17. Department of Conservation, Wellington.
- Gales NJ (2008). *Phocarctos hookeri*. In: IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4. [www.iucnredlist.org](http://www.iucnredlist.org). Downloaded on 1 June 2011.
- Gales NJ; Fletcher DJ (1999). Abundance, distribution and status of the New Zealand sea lion, *Phocarctos hookeri*. *Wildlife Research* 26: 35-52.
- Gales NJ; Mattlin RH (1997). Summer diving behaviour of lactating female New Zealand sea lions. *Canadian Journal of Zoology* 75:1695-1706.
- Gilbert DJ; Chilvers BL (2008). Final report on New Zealand sea lion pupping rate. POP2006-01 Objective 3. Analysis from sea lion database to estimate pupping rate and associated parameters. NIWA Report WLG2008-77. 26p.
- Gill BJ (1998). Prehistoric breeding sites of New Zealand sea lions (*Phocarctos hookeri*, Carnivora: Otariidae) at North Cape. *Records of the Auckland Museum* 35: 55-64.
- Hanke W; Witte M; Miersch L; Brede M; Oeffner J; Michael M; Hanke F; Leder A; Dehnhardt G (2010). Harbor seal vibrissa morphology suppresses vortex-induced vibrations. *Journal of Experimental Biology* 213: 2665-2672.
- IUCN, 2010. IUCN Red List of Threatened Species. Version 2010.4. <http://www.iucnredlist.org>. Downloaded on 1 June 2011.
- Katsanevakis S (2008). Marine debris, a growing problem: Sources, distribution, composition, and impacts. In: Hofer TN (ed) *Marine Pollution: New Research*. Nova Science Publishers, New York. pp. 53-100.
- Lalas C; Bradshaw CJA (2003). Expectations for population growth at new breeding locations for the vulnerable New Zealand sea lion using a simulation model. *Biological Conservation* 114:67-78.
- Leung ES; Chilvers BL; Moore AB; Nakagawa S; Robertson BC (2012). Sexual segregation in juvenile New Zealand sea lion foraging ranges: implications for intraspecific competition, population dynamics and conservation. *PLoS ONE*. 7(9): e45389.
- Lyle JM (2011). SRP2010-03: Fur seal interactions with SED excluder device. Unpublished Final Research Report for Ministry of Fisheries, 26 July 2011. 20 pp.
- Lyle JM; Willcox ST (2008). Dolphin and seal interactions with mid-water trawling in the Commonwealth small pelagic fishery, including an assessment of bycatch mitigation. Final Report Project R05/0996. Australian Fisheries Management Authority, National Library of Australia. 39p.
- MacKenzie DI (2011). Estimation of demographic parameters for New Zealand sea lions breeding on the Auckland Islands. Final Report November 2012, Contract Number: Objective 3: POP 2011/01. Department of Conservation, Wellington. 88 p.
- Maloney A; Chilvers BL; Haley M; Muller CG; Roe WD; Debski I (2009). Distribution, pup production and mortality of New Zealand sea lion (*Phocarctos hookeri*) on Campbell Island / Motu Ihupuku, 2008. *New Zealand Journal of Ecology* 33: 97-105.
- Maloney A; Chilvers BL; Muller CG; Haley M (2012). Increasing pup production of New Zealand sea lions at Campbell Island/Motu Ihupuku: can it continue? *New Zealand Journal of Zoology* 39:19-29.
- Marshall CD (2008). Feeding Morphology. pp. 406-413. In W. F. Perrin, B. Wursig, and J. G. M. Thewissen (eds.). *Encyclopedia of Marine Mammals*. Academic Press.
- Mattlin RH (2004). QMA SQU6T New Zealand sea lion incidental catch and necropsy data for the fishing years 2000-01, 2001-02 and 2002-03. Report prepared for the NZ Ministry of Fisheries, Wellington, NZ. July 2004. 21p.
- Marlow BJ (1975). The comparative behaviour of the Australasian sea lions *Neophoca cinerea* and *Phocarctos hookeri*. *Mammalia* 39:159-230.
- McConkey S; McConnell H; Lalas C; Heinrich S; Ludmerer A; McNally N; Parker E; Borofsky C; Schimanski K; McIntosh M (2002). A northward spread in the breeding distribution of the New Zealand sea lion. *Australian Mammalogy* 24:97-106.
- McNally N; Heinrich S; Childerhouse SJ (2001). Distribution and breeding of New Zealand sea lions *Phocarctos hookeri* on Campbell Island. *New Zealand Journal of Zoology* 28: 79-87.
- Meynier L (2010). New Zealand sea lion bio-energetic modelling: Final Report for Project IPA2009-09. Presented to AEWG May 17 2011, Wellington, New Zealand. 34p.
- Meynier L; Morel PCH; Chilvers BL; Mackenzie DDS; MacGibbon A; Duignan PJ (2008). Temporal and sex differences in the blubber fatty acid profiles of the New Zealand sea lion *Phocarctos hookeri*. *Marine Ecology Progress Series* 366: 271-278.
- Meynier L; Mackenzie DDS; Duignan PJ; Chilvers BL; Morel PCH (2009). Variability in the diet of New Zealand sea lion (*Phocarctos hookeri*) at the Auckland Islands, New Zealand. *Marine Mammal Science* 25: 302-326.

- Ministry of Fisheries. (2011). Report from the Fisheries Assessment Plenary, May 2011: stock assessments and yield estimates. Ministry of Fisheries, Wellington, New Zealand. 1178p.
- Moore JM; Wallace B; Lewison RL; Zydelis R; Cox T; Crowder L (2009). A review of marine mammal, sea turtle and seabird bycatch in USA fisheries and the role of policy in shaping management. *Marine Policy* 33:435–451.
- Moore PJ; Moffatt RD (1990). Research and management projects on Campbell Island 1987–88. Science and Research Internal Report Series 57. Wellington, Department of Conservation. 101 p.
- Nagaoka L (2001). Using Diversity Indices to Measure Changes in Prey Choice at the Shag River Mouth Site, Southern New Zealand. *International Journal of Osteoarchaeology* 11: 101–111.
- Nagaoka L (2006). Prehistoric seal carcass exploitation at the Shag Mouth site, New Zealand. *Journal of Archaeological Science* 33: 1474–1481.
- Ponte G; van den Berg A; Anderson RWG (2010). Impact characteristics of the New Zealand Fisheries sea lion exclusion device stainless steel grid. Final Research Report for Ministry of Fisheries project IPA2009-06, Oct. 2010. 24p.
- Ponte G; van den Berg A; Anderson RWG (2011). Further analysis of the impact characteristics of the New Zealand Fisheries sea lion exclusion device stainless steel grid. Final Research Report for Ministry of Fisheries project SRP2010-05, Sept. 2011. 36 p.
- Ray GC (1963). Locomotion in pinnipeds. *Natural History* 72: 10–21.
- Read AJ; Drinker P; Northridge S (2006). Bycatch of marine mammals in U.S. and global fisheries. *Conservation Biology* 20:163–169.
- Richard Y; Abraham ER; Filippi D (2011). Assessment of the risk to seabird populations from New Zealand commercial fisheries. Final Research Report for research projects IPA2009-19 and IPA2009-20. (Unpublished report held by Ministry of Fisheries, Wellington.). 66 p.
- Riet-Sapirza FG (2007). Milk composition of the New Zealand sea lion and factors that influence it. PhD thesis, Massey University, Palmerston North, New Zealand.
- Riet-Sapirza FG; Lopez-Villalobos N; Mackenzie DDS; Duignan PJ; Macgibbon A; Chilvers BL (2012). Assimilation and deposition of fatty acid biomarkers in blood serum and milk of lactating New Zealand sea lions, *Phocartos hookeri*. *Canadian Journal of Zoology*
- Robertson BC; Chilvers BL (2011). The population decline of the New Zealand sea lion *Phocartos hookeri*: a review of possible causes. *Mammal Review* 41: 253–275.
- Robertson BC; Chilvers BL; Duignan PJ; Wilkinson IS; Gemmel NJ (2006). Dispersal of breeding adult male *Phocartos hookeri*: implications for disease transmission, population management and species recovery. *Biological Conservation* 127:227–236.
- Roe WD (2010). External review of NZ sea lion bycatch necropsy data and methods. Report prepared for the NZ Ministry of Fisheries, Wellington. 8p.
- Roe WD; Meynier L (2010). Review of Necropsy Records for Bycaught NZ sea lions (*Phocartos hookeri*), 2000–2008. Report for NZ Ministry of Fisheries project PRO2008-03, Wellington. 46p.
- Smith IWG (1989). Maori Impact on the Marine Magafauna: Pre-European Distributions of New Zealand Sea Mammals. In Sutton, D. G. (ed.) *Saying so doesn't make it so*. Papers in Honour of B. Foss Leach. *New Zealand Archaeological Association Monograph* 17:76–108.
- Smith IWG (2011). Estimating the magnitude of pre-European Maori marine harvest in two New Zealand study areas. *New Zealand Aquatic Environment and Biodiversity Report* No. 82.
- Smith MH; Baird SJ (2005). Factors that may influence the level of incidental mortality of New Zealand sea lions (*Phocartos hookeri*) in the squid (*Nototodarus* spp.) trawl fishery in SQU 6T. *New Zealand Fisheries Assessment Report* 2005/20. 35 p.
- Smith MH; Baird SJ (2007a). Estimation of incidental captures of New Zealand sea lions (*Phocartos hookeri*) in New Zealand fisheries in 2003–04, with particular reference to the SQU 6T squid trawl fishery. *New Zealand Fisheries Assessment Report* 2007/7. 32 p.
- Smith MH; Baird SJ (2007b). Estimation of incidental captures of New Zealand sea lions (*Phocartos hookeri*) in New Zealand fisheries in 2004–05, with particular reference to the SQU 6T squid trawl fishery. *New Zealand Aquatic Environment and Biodiversity Report* No. 12. 31 p.
- Thompson FN; Abraham ER (2010). Estimation of the capture of New Zealand sea lions (*Phocartos hookeri*) in trawl fisheries, from 1995–96 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report* No. 66, Wellington.
- Thompson FN; Abraham ER; Berkenbusch K (2011). Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2009–10. Draft Final Research Report for Ministry for Primary Industries project PRO2010-01 (Unpublished report held by the Ministry for Primary Industries, Wellington). 80 pages.
- Thompson FN; Abraham ER; Berkenbusch K (2012). Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2010–11. Draft Final Research Report for Ministry for Primary Industries project PRO2010-01 (Unpublished report held by the Ministry for Primary Industries, Wellington). 90 pages.
- Townsend AJ; de Lange PJ; Duffy CAJ; Miskelly CM; Molloy J; Norton D (2008). New Zealand Threat Classification System Manual. Department of Conservation, Wellington, New Zealand.
- Tuck ID (2009). Characterisation of scampi fisheries and the examination of catch at length and spatial distribution of scampi in SCI 1, 2, 3, 4A and 6A. *New Zealand Fisheries Assessment Report* 2009/27:102 pages.
- Wade PR (1998). Calculating limits to the allowable human caused mortality of cetaceans and pinnipeds. *Marine Mammal Science* 14:1–37.
- Wade PR; Angliss R (1997). Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop. *NOAA Technical Memo NMFS-OPR-12*.
- Walker GE; Ling JK (1981). New Zealand sea lion. In Ridway S.H., Harrison R.J. (eds.). *Handbook of Marine Mammal*, Vol. 1. Academic Press Inc., London, United Kingdom.
- Wilkinson IS; Burgess J; Cawthorn MW (2003). New Zealand sea lions and squid—managing fisheries impacts on a threatened marine mammal. pp 192–207 In: N. Gales, M. Hindell, and R. Kirkwood (eds.). *Marine mammals: Fisheries, tourism and management issues*. CSIRO Publishing, Melbourne.
- Wilkinson IS; Childerhouse SJ; Duignan PJ; Gulland FMD (2000). Infanticide and cannibalism in the New Zealand sea lion. *Marine Mammal Science* 16:495–500.
- Wilkinson IS; Duignan PJ; Grinberg A; Chilvers BL; Robertson BC (2006). *Klebsiella pneumoniae* epidemics: Possible impact on New Zealand sea lion recruitment. pp. 455–471. In: A. Trites, S. Atkinson, D. DeMaster, L. Fritz, T. Gelatt, L. Rea, and K. Wynne. (eds.). *Sea lions of the world*. Alaska Sea Grant College Program, University of Alaska Fairbanks.
- Woodley TH; Lavigne DM (1991). Incidental capture of pinnipeds in commercial fishing gear. *International Marine Mammal Association Technical Report* 91-01: 35 p.

## 4. New Zealand fur seal (*Arctocephalus forsteri*)

<i>Scope of chapter</i>	This chapter outlines the biology New Zealand fur seals ( <i>Arctocephalus forsteri</i> ), the nature of any fishing interactions, the management approach, trends in key indicators of fishing effects and major sources of uncertainty.
<i>Area</i>	All of the New Zealand EEZ and territorial sea.
<i>Focal localities</i>	Areas with significant fisheries interactions include waters over or close to the continental shelf surrounding the South Island and southern offshore islands, notably Cook Strait, West Coast South Island, Banks Peninsula and the Bounty Islands, plus offshore of Bay of Plenty-East Cape.
<i>Key issues</i>	Improving estimates of incidental bycatch in some fisheries, and assessing the potential for populations to sustain the present levels of bycatch.
<i>Emerging issues</i>	Improving data and information sources for future ecological risk assessments.
<i>MPI Research (current)</i>	PRO2010-01 <i>Estimating the nature &amp; extent of incidental captures of seabirds, marine mammals &amp; turtles in New Zealand commercial fisheries</i> ; PRO2012-02 <i>Assess the risk posed to marine mammal populations from New Zealand fisheries</i> .
<i>Other Govt Research (current)</i>	<u>DOC Marine Conservation Services Programme (CSP)</u> : INT2012-01 <i>To understand the nature and extent of protected species interactions with New Zealand commercial fishing activities</i> .
<i>Links to 2030 objectives</i>	Objective 6: Manage impacts of fishing and aquaculture. Strategic Action 6.2: Set and monitor environmental standards, including for threatened and protected species and seabed impacts
<i>Related issues/chapters</i>	See the New Zealand sea lion chapter.

### 4.1. Context

Management of fisheries impacts on New Zealand (NZ) fur seals is legislated under the Marine Mammals Protection Act (MMPA) 1978 and the Fisheries Act (FA) 1996. Under s.3E of the MMPA, the Minister of Conservation, with the concurrence of the Minister for Primary Industries (formerly the Minister of Fisheries), may approve a population management plan (PMP). There is no PMP in place for NZ fur seals.

In the absence of a PMP, the Ministry for Primary Industries (MPI) manages fishing-related mortality of NZ fur seals under s.15(2) of the FA “to avoid, remedy, or mitigate the effect of fishing-related mortality on any protected species, and such measures may include setting a limit on fishing-related mortality.”

All marine mammal species are designated as protected species under s.2(1) of the FA. In 2005, the Minister of Conservation approved the Conservation General Policy, which specifies in Policy 4.4 (f) that “Protected marine species should be managed for their long-term viability and recovery throughout their natural range.” DOC’s Regional Conservation Management Strategies outline specific policies and objectives for protected marine species at a regional level.

In 2004, DOC approved the *Department of Conservation Marine Mammal Action Plan for 2005–2010*<sup>20</sup> (Suisted and Neale 2009). The plan specifies a number of species-specific key objectives for NZ fur seals, of which the following is most relevant for fisheries interactions: “*To control/mitigate fishing-related mortality of NZ fur seals in trawl fisheries (including the WCSI hoki and Bounty Island southern blue whiting fisheries).*”

Management of NZ fur seal incidental captures aligns with Fisheries 2030 Objective 6: *Manage impacts of fishing and aquaculture*. Further, the management actions follow Strategic Action 6.2: *Set and monitor environmental standards, including for threatened and protected species and seabed impacts*.

All National Fisheries Plans except those for inshore shellfish and freshwater fisheries are relevant to the management of fishing-related mortality of NZ fur seals.

Under the National Deepwater Plan, the objective most relevant for management of NZ fur seals is Management Objective 2.5: *Manage deepwater and middle-depth fisheries to avoid or minimise adverse effects on the long-term viability of endangered, threatened and protected species*.

Specific objectives for the management of NZ fur seals bycatch are to be outlined in the fishery-specific chapters of the National Deepwater Plan for the fisheries with which NZ fur seals are most likely to interact. These fisheries include hoki (HOK), southern blue whiting (SBW), hake (HAK) and jack mackerel (JMA). The HOK chapter of the National Deepwater Plan is complete and includes Operational Objective 2.11: *Ensure that incidental marine mammal captures in the hoki fishery are avoided and minimised to acceptable levels (which may include standards) by 2012*. The SBW chapter is nearing completion while the timeframes for the HAK and JMA chapters are yet to be confirmed.

Management Objective 7 of the National Fisheries Plan for Highly Migratory Species (HMS) is to “*Implement an ecosystem approach to fisheries management, taking into account associated and dependent species.*” This comprises four components: Avoid, remedy, or mitigate the adverse effects of fishing on associated and dependent species, including through maintaining foodchain relationships; Minimise unwanted bycatch and maximise survival of incidental catches of protected species in HMS fisheries, using a risk management approach; Increase the level and quality of information available on the capture of protected species; and Recognise the intrinsic values of HMS and their ecosystems, comprising predators, prey, and protected species.

The Environment Objective is the same for all groups of fisheries in the draft National Fisheries Plan for Inshore Finfish, to “Minimise adverse effects of fishing on the aquatic environment, including on biological diversity”. The draft National Fisheries Plans for Inshore Shellfish and Freshwater have the same objective but are unlikely to be relevant to management of fishing-related mortality of NZ fur seals.

## **4.2. Biology**

### **4.2.1. Taxonomy**

The NZ fur seal (*Arctocephalus forsteri* (Lesson, 1828)) is one of only two species of otariid (eared seals, includes fur seals and sea lions) native to New Zealand, the other being the New Zealand sea lion (*Phocarctos hookeri* (Gray, 1844)).

---

<sup>20</sup> DOC has confirmed that the Marine Mammal Action Plan for 2005–2010 still reflects DOC’s priorities for marine mammal conservation.

#### 4.2.2. Distribution

Pre-European archaeological evidence suggests that NZ fur seals were present along much of the east coasts of the North Island (except the less rocky coastline of Bay of Plenty and Hawke Bay) and the South Island, and, to a lesser extent, on the west coasts, where fewer areas of suitable habitat were available (Smith 1989, 2005, 2011). A combination of subsistence hunting and commercial harvest resulted contraction of the species' range and in population decline almost to the point of extinction (Smith 1989, 2005, 2011, Ling 2002, Lalas 2008). NZ fur seals became fully protected in the 1890's and, with the exception of one year of licenced harvest in the 1950's, have remained protected since.

Currently, NZ fur seals are dispersed throughout New Zealand waters, especially in waters south of about 40° S to Macquarie Island. On land, NZ fur seals are distributed around the New Zealand coastline, on offshore islands, and on sub-Antarctic islands (Crawley and Wilson 1976, Wilson 1981, Mattlin 1987). The recolonisation of the coastline by NZ fur seals has resulted in the northward expansion of the distribution of breeding colonies and haulouts (Lalas and Bradshaw 2001), and breeding colonies present on many exposed rocky areas (Baird 2011). The extent of breeding colony distribution in New Zealand waters is bounded to the north by a very small (space-limited) colony at Gannet Island off the North Island west coast (latitude 38° S), to the east by colonies of unknown sizes at the Chatham Islands group, to the west by colonies of unknown size on Fiordland offshore islands, and to the south by unknown numbers on Campbell Island. Outside New Zealand waters, breeding populations exist in South and Western Australia (Shaughnessy *et al.* 1994, Shaughnessy 1999, Goldsworthy *et al.* 2003).

The seasonal distribution of the NZ fur seals is determined by the sex and maturity of each animal. Males are generally at the breeding colonies from late October to late January then move to haulout areas around the New Zealand coastline (see Bradshaw *et al.* 1999), with peak density of males and sub-adult males at haulouts during July–August and lowest densities in September–October (Crawley and Wilson 1976). Females arrive at the breeding colony from November and lactating females remain at the colony (apart from short foraging trips) for about 10 months until the pups are weaned, usually during August–September (Crawley and Wilson 1976).

#### 4.2.3. Foraging ecology

Most foraging research in New Zealand has focused on lactating NZ fur seals at Open Bay Islands off the South Island west coast (Mattlin *et al.* 1998), Otago Peninsula (Harcourt *et al.* 2002), and Ohau Point, Kaikoura (Boren 2005), using time-depth-recorders, satellite-tracking, or very-high-frequency transmitters. Individual females show distinct dive pattern behaviour and may be relatively shallow or deep divers, but most forage at night and in depths shallower than 200 m. At Open Bay Islands, dives were generally deeper and longer in duration during autumn and winter. Females can dive to at least 274 m (for a 5.67 min dive in autumn) and remain near the bottom at over 237 m for up to 11.17 min in winter (Mattlin *et al.* 1998). Females in some locations undertook longer dive trips, with some to deeper waters, in autumn (in over 1000 m beyond the continental shelf; Harcourt *et al.* 2002).

The relatively shallow dives and nocturnal feeding during summer suggested that seals fed on pelagic and vertical migrating prey species (for example, arrow squid, *Nototodarus sloanii*). Conversely, the deeper dives and increased number of dives in daylight during autumn and winter suggested that the prey species may include benthic, demersal, and pelagic species (Mattlin *et al.* 1998, Harcourt *et al.* 2002). The deeper dives enabled seals to forage along or off the continental shelf (within 10 km) of the colony studied (at Open Bay Islands). These deeper dives may be to the benthos or to depths in the water column where spawning hoki are concentrated.

Methods to analyse NZ fur seal diets have included investigation of freshly killed animals (Sorensen 1969), scats, and regurgitates (e.g. Allum and Maddigan 2012). Fish prey items can be recognised by the presence of otoliths, bones, scales, and lenses, while cephalopods are indicated by beaks and pens. Foraging appears to be specific to individuals and different diets may be represented in the scats and regurgitations of males and females as well as juveniles from one colony. These analyses can be biased, however, particularly if only one collection method is used, and this limits fully quantitative assessment of prey species composition.

Dietary studies of NZ fur seals have been conducted at colonies in Nelson-Marlborough, west coast South Island, Otago Peninsula, Kaikoura, Banks Peninsula, Snares Islands, and off Stewart Island, and summaries are provided by Carey (1992), Harcourt (2001), Boren (2010), and Baird (2011).

NZ fur seals are opportunistic foragers and, depending on the time of year, method of analysis, and location, their diet includes at least 61 taxa (Holborow 1999) of mainly fish (particularly lanternfish (myctophids) in all studied colonies except Tonga Island (in Golden Bay, Willis *et al.* 2008), as well as anchovy (*Engraulis australis*), aruhu (*Auchenoceros punctatus*), barracouta (*Thrysites atun*), hoki (*Macruronus novaezelandiae*), jack mackerel (*Trachurus* spp.), pilchard (*Sardinops sagax*), red cod (*Pseudophycis bachus*), red gurnard (*Chelidonichthys kumu*), silverside (*Argentina elongate*), sprat (*Sprattus* spp.) and cephalopods (octopus (*Macroctopus maorum*), squid (*Nototodarus sloanii*, *Sepioteuthis bilineata*)). For example, myctophids were present in Otago scats throughout the year (representing offshore foraging), but aruhu, sprat, and juvenile red cod were present only during winter-spring (Fea *et al.* 1999). Medium-large arrow squid predominated in summer and autumn. Jack mackerel species, barracouta, and octopus were dominant in winter and spring. Prey such as lanternfish and arrow squid rise in the water column at night, the time when NZ fur seals exhibit shallow foraging (Harcourt *et al.* 1995, Mattlin *et al.* 1998, Fea *et al.* 1999).

#### 4.2.4. Reproductive biology

NZ fur seals are sexually dimorphic and polygynous (Crawley and Wilson 1976); males may weigh up to 160 kg, whereas females weigh up to about 50 kg (Miller 1975; Mattlin 1978a, 1987; Troy *et al.* 1999). Adult males are much larger around the neck and shoulders than females and breeding males are on average 3.5 times the weight of breeding females (Crawley and Wilson 1976). Females are philopatric and are sexually mature at 4–6 years, whereas males mature at 5–9 years (Mattlin 1987, Dickie and Dawson 2003). The maximum age recorded for NZ fur seals in New Zealand waters is 22 years for females (Dickie and Dawson 2003) and 15 years for males (Mattlin 1978).

NZ fur seals are annual breeders and generally produce one pup after a gestation period of about 10 months (Crawley and Wilson 1976). Twinning can occur and females may foster a pup (Dowell *et al.* 2008), although both are rare. Breeding animals come ashore to mate after a period of sustained feeding at sea. Breeding males arrive at the colonies to establish territories during October–November. Breeding females arrive at the colony from late November and give birth shortly after. Peak pupping occurs in mid December (Crawley and Wilson 1976).

Females remain at the colony with their newborn pups for about 10 days, by which time they have usually mated. Females then leave the colony on short foraging trips of 3–5 days before returning to suckle pups for 2–4 days (Crawley and Wilson 1976). As the pups grow, these foraging trips are progressively longer in duration. Pups remain at the breeding colony from birth until weaning (at 8–12 months of age).

Breeding males generally disperse after mating to feed and occupy haulout areas, often in more northern areas (Crawley and Wilson 1976). This movement of breeding adults away from the colony area during January allows for an influx of sub-adults from nearby areas. Little is described about the ratio of males to females on breeding colonies (Crawley and Wilson 1976), or the reproductive

success. Boren (2005) reported a fecundity rate of 62% for a Kaikoura colony, based on two annual samples of between about 5 and 8% of the breeding female population. This rate is similar to the 67% estimated by Goldsworthy and Shaughnessy (1994) for a South Australian colony.

Newborn pups are about 55 cm long and weigh about 3.5 kg (Crawley and Wilson 1976). Male pups are generally heavier than female pups at birth and throughout their growth (Crawley and Wilson 1976, Mattlin 1981, Chilvers *et al.* 1995, Bradshaw *et al.* 2003b, Boren 2005). Pup growth rates may vary by colony (see Harcourt 2001). The proximity of a colony to easily accessible rich food sources will vary, and pup condition at a colony can vary markedly between years (Mattlin 1981, Bradshaw *et al.* 2000, Boren 2005). Food availability may be affected by climate variation, and pup growth rates probably represent variation in the ability of mothers to provision their pups from year to year. The sex ratio of pups at a colony may vary by season (Bradshaw *et al.* 2003a, 2003b, Boren 2005), and in years of high food resource availability, more mothers may produce males or more males may survive (Bradshaw *et al.* 2003a, 2003b).

#### 4.2.5. Population biology

Historically, the population of NZ fur seals in New Zealand was thought to number above 1.25 million animals (possibly as high as 1.5 to 2 million) before the extensive sealing of the early 19<sup>th</sup> century (Richards 1994). Present day population estimates for NZ fur seals in New Zealand are few and highly localised. In the most comprehensive attempt to quantify the total NZ fur seal population, Wilson (1981) summarised population surveys of mainland New Zealand and offshore islands undertaken in the 1970s and estimated the population size within the New Zealand region at between 30,000 and 50,000 animals. Since then, several authors have suggested a population size of ~100,000 animals (Taylor 1990, see Harcourt 2001), but this estimate is very much an approximation and its accuracy is difficult to assess in the absence of comprehensive surveys.

Fur seal colonies provide the best data for consistent estimates of population numbers, generally based on pup production in a season (see Shaughnessy *et al.* 1994). Data used to provide colony population estimates of NZ fur seals have been, and generally continue to be, collected in an *ad hoc* fashion. Regular pup counts are made at some discrete populations. A 20-year time series of Otago Peninsula colony data is updated, maintained, and published primarily by Chris Lalas (assisted by Sanford (South Island) Limited), and the most recent estimate is 20,000–30,000 animals (Lalas 2008). A 20-year plus time series of pup counts exists for three west coast South Island colonies (Cape Foulwind, Wekekeura Point, and Open Bay Islands; Best 2011). Recent Kaikoura work by Boren (2005) covered four seasons and unpublished data are available for the subsequent seasons.

Other studies of breeding colonies generally provide estimates for one or two seasons, but many of these are more than 10 years old. Published estimates suggest that populations have stabilised at the Snares Islands after a period of growth in the 1950s and 1960s (Carey 1998) and increased at the Bounty Islands (Taylor 1996), Nelson-Marlborough region (Taylor *et al.* 1995), Kaikoura (Boren 2005), Otago (Lalas and Harcourt 1995, Lalas and Murphy 1998, Lalas 2008), and near Wellington (Dix 1993).

For many areas where colonies or haulouts exist, count data have been collected opportunistically (generally by Department of Conservation staff during their field activities) and thus data are not often comparable because counts may represent different life stages, different assessment methods, and different seasons (see Baird 2011).

Baker *et al.* (2010a) conducted an aerial survey of the South Island west coast from Farewell Spit to Puysegur Point and Solander Island in 2009 but were their counts were quite different from ground counts collected at a similar time at the main colonies (Melina and Cawthorn 2009). This discrepancy was thought to be a result mainly of the survey design and the nature of the terrain. However, the

aerial survey confirmed the localities shown by Wilson (1981) of potentially large numbers of pups at sites such as Cascade Point, Yates Point, Chalky Island, and Solander Island.

Population numbers for some areas, especially more isolated ones, are not well known. The most recent counts for the Chatham Islands were collected in the 1970s (Wilson 1981), and the most recent for the Bounty Islands in 1993–94. Taylor (1996) reported an increase in pup production at the Bounty Islands since 1980, and estimated that the total population was at least 21 500, occupying over 50% of the available area. Information is sparse for populations at Campbell Island, the Auckland Islands group and the Antipodes Islands

Little is reported about the natural mortality of NZ fur seals, other than reports of sources and estimates of pup mortality for some breeding colonies. Estimates of pup mortality or pup survival vary in the manner in which they were determined and in the number of seasons they represent, and are not directly comparable. Each colony will be affected by different sources of mortality related to habitat, location, food availability, environment, and year, as well as the ability of observers to count all the dead pups (may be limited by terrain, weather, or time of day).

Reported pup mortality rates vary: 8% for Otago Peninsula pups up to 30 days old and 23% for pups up to 66 days old (Lalas and Harcourt 1995); 20% from birth to 50 days and about 40% from birth to 300 days for Taumaka Island, Open Bay Islands pups (Mattlin 1978b); and in one year, 3% of Kaikoura pups before the age of 50 days (Boren 2005). Starvation was the major cause of death, although stillbirth, suffocation, trampling, drowning, predation, and human disturbance also occur. Pup survival of at least 85% was estimated for a mean 47 day interval for three Otago colonies, incorporating data such as pup body mass (Bradshaw *et al.* 2003b), though pup mortality before the first capture effort was unknown. Other sources of natural mortality for NZ fur seals include predators such as sharks and NZ sea lions (Mattlin 1978b, Bradshaw *et al.* 1998).

Human-induced sources of mortality include: fishing, for example, entanglement or capture in fishing gear; vehicle-related deaths (Lalas and Bradshaw 2001, Boren 2005, Boren *et al.* 2006, 2008); and mortality through shooting, bludgeoning, and dog attacks. NZ fur seals are vulnerable to certain bacterial diseases and parasites and environmental contaminants, though it is not clear how life-threatening these are. The more obvious problems include tuberculosis infections, *Salmonella*, hookworm enteritis, phocine distemper, and septicaemia (associated with abortion) (Duignan 2003, Duignan and Jones 2007). Low food availability and persistent organohalogen compounds (which can affect the immune and the reproductive systems) may also affect NZ fur seal health.

Various authors have investigated fur seals genetic differentiation among colonies and regions in New Zealand (Lento *et al.* 1994; Robertson and Gemmell (2005). Lento *et al.* (1994) described the geographic distribution of mitochondrial cytochrome *b* DNA haplotypes, whereas Robertson and Gemmell (2005) described low levels of genetic differentiation (consistent with homogenising gene flow between colonies and an expanding population) based on genetic material from NZ fur seal pups from seven colonies. One aim of the work is to determine the provenance of animals captured during fishing activities, through the identification and isolation of any colony genetic differences.

#### **4.2.6. Conservation biology and threat classification**

Threat classification is an established approach for identifying species at risk of extinction (IUCN 2010). The risk of extinction for NZ fur seals has been assessed under two threat classification systems: the New Zealand Threat Classification System (Townsend *et al.* 2008) and the International Union for the Conservation of Nature (IUCN) Red List of Threatened Species (IUCN 2010).

In 2008, the IUCN updated the Red List status of NZ fur seals, listing them as Least Concern on the basis of their large and apparently increasing population size (Goldsworthy and Gales 2008). In 2010,

DOC updated the New Zealand Threat Classification status of all NZ marine mammals (Baker *et al.* 2010b). In the revised list, NZ fur seals were classified as Not Threatened with the qualifiers increasing (Inc) and secure overseas (SO) (Baker *et al.* 2010b).

#### **4.3. Global understanding of fisheries interactions**

NZ fur seals are found in both Australian and New Zealand waters. Overall abundance has been suggested to be as high as 200 000, with about half of the population in Australian waters (Goldsworthy and Gales 2008). However, this figure is very much an approximation, and its accuracy is difficult to assess in the absence of comprehensive surveys.

Pinnipeds are caught incidentally in a variety of fisheries worldwide (Read *et al.* 2006), including: NZ fur seals, Australian fur seals, and Australian sea lions in Australian trawl and inshore fisheries (e.g., Shaughnessy 1999, Norman 2000); Cape fur seals in South African fisheries (Shaughnessy and Payne 1979); South American sea lions in trawl fisheries off Patagonia (Dans *et al.* 2003); and seals and sea lions in United States waters (Moore *et al.* 2009).

#### **4.4. State of knowledge in New Zealand**

NZ fur seals are attracted to feeding opportunities in various fishing gears and anecdotal evidence suggests that the sound of winches as trawlers haul their gear acts as a ‘dinner gong’. The attraction of fish in a trawl net, on longline hooks, or caught in a setnet provide opportunities for NZ fur seals to interact with fishing gear, which can result in capture and, potentially, death via drowning or injury.

Most captures occur in trawl fisheries and NZ fur seals are most at risk from capture during shooting and hauling (Shaughnessy and Payne 1979), when the net mouth is within diving depths. Once in the net some animals may have difficulty in finding their way out within their maximum breath-hold time (Shaughnessy and Davenport 1996). The operational aspects that are associated with NZ fur seal captures on trawlers include factors that attract the NZ fur seals, such as the presence of offal and discards, the sound of the winches, vessel lights, and the presence of ‘stickers’ in the net (Baird 2005). NZ fur seals are at particular risk of capture when a vessel partially hauls the net during a tow and executes a turn with the gear close to the surface. At the haul, NZ fur seals often attempt to feed from the codend as it is hauled and dive after fish that come loose and escape from the net (Baird 2005).

Factors identified as important influences on the potential capture of NZ fur seals in trawl gear include the year or season, the fishery area, gear type and fishing strategies (often specific to certain nationalities within the fleet), time of day, and distance to shore (Baird and Bradford 2000, Mormede *et al.* 2008, Smith and Baird 2009). These analyses did not include any information on NZ fur seal numbers or activity in the water at the stern of the vessel. Other influences on NZ fur seal capture rate (of Australian and NZ fur seals) may include inclement weather and sea state, vessel speed, increased numbers of vessels and trawl frequency, and potentially the weight of the fish catch and the presence of certain bycatch fish species (Hamer and Goldsworthy 2006). This Australian study found similar mortality rates for tows with and without Seal Exclusion Devices (see also Hooper *et al.* 2005).

The spatial and temporal overlap of commercial fishing grounds and NZ fur seal foraging areas has resulted in NZ fur seal captures in fishing gear (Matlin 1987, Rowe 2009). Most fisheries with observed captures occur in waters over or close to the continental shelf. Because the topography around much of the South Island and offshore islands slopes steeply to deeper waters, most captures occur close to colonies and haulouts (Figures 4.1 and 4.2).

Observed NZ fur seal captures are mainly from trawls in defined seasons in areas where fishing occurs relatively close to NZ fur seal colonies or haulouts. Winter hoki fisheries attract NZ fur seals

off the west coast South Island and in Cook Strait between late June and September (Table 4.1). In August–October, NZ fur seals are caught in southern blue whiting effort near the Bounty Islands and Campbell Island. In September–October captures may occur in hoki and ling fisheries off Puysegur Point on the southwestern coast of the South Island. Captures are also reported from the Stewart-Snares shelf fisheries that operate during summer months, mainly for hoki and other middle depths species and squid, and from fisheries throughout the year on the Chatham Rise though captures have not been observed east of longitude 180° on the Chatham Rise.

Captures were reported from trawl fisheries for species such as hoki, hake (*Merluccius australis*), ling (*Genypterus blacodes*), squid, southern blue whiting, Jack mackerel, and barracouta (Baird and Smith 2007, Abraham *et al.* 2010a). Between 1 and 3% of observed tows targeting middle depths fish species catch NZ fur seals compared with about 1% for squid tows, and under 1% of observed tows targeting deepwater species such as orange roughy (*Hoplostethus atlanticus*) and oreo species (for example, *Allocyttus niger*, *Pseudocottus maculatus*) (Baird and Smith 2007). The main fishery areas that contribute to the estimated annual catch of NZ fur seals (modelled from observed captures) in middle depths and deepwater trawl fisheries are Cook Strait hoki, west coast South Island middle depths fisheries (mainly hoki), western Chatham Rise hoki, and the Bounty Islands southern blue whiting fishery (Baird and Smith 2007, Thompson and Abraham 2010). Captures on longlines occur when the NZ fur seals attempt to feed on the fish catch during hauling. Most NZ fur seals are released alive from surface and bottom longlines, typically with a hook and short snood or trace still attached.

**Table 4.1: Monthly distribution of NZ fur seal activity and the main trawl and longline fisheries with observed reports of NZ fur seal incidental captures.**

NZ fur seals	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep
Breeding males	At breeding colony				Dispersed at sea or at haulouts							
Breeding females	At sea	At breeding colony		At breeding colony and at-sea foraging and suckling							At sea	
Pups	At sea		At breeding colony								At sea	
Non-breeders	Dispersed at sea, at haulouts, or breeding colony periphery											
Major fisheries	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep
Hoki trawl	Puysegur		Chatham Rise						Cook Strait, west coast South Island			
Squid trawl			Stewart-Snares shelf, Auckland Is. Shelf, East Coast South Island									
Southern blue whiting trawl	Campbell Rise										Bounty Is., Pukaki Rise	
Southern bluefin tuna longline						Fiordland						




Figure 4.1: Distribution of trawl fishing effort and observed NZ fur seal captures, 2002-03 to 2010-11 (for more information see <http://data.dragonfly.co.nz/psc/>). Fishing effort is mapped into 0.2-degree cells, with the colour of each cell being related to the amount of effort. Observed fishing events are indicated by black dots, and observed captures are indicated by red dots. Fishing is only shown if the effort could be assigned a latitude and longitude, and if there were three or more vessels fishing within a cell. In this case, 96.0% of the effort is shown.




Figure 4.2: Distribution of surface longline fishing effort and observed NZ fur seal captures, 2002-03 to 2010-11 (for more information see <http://data.dragonfly.co.nz/psc/>). Fishing effort is mapped into 0.2-degree cells, with the colour of each cell being related to the amount of effort. Observed fishing events are indicated by black dots, and observed captures are indicated by red dots. Fishing is only shown if the effort could be assigned a latitude and longitude, and if there were three or more vessels fishing within a cell. In this case, 75.3% of the effort is shown.

#### 4.4.1. Quantifying fisheries interactions

Observer data and commercial effort data have been used historically to characterise the incidental captures and estimate the total numbers caught (Baird and Smith 2007, Smith and Baird 2009, Thompson and Abraham 2010, Abraham and Thompson 2011). This approach is currently applied using information collected under DOC project INT2012-01 and analysed under MPI project PRO2010-01 (Thompson *et al.* 2011, Thompson *et al.* 2012). The analytical methods used to estimate capture numbers across the commercial fisheries have depended on the quantity and quality of the data, in terms of the numbers observed captured and the representativeness of the observer coverage. Initially, stratified ratio estimates were provided for the main trawl fisheries, starting in the late 1980s, after scientific observers reported 198 NZ fur seal deaths during the July to September west coast South Island spawning hoki fishery (Mattlin 1994a, 1994b). In the following years, ratio estimation was used to estimate NZ fur seal captures in the Taranaki Bight jack mackerel fisheries and Bounty Platform, Pukaki Rise, and Campbell Rise southern blue whiting fisheries, based on observed catches and stratified by area, season, and gear type (Baird 1994).

In the last 10 years, model-based estimates of captures have been developed for all trawl fisheries in waters south of 40° S (Baird and Smith 2007, Smith and Baird 2009, Thompson and Abraham 2010, Abraham and Thompson 2011, Thompson *et al.* 2011, Thompson *et al.* 2012). These models use the observed and unobserved data in an hierarchical Bayesian approach that combines season and vessel-season random effects with covariates (for example, day of fishing year, time of day, tow duration, distance from shore, gear type, target) to model variation in capture rates among tows. This method compensates in part for the lack of representativeness of the observer coverage and includes the contribution from correlation in the capture rate among tows by the same vessel. The method is limited by the very large differences in the observed and non-observed proportions of data for the different vessel sizes; most observer coverage is on larger vessels that generally operate in waters deeper than 200 m. The operation of inshore vessels in terms of the location of effort, gear, and the fishing strategies used is also relatively unknown compared with the deeper water fisheries although changes to reporting requirements means that data is now improving and inshore trawl effort (not including flatfish trawl effort) is now able to be included in the modelling (Thompson *et al.* 2012, see also description of the Trawl Catch Effort Return, TCER, in use since 2007/08, in Chapter 7 on benthic effects).


Since 2005, there has been a small downward trend in estimated capture rates, and annual estimated NZ fur seal captures (Smith and Baird 2009, Thompson and Abraham 2010, Abraham and Thompson 2011, Thompson *et al.* 2011, Thompson *et al.* 2012, Figure 4.3). This probably reflects efforts to reduce bycatch combined with a reduction in fishing effort since the late 1990s. Similar modelling methods were used to produce the most recent set of estimated NZ fur seal captures in trawl fisheries (Thompson and Abraham 2010, Abraham and Thompson 2011, Thompson *et al.* 2011, Thompson *et al.* 2012). The overall downward trend in estimated annual captures for trawl fisheries has continued (see Table 4.2), as a result of the continued decrease in total tows made each year and a concurrent decrease in capture rate. Note these capture rates include animals that are released alive (7% of observed trawl capture in 2008-09, Thompson and Abraham 2010).

Ratio estimation was used to calculate total captures in longline fisheries by target fishery fleet and area (Baird 2008) and by all fishing methods (Abraham *et al.* 2010a). NZ fur seal captures in surface longline fisheries have been generally observed in waters south and west of Fiordland, but also in the Bay of Plenty and off East Cape. Estimated numbers range from 127 (95% c.i. 121–133) in 1998–99 to 25 (14–39) in 2007–08 during southern bluefin tuna fishing by chartered and domestic vessels (Abraham *et al.* 2010a). These capture rates include animals that are released alive (100% of observed surface longline capture in 2008-09, Thompson and Abraham 2010). Captures of NZ fur seals have also been recorded in other fisheries; 8 in setnets and 2 in bottom longline fisheries since 2002-03

(Thompson *et al.* 2012). Captures associated with recreational fishing activities are poorly known (Abraham *et al.* 2010b).

**Table 4.2: Effort, observed and estimated NZ fur seal captures in trawl and surface longline fisheries by fishing year in the New Zealand EEZ (Abraham and Thompson 2011 and <http://data.dragonfly.co.nz/psc/>). For each fishing year, the table gives the the total number of tows or hooks; the observer coverage (the percentage of tows or hooks that were observed); the number of observed captures (both dead and alive); the capture rate (captures per hundred tows or per thousand hooks); the estimation method used (model or ratio); and the mean number of estimated total captures (with 95% confidence interval). For more information on the methods used to prepare the data, see Abraham and Thompson (2011).**

Fishing year	Fishing effort		Observed captures		Estimated captures		
	All effort	% observed	Number	Rate	Type	Mean	95% c.i.
<i>Trawl fisheries</i>							
1998–1999	153 412	4.7	190	2.62	Ratio	1 591	1454–1744
1999–2000	139 057	5.5	203	2.65	Ratio	1 539	1400–1693
2000–2001	134 243	6.8	170	1.87	Ratio	1 490	1348–1649
2001–2002	127 883	6.0	157	2.03	Ratio	1 273	1164–1394
2002–2003	130 344	5.2	68	1.00	Model	841	503 – 1380
2003–2004	121 494	5.4	84	1.28	Model	1 052	635 – 1728
2004–2005	120 590	6.4	200	2.59	Model	1 471	914 – 2392
2005–2006	110 230	5.9	143	2.18	Model	917	577 – 1479
2006–2007	103 529	7.7	73	0.92	Model	533	324 – 871
2007–2008	89 537	10.1	141	1.56	Model	765	476 – 1348
2008–2009	87 587	11.2	72	0.73	Model	546	308 – 961
2009–2010	92 886	9.7	72	0.80	Model	472	269 – 914
2010–2011	86 073	8.6	69	0.93	Model	376	221 – 668
<i>Surface longline fisheries</i>							
1998–1999	6 855 124	18.9	102	0.08	Ratio	138	120–160
1999–2000	8 258 537	10.4	42	0.05	Ratio	67	54–83
2000–2001	9 698 805	10.8	43	0.04	Ratio	64	51–83
2001–2002	10 833 533	9.1	44	0.04	Ratio	75	61–93
2002–2003	10 764 588	20.4	56	0.03	Ratio	73	63–87
2003–2004	7 380 779	21.8	40	0.02	Ratio	107	61–189
2004–2005	3 676 365	21.3	20	0.03	Ratio	46	26–71
2005–2006	3 687 339	19.1	12	0.02	Ratio	59	28–100
2006–2007	3 738 362	27.8	10	0.01	Ratio	31	18–49
2007–2008	2 244 339	19.0	10	0.02	Ratio	29	17–46
2008–2009	3 115 633	30.1	22	0.02	Ratio	48	29–75
2009–2010	2 992 285	22.3	19	0.03	Ratio	65	35–103
2010–2011	3 164 159	21.3	17	0.03	Ratio	57	26–99



**Figure 4.3: Observed captures of NZ fur seals in trawl fisheries (both dead and alive), the capture rate (captures per hundred tows) and the mean number of estimated total captures (with 95% confidence interval) by fishing year for regions with more than 50 observed captures since 2002-03: (a) the New Zealand EEZ; (b) the Cook Strait area; (c) the East Coast South Island area; (d) the Stewart Snares Shelf area; (e) the Subantarctic area; and (f) the West Coast South Island area (Abraham and Thompson 2011 and <http://data.dragonfly.co.nz/psc/>). For more information on the methods used to prepare the data, see Abraham and Thompson (2011).**

#### 4.4.2. Managing fisheries interactions

The impact of fishing related captures on the NZ fur seal population is presently unknown. However, fishing interactions are considered unlikely to have adverse population-level consequences for NZ fur seals given: the scale of bycatch relative to overall NZ fur seal abundance; the apparently increasing population and range; and the NZ and IUCN threat status of the species. The consequences of fishing related mortality for some individual colonies may be more or less severe.

Management has focused on encouraging vessel operators to alter fishing practices to reduce captures, and monitoring captures via the observer programme. A marine mammal operating procedure (MMOP) has been developed by the deepwater sector to reduce the risk of marine mammal captures and is currently applied to trawlers greater than 28 m LOA and is supported by annual training. It includes a number of mitigation measures, such as managing offal discharge and refraining from shooting and hauling the gear when NZ fur seals are congregating around the vessel. Its major focus is reducing the time gear is at or near the surface when it poses the greatest risk. MPI monitors and audits vessel performance against this procedure (see the MPI National Deepwater Plan for further details).

Research into methods to minimise or mitigate NZ fur seal captures in commercial fisheries has focused on fisheries in which NZ fur seals are more likely to be captured (trawl fisheries, see Clement and Associates 2009). Finding ways to mitigate captures has proven difficult because the animals are free swimming, can easily dive to the depths of the net when it is being deployed, hauled, or brought to the surface during a turn, and are known to deliberately enter nets to feed. Further, any measures also need to ensure that the catch is not greatly compromised, either in terms of the amount of fish or their condition. This is one potential drawback of using seal exclusion devices (see Rowe 2007). Adhering to current risk mitigation methods (e.g. MMOP) will help to minimise the level of impacts, however rates may fluctuate depending on fleet deployment, NZ fur seal abundance and local feeding conditions.

#### 4.4.3. Modelling population-level impacts of fisheries interactions

The uncertainty about the size of the NZ fur seal population has restricted the potential to investigate any effects that NZ fur seal deaths through fishing may have on the population as a whole or on the viability of colonies or groups of colonies. The provenance of NZ fur seals caught during fishing is presently unknown, although proposed genetic research potentially could identify which animals belonged to a specific colony (Robertson and Gemmell 2005).

In response to the requirements for the Marine Stewardship Council certification of the hoki fishery (one target fishery contributing to NZ fur seal mortality), expert knowledge about NZ fur seals and their interactions with trawl gear (including some comparisons of annual capture estimates) have been used for an expert-based qualitative ecological risk assessment (ERAs). The results of this study have not been reviewed by the AEWG or DOC's CSP-TWG.

The impact of fisheries interactions on NZ fur seal populations (and other marine mammal populations) will be assessed in the marine mammal risk assessment project (PRO2012-02) due to be commissioned in 2013. The goal of this project is to assess the risk posed to marine mammal populations by New Zealand fisheries by applying a similar approach to the recent seabird risk assessment (Richard *et al.* 2011). In this approach, risk is defined as the ratio of total estimated annual fatalities due to bycatch in fisheries, to the level of Potential Biological Removal (PBR, Wade 1998). The results should be available in 2014.



#### 4.4.4. Sources of uncertainty

Any measure of the effect of NZ fur seal mortality from commercial fisheries on NZ fur seal populations requires adequate information on the size of the populations at different colonies. Although there is reasonable information about where the main NZ fur seal breeding colonies exist, the size and dynamics of the overall populations are poorly understood. At present, the main sources of uncertainty are the lack of consistent data on: abundance by colony and in total; population demographic parameters; and at-sea distribution (which would ideally be available at the level of a colony or wider geographic area where several colonies are close together) (Baird 2011). Collation and analysis of existing data, such as that for the west coast South Island, would fill some of these gaps; there is a 20-year time series of pup production from three west coast South Island colonies, a reasonably long data series from the Otago Peninsula, and another from Kaikoura. Maximum benefit could be gained through the use of all available data, as shown by the monitoring of certain colonies of NZ fur seals in Australia to provide a measure of overall population stability (see Shaughnessy *et al.* 1994, Goldsworthy *et al.* 2003).

Fur seals may forage in waters near a colony or haulout, or may range widely, depending on the sex, age, and individual preferences of the animal (Baird 2011). It is not known whether the NZ fur seals around a fishing vessel are from colonies nearby. Some genetic work is proposed to test the potential to differentiate between colonies so that in the future NZ fur seals drowned by fishing gear may be identified as being from a certain colony (Robertson and Gemmell 2005).

The low to moderate levels of observer coverage in some fishery-area strata adds uncertainty to the total estimated captures. However, the main source of uncertainty in the level of bycatch is the paucity of information from the inshore fishing fleets using a variety of methods. Recent increases in observer coverage enabled fur seal capture estimates to include inshore fishing effort. Further increases in coverage, particularly for inshore fisheries, would provide better data on the life stage, sex, and size of captured animals, as well as samples for fatty acid or stable isotope analysis to assess diet and to determine provenance. Information on the aspects of fishing operations that lead to capture in inshore fisheries would also be useful to design mitigation.

#### 4.5. Indicators and trends

Population size	Unknown, but potentially ~100 000 in the New Zealand EEZ <sup>21</sup> .
Population trend	Increasing at some mainland colonies but unknown for offshore island colonies. Range is thought to be increasing.
Threat status	NZ: Not Threatened, Increasing, Secure Overseas, in 2010 <sup>22</sup> . IUCN: Least Concern, in 2008 <sup>23</sup> .
Number of interactions <sup>24</sup>	376 estimated captures (95%CI: 221-668) in trawl fisheries in 2010-11 57 estimated captures (95%CI: 26-99) in surface longline fisheries in 2010-11 69 observed captures in trawl fisheries in 2010-11 17 observed captures in surface longline fisheries in 2010-11
Trends in interactions	<p>Trawl fisheries:</p>  <p>Surface longline fisheries:</p> 

<sup>21</sup> Taylor (1990), Harcourt (2001).

<sup>22</sup> Baker *et al.* (2010b).

<sup>23</sup> Goldsworthy and Gales (2008).

<sup>24</sup> For more information, see: <http://data.dragonfly.co.nz/psc/>.

## 4.6. References

- Abraham, E.R.; Thompson, F.N. (2011). Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2008–09. Final Research Report prepared for Ministry of Fisheries project PRO2007/01. 170 pages. (Unpublished report held by the Ministry of Fisheries, Wellington.)
- Abraham, E.R.; Thompson, F.N.; Oliver, M.D. (2010a). Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 45*. 148 p.
- Abraham, E.R.; Berkenbusch, K.N.; Richard, Y. (2010b). The capture of seabirds and marine mammals in New Zealand non-commercial fisheries. *New Zealand Aquatic Environment and Biodiversity Report No. 64*. 52 p.
- Allum, L.L.; Maddigan, F.W. (2012). Unusual stability of diet of the New Zealand fur seal (*Arctocephalus forsteri*) at Banks Peninsula, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 46: 91–96.
- Baird, S.J. (Comp.) (1994). Nonfish species and fisheries interactions working group report. New Zealand Fisheries Assessment Working Group Report 94/1. Ministry of Fisheries. [Unpublished report held in the NIWA library.] 54 p.
- Baird, S.J. (2005). Review of observer comments that relate to captures of New Zealand fur seals (*Arctocephalus forsteri*) during hoki (*Macruronus novaezelandiae*) trawl fishery operations. *NIWA Client Report WLG205-43*. (Report prepared for the Hoki Fishery Management Company) 27 p.
- Baird, S.J. (2008). Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in longline fisheries in New Zealand waters, 1994–95 to 2005–06. *New Zealand Aquatic Environment and Biodiversity Report No. 20*. 21 p.
- Baird, S.J. (2011). New Zealand fur seals — summary of current knowledge. *New Zealand Aquatic Environment and Biodiversity Report No. 72*. 50 p.
- Baird, S.J.; Bradford, E. (2000). Factors that may have influenced the capture of New Zealand fur seals (*Arctocephalus forsteri*) in the west coast South Island hoki fishery, 1991–98. *NIWA Technical Report 92*. 35 p.
- Baird, S.J.; Smith, M.H. (2007). Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in commercial fisheries in New Zealand waters, 2003–04 to 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No. 14*. 98 p.
- Baker, B.; Jensz, K.; Cawthorn, M.; Cunningham, R. (2010a). Census of New Zealand fur seals on the west coast of New Zealand's South Island. [Report to the Deepwater Group Ltd., available at [www.latitude-42.com.au](http://www.latitude-42.com.au).] 22 p.
- Baker, C.S., B.L. Chilvers, B.L., R. Constantine, R.; S. DuFresne, S.; R. Mattlin, R.; A. van Helden, A.; and R. Hitchmough, R. (2010b). Conservation status of New Zealand marine mammals (suborders Cetacea and Pinnipedia), 2009. *New Zealand Journal of Marine and Freshwater Research* 44: 101–115.
- Best, H. (2011). West Coast South Island monitoring of NZ fur seal pup numbers, 1991–2010 – a summary. Unpublished report for the Department of Conservation. 5 p.
- Boren, L.J. (2005). New Zealand fur seals in the Kaikoura region: colony dynamics, maternal investment and health. A thesis submitted in partial fulfillment of the requirements for the degree of Doctor of Philosophy in Biological Sciences at the University of Canterbury, University of Canterbury. 261 p.
- Boren, L. (2010). Diet of New Zealand fur seals (*Arctocephalus forsteri*): a summary. *DOC Research & Development Series* 319. Department of Conservation, Wellington. 19 p.
- Boren, L.; Morrissey, M.; Gemmell, N.J. (2008). Motor vehicle collisions and the New Zealand fur seal in the Kaikoura region. *Marine Mammal Science* 24(1): 235–238.
- Boren, L.J.; Morrissey, M.; Muller, C.G.; Gemmell, N.J. (2006). Entanglement of New Zealand fur seals in man-made debris at Kaikoura, New Zealand. *Marine Pollution Bulletin* 52: 442–446.
- Bradshaw, C.J.A.; Barker, R.J.; Harcourt, R.G.; Davis, L.S. (2003a). Estimating survival and capture probability of fur seal pups using multistate mark-recapture models. *Journal of Mammalogy* 84(1): 65–80.
- Bradshaw, C.J.A.; Davis, L.S.; Lalas, C.; Harcourt, R.G. (2000). Geographic and temporal variation in the condition of pups of the New Zealand fur seal (*Arctocephalus forsteri*): evidence for density dependence and differences in the marine environment. *Journal of Zoology* 252: 41–51.
- Bradshaw, C.J.A.; Harcourt, R.G.; Davis, L.S. (2003b). Male-biased sex ratios in New Zealand fur seal pups relative to environmental variation. *Behavioral Ecology and Sociobiology* 53(5): 297–307.
- Bradshaw, C.J.A.; Lalas, C.; McConkey, S. (1998). New Zealand sea lion predation on New Zealand fur seals. *New Zealand Journal of Marine and Freshwater Research* 32: 101–104.
- Bradshaw, C.J.A.; Thompson, C. M.; Davis, L.S.; Lalas, C. (1999). Pup density related to terrestrial habitat use by New Zealand fur seals. *Canadian Journal of Zoology* 77(10): 1579–1586.
- Carey, P. W. (1992). Fish prey species of the New Zealand fur seal (*Arctocephalus forsteri* Lesson). *New Zealand Journal of Ecology* 16(1): 41–46.
- Carey, P.W. (1998). New Zealand fur seals (*Arctocephalus forsteri*) at the Snares Islands: a stabilised population? *New Zealand Journal of Marine and Freshwater Research* 32(1): 113–118.
- Chilvers, B.L.; Wilson, K.J.; Hickling, G.J. (1995). Suckling behaviours and growth rates of New Zealand fur seals, *Arctocephalus forsteri*, at Cape Foulwind, New Zealand. *New Zealand Journal of Zoology* 22: 263–270.
- Clement and Associates (2009). Mitigating incidental captures of fur seals in trawl fisheries. Final Report prepared for Department of Conservation project MIT2006/09. 45 p.
- Crawley, M.C.; Wilson, G.J. (1976). The natural history and behaviour of the New Zealand fur seal (*Arctocephalus forsteri*). *Tuatara* 22(1): 1–29.
- Dickie, G.S.; Dawson, S.M. (2003). Age, growth, and reproduction in New Zealand fur seals. *Marine Mammal Science* 19(1): 173–185.
- Dans, S.L.; Alonso, M.K.; Crespo, E.A.; Pedraza, S.N.; Garcia, N.A. (2003). Interactions between marine mammals and high sea fisheries in Patagonia: an integrated approach. pp. 100–115, in Gales, N.; Hindell, M.; Kirkwood, R. (Eds.) *Marine mammals fisheries tourism and management issues*. CSIRO Publishing. 446p.
- Dix, B. (1993) A new record this century of a breeding colony in the North Island for the New Zealand fur seal *Arctocephalus forsteri*. *Journal of the Royal Society of New Zealand* 23: 1–4.
- Dowell, S.A.; Boren, L.J.; Negro, S.; Muller, C.G.; Caudron, A.K.; Gemmell, N.J. (2008). Rearing two New Zealand fur seal (*Arctocephalus forsteri*) pups to weaning. *Australian Journal of Zoology* 56: 33–39.
- Duignan, P.J. (2003). Disease investigations in stranded marine mammals, 1999–2002. *DOC Science Internal Series 104*. Department of Conservation, Wellington. 32 p.
- Duignan, P.J.; Jones, G.W. (2007). Autopsy of pinnipeds incidentally caught in commercial fisheries, 2002/03 and 2003/04. *DOC Research & Development Series* 280. Department of Conservation, Wellington. 41 p.

- Fea, N.I.; Harcourt, R.; Lalas, C. (1999). Seasonal variation in the diet of New Zealand fur seals (*Arctocephalus forsteri*) at Otago Peninsula, New Zealand. *Wildlife Research* 26: 147–160.
- D.; Bulman, C.; He, X.; Lecombe, J.; Litman, C. (2003). Trophic interactions between marine mammals and Australian fisheries: an ecosystem approach, pp. 62–99. In Gales, N.; Hindell, M.; Kirkwood, R. (Eds.), *Marine mammals — fisheries, tourism and management issues*. CSIRO Publishing. 446 p.
- Goldsworthy, S.D.; Shaughnessy, P.D. (1994). Breeding biology and haul-out pattern of the New Zealand fur seal, *Arctocephalus forsteri*, at Cape Gantheaume, South Australia. *Wildlife Research* 21: 365–376.
- Goldsworthy, S.; Gales, N. (2008). *Arctocephalus forsteri*. In: IUCN 2010. IUCN Red List of Threatened Species. Version 2011.1. www.iucnredlist.org. Downloaded on 21 June 2011.
- Hamer, D.J.; Goldsworthy, S.D. (2006). Seal–fishery operational interactions: Identifying the environmental and operational aspects of a trawl fishery that contribute to bycatch and mortality of Australian fur seals (*Arctocephalus pusillus doriferus*). *Biological Conservation* 130: 517–529.
- Harcourt, R.G. (2001). Advances in New Zealand mammalogy 1990–2000: Pinnipeds. *Journal of the Royal Society of New Zealand* 31(1): 135–160.
- Harcourt, R. G.; Bradshaw, C.J.A.; Dickson, K.; Davis, L.S. (2002). Foraging ecology of a generalist predator, the female New Zealand fur seal. *Marine Ecology Progress Series* 227: 11–24.
- Harcourt, R.G.; Schulman, A.; Davis, L.S.; Trillmich, F (1995). Summer foraging by lactating female New Zealand fur seals (*Arctocephalus forsteri*) off Otago Peninsula, New Zealand. *Canadian Journal of Zoology* 73: 678–690.
- Holborrow, J. (1999). The diet of New Zealand fur seals (*Arctocephalus forsteri*) in Southern New Zealand. [Unpublished MSc thesis, University of Otago, New Zealand.]
- Hooper, J., J.M. Clark, C. Charman and D. Agnew. (2005). Sea mitigation measures on trawl vessels fishing for krill in CCAMLR Subarea 48.3. *CCAMLR Science* 12: 195–205.
- IUCN, (2010). IUCN Red List of Threatened Species. Version 2011.1. www.iucnredlist.org. Downloaded on 21 June 2011.
- Lalas, C. (2008). Recolonisation of Otago, southern New Zealand, by fur seals and sea lions: unexpected patterns and consequences, 15 p. In Clarkson, B.; Kurian, P.; Nachowitz, T.; Rennie, H. (Eds.), *Proceedings of the Conser-Vision Conference*, University of Waikato, available at www.waikato.ac.nz/wfass/conserv-vision.
- Lalas, C.; Bradshaw, C.J.A. (2001). Folklore and chimerical numbers: review of a millennium of interaction between fur seals and humans in the New Zealand region. *New Zealand Journal of Marine and Freshwater Research* 35(3): 477–497.
- Lalas, C.; Harcourt, R. (1995). Pup production of the New Zealand fur seal on the Otago Peninsula, New Zealand. *Journal of the Royal Society of New Zealand* 25(1): 81–88.
- Lalas, C.; Murphy, B. (1998). Increase in the abundance of New Zealand fur seals at the Catlins, South Island, New Zealand. *Journal of the Royal Society of New Zealand* 28(2): 287–294.
- Lento, G.M., R.H. Mattlin. G.K. Chambers and C. S. Baker. (1994). Geographic distribution of mitochondrial cytochrome b DNA haplotypes in New Zealand fur seals (*Arctocephalus forsteri*). *Canadian Journal of Zoology* 72: 293 – 299.
- Ling JK, (2002). Impact of colonial sealing on seal stocks around Australia, New Zealand and subantarctic islands between 150 and 170 degrees East. *Australian Mammalogy* 24: 117–126.
- Mattlin, R.H. (1978a). Population biology, thermoregulation and site preference of the New Zealand fur seal (*Arctocephalus forsteri* (Lesson, 1828) on the Open Bay Islands, New Zealand. Unpubl. PhD Thesis, Christchurch, University of Canterbury.
- Mattlin, R.H. (1978b). Pup mortality of the New Zealand fur seal (*Arctocephalus forsteri* Lesson). *New Zealand Journal of Ecology* 1: 138–144.
- Mattlin, R.H. (1981). Pup growth of the New Zealand fur seal, *Arctocephalus forsteri*, on the Open Bay Islands, New Zealand. *Journal of Zoology (London)* 193: 305–314.
- Mattlin, R.H. (1987). New Zealand fur seal, *Arctocephalus forsteri*, within the New Zealand region. In Croxall, J.P.; Gentry, R.L. Status, biology, and ecology of fur seals: Proceedings of an international symposium and workshop, Cambridge, England, 23–27 April 1984. *NOAA Technical Report NMFS-51*.
- Mattlin, R.H. (1994a). Incidental catch of fur seals in the west coast South Island hoki trawl fishery, 1989–92. New Zealand Fisheries Assessment Research Document 93/19. 18 p. (Unpublished report available at NIWA library, Wellington.)
- Mattlin, R.H. (Comp. & Ed.)(1994b). Seals and sea birds–fisheries interactions: report of a workshop, Wellington, 1992. *New Zealand Fisheries Occasional Publication No. 8*. 18 p. plus appendices.
- Mattlin, R.H.; Gales N.J.; Costa, D.P. (1998). Seasonal dive behaviour of lactating New Zealand fur seals (*Arctocephalus forsteri*). *Canadian Journal of Zoology* 76(2): 350–360.
- Mellina, E.; Cawthorn, M. (2009). New Zealand fur seal (*Arctocephalus forsteri*) population assessment in Fiordland Sounds. Draft Report. Presented to AEWG July 7 2009, Wellington, New Zealand. 31p.
- Miller, E.H. 1975. Body and organ measurements of fur seals, *Arctocephalus forsteri* (Lesson), from New Zealand. *Journal of Mammalogy* 56(2): 511–513.
- Moore, J.E.; Wallace, B.P.; Lewison, R.L.; Žydelis, R.; Cox, T.M.; Crowder, L.B. (2009). A review of marine mammal, sea turtle and seabird bycatch in USA fisheries and the role of policy in shaping management. *Marine Policy* 33: 435–451.
- Mormede, S.; Baird, S.J.; Smith, M.H. (2008). Factors that may influence the probability of fur seal capture in selected New Zealand fisheries. *New Zealand Aquatic Environment and Biodiversity Report No. 19*. 42 p.
- Norman, F.I. (2000). Preliminary investigation of the bycatch of marine birds and mammals in inshore commercial Fisheries, Victoria, Australia. *Biological Conservation* 92:217–226.
- Richard, Y., Abraham, E. R., & Filippi, D. (2011). Assessment of the risk to seabird populations from New Zealand commercial fisheries. Final Research Report for research projects IPA2009-19 and IPA2009-20. (Unpublished report held by Ministry of Fisheries, Wellington.). 66 pages.
- Richards, R. 1994: "The upland seal" of the Antipodes and Macquarie Islands: a historian's perspective. *Journal of The Royal Society of New Zealand* 24:289–295.
- Robertson, B.C.; Gemmell, N.J. (2005). Microsatellite DNA markers for the study of population structure in the New Zealand fur seal *Arctocephalus forsteri*. *DOC Science Internal Series* 196. Department of Conservation, Wellington. 18 p.
- Rowe, S.J. (2007). A review of methodologies for mitigating incidental catch of protected marine mammals. *DOC Research and Development Series* 283. 47p. Science and Technical Publishing . DOC Wellington.
- Rowe, S.J. (2009). Conservation Services Programme observer report: 1 July 2004 to 30 June 2007. *DOC Marine Conservation Services Series 1*. Department of Conservation, Wellington. 93 p.
- Shaughnessy, P.D. (1999). The action plan for Australian seals. Environment Australia. 62 p.
- Shaughnessy, P.D.; Davenport, S.R. (1996). Underwater videographic observations and incidental mortality of fur seals around fishing equipment in south-eastern Australia. *Marine and Freshwater Research* 47: 553–556.

## AEBAR 2012: Protected species: Fur seals

- Shaughnessy, P.D.; Gales, N.J.; Dennis, T.E.; Goldsworthy, S.D. (1994). Distribution and abundance of New Zealand fur seals, *Arctocephalus forsteri*, in South Australia and Western Australia. *Wildlife Research* 21(6): 667–695.
- Shaughnessy, P.D.; Payne, A.I.L (1979). Incidental mortality of Cape fur seals during trawl fishing activities in South African waters. *Fisheries Bulletin, South Africa* 12: 20–25.
- Smith, I.W.G. (1989). Maori impact on the marine megafauna: pre-European distributions of New Zealand sea mammals, pp. 76–108. In Sutton, D.G. (Ed.) “Saying so doesn’t make it so”, papers in honour of B. Foss Leach. New Zealand Archaeological Association, Dunedin.
- Smith, I.W.G. (2005). Retreat and resilience: fur seals and human settlement in New Zealand, pp. 6–18. In Monks, G. (Ed.), “The Exploitation and Cultural Importance of Sea Mammals”. Oxbow Books, Oxford. 173 p.
- Smith, I.W.G. (2011). Estimating the magnitude of pre-European Maori marine harvest in two New Zealand study areas. *New Zealand Aquatic Environment and Biodiversity Report* No. 82.
- Smith, M.H.; Baird, S.J. (2009). Model-based estimation of New Zealand fur seal (*Arctocephalus forsteri*) incidental captures and strike rates for trawl fishing in New Zealand waters for the years 1994–95 to 2005–06. *New Zealand Aquatic Environment and Biodiversity Report* No. 40. 90 p.
- Sorensen, J.H. (1969). New Zealand seals with special reference to the fur seal. *Fisheries Technical Report* No. 39. N.Z. Marine Department. 35 p.
- Suisted, R.; Neale, D. M. (2004). *Department of Conservation Marine Mammal Action Plan for 2005-2010*. Department of Conservation: Wellington.
- Taylor, R.H. (1990). Records of subantarctic fur seals in New Zealand (Note). *New Zealand Journal of Marine and Freshwater Research* 24: 499–502.
- Taylor, R.H. (1996). Distribution, abundance and pup production of the New Zealand fur seal (*Arctocephalus forsteri* Lesson) at the Bounty Islands. *Science for Conservation* No. 32. 14 p.
- Taylor, R.H.; Barton, K.J.; Wilson, P.R.; Thomas, B.W.; Karl, B.J. (1995). Population status and breeding of New Zealand fur seals (*Arctocephalus forsteri*) in the Nelson-northern Marlborough region. *New Zealand Journal of Marine and Freshwater Research* 29(2): 223–234.
- Thompson, F.N.; Abraham, E.R. (2010). Estimation of fur seal (*Arctocephalus forsteri*) bycatch in New Zealand trawl fisheries, 2002–03 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report* No. 61. 37 p.
- Thompson, F.N., E.R. Abraham., and K. Berkenbusch. 2011. Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2009–10. Draft Final Research Report for Ministry for Primary Industries project PRO2010-01 (Unpublished report held by the Ministry for Primary Industries, Wellington). 80 pages.
- Thompson, F.N., E.R. Abraham., and K. Berkenbusch. 2012. Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2010–11. Draft Final Research Report for Ministry for Primary Industries project PRO2010-01 (Unpublished report held by the Ministry for Primary Industries, Wellington). 90 pages.
- Troy, S.K., R. Mattlin, P.D. Shaughnessy and P.S. Davie. (1999). Morphology, age and survival of adult male New Zealand fur seals, *Arctocephalus forsteri*, in South Australia. *Wildlife Research* 26: 21–34.
- Wade, P.R. 1998. Calculating limits to the allowable human caused mortality of cetaceans and pinnipeds. *Marine Mammal Science* 14: 1–37.
- Wilson, G. J. (1981). Distribution and abundance of the New Zealand fur seal, *Arctocephalus forsteri*. *Fisheries Research Division Occasional Publication* No. 20. 39 p.
- Willis, T.J.; Triossi, F.; Meynier, L. (2008). Diet of fur seals *Arctocephalus forsteri* at Tonga Island, Abel Tasman National Park. NIWA Client Report: NEL2008-011 12 p. prepared for the Department of Conservation and available at <http://www.doc.govt.nz/upload/documents/conservation/marine-and-coastal/marine-protected-areas/tonga-island-seal-diet.pdf>.

## 5. New Zealand seabirds

<i>Scope of chapter</i>	This chapter focuses on estimates of captures and risk assessments conducted for seabirds that breed in New Zealand waters. Also included are descriptions of the nature of fishing interactions, the management context and approach, trends in key indicators and major sources of uncertainty. It does not include detail on the biology or response of individual seabird species other than those four taxa for which quantitative population modelling has been conducted.
<i>Area</i>	New Zealand EEZ and Territorial Sea (noting that many seabirds are highly migratory and spend prolonged periods outside the NZ EEZ; on the high seas these effects are considered by CCSBT, WCPFC, CCAMLR, SPRFMO, etc. and New Zealand capture estimates are reported to those organisations).
<i>Focal localities</i>	Interactions with fisheries occur in many parts of the EEZ and TS.
<i>Key issues</i>	Quantitative and semi-quantitative risk assessments can be improved through better estimates of: incidental captures in fisheries that are poorly or un-observed; species identity, especially of birds released alive; cryptic mortality rates; survival of birds released alive; and the ability of seabird populations to sustain given levels of bycatch, especially given fisheries interactions and captures outside the New Zealand EEZ and in non-commercial fisheries. Consolidating qualitative and (semi) quantitative risk assessments is a key challenge.
<i>Emerging issues</i>	Assessing fisheries impacts in the context of other factors influencing seabird survival and reproduction, including other anthropogenic effects. Magnitude of “deck strike” mortality.
<i>MFish Research (current)</i>	PRO2006-01 <i>Demographic, distributional and trophic information on selected seabird species</i> ; PRO2006-02 <i>Modelling the effects of fishing on selected seabird species</i> ; PRO2010-01 <i>Estimating incidental captures of protected species</i> ; PRO2010-02 <i>Addressing key areas of uncertainty (including in risk assessments) for a revised NPOA-seabirds</i> .
<i>Other Govt Research (current)</i>	DOC Conservation Services Programme (CSP) projects: INT2012-01, <i>Observing commercial fisheries</i> ; INT2010-02, <i>Identification of seabirds captured in New Zealand fisheries</i> ; POP2011-02, <i>Flesh-footed shearwater population study trial and at-sea distribution</i> ; POP2012-03, <i>Black petrel at-sea distribution and population estimate</i> ; POP2012-04, <i>Campbell Island and grey-headed albatrosses population estimates</i> ; POP2012-05, <i>White-capped albatross population estimate</i> ; POP2012-06, <i>Salvin’s albatross population estimate and at-sea distribution</i> ; POP2012-07, <i>Gibson’s albatross population estimate</i> ; POP2012-08, <i>Pitt Island shags foraging ecology</i> ; MIT2012-01, <i>Inshore bottom longline seabird mitigation design and analysis</i> ; MIT2012-02, <i>Inshore trawl warp-strike mitigation analysis of effectiveness</i> ; MIT2012-03, <i>Review of mitigation techniques in setnet fisheries</i> ; MIT2012-04, <i>Surface longline seabird mitigation</i> ; MIT2012-05, <i>Protected species bycatch newsletter</i>
<i>Links to 2030 objectives</i>	Objective 6: Manage impacts of fishing and aquaculture. Strategic Action 6.2: Set and monitor environmental standards, including for threatened and protected species and seabed impacts.
<i>Related chapters/issues</i>	National Plan of Action to reduce the incidental catch of seabirds in New Zealand fisheries

## 5.1. Context

Seabird names and taxonomy in this document generally follow that adopted by the Ornithological Society of New Zealand (OSNZ 2010) except where a different classification has been agreed by the parties to the Agreement for the Conservation of Albatrosses and Petrels, ACAP, or the New Zealand Threat Classification Scheme (NZTCS) classifies multiple taxa within a single OSNZ species (Table 5.1). The key exceptions to the OSNZ (2010) classification are for: white-capped albatross (OSNZ cites a subspecies *Thalassarche cauta steadi* whereas full species status is used here following ACAP); blue penguins (OSNZ cites a single species, little penguin *Eudyptula minor*, whereas multiple sub-species are used here to reflect NZTCS); and OSNZ (2010) and white-fronted tern (OSNZ cite a single species *Sterna striata*, whereas multiple sub-species are use here to reflect NZTCS). Southern and northern Buller's albatrosses are treated as separate taxa here, although ACAP lists a single species "Buller's albatross". The taxonomy and common names adopted here will, therefore, differ in some instances from those used in legislation or other documents.

There are about 140 000 bird species worldwide, but fewer than 400 are classified as seabirds (being specialised marine foragers). All but seven seabird taxa in New Zealand are absolutely protected under s.3 of the Wildlife Act 1953, meaning that it is an offence to hunt or kill them. Southern black-backed gull, *Larus dominicanus*, is the only species that is not protected. Black shag, *Phalacrocorax carbo*, and sea hawk, *Catharacta lonnbergi*, are partially protected, and sooty shearwater, *Puffinus griseus*, grey-faced petrel, *Pterodroma macroptera*, little shag, *Phalacrocorax melanoleucus brevirostris*, and pied shag, *Phalacrocorax varius*, may be hunted or killed subject to Minister's notification. Of the 85 seabird taxa that breed in New Zealand waters, 47 are considered threatened (by far the largest number on the world). For albatrosses and petrels, a key threat is injury or death in fishing operations, although the Wildlife Act provides defences if the death or injury took place as part of a fishing operation or if all reasonable steps to avoid the death or injury were taken, as long as the interaction is reported. Commercial fishers are required to complete a Non-Fish and Protected Species Catch Return (NFPSR, s11E of the Fisheries (Reporting) Regulations 2001).

Relevant, high level guidance from the 2005 statement of General Policy under the Conservation Act 1987 and Wildlife Act 1953 includes the following stated policies:

- 4.4 (f) Marine protected species should be managed for their long-term viability and recovery throughout their natural range.
- 4.4 (g) Where unprotected marine species are identified as threatened, consideration will be given to amending the Wildlife Act 1953 schedules to declare such species absolutely protected.
- 4.4 (j) Human interactions with marine mammals and other marine protected species should be managed to avoid or minimise adverse effects on populations and individuals.
- 4.4 (l) The Department should work with other agencies and interests to protect marine species.

The Minister of Conservation may approve a Population Management Plan (PMP) for one or more species under s.14F of the Wildlife Act and a PMP can include a maximum allowable level of fishing-related mortality for a species (MALFiRM). Such a limit would apply to New Zealand fisheries waters and would be for the purpose of enabling a threatened species to achieve a non-threatened status as soon as reasonably practicable or, in the case of non-threatened species, neither cause a net reduction in the size of the population nor seriously threaten the reproductive capacity of the species (s.14G). No PMPs are in place for seabirds but, in the absence of a PMP, the Minister of Fisheries (Primary Industries) may, after consultation with the Minister of Conservation, take such measures as they consider necessary to avoid, remedy, or mitigate the effect of fishing-related mortality on any protected species (s.15(2) of the Fisheries Act).

New Zealand is a signatory to a number of international conventions and agreements to provide for the management of threats to seabirds, including:

- the United Nations Convention on the Law of the Sea (UNCLOS);
- the United Nations Fish Stocks Agreement (insofar as it relates to the conservation of non-target, associated and dependent species);
- the Convention on Biological Diversity (CBD);
- the Convention on Migratory Species (CMS);
- the Food and Agriculture Organisation's (FAO) International Plan of Action for Reducing the Incidental Catch of Seabirds in Longline Fisheries (IPOA);
- the FAO Code of Conduct for Responsible Fisheries and the interpretive Best Practice Technical Guidelines;
- the Agreement on the Conservation of Albatrosses and Petrels (ACAP)

The ACAP agreement requires that parties achieve and maintain a favourable conservation status for a number of albatross and petrel taxa. Under the IPOA-seabirds, New Zealand developed a National Plan of Action (NPOA) to reduce the incidental catch of seabirds in New Zealand fisheries in 2004 (MFish and DOC 2004) and recently (2012) consulted on a revised NPOA-seabirds (<http://www.fish.govt.nz/en-nz/Consultations/npoa+seabirds/default.htm>). The scopes of the 2004 NPOA (and the 2012 draft) are broader than the original IPOA to facilitate a co-ordinated and long-term approach to reducing the impact of fishing activity on seabirds.

Management of fishing-related mortality of seabirds is consistent with Fisheries 2030 Objective 6: *Manage impacts of fishing and aquaculture*. Further, the management actions follow Strategic Action 6.2: *Set and monitor environmental standards, including for threatened and protected species and seabed impacts*.

All National Fisheries Plans except that for freshwater fisheries are relevant to the management of fishing-related mortality of seabirds.

Under the National Deepwater Plan, the objective most relevant for management of seabirds is Management Objective 2.5: *Manage deepwater and middle-depth fisheries to avoid or minimise adverse effects on the long-term viability of endangered, threatened and protected species*.

Management objective 7 of the National Fisheries Plan for Highly Migratory Species (HMS) is to *“Implement an ecosystem approach to fisheries management, taking into account associated and dependent species”*. This comprises four components: *Avoid, remedy, or mitigate the adverse effects of fishing on associated and dependent species, including through maintaining food-chain relationships; Minimise unwanted bycatch and maximise survival of incidental catches of protected species in HMS fisheries, using a risk management approach; Increase the level and quality of information available on the capture of protected species; and Recognise the intrinsic values of HMS and their ecosystems, comprising predators, prey, and protected species*.

The Environment Objective is the same for all groups of fisheries in the draft National Fisheries Plan for Inshore Finfish and the draft National Fisheries Plan for Inshore Shellfish, to *“Minimise adverse effects of fishing on the aquatic environment, including on biological diversity”*. The draft National Fisheries Plan for Freshwater has the same objective but is unlikely to be relevant to management of fishing-related mortality of seabirds.

**Table 5.1: List of New Zealand seabird taxa, excluding occasional visitors and vagrants, according to the Ornithological Society of New Zealand (OSNZ 2010) unless otherwise indicated (all taxa under the New Zealand Threat Classification System are listed and ACAP taxonomy generally takes precedence). Broad categories of threat status are listed, but comprehensive threat classifications are given by IUCN (<http://www.iucnredlist.org/>) and DOC (<http://www.doc.govt.nz/publications/conservation/nz-threat-classification-system/nz-threat-classification-system-lists-2008-2011/>, see also Miskelly *et al.* 2008, to be updated shortly).**

Common name	Scientific name	DOC category
Wandering albatross	<i>Diomedea exulans</i>	–
Antipodean albatross	<i>Diomedea antipodensis antipodensis</i>	Threatened
Gibson's albatross	<i>Diomedea antipodensis gibsonii</i>	Threatened
Southern royal albatross	<i>Diomedea epomophora</i>	At Risk
Northern royal albatross	<i>Diomedea sanfordi</i>	At Risk
Black-browed albatross	<i>Thalassarche melanophrys</i>	–
Campbell black-browed albatross	<i>Thalassarche impavida</i>	At Risk
Southern Buller's albatross	<i>Thalassarche bulleri</i>	At Risk
Northern Buller's albatross	<i>Thalassarche bulleri platei</i>	At Risk
White-capped albatross	<i>Thalassarche steadi</i> *	Threatened
Salvin's albatross	<i>Thalassarche salvini</i>	Threatened
Chatham Island albatross	<i>Thalassarche eremita</i>	At Risk
Indian yellow-nosed albatross	<i>Thalassarche carteri</i>	–
Grey-headed albatross	<i>Thalassarche chrysostoma</i>	Threatened
Light mantled sooty albatross	<i>Phoebastria palpebrata</i>	At Risk
Flesh-footed shearwater	<i>Puffinus carneipes</i>	Threatened
Wedge-tailed shearwater	<i>Puffinus pacificus</i>	At Risk
Buller's shearwater	<i>Puffinus bulleri</i>	At Risk
Sooty shearwater	<i>Puffinus griseus</i>	At Risk
Short-tailed shearwater	<i>Puffinus tenuirostris</i>	–
Fluttering shearwater	<i>Puffinus gavia</i>	At Risk
Hutton's shearwater	<i>Puffinus huttoni</i>	At Risk
Kermadec little shearwater	<i>Puffinus assimilis kermadecensis</i>	At Risk
North Island little shearwater	<i>Puffinus assimilis haurakiensis</i>	At Risk
Subantarctic little shearwater	<i>Puffinus elegans</i>	At Risk
Northern diving petrel	<i>Pelecanoides urinatrix urinatrix</i>	At Risk
Southern diving petrel	<i>Pelecanoides urinatrix chathamensis</i>	At Risk
Subantarctic diving petrel	<i>Pelecanoides urinatrix exsul</i>	–
South Georgian diving petrel	<i>Pelecanoides georgicus</i>	Threatened
Grey petrel	<i>Procellaria cinerea</i>	At Risk
Black (Parkinson's) petrel	<i>Procellaria parkinsoni</i>	Threatened
Westland petrel	<i>Procellaria westlandica</i>	At Risk
White-chinned petrel	<i>Procellaria aequinoctialis</i>	At Risk
Kerguelen petrel	<i>Lugensa brevirostris</i>	–
Southern Cape petrel	<i>Daption capense capense</i>	–
Snares Cape petrel	<i>Daption capense australe</i>	At Risk
Antarctic fulmar	<i>Fulmarus glacialis</i>	–
Southern giant petrel	<i>Macronectes giganteus</i>	–
Northern giant petrel	<i>Macronectes halli</i>	At Risk
Fairy prion	<i>Pachyptila turtur</i>	At Risk
Chatham fulmar prion	<i>Pachyptila crassirostris crassirostris</i>	At Risk
Lesser fulmar prion	<i>Pachyptila crassirostris flemingi</i>	At Risk
Thin-billed prion	<i>Pachyptila belcheri</i>	–
Antarctic prion	<i>Pachyptila desolata</i>	At Risk
Salvin's prion	<i>Pachyptila salvini</i>	–
Broad-billed prion	<i>Pachyptila vittata</i>	At Risk
Blue petrel	<i>Halobaena caerulea</i>	–
Pycroft's petrel	<i>Pterodroma pycrofti</i>	At Risk
Cook's petrel	<i>Pterodroma cookii</i>	At Risk
Black-winged petrel	<i>Pterodroma nigripennis</i>	–
Chatham petrel	<i>Pterodroma axillaris</i>	Threatened
Mottled petrel	<i>Pterodroma inexpectata</i>	At Risk
White-naped petrel	<i>Pterodroma cervicalis</i>	At Risk
Kermadec petrel	<i>Pterodroma neglecta</i>	At Risk
Grey-faced petrel	<i>Pterodroma macroptera gouldi</i>	–
Chatham Island taiko	<i>Pterodroma magentae</i>	Threatened

# AEBAR 2012: Protected species: Seabirds

White-headed petrel	<i>Pterodroma lessonii</i>	–
Soft-plumaged petrel	<i>Pterodroma mollis</i>	–
Wilson's storm petrel	<i>Oceanites oceanicus</i>	–
Kermadec storm petrel	<i>Pelagodroma albicunius</i>	Threatened
New Zealand storm petrel	<i>Pealeornis maoriana</i>	Threatened
Grey-backed storm petrel	<i>Garrodia nereis</i>	At Risk
New Zealand white-faced storm petrel	<i>Pelagodroma marina maoriana</i>	At Risk
Black-bellied storm petrel	<i>Fregetta tropica</i>	–
White-bellied storm petrel	<i>Fregetta grallaria grallaria</i>	Threatened
Yellow-eyed penguin	<i>Megadyptes antipodes</i>	Threatened
Northern blue penguin**	<i>Eudyptula minor iredalei**</i>	At Risk
Southern blue penguin**	<i>Eudyptula minor minor**</i>	At Risk
Chatham Island blue penguin**	<i>Eudyptula minor chathamensis**</i>	At Risk
White-flipped blue penguin**	<i>Eudyptula minor albosignata**</i>	Threatened
Eastern rockhopper penguin	<i>Eudyptes filholi</i>	Threatened
Fiordland crested penguin	<i>Eudyptes pachyrhynchus</i>	Threatened
Snares crested penguin	<i>Eudyptes robustus</i>	At Risk
Erect-crested penguin	<i>Eudyptes sclateri</i>	At Risk
Red-tailed tropicbird	<i>Phaethon rubricauda</i>	Threatened
Australasian gannet	<i>Morus serrator</i>	–
Masked booby	<i>Sula dactylatra fullageri</i>	Threatened
Black shag	<i>Phalacrocorax carbo novaehollandiae</i>	At Risk
Pied shag	<i>Phalacrocorax varius varius</i>	Threatened
Little black shag	<i>Phalacrocorax sulcirostris</i>	At Risk
Little shag	<i>Phalacrocorax melanoleucos brevirostris</i>	–
Stewart Island shag	<i>Leucocarbo chalconotus</i>	Threatened
King shag	<i>Leucocarbo carunculatus</i>	Threatened
Chatham Island shag	<i>Leucocarbo onslowi</i>	Threatened
Bounty Island shag	<i>Leucocarbo ranfurlyi</i>	Threatened
Auckland Island shag	<i>Leucocarbo colensoi</i>	Threatened
Campbell Island shag	<i>Leucocarbo campbelli</i>	At Risk
Spotted shag	<i>Stictocarbo punctatus punctatus</i>	–
Blue shag	<i>Stictocarbo punctatus oliveri</i>	At Risk
Pitt Island shag	<i>Stictocarbo featherstoni</i>	Threatened
Subantarctic skua	<i>Catharacta antarctica lonnbergi</i>	At Risk
South Polar skua	<i>Catharacta maccormicki</i>	–
Pomarine skua	<i>Stercorarius pomarinus</i>	–
Arctic skua	<i>Stercorarius parasiticus</i>	–
Long-tailed skua	<i>Stercorarius longicaudus</i>	–
Southern black-backed gull	<i>Larus dominicanus dominicanus</i>	–
Red-billed gull	<i>Larus novaehollandiae scopulinus</i>	Threatened
Black-billed gull	<i>Larus bulleri</i>	Threatened
Caspian tern	<i>Hydroprogne caspia</i>	Threatened
White-fronted tern***	<i>Sterna striata striata***</i>	At Risk
Southern white-fronted tern***	<i>Sterna striata aucklandornia***</i>	Threatened
Arctic tern	<i>Sterna paradisaea</i>	–
New Zealand Antarctic tern	<i>Sterna vittata bethunei</i>	At Risk
Eastern little tern	<i>Sternula albifrons sinensis</i>	–
New Zealand fairy tern	<i>Sternula nereis davisae</i>	Threatened
Sooty tern	<i>Onychoprion fuscata serratus</i>	At Risk
Black-fronted tern	<i>Chlidonias albostratus</i>	Threatened
White-winged black tern	<i>Chlidonias leucopterus</i>	–
Brown noddy	<i>Anous stolidus pileatus</i>	–
Black noddy	<i>Anous tenuirostris minutus</i>	At Risk
Grey noddy	<i>Procelsterna cerulea albivittata</i>	At Risk
White tern	<i>Gygis alba candida</i>	Threatened

## Notes:

\* OSNZ (2010) classify New Zealand white-capped albatross as a subspecies *Thalassarche cauta steadi*. Full species status is used here following ACAP.

\*\* OSNZ (2010) classify a single species, little penguin *Eudyptula minor*. Multiple taxa are included here to reflect classification in the New Zealand Threat Classification Scheme.

\*\*\* OSNZ (2010) classify a single species, white-fronted tern *Sterna striata*. Multiple taxa are included here to reflect classification in the New Zealand Threat Classification Scheme.

## **5.2. Biology**

Taylor (2000) provided an excellent summary of the characteristics, ecology, and life history traits of seabirds (defined for the purpose of this document by the list in Table 5.1) which is further summarised here.

All seabirds spend part of their life cycle feeding over the open sea. They have webbed feet, water-resistant feathering to enable them to fully immerse in salt water, and powerful wings or flippers. All have bills with sharp hooks, points, or filters which enable them to catch fish, cephalopods, crustaceans, and plankton. Seabirds can drink saltwater and have physiological adaptations to remove excess salt.

Most seabird taxa are relatively long-lived; most live to 20 years and 30–40 years is typical for the oldest individuals. A few groups, notably albatrosses, can live for 50–60 years. Most taxa have relatively late sexual maturity. Red-billed gull and blue penguin have been recorded nesting as yearlings and diving petrels and yellow-eyed penguins can begin as 2-year-olds, but most seabirds start nesting only at age 3–6 years, and some albatross and petrel taxa delay nesting until 8–15 years old. In these late developers, individuals first return to colonies at 2–6 years old. Richard *et al.* (2011) list values for several demographic parameters that they used for a comprehensive seabird risk assessment. Most seabirds, and especially albatrosses and some petrels, usually return to the breeding colony where they were reared, or nest close-by. Seabirds also have a tendency to mate for long periods with the same partner, and albatross pairs almost always remain together unless one partner fails to return to the colony.

The number of eggs laid varies among families. Albatrosses and petrels lay only one egg per year (sometimes nesting every other year) and do not replace it if it is damaged or lost. Other taxa such as gannets lay one egg but can replace it if the egg is lost. Most penguins lay two eggs but some raise only one chick and eject the second egg; replacement laying is uncommon. Blue penguins, gulls, and terns lay 1–3 eggs and can lay up to three clutches in a year if eggs are damaged or lost. Shags lay 2–5 eggs, can replace clutches, and have several breeding seasons in a year. Incubation in albatrosses and petrels lasts 40–75 days and chick rearing 50–280 days. In gulls and terns, incubation is completed in 20–25 days and chicks fledge in 20–40 days. In general, the lower the potential reproductive output of a taxon, the higher the adult survival rates and longevity.

Some seabirds such as shags, blue penguins, and yellow-eyed penguins live their lives and forage relatively close to where they breed, but many, including most albatrosses and petrels, spend large parts of their lives in international waters or in the waters of other nations far away from their breeding locations. They can travel great distances across oceans during foraging flights and migratory journeys.

## **5.3. Global understanding of fisheries interactions**

Fishing related mortality of seabirds has been recognised as a serious, worldwide issue for only about 20 years (Bartle 1991, Brothers 1991, Brothers *et al.* 1999, Croxall 2008) and the Food & Agriculture Organization of the United Nations (FAO) released its International Plan of Action for reducing incidental catch of seabirds in longline fisheries (IPOA-seabirds) in 1999 (FAO 1999). The IPOA-Seabirds called on countries with (longline) fisheries that interact with seabirds to assess their fisheries to determine if a problem exists and, if so, to develop national plans (NPOA-seabirds) to reduce the incidental seabird catch in their fisheries. Lewison *et al.* (2004) noted that, in spite of the recognition of the problem, few comprehensive assessments of the effects of fishing-related mortality had been conducted in the decade or so after the problem was recognised. They reasoned that: many vulnerable species live in pelagic habitats, making surveys logistically complex and expensive; capture data are sparse; and understanding of the potential for affected populations to sustain

additional mortality is poor. Soykan *et al.* (2008) identified similar questions in a Theme Section published in Endangered Species Research, including: Where is bycatch most prevalent? Which species are taken as bycatch? Which fisheries and gear types result in the highest bycatch of marine megafauna? What are the population-level effects on bycatch species? How can bycatch be reduced?

There has been substantial progress on these questions since 2004. Croxall *et al.* (2012) reviewed the threats to 346 seabird taxa and concluded that: seabirds are more threatened than other comparable groups of birds; that their status has deteriorated faster over recent decades; and that fishing-related mortality is the most pervasive and immediate threat to many albatross and petrels. They listed the principal threats while at sea were posed by commercial fisheries (through competition and mortality on fishing gear) and pollution, and those on land were alien predators, habitat degradation and human disturbance. Direct exploitation, impacts of aquaculture, energy generation operations, and climate change were listed as threats for some taxa or areas where understanding was particularly poor.

Croxall *et al.* (2012) categorise responses to the issue of fishing-related mortality as

- using long-term demographic studies of relevant seabird species, linked to observational and recovery data to identify the cause of population declines (e.g. Croxall *et al.* 1998, Tuck *et al.* 2004, Poncet *et al.* 2006);
- risk assessments, based on spatiotemporal overlap between seabird species susceptible to bycatch and effort data for fisheries likely to catch them (e.g. Waugh *et al.* 2008; Filippi *et al.* 2010; Tuck *et al.* 2011);
- working with multinational and international bodies (e.g. FAO and RFMOs) to develop and implement appropriate regulations for the use of best-practice techniques to reduce or eliminate seabird bycatch and;
- working with fishers (and national fishery organisations) to assist cost-effective implementation of these mitigation techniques.

Seabirds are ranked by the International Union for the Conservation of Nature (IUCN) as the world's most threatened bird grouping (Croxall *et al.* 2012). Globally they face a number of threats to their long term viability, both at their breeding sites and while foraging at sea. Work at the global level on reducing threats at breeding sites is a major focus of the Agreement on the Conservation of Albatrosses and Petrels (ACAP) and, in New Zealand, is a DOC responsibility, but the key threat to seabirds at sea, especially albatrosses and petrels, is incidental capture and death through fishing operations.

Some seabirds do not range far from their breeding or roosting sites and incidental captures of these taxa can be managed by a single jurisdiction. Conversely, conservation of highly migratory taxa such as albatrosses and petrels cannot be achieved by one country acting independently of other nations which share the same populations (e.g., ACAP). Because of this, in recent years countries which share populations of threatened seabirds have sought to take actions on an international level to complement policy and actions taken within their own jurisdictions.

The ICES Working Group on Seabird Ecology agreed (WGSE 2011) that the three most important indirect effects of fisheries on seabird populations were: the harvesting of seabird food; discards as food subsidies; and modification of marine habitats by dredges and trawls. Many seabird prey species are fished commercially (e.g., Furness 2003) or can be impacted indirectly by fishing of larger predators. These relationships are complex and poorly understood but WGSE (2011) agreed that impacts on populations of seabirds were inevitable. Fishery discards and offal have the potential to benefit seabird species, especially those that ordinarily scavenge (Furness *et al.* 1992, Wagner and Boersma 2011). However, discarding can also modify the way in which birds forage for food (e.g., Bartumeus *et al.* 2010; Louzao *et al.* 2011), sometimes with farther-reaching behavioural consequences with negative as well as positive effects. Louzao *et al.* (2011) stated that discards can affect movement patterns (Arcos and Oro 1996), improve reproductive performance (Oro *et al.*, 1997;

1999) and increase survival (Oro and Furness, 2002; Oro *et al.* 2004). Benefits for scavengers and kleptoparasitic taxa (those that obtain food by stealing from other animals) feeding on discards can also have consequent negative impacts on other species, especially diving species, that share breeding sites or are subject to displacement (Wagner and Boersma 2011). Dredging and bottom trawling both affect benthic habitat and fauna (see Rice 2006 and the benthic effects chapter in this document) and WGSE (2011) agreed that this probably affects some seabird populations, although little work has been done in this area.

#### 5.4. State of knowledge in New Zealand

Before the arrival of humans, the absence of mammalian predators in New Zealand made it a relatively safe breeding place for seabirds and large numbers of a wide variety of taxa bred here, including substantial numbers on the main North and South Islands. Today, New Zealand's extensive coastline, numerous inshore and offshore islands (many of them predator free) and surrounding seas and oceans continue to make it an important foraging and breeding ground for about 145 seabird taxa, second only to the USA (GA Taylor, Department of Conservation, personal communication). Roughly 95 of these taxa breed in New Zealand (Figures 5.1 and 5.2; Table 5.2), including the greatest number of albatrosses (14), petrels (32), shags (13) and penguins (9) of any area in the world (Miskelly *et al.* 2008). More than a third are endemic (i.e. breed nowhere else in the world), giving New Zealand by far the largest number of endemic seabird taxa in the world.




Figure 5.1 (after Croxall *et al* 2012). Number of endemic breeding seabird taxa by country.


Some seabirds use New Zealand waters but do not breed here. Some visit here occasionally to feed (e.g. Indian Ocean yellow-nosed albatross and snowy wandering albatross), whereas others are frequent visitors (e.g. short-tailed shearwater and Wilson's storm petrel), sometimes for extended durations (e.g. juvenile giant petrels).

Taylor (2000) lists a wide range of threats to New Zealand seabird taxa including introduced mammals, avian predators (weka), disease, fire, weeds, loss of nesting habitat, competition for nest sites, coastal development, human disturbance, commercial and cultural harvesting, volcanic eruptions, pollution, plastics and marine debris, oil spills and exploration, heavy metals or chemical contaminants, global sea temperature changes, marine biotoxins, and fisheries interactions. Seabirds are caught in trawl, longline, set-net, and, occasionally, other fisheries (e.g. annual assessments by SJ Baird from 1994 to 2005, Baird & Smith 2008, Waugh *et al.* 2008, Abraham *et al.* 2010) and New Zealand released its National Plan of Action to reduce the incidental catch of seabirds (NPOA-seabirds) in 2004. This stated there was, at that time, limited information about the level of incidental

catch and population characteristics of different seabird taxa, and that this made quantifying the overall impact of fishing difficult. A key objective of New Zealand's NPOA-seabirds was to improve this information and gain a better understanding of the impact of incidental catch on seabird taxa. Seabird taxa caught in New Zealand fisheries range in IUCN threat ranking from critically endangered (e.g. Chatham Island shag), to least concern (e.g. flesh-footed shearwater) (e.g., Vie *et al.* 2009).

**Table 5.2 (after Taylor 2000): Number of species (spp.) and taxa of seabirds of different families in New Zealand and worldwide in 2000. Additional taxa may have been recorded since.**

Family	Common name	World breeding		NZ breeding		NZ visitors, vagrants	
		N spp.	N taxa	N spp.	N taxa	N spp.	N taxa
Spheniscidae	Penguins	17	26	6	10	8	10
Gaviidae	Divers, loons	4	6	—	—	—	—
Podicipedidae	Grebes	10	20	2	2	—	—
Diomedidae	Albatrosses	24	24	13	13	7	7
Procellariidae	Petrels, shearwaters	70	109	28	31	20	23
Hydrobatidae	Storm-petrels	20	36	4	5	2	3
Pelecanoididae	Diving petrels	4	9	2	4	—	—
Phaethontidae	Tropicbirds	3	12	1	1	1	1
Pelecanidae	Pelicans	7	12	—	—	1	1
Sulidae	Gannets	9	19	2	2	1	1
Phalacrocoracidae	Shags	39	57	12	13	—	—
Fregatidae	Frigatebirds	5	11	—	—	2	2
Anatidae	Marine ducks	18	27	—	—	—	—
Scolopacidae	Phalaropes	2	2	—	—	2	2
Chionidae	Shearwaters	2	5	—	—	—	—
Stercorariidae	Skuas	7	10	1	1	4	4
Laridae	Gulls	51	78	3	3	—	—
Sternidae	Terns, noddies	43	121	10	11	8	8
Rynchopidae	Skimmers	2	4	—	—	—	—
Alcidae	Auks, puffins	22	45	—	—	—	—
	Total	359	633	84	96	56	62



**Figure 5.2 (from Croxall *et al.* 2012, supplementary material): The number of breeding and resident seabird species by country in each IUCN category (excluding Least Concern). FST, French Southern Territories; SGSSI, South Georgia & South Sandwich Islands; FI(M), Falkland Islands (Malvinas); H&M, Heard Island & McDonald Islands.**

Different taxa and populations face different threats from fishing operations depending on their biological characteristics and foraging behaviours. Biological traits such as diving ability, agility, size, sense of smell, eyesight and diet, foraging factors such as the season and areas they forage, their aggressiveness, the boldness (or shyness) they display in their attraction to fishing activity can all determine their susceptibility to capture, injury, or death from fishing operations. Some fishing methods pose particular threats to some guilds or types of seabirds. For example, penguins are particularly vulnerable to set net operations and large albatrosses appear to be vulnerable to all forms of longlining. The nature and extent of interactions differs spatially, temporally, seasonally and diurnally between sectors, fisheries and between fleets and vessels within fisheries. In 2010/11 the taxa most frequently observed caught in New Zealand commercial fisheries in descending order were white-chinned petrel, sooty shearwater, southern Buller's albatross, white-capped albatross, Salvin's albatross, and flesh footed shearwater, grey petrel, cape petrel, storm petrels, and black petrel.

The management of fisheries to ensure the long-term viability of seabird populations requires an understanding of the risks posed by fishing and other anthropogenic drivers. Several studies have already estimated the number of seabirds caught annually within the New Zealand Exclusive Economic Zone (EEZ) in a range of fisheries (e.g., Baird & Smith 2008, Waugh *et al.* 2008, Abraham *et al.* 2010). In order to evaluate whether the viability of seabird populations is jeopardised by incidental mortality from commercial fishing, the number of annual fatalities needs to be compared with the capacity of the populations to replace those losses; this depends on the size and productivity of each population. Seabirds that breed in New Zealand die as a result of interactions with commercial or recreational fishing operations in waters under New Zealand jurisdiction, through interactions with New Zealand vessels or other nations' vessels on the High Seas and through interactions with commercial, recreational or artisanal fishing operations in waters under the jurisdiction of other states.

Unfortunately, sufficient data to build fully quantitative population models to assess risks and explore the likely results of different management approaches are available for only very few taxa (e.g., Fletcher *et al.* 2008, Francis and Bell 2010, Francis *et al.* 2008, Dillingham and Fletcher 2011). For this reason, broad seabird risk assessments need to rely on expert knowledge (level-1) or to be semi-quantitative (level-2) (Hobday *et al.* 2007). Rowe 2010b described a level-1 seabird risk assessment and Baird and Gilbert (2010) described a semi-quantitative assessment for seabird taxa for which reasonable numbers of observed captures were available. These assessments were based on expert knowledge or not comprehensive and could not be used directly to assess risk for all seabird taxa and fisheries.

#### **5.4.1. Quantifying fisheries interactions**

Information with which to characterise seabird interactions with fisheries comes from a variety of sources. Some is opportunistically collected, whilst other information collection is targeted at specifically describing the nature and extent of seabird captures in fisheries. This section is focussed on the targeted information collection.

Many New Zealand commercial fisheries have MPI observer coverage, much of which is funded by DOC's CSP programme (e.g., Rowe 2009, 2010, Ramm 2011, 2012). Observers generate independent data on the number of captures of seabirds, the number of fishing events observed, and at-sea identification of the seabirds for these fisheries. Commercial fishers are required to provide effort data allowing estimation of the total number of fishing events in a fishery. In combination these data have been used for many years to assess the nature and extent of seabird captures in fisheries (e.g., Abraham *et al.* 2010, Abraham and Thompson 2009a, 2010, 2011 a&b, Ayers *et al.* 2004, Baird 1994, 1995, 1996, 1997, 1999, 2000 a&b, 2001 a&b, 2003, 2004 a–c, 2005, Baird *et al.* 1998, 1999, Baird & Griggs 2004, Thompson and Abraham 2009). Fisher-reported captures (on NFPSR forms available since 1 October 2008) have not been used to estimate total captures because the reported capture rates

are much lower than those reported by independent observers (Abraham and Thompson 2011) and the species identification is less certain. Specimens and photographs (especially for birds released alive) are also collected allowing verification of at-sea identifications (from carcasses or photographs) and description of biological characters (sex, age, condition, etc., available only from carcasses).

In some fisheries observer data are temporally and spatially well stratified, whilst in others data are only available from a spatially select part of the fishery, or a limited part of the year. Where sufficient observer data are available, estimates of total seabird captures in the fishery are calculated. The methods currently used in estimating seabird captures in New Zealand fisheries are described in Abraham and Thompson (2011a). In this context, captures include all seabirds recovered on a fishing vessel except birds that simply land on the deck or collide with a vessel's superstructure, decomposing animals, records of tissue fragments, and birds caught during trips carried out under special permit (e.g., for trials of mitigation methods). Observer coverage has been highly heterogeneous in that some fisheries and areas have had much higher coverage than others. This complicates estimation of the total number of seabirds captured, especially when estimates include more than one fishery, because the distribution of birds and captures is heterogeneous (Figure 5.3).

Abraham and Thompson (2011, available at: [http://fs.fish.govt.nz/Doc/22872/AEBR\\_79.pdf.ashx](http://fs.fish.govt.nz/Doc/22872/AEBR_79.pdf.ashx)) made model-based estimates of captures in New Zealand trawl and longline fisheries for the following taxa or groups: sooty shearwater (*Puffinus griseus*); white-chinned petrel (*Procellaria aequinoctialis*); white-capped albatross (*Thalassarche steadi*); other albatrosses; and all other birds. The three individual species were chosen because they are the most frequently caught in trawl and longline fisheries. Captures of other albatrosses are mostly Salvin's, southern Buller's, Gibson's or Antipodean wandering albatrosses, or Campbell albatrosses. The other birds category includes many taxa but grey, black, great-winged, and Cape petrels, flesh-footed shearwater, and spotted shag are relatively common observed captures (the latter based on few observations that included 31 captures in one event). Estimated captures up to and including the 2010/11 year are shown in Tables 5.3 and 5.4.

Observed captures of seabirds in trawl fisheries were most common off both coasts of the South Island, along the Chatham Rise, on the fringes of the Stewart-Snares shelf, and around the Auckland Islands (Figure 5.4). This largely reflects the distribution of the major commercial fisheries for squid, hoki, and middle-depth species which have tended to have relatively high observer coverage. White-capped, Salvin's, and southern Buller's have been the most frequently observed captured albatrosses, and sooty shearwater and white chinned petrel have been the other species most frequently observed (Table 5.5). About 42% of observed captures were albatrosses.

Observed captures of seabirds in surface longline fisheries were most common off the southwest coast of the South Island and the northeast coast of the North Island (Figure 5.5), again largely reflecting the distribution of the major commercial fisheries (for southern bluefin and other tunas). The charter fleet targeting tuna has historically had much higher observer coverage than the domestic fleet. Southern Buller's and white-capped have been the most frequently observed captured albatrosses, and grey, white-chinned, and black petrels have been the other species most frequently observed (Table 5.6). About 77% of observed captures were albatrosses.

Observed captures of seabirds in bottom longline fisheries were most common off the south coast of the South Island, along the Chatham Rise, scattered throughout the SubAntarctic, and off the northeast coast of the North Island, especially around the Hauraki Gulf (Figure 5.6). This distribution largely reflects the distribution of the ling and snapper longline fisheries that have received most observer coverage; other bottom longline fisheries have had much less coverage. Salvin's and Chatham have been the most frequently observed captured albatrosses, and white chinned petrel, grey petrel, sooty shearwater, and black petrels have been the other species most frequently observed (Table 5.7). Only about 14% of observed captures were albatrosses.




Figure 5.3 (reproduced from Abraham and Thompson 2011): All observed seabird captures in trawl, surface longline, and bottom longline fishing within the New Zealand region, between October 2008 and September 2009. The colour within each 0.2 degree cell indicates the number of fishing events (tows and sets, darker colours indicate more fishing) and the black dots indicate the number of observed events (larger dots indicate more observations). The coloured symbols indicate the location of observed seabird captures, randomly jittered by 0.2 degrees. The 500 m and 1000 m depth contours are shown.

**Table 5.3: Summary of observed and model-estimated total captures of all seabirds (top half) and white-capped albatross (bottom half) by October fishing year in trawl (BT, effort in tows), surface longline (SLL, effort in hooks) and bottom longline (BLL, effort in hooks) fisheries between 2002–30 and 2010–11. Observed and modelled rates are per 100 trawl tows or 1000 longline hooks. Caps, observed captures; % obs, percentage of effort observed; % incl, percentage of total effort included in the model. Data version v20121101.**

Models for all seabirds		Fishing effort		Seabirds			Model estimates		
Year	All effort	Observed	% obs	Caps	Rate	Mean	95% c.i.	% incl	Rate
BT 2002–03	130 344	6 834	5.2	269	3.94	3126	2451–4045	100.0	2.40
BT 2003–04	121 498	6 546	5.4	262	4.00	2624	2034–3456	100.0	2.16
BT 2004–05	120 585	7 709	6.4	483	6.27	4337	3358–5861	100.0	3.60
BT 2005–06	110 234	6 553	5.9	356	5.43	3424	2696–4363	100.0	3.11
BT 2006–07	103 529	7 928	7.7	211	2.66	2027	1559–2678	100.0	1.96
BT 2007–08	89 537	9 046	10.1	234	2.59	1976	1515–2574	100.0	2.21
BT 2008–09	87 589	9 804	11.2	469	4.78	2505	2059–3140	100.0	2.86
BT 2009–10	92 886	9 006	9.7	256	2.85	2176	1672–2882	100.0	2.34
BT 2010–11	86 074	7 445	8.6	370	4.97	2788	2172–3611	100.0	3.24
SLL 2002–03	10764 588	2 195 152	20.4	115	0.05	2349	1735–3271	100.0	0.022
SLL 2003–04	7 380 779	1 607 304	21.8	71	0.04	1582	1212–2064	100.0	0.021
SLL 2004–05	3 676 365	783 812	21.3	41	0.05	660	499–885	100.0	0.018
SLL 2005–06	3 687 339	705 945	19.1	37	0.05	785	589–1062	100.0	0.021
SLL 2006–07	3 738 362	1 040 948	27.8	187	0.18	923	720–1239	100.0	0.025
SLL 2007–08	2 244 339	426 310	19.0	41	0.10	509	397–650	100.0	0.023
SLL 2008–09	3 115 633	937 233	30.1	57	0.06	642	502–814	100.0	0.021
SLL 2009–10	2 992 285	665 883	22.3	135	0.20	903	702–1191	100.0	0.030
SLL 2010–11	3 164 159	674 522	21.3	47	0.07	740	547–1019	100.0	0.023
BLL 2002–03	37 671 038	10 772 020	28.6	296	0.03	1718	1250–2268	89.2	0.005
BLL 2003–04	43 397 540	5 162 608	11.9	54	0.01	1151	761–1604	90.2	0.003
BLL 2004–05	41 818 638	2 883 725	6.9	30	0.01	1191	802–1661	88.0	0.003
BLL 2005–06	37 126 833	3 802 951	10.2	41	0.01	1037	701–1431	87.3	0.003
BLL 2006–07	38 124 470	2 315 772	6.1	58	0.03	1236	833–1716	86.2	0.003
BLL 2007–08	41 464 276	3 589 511	8.7	40	0.01	1193	824–1621	86.0	0.003
BLL 2008–09	37 389 512	4 024 816	10.8	33	0.01	1037	699–1458	86.5	0.003
BLL 2009–10	40 413 281	2 271 623	5.6	68	0.03	1062	716–1474	86.1	0.003
BLL 2010–11	40 826 726	1 730 585	4.2	29	0.02	1403	955–1967	85.8	0.003
<b>White-capped albatross models</b>									
Year	All effort	Observed	% obs	Caps	Rate	Mean	95% c.i.	% incl	Rate
BT 2002–03	130 344	6 834	5.2	85	1.24	861	648–1119	100.0	0.66
BT 2003–04	121 498	6 546	5.4	148	2.26	905	701–1144	100.0	0.74
BT 2004–05	120 585	7 709	6.4	243	3.15	1200	976–1502	100.0	1.00
BT 2005–06	110 234	6 553	5.9	69	1.05	609	439–826	100.0	0.55
BT 2006–07	103 529	7 928	7.7	57	0.72	437	315–591	100.0	0.42
BT 2007–08	89 537	9 046	10.1	42	0.46	312	205–443	100.0	0.35
BT 2008–09	87 589	9 804	11.2	97	0.99	471	352–625	100.0	0.54
BT 2009–10	92 886	9 006	9.7	48	0.53	381	266–527	100.0	0.41
BT 2010–11	86 074	7 445	8.6	39	0.52	356	236–496	100.0	0.41
SLL 2002–03	10764 588	2 195 152	20.4	2	0.00	101	63–149	100.0	0.001
SLL 2003–04	7 380 779	1 607 304	21.8	17	0.01	228	148–325	100.0	0.003
SLL 2004–05	3 676 365	783 812	21.3	3	0.00	58	35–86	100.0	0.002
SLL 2005–06	3 687 339	705 945	19.1	2	0.00	54	32–81	100.0	0.001
SLL 2006–07	3 738 362	1 040 948	27.8	28	0.03	42	32–55	100.0	0.001
SLL 2007–08	2 244 339	426 310	19.0	4	0.01	55	33–81	100.0	0.002
SLL 2008–09	3 115 633	937 233	30.1	3	0.00	78	48–114	100.0	0.003
SLL 2009–10	2 992 285	665 883	22.3	31	0.05	135	94–185	100.0	0.005
SLL 2010–11	3 164 159	674 522	21.3	3	0.00	84	52–123	100.0	0.003
BLL 2002–03	37 671 038	10 772 020	28.6	0	0.00	1	0–4	44.8	0.000
BLL 2003–04	43 397 540	5 162 608	11.9	1	0.00	3	0–7	50.3	0.000
BLL 2004–05	41 818 638	2 883 725	6.9	0	0.00	2	0–6	39.6	0.000
BLL 2005–06	37 126 833	3 802 951	10.2	1	0.00	3	1–6	36.4	0.000
BLL 2006–07	38 124 470	2 315 772	6.1	0	0.00	2	0–5	30.6	0.000
BLL 2007–08	41 464 276	3 589 511	8.7	0	0.00	2	0–6	29.9	0.000
BLL 2008–09	37 389 512	4 024 816	10.8	0	0.00	2	0–5	32.1	0.000
BLL 2009–10	40 413 281	2 271 623	5.6	0	0.00	2	0–6	30.1	0.000
BLL 2010–11	40 826 726	1 730 585	4.2	0	0.00	2	0–5	28.6	0.000

**Table 5.4: Summary of observed and model-estimated total captures of sooty shearwater (top half) and white-chinned petrel (bottom half) by October fishing year in trawl (BT, effort in tows), surface longline (SLL, effort in hooks) and bottom longline (BLL, effort in hooks) fisheries between 2002–30 and 2010–11. Observed and modelled rates are per 100 trawl tows or 1000 longline hooks. Caps, observed captures; % obs, percentage of effort observed; % incl, percentage of total effort included in the model. Data version v20121101.**

Sooty shearwater models		Fishing effort		Seabirds			Model estimates		
Year	All effort	Observed	% obs	Caps	Rate	Mean	95% c.i.	% incl	Rate
BT 2002–03	130 344	6 834	5.2	120	1.76	999	642–1523	100.0	0.77
BT 2003–04	121 498	6 546	5.4	54	0.82	370	224–590	100.0	0.30
BT 2004–05	120 585	7 709	6.4	74	0.96	494	319–758	100.0	0.41
BT 2005–06	110 234	6 553	5.9	169	2.58	976	657–1456	100.0	0.89
BT 2006–07	103 529	7 928	7.7	84	1.06	497	328–748	100.0	0.48
BT 2007–08	89 537	9 046	10.1	82	0.91	416	276–627	100.0	0.46
BT 2008–09	87 589	9 804	11.2	152	1.55	521	371–744	100.0	0.59
BT 2009–10	92 886	9 006	9.7	43	0.48	260	159–409	100.0	0.28
BT 2010–11	86 074	7 445	8.6	110	1.48	488	331–722	100.0	0.57
SLL 2002–03	10 771 388	2 195 152	20.4	8	0.00	14	8–31	100.0	0.000
SLL 2003–04	7 386 864	1 607 304	21.8	3	0.00	7	3–19	100.0	0.000
SLL 2004–05	3 679 865	783 812	21.3	0	0.00	2	0–9	100.0	0.000
SLL 2005–06	3 689 879	705 945	19.1	0	0.00	2	0–9	100.0	0.000
SLL 2006–07	3 739 962	1 040 948	27.8	2	0.00	4	2–10	100.0	0.000
SLL 2007–08	2 245 939	426 310	19.0	0	0.00	2	0–6	100.0	0.000
SLL 2008–09	3 115 633	937 233	30.1	0	0.00	2	0–8	100.0	0.000
SLL 2009–10	2 992 285	665 883	22.3	0	0.00	2	0–7	100.0	0.000
SLL 2010–11	3 166 559	674 522	21.3	0	0.00	2	0–9	100.0	0.000
BLL 2002–03	37 789 058	10 772 020	28.6	32	0.00	97	45–216	100.0	0.000
BLL 2003–04	43 493 500	5 162 608	11.9	17	0.00	82	30–202	100.0	0.000
BLL 2004–05	41 868 788	2 883 725	6.9	3	0.00	81	20–213	100.0	0.000
BLL 2005–06	37 138 783	3 802 951	10.2	3	0.00	46	6–151	100.0	0.000
BLL 2006–07	38 150 820	2 315 772	6.1	1	0.00	53	7–169	100.0	0.000
BLL 2007–08	41 502 096	3 589 511	8.7	6	0.00	61	17–157	100.0	0.000
BLL 2008–09	37 424 356	4 023 916	10.8	0	0.00	54	6–169	100.0	0.000
BLL 2009–10	40 445 221	2 279 233	5.6	7	0.00	53	10–165	100.0	0.000
BLL 2010–11	40 878 991	1 728 765	4.2	0	0.00	69	6–235	100.0	0.000
White-chinned petrel models									
Year	All effort	Observed	% obs	Caps	Rate	Mean	95% c.i.	% incl	Rate
BT 2002–03	130 344	6 834	5.2	13	0.19	148	67–280	100.0	0.11
BT 2003–04	121 498	6 546	5.4	18	0.27	117	61–207	100.0	0.10
BT 2004–05	120 585	7 709	6.4	55	0.71	266	169–403	100.0	0.22
BT 2005–06	110 234	6 553	5.9	70	1.07	436	270–688	100.0	0.40
BT 2006–07	103 529	7 928	7.7	29	0.37	135	82–216	100.0	0.13
BT 2007–08	89 537	9 046	10.1	59	0.65	271	168–430	100.0	0.30
BT 2008–09	87 589	9 804	11.2	104	1.06	316	222–453	100.0	0.36
BT 2009–10	92 886	9 006	9.7	74	0.82	295	189–461	100.0	0.32
BT 2010–11	86 074	7 445	8.6	130	1.75	540	359–817	100.0	0.63
SLL 2002–03	10764 588	2 195 152	20.4	4	0.00	79	43–128	100.0	0.001
SLL 2003–04	7 380 779	1 607 304	21.8	2	0.00	53	27–87	100.0	0.001
SLL 2004–05	3 676 365	783 812	21.3	3	0.00	30	14–49	100.0	0.001
SLL 2005–06	3 687 339	705 945	19.1	1	0.00	30	14–50	100.0	0.001
SLL 2006–07	3 738 362	1 040 948	27.8	5	0.00	30	16–48	100.0	0.001
SLL 2007–08	2 244 339	426 310	19.0	4	0.01	22	11–35	100.0	0.001
SLL 2008–09	3 115 633	937 233	30.1	3	0.00	26	13–43	100.0	0.001
SLL 2009–10	2 992 285	665 883	22.3	3	0.00	25	12–41	100.0	0.001
SLL 2010–11	3 164 159	674 522	21.3	8	0.01	34	19–52	100.0	0.001
BLL 2002–03	37 671 038	10 772 020	28.6	132	0.01	350	246–540	100.0	0.001
BLL 2003–04	43 397 540	5 162 608	11.9	15	0.00	139	81–215	100.0	0.000
BLL 2004–05	41 818 638	2 883 725	6.9	11	0.00	188	105–290	100.0	0.000
BLL 2005–06	37 126 833	3 802 951	10.2	13	0.00	189	108–303	100.0	0.001
BLL 2006–07	38 124 470	2 315 772	6.1	12	0.01	225	123–364	100.0	0.001
BLL 2007–08	41 464 276	3 589 511	8.7	10	0.00	261	143–423	100.0	0.001
BLL 2008–09	37 389 512	4 024 816	10.8	1	0.00	204	97–380	100.0	0.001
BLL 2009–10	40 413 281	2 271 623	5.6	1	0.00	172	86–282	100.0	0.000
BLL 2010–11	40 826 726	1 730 585	4.2	24	0.01	422	225–769	100.0	0.001





Figure 5.4: Map of trawl fishing effort and all observed seabird captures in trawls, October 2003 to September 2011. Fishing effort is mapped into 0.2-degree cells, with the colour of each cell being related to the amount of effort (events). Observed fishing events are indicated by black dots, and observed captures are indicated by red dots. Fishing is shown only if the effort could be assigned a latitude and longitude, and if there were three or more vessels fishing within a cell (here, 96% of effort is displayed).

**Table 5.5: Summary of seabirds observed captured in trawl fisheries 2002–03 to 2010–11. Declared target species are: SQU, arrow squid; HOK+, hoki, hake, ling; Mid., other middle depth species silver, white, and common warehou, barracouta, alfonsinos, stargazer; SCI, scampi; ORH+, orange roughy and oreos; SBW, southern blue whiting; JMA, Jack mackerels; Ins., other inshore species for which one or more captures have been observed; tarakihi, red cod, spiny dogfish, John dory, snapper; FLA, flatfishes. Data version v20121101.**


Species or group	Declared target species									
	SQU	HOK+	Mid.	SCI	ORH+	SBW	JMA	Ins.	FLA	Total
White capped albatross	679	54	52	15	6	0	1	22	0	829
Salvin's albatross	18	87	25	29	16	2	0	20	0	197
Southern Buller's	49	41	19	4	3	0	1	1	0	118
Campbell albatross	2	5	0	1	0	1	0	0	0	9
Chatham Island albatross	0	0	0	1	8	0	0	0	0	9
Southern royal albatross	5	0	0	0	0	1	0	0	0	6
Southern black-browed	1	2	0	0	0	0	0	2	0	5
Gibson's albatross	0	0	0	0	1	0	0	0	0	1
Northern royal albatross	0	0	0	0	1	0	0	0	0	1
Albatross indet.	10	10	1	5	0	4	1	1	0	32
All albatrosses	764	199	97	55	35	8	3	46	0	1207
Sooty shearwater	540	181	119	37	5	0	5	1	0	888
White chinned petrel	387	43	42	48	1	0	9	0	0	530
Cape petrels	1	34	1	3	19	1	2	0	0	61
Flesh footed shearwater	0	1	0	35	0	0	0	2	0	38
Spotted shag	0	0	0	0	0	0	0	0	32	32
Grey petrel	1	2	0	0	3	22	0	0	0	28
Common diving petrel	5	5	0	1	2	0	1	0	0	14
Westland petrel	0	11	1	0	0	0	1	0	0	13
Fairy prion	0	4	0	0	0	0	5	0	0	9
Antarctic prion	7	0	0	0	0	0	0	0	0	7
Northern giant petrel	0	3	1	1	1	0	0	0	0	6
Giant petrel	3	1	0	0	0	0	0	0	0	4
Grey-backed storm petrel	3	1	0	0	0	0	0	0	0	4
Fulmar prion	0	0	0	0	0	0	3	0	0	3
Black petrel	0	0	0	1	0	0	0	1	0	2
Black-bellied storm petrel	1	1	0	0	0	0	0	0	0	2
White-faced storm petrel	0	0	0	0	2	0	0	0	0	2
Black backed gull	0	0	0	0	0	0	0	0	1	1
Short tailed shearwater	0	0	1	0	0	0	0	0	0	1
White headed petrel	1	0	0	0	0	0	0	0	0	1
Other bird indet.	11	5	3	2	1	5	0	2	2	31
All other birds	960	292	168	128	34	28	26	6	35	1677
All observed birds	1724	491	265	183	69	36	29	52	35	2884
Approx. proportion obs	0.23	0.14	0.06	0.09	0.26	0.35	0.25	0.01	0.01	0.08



**Figure 5.5:** Map of surface longline fishing effort and all observed seabird captures by surface longlines, October 2003 to September 2011. Fishing effort is mapped into 0.2-degree cells, with the colour of each cell being related to the amount of effort (events). Observed fishing events are indicated by black dots, and observed captures are indicated by red dots. Fishing is shown only if the effort could be assigned a latitude and longitude, and if there were three or more vessels fishing within a cell (here, 75.3% of effort is displayed).

**Table 5.6: Summary of seabirds observed captured in surface longline fisheries 2002–03 to 2010–11. Declared target species are: SBT, southern bluefin tuna; BIG, bigeye tuna; SWO, broadbill swordfish; ALB, albacore tuna. Data version v20121101.**

Species or group	Declared target species				
	SBT	BIG	SWO	ALB	Total
Southern Buller's albatross	296	7	1	8	312
White capped albatross	91	1	1	0	93
Campbell albatross	18	3	2	17	40
Antipodean albatross	4	8	15	3	30
Gibson's albatross	8	6	9	7	30
Wandering albatrosses	8	3	0	0	11
Salvin's albatross	3	4	0	1	8
Antipodean / Gibson's	0	2	5	0	7
Black browed albatrosses	0	2	2	0	4
Southern royal albatross	4	0	0	0	4
Southern black-browed	2	0	0	0	2
Light-mantled sooty	1	0	0	0	1
Northern royal albatross	0	1	0	0	1
Pacific albatross	1	0	0	0	1
Albatrosses indet.	2	1	33	0	36
Total albatrosses	438	38	68	36	580
Grey petrel	38	0	3	5	46
White chinned petrel	21	8	2	2	33
Black petrel	0	23	2	1	26
Great winged petrel	0	1	2	17	20
Sooty shearwater	4	0	1	8	13
Flesh footed shearwater	0	11	1	0	12
Westland petrel	6	0	0	2	8
Cape petrels	2	0	0	0	2
Southern giant petrel	2	0	0	0	2
White headed petrel	0	0	0	2	2
Petrels indet.	0	1	0	0	1
Total other birds	73	44	11	37	165
All observed birds	511	82	79	73	745
Approx. proportion obs	0.42	0.03	0.10	0.38	0.22



**Figure 5.6: Map of bottom longline fishing effort and all observed seabird captures by bottom longlines, October 2003 to September 2011. Fishing effort is mapped into 0.2-degree cells, with the colour of each cell being related to the amount of effort (events). Observed fishing events are indicated by black dots, and observed captures are indicated by red dots. Fishing is shown only if the effort could be assigned a latitude and longitude, and if there were three or more vessels fishing within a cell (here, 79.3% of effort is displayed).**

**Table 5.7: Summary of seabirds observed captured in bottom longline fisheries 2002–03 to 2010–11. Declared target species are: LIN, ling; SNA, snapper; BNS, bluenose; HPB, hapuku or bass. Data version v20121101.**

Species or group	Declared target species				
	LIN	SNA	BNS	HPB	Total
Salvin's albatross	51	0	0	0	51
Chatham Island albatross	18	0	0	0	18
Southern Buller's albatross	4	0	3	0	7
Campbell albatross	0	0	2	1	3
Wandering albatrosses	2	0	1	0	3
White capped albatross	2	0	0	0	2
Black browed albatrosses	1	0	0	0	1
Indian yellow-nosed albatross	1	0	0	0	1
Southern royal albatross	1	0	0	0	1
Albatross indet.	2	0	0	0	2
All albatrosses	82	0	6	1	89
White chinned petrel	217	0	2	0	219
Grey petrel	79	0	0	0	79
Sooty shearwater	68	0	0	1	69
Black petrel	0	28	14	7	51
Flesh footed shearwater	0	36	0	3	39
Cape petrels	24	0	0	0	24
Common diving petrel	23	0	0	0	23
Great winged petrel	0	0	0	6	6
Fluttering shearwater	0	4	0	0	4
Northern giant petrel	4	0	0	0	4
Prions	4	0	0	0	4
Storm petrels	3	0	0	0	3
Gannets	0	2	0	0	2
Pied shag	0	2	0	0	2
Black backed gull	0	1	0	0	1
Buller's shearwater	0	1	0	0	1
Crested penguins	1	0	0	0	1
Giant petrel	1	0	0	0	1
Red billed gull	0	1	0	0	1
Other birds indet	1	10	0	0	11
All other birds	425	85	16	17	545
All birds observed	507	85	22	18	634
Approx. proportion obs	0.20	0.01	0.01	0.01	0.10

Model-based estimates of captures can be combined across trawl and longline fisheries (Figure 5.7). Summed across all bird taxa, trawl, surface longline, and bottom longline fisheries account for 55%, 21%, and 24% of captures, respectively, but there are substantial differences in these proportions among seabird taxa. A high proportion (87% between 2003 and 2011) of white-capped albatross captures are taken in trawl fisheries with almost all of the remainder taken in surface longline fisheries. The trawl fishery also accounts for 89% of sooty shearwaters captured, with most of the remainder taken by bottom longliners. The proportion captured by trawl fisheries reduces to 53% for all other albatrosses combined, with 30% and 17% taken in surface and bottom longline fisheries, respectively. Bottom longline and trawl take similar proportions of the white-chinned petrels captured (43% and 50%, respectively).

Over the 2003 to 2011 period, there appear to have been downward trends (across all fisheries) in the estimated captures of all birds combined, white-capped albatross, and non-albatross taxa other than sooty shearwaters and white-chinned petrel (Figure 5.7). Estimated captures of other albatrosses, sooty shearwaters, and white-chinned petrel appear to have fluctuated without much trend, although there is some evidence for an increasing trend for white-chinned petrel, especially in trawl fisheries.


Because fishing effort often changes with time, estimates of total captures may not be the only index required for comprehensive monitoring. The number of captures (with certain caveats, see later) is clearly more biologically relevant for the birds, but capture rates by fishery may be more useful measures to assess fishery performance and the effectiveness of mitigation approaches. Dividing modelled catch estimates by the number of tows or hooks set in a particular fishery in each year provides catch rate indices by fishery. These are typically reported as the number of birds captured per 100 trawl tows or per 1000 longline hooks (Figures 5.8 to 5.10).



**Figure 5.7: Model-based estimates of captures of the most numerous seabird taxa observed captured in trawl, surface longline, and bottom longline fisheries between 2002/03 and 2010/11. For confidence limits see Tables 3 and 4. Note this level of aggregation conceals any different trends within a fishing method (e.g., deepwater vs. inshore and flatfish trawl or large vs. small longliners).**


For white-capped albatross, captures rates declined between 2002/03 and 2010/11, and especially up to 2006/07, in the major offshore trawl fisheries for squid and hoki (Figure 5.8) but showed no trend for inshore trawlers and increased for surface longliners targeting southern bluefin tuna. Together, these fisheries account for 82% of all estimated captures of white-capped albatross in these years.

## White-capped albatross captures and capture rates




**Figure 5.8:** Model-based estimates of captures (left panels) and capture rates (right panels, captures per 100 trawl tows or 1000 longline hooks) of white capped albatross in the four fisheries estimated to have taken the most captures between 2002/03 and 2010/11 (cumulatively, 82% of all white-capped albatross captures). Data version v20121101.

For white-chinned petrel, captures rates increased between 2002/03 and 2010/11 in squid and scampi trawlers (Figure 5.9) but showed no trend for bottom longliners targeting ling and bluenose. Together, these fisheries account for 81% of all estimated captures of white-chinned petrel in these years.



**Figure 5.9:** Model-based estimates of captures (left panels) and capture rates (right panels, captures per 100 trawl tows or 1000 longline hooks) of white chinned petrels in the four fisheries estimated to have taken the most captures between 2002/03 and 2010/11 (cumulatively, 81% of all white-chinned petrel captures). Data version v20121101.

For sooty shearwaters, captures rates decreased between 2002/03 and 2010/11 for bottom longliners targeting ling, but showed no trend in squid, middle-depth, and hoki trawlers (Figure 5.10). Together, these fisheries account for 80% of all estimated captures of sooty shearwaters in these years.



**Figure 5.10: Model-based estimates of captures (left panels) and capture rates (right panels, captures per 100 trawl tows or 1000 longline hooks) of sooty shearwaters in the four fisheries estimated to have taken the most captures between 2002/03 and 2010/11 (cumulatively, 80% of all sooty shearwater captures). Data version v20121101.**

On-board captures recorded by observers represent the most reliable source of information for monitoring trends in total captures and capture rates, but these data have three main deficiencies with respect to estimating total fatalities, especially to species level. First, some captured seabirds are released alive (23% in trawl fisheries between 2002–03 and 2010–11, 29% in surface longline fisheries, and 25% in bottom longline fisheries), meaning that, all else being equal, estimates of captures may overestimate total fatalities, depending on the survival rate of those released. Second, identifications by observers are not completely reliable and sometimes use generic codes rather than species codes. A high proportion of dead captures are returned for necropsy and formal identification (87% in trawl fisheries between 2002–03 and 2010–11, 83% in surface longline fisheries, and 89% in bottom longline fisheries), but there remains uncertainty in the identity of 11–17% of dead captures and 100% of those released alive. Third, not all birds killed or mortally wounded by fishing gear are recovered on a fishing vessel. Some birds caught on longline hooks fall off before being recovered, and birds that collide with trawl warps may be dragged under the water and drowned or injured to the extent that they are unable to fly or feed. Excluding this “cryptic” mortality means that, all else being equal, estimates of captures will underestimate total fatalities. These deficiencies do not greatly affect the suitability of estimates of captures and capture rates for monitoring purposes, but they have necessitated the development of alternative measures for assessing risk and population consequences.

#### 5.4.2. Managing fisheries interactions

New Zealand had taken steps to reduce incidental captures of seabirds before the advent of the IPOA in 1999 and the NPOA in 2004. For example, regulations were put in place under the Fisheries Act to prohibit drift net fishing in 1991 and prohibit the use of netsonde monitoring cables (“third wires”) in trawl fisheries in 1992. The use of tori lines (streamer lines designed to scare seabirds away from baited hooks) was made mandatory in all tuna longline fisheries in 1992.

The fishing industry also undertook several initiatives to reduce captures include funding research into new or improved mitigation measures, and adopting voluntary codes of practice and best practice fishing methods. Codes of practice have been in place in the joint venture tuna longline fishery since 1997–98, requiring, *inter alia*, longlines to be set at night and voluntary upper limits on the incidental catch of seabirds. That limit was steadily reduced from 160 “at risk” seabirds in 1997–98, to 75 in 2003–04. Most vessels in the domestic longline tuna fishery had also voluntarily adopted night setting, by 2004. A code of practice was in place for the ling auto-line fishery by 2002–03. Other early initiatives included reduced deck lighting, the use of thawed rather than frozen baits, sound deterrents, discharging of offal away from setting and hauling, weighted branch lines, different gear hauling techniques and line shooters. Current regulated and voluntary initiatives are summarised by fishery in Table 5.8.

In 2002, MFish, DOC, and stakeholders began working with other countries to reduce the incidental catch of seabirds. As a result, a group called Southern Seabird Solutions was formed and formally established as a Trust in 2003 (<http://www.southernseabirds.org/>) and received royal patronage in 2012. Southern Seabird Solutions exists to promote responsible fishing practices that avoid the incidental capture of seabirds in New Zealand and the southern ocean. Membership includes representatives from the commercial fishing industry, environmental and conservation groups, and government departments. The Trust’s vision is that: *All fishers in the Southern Hemisphere avoid the capture of seabirds*, and this is underpinned by the strategic goals on: Culture Change; Supporting Collaboration; Mitigation Development and Knowledge Transfer; Recognising Success; and Strengthening the Trust.

Building on these initiatives, New Zealand’s 2004 NPOA established a more comprehensive framework to reducing incidental captures approach across all fisheries (because focussing on longline fisheries like the IPOA was considered neither equitable nor sufficient).

It included two goals that set the overall direction:

1. To ensure that the long-term viability of protected seabird species is not threatened by their incidental catch in New Zealand fisheries waters or by New Zealand flagged vessels in high seas fisheries; and
2. To further reduce incidental catch of protected seabird species as far as possible, taking into account advances in technology, knowledge and financial implications.

Together the two goals established the NPOA as a long-term strategy. The second goal was designed to build on the first goal by promoting and encouraging the reduction of incidental catch beyond the level that is necessary to ensure long term viability. The goals recognised that, although seabird deaths may be accidentally caused by fishing, most seabirds are absolutely protected under the Wildlife Act. The second goal balances the need to continue reducing incidental catch against the factors that influence how this can be achieved in practice (e.g., advances in technology and the costs of mitigation). The scope of the NPOA included:

- all seabird species absolutely or partially protected under the Wildlife Act;
- commercial and non-commercial fisheries;
- all New Zealand fisheries waters; and
- high seas fisheries in which New Zealand flagged vessels participate, or where foreign flagged vessels catch protected seabird species.

Specific objectives were established in the NPOA as follows:

1. Implement efficient and effective management measures to achieve the goals of the NPOA, using best practice measures where possible;
2. Ensure that appropriate incentives and penalties are in place so that fishers comply with management measures;
3. Establish mandatory bycatch limits for seabird species where they are assessed to be an efficient and effective management measure and there is sufficient information to enable an appropriate limit to be set;
4. Ensure that there is sufficient, reliable information available for the effective implementation and monitoring of management measures;
5. Establish a transparent process for monitoring progress against management measures;
6. Ensure that management measures are regularly reviewed and updated to reflect new information and developments, and to ensure the achievement of the goals of the NPOA;
7. Encourage and facilitate research into affected seabird species and their interactions with fisheries;
8. Encourage and facilitate research into new and innovative ways to reduce incidental catch;
9. Provide mechanisms to enable all interested parties to be involved in the reduction of incidental catch;
10. Promote education and awareness programmes to ensure that all fishers are aware of the need to reduce incidental catch and the measures available to achieve a reduction.

The NPOA-seabirds sets out the mix of voluntary and mandatory measures that would be used to help reduce incidental captures of seabirds, noted research into the extent of the problem and the techniques for mitigating it, and outlined mechanisms to oversee, monitor and review the effectiveness of these measures. It was not within the scope of the NPOA to address threats to seabirds other than fishing. Such threats are identified in DOC's Action Plan for Seabird Conservation in New Zealand (Taylor 2000) and their management is undertaken by DOC.

Since publication of the NPOA in 2004, more progress has been made in the commercial fishing sector, including:

- in the deepwater fishing sector;
  - industry has implemented vessel specific risk management plans (VMPs) comprising non-mandatory seabird scaring devices offal management and other measures to reduce risks to seabirds,
  - Government has implemented mandatory measures to reduce risk to seabirds (e.g., use and deployment of seabird scaring devices), and
  - industry has taken a proactive stance in resourcing a 24/7 liaison officer to undertake incident response actions, mentoring, VMP and regime development and reviewing, and fleet wide training;
- in the bottom and surface long-line sectors, Government has implemented mandatory measures including tori lines, night setting, line weighting and offal management;
- a number of research projects have been or are currently being undertaken by government and industry into offal discharge, efficacy of seabird scaring devices, line weighting and longline setting devices; and
- workshops organised by both industry bodies and Southern Seabird Solutions are being held for the inshore trawl and longline sectors.

Areas still requiring progress identified in MPI's 2012 consultation documents for a revision to the NPOA-seabirds included:

- development and implementation of mitigation measures, and education, training and outreach in commercial set net fisheries and inshore trawl fisheries;
- implementation of spatially and temporally representative at-sea data collection in inshore and some HMS fisheries;
- development and implementation of mitigation measures for net captures in trawl fisheries;
- development and implementation of mitigation measures, education, training and outreach in, and risk assessment of non-commercial fisheries (especially setnet and line fisheries).

Mitigation has developed substantially since FAO's IPOA was published and a number of recent reviews consider the effectiveness of different methods (Bull 2007, 2009) and summarise currently accepted best practice (ACAP 2011). In December 2010, FAO held a Technical Consultation where International Guidelines on bycatch management and reduction of discards were adopted (FAO2010). The text included an agreement that the guidelines should complement appropriate bycatch measures addressed in the IPOA-Seabirds and its Best Practice Technical Guidelines (FAO 2009). The Guidelines were subsequently adopted by FAO in January 2011.

The most important factor influencing contacts between seabirds and trawl warp cables is the discharge of offal (Wienecke and Robertson 2002; Sullivan *et al.* 2006, ACAP 2011). Offal management methods used to reduce the attraction of seabirds to vessels include mealings, mincing, and batching. ACAP recommends (ACAP 2011) full retention of all waste material where practicable because this significantly reduced the number of seabirds feeding behind vessels compared with the discharge of unprocessed fish waste (Abraham 2009; Wienecke and Robertson 2002; Favero *et al.* 2010) or minced waste (Melvin *et al.* 2010). Offal management has been found to be a key driver of seabird bycatch in New Zealand trawl fisheries (Abraham 2007; Abraham and Thompson 2009b; Abraham *et al.* 2009; Abraham 2010; Pierre *et al.* 2010, 2012 a&b). Other best practice recommendations (ACAP 2011) are the use of bird-scaring lines to deter birds from foraging near the trawl warps, use of snatch blocks to reduce the aerial extent of trawl warps, cleaning fish and benthic material from nets before shooting, minimising the time the trawl net is on the surface during hauling, and binding of large meshes in pelagic trawl before shooting.

**Table 5.8 (after MPI 2012, consultation documents for a revised NPOA-seabirds): summary of current mitigation measures applied to New Zealand vessels fishing in New Zealand waters to avoid incidental seabird captures. R, regulated; SM, required via a self-managed regime (non-regulatory, but required by industry organisation and audited independently by Government); V, voluntary with at least some use known; N/A, measure not relevant to the fishery; years in parentheses indicate year of implementation; \*, part of a vessel management plan (VMP). Note, this table may not capture all voluntary measures adopted by fishers.**

Mitigation Measure	Surface longline	Bottom longline	Trawl >=28 m	Trawl <28 m	Set net	Notes
Netsonde cable prohibition	N/A	N/A	R (1992)	R (1992)	N/A	Netsonde cables also called third wires
Streamer (tori) lines	R	R	N/A	N/A	N/A	
Additional streamer line	–	–	N/A	N/A	N/A	
Night setting	R (or line weighting)	R (or line weighting)	–	–	–	 <p>Longlines must use night setting if not line weighting, or <i>vice-versa</i> To prevent warp captures and collisions</p>
Line weighting	R (or night setting)	R (or night setting)	N/A	N/A	N/A	
Seabird scaring device	N/A	N/A	R (2006)	R?	N/A	
Additional bird scaring device	N/A	N/A	SM (2008)*	–	N/A	
Dyed bait	V	–	N/A	N/A	N/A	
Offal management	V	R	SM (2008)*	–	–	
VMPs			SM (2008)	V	–	Some VMPs developed for vessels < 28m
Code of Practice	V	–	VMP	–	–	

Note: A vessel management plan (VMP) is a vessel-specific seabird risk management plan which specifies seabird mitigation devices to be used, operational management requirements to minimise the attraction of seabirds to vessels, and incident response requirements and other techniques or processes in place to minimise risk to seabirds from fishing operations.

In New Zealand, the three legally permitted devices used for mitigation by trawlers are tori lines (e.g., Sullivan *et al.* 2006), bird bafflers (Crysel 2002), and warp scarers (Carey 2005). Middleton and Abraham (2007) reported experimental trials of mitigation devices designed to reduce the frequency of collisions between seabirds and trawl warps on 18 observed vessels in squid trawl fishery in 2006. The frequencies of birds striking either warps or one of three mitigation devices (tori lines, 4-boom bird bafflers, and warp scarers) were assessed using standardised protocols during commercial fishing. Different warp strike mitigation treatments were used on different tows according to a randomised experimental design. Middleton and Abraham (2007) confirmed that the discharge of offal was the main factor influencing seabird strikes; almost no strikes were recorded when there was no discharge, and strike rates were low when only sump water was discharged (see also Abraham *et al.* 2009). In addition to this effect, tori lines were shown to be most effective mitigation approach and reduced warp strikes by 80–95% of their frequency without mitigation. Other mitigation approaches were only 10–65% effective. Seabirds struck tori lines about as frequently as they did the trawl warps in the absence of mitigation but the consequences are unknown.

Recommended best practice for surface (pelagic) longline fisheries and bottom (demersal) longlines (ACAP 2011) includes weighting of lines to ensure rapid sinking of baits (including integrated weighted line for bottom longlines), setting lines at night when most vulnerable birds are less active, and the proper deployment of bird scaring lines (tori lines) over baits being set, and offal management (especially for bottom longlines). A range of other measures are offered for consideration.

### **5.4.3. Modelling fisheries interactions and estimating risk**

#### **5.4.3.1. Hierarchical structure of risk assessments**

Hobday *et al.* (2007) described a hierarchical framework for ecological risk assessment in fisheries (see Figure 5.11). The hierarchy included three levels: Level 1 qualitative, expert-based assessments (often based on a Scale, Intensity, Consequence Analysis, SICA); Level 2 semi-quantitative analysis (often using some variant of Productivity Susceptibility Analysis, PSA); and Level 3 fully quantitative modelling including uncertainty analysis. The hierarchical structure is designed to “screen out” potential effects that pose little or low risk for the least investment in data collection and analysis, escalating to risk treatment or higher levels in the hierarchy only for those potential effects that pose non-negligible risk. This structure relies for its effectiveness on a low potential for false negatives at each stage, thereby identifying and screening out activities that are ‘low risk’ with high certainty. This focuses effort on remaining higher risk activities. In statistical terms, risk assessment tolerates Type I errors (false positives, i.e. not screening out activities that may actually present a low risk) in order to avoid Type II errors (false negatives, i.e. incorrectly screening out activities that actually constitute high risk), and it is important to distinguish this approach from normal estimation methods. Whereas normal estimation strives for a lack of bias and a balance of Type I and Type II errors, risk assessment is designed to answer the question “how bad could it be?” The divergence between the risk assessment approach and normal, unbiased estimation approaches should diminish at higher levels in the risk assessment hierarchy, where the assessment process should be informed by good data that support robust estimation.




Figure 5.11 (from Hobday *et al.*): Diagrammatic representation of the hierarchical risk assessment process where activities that present low risk are progressively screened out by assessments of increasingly high data content, sophistication, and cost.

#### 5.4.3.2. Qualitative (Level 1) risk assessment

Rowe (2010) summarised an expert-based, qualitative (Level 1) risk assessment, commissioned by DOC, for the incidental mortality of seabirds caused by New Zealand fisheries. The main focus was on fisheries operating within the NZ EEZ and on all seabirds absolutely or partially protected under the Wildlife Act 1953. New Zealand flagged vessels fishing outside the EEZ were included, but risk from non-NZ fisheries and other human causes were not included.

The panel of experts who conducted the Level 1 risk assessment assessed the threat to each of 101 taxa posed by 26 fishery groups, scoring exposure and consequence independently according to the schemas in Tables 5.9 and 5.10 (details in Rowe 2010b). The risk for a given taxon posed by a given fishery was calculated as the product of exposure and consequence scores. Potential risk was estimated as the risk posed by a fishery assuming no mitigation was in place, and residual risk (called “optimum risk” by Rowe 2010b) was estimated assuming that mitigation was in place throughout a given fishery and deployed correctly. The panel also agreed a confidence score for each taxon-fishery interaction using the schema in Table 5.11.

Table 5.9: Exposure scores used by Rowe (2010) (modified from Fletcher 2005, Hobday *et al* 2007)

Score	Descriptor	Description
0	Remote	The species will not interact directly with the fishery
1	Rare	Interactions may occur in exceptional circumstances
2	Unlikely	Evidence to suggest interactions possible
3	Possible	Evidence to suggest interactions occur, but are uncommon
4	Occasional	Interactions likely to occur on occasion
5	Likely	Interactions are expected to occur

**Table 5.10: Consequence scores used by Rowe (2010) (modified from Fletcher 2005, Campbell & Gallagher 2007, Hobday *et al.* 2007)**

Score	Descriptor	Description
1	Negligible	Some or one individual/s impacted, no population impact.
2	Minor	Some individuals are impacted, but minimal impact on population structure or dynamics. In the absence of further impact, rapid recovery would occur
3	Moderate	The level of interaction / impact is at the maximum acceptable level that still meets an objective. In the absence of further impact, recovery is expected in years
4	Major	Wider and longer term impacts; loss of individuals; potential loss of genetic diversity. Level of impact is above the maximum acceptable level. In the absence of further impact, recovery is expected in multiple years
5	Severe	Very serious impacts occurring, loss of seabird populations causing local extinction; decline in species with single breeding population, measurable loss of genetic diversity. In the absence of further impact, recovery is expected in years to decades
6	Intolerable	Widespread and permanent / irreversible damage or loss occurring; local extinction of multiple seabird populations; serious decline of a species with a single breeding population, significant loss of genetic diversity. Even in the absence of further impact, long-term recovery period to acceptable levels will be greater than decades or may never occur

**Table 5.11: Confidence scores used by Rowe (2010) (after Hobday 2007)**

Score	Descriptor	Rationale for confidence score
1a	Low	Data exists, but is considered poor or conflicting.
1b		No data exists.
1c		Agreement between experts, but with low confidence
1d		Disagreement between experts
2a	High	Data exists and is considered sound.
2b		Consensus between experts
2c		High confidence exposure to impact can not occur (e.g. no spatial overlap of fishing activity and at-sea seabird distribution)

Total potential and residual risk for a seabird taxon was estimated by summing the scores across all fisheries (Table 5.12 shows taxa with an aggregate score of 30 or higher), and total potential and residual risk posed by a fishery group was estimated by summing the scores across all seabird taxa (Table 5.13 shows the results for all 26 fishery groups).

White-chinned petrel, Sooty shearwater, Black (Parkinson's) petrel, Salvin's albatross, White-capped albatross, and Flesh-footed shearwater were all estimated by this procedure to have an aggregate risk score of 90 or higher (range 92 to 123) even if mitigation was in place and deployed properly across all fisheries. Of the 101 seabird taxa considered, the aggregate risk score was less than 30 for 70 taxa with respect to potential risk and for 72 taxa with respect to residual risk.

**Table 5.12: Potential and residual risk scores for each seabird taxon with a potential risk score of  $\geq 30$  in Rowe (2010). Residual risk (“optimal risk” in Rowe 2010b, not tabulated therein for grey-faced petrel or light-mantled albatross) is estimated assuming mitigation is deployed and correctly used throughout all interacting fisheries.**

<b>Taxon</b>	<b>Potential score</b>	<b>Residual score</b>	<b>Percent reduction</b>
White-chinned petrel	159	123	23
Sooty shearwater	126	108	14
Black (Parkinson's) petrel	139	106	24
Salvin's albatross	161	106	34
White-capped albatross	141	94	33
Flesh-footed shearwater	117	92	21
Southern Buller's albatross	123	85	31
Grey petrel	123	84	32
Black-browed albatross	114	80	30
Northern Buller's albatross	107	72	33
Chatham albatross	114	71	38
Campbell albatross	97	66	32
Westland petrel	89	59	34
Antipodean albatross	89	55	38
Gibson's albatross	89	55	38
Wandering albatross	89	55	38
Southern royal albatross	79	49	38
King shag	48	48	0
Pitt Island shag	46	46	0
Chatham Island shag	45	45	0
Hutton's shearwater	37	35	5
Northern giant petrel	62	35	44
Pied shag	35	35	0
Indian yellow-nosed albatross	58	34	41
Southern giant petrel	61	34	44
Fluttering shearwater	34	32	6
Spotted shag	31	31	0
Stewart Island shag	31	31	0
Yellow-eyed penguin	30	30	0
Grey-faced petrel	31	–	–
Light-mantled albatross	30	–	–

Setnet and inshore trawl fisheries groups posed the greatest residual risk to seabirds (summed across all taxa); both had aggregate scores of over 200 and had no substantive mitigation. Surface and bottom longline fisheries and middle-depth trawl fisheries for finfish and squid also had aggregate risk scores of 100 or more. These risk scores were substantially reduced if mitigation was assumed to be deployed throughout these fisheries (reductions of 24 to 56%), but all remained above 100. Trawling for southern blue whiting and deep-water species, inshore drift net, various seine methods, ring net, diving, dredging, and hand gathering all had aggregate risk scores of 40 or less if mitigation was assumed to be deployed throughout these fisheries. Diving, dredging, and hand gathering were all judged by the panel to pose essentially no risk to seabirds.

**Table 5.13: Cumulative potential risk and residual risk scores across all seabird taxa for each fishery from Rowe (2010). Residual risk (“optimal risk” in Rowe 2010b) is estimated assuming mitigation is deployed and correctly used throughout a given fishery.**

<b>Fishery group</b>	<b>No. taxa</b>	<b>Potential risk</b>	<b>Residual risk</b>	<b>Percent reduction</b>
Setnet	42	374	374	0
Inshore trawl	44	225	225	0
Surface longline: charter	25	313	191	39
Surface longline: domestic	25	302	184	39
Bottom longline: small	33	354	154	56
Bottom longline: large	32	311	139	55
Mid-depth trawl: finfish	22	160	122	24
Mid-depth trawl: squid	21	156	118	24
Mid-depth trawl: scampi	23	94	94	0
Hand line	27	68	68	0
Squid jig	44	62	62	0
Dahn line	29	61	61	0
Pots, traps	17	61	61	0
Trot line	29	61	61	0
Pelagic trawl	27	63	51	19
Troll	23	50	50	0
Mid-depth trawl: southern blue whiting	21	53	40	25
Deep water trawl	21	46	35	24
Inshore drift net	12	33	33	0
Danish seine	15	32	32	0
Beach seine	16	29	29	0
Purse seine	11	22	22	0
Ring net	12	13	13	0
Diving	0	0	0	–
Dredge	0	0	0	–
Hand gather	0	0	0	–

### 5.4.3.3. Semi-quantitative (Level 2) risk assessment

The level 2 method developed by MPI arose initially from an expert workshop hosted by the Ministry of Fisheries in 2008 and attended by experts with specialist knowledge of New Zealand fisheries, seabird-fishery interactions, seabird biology, population modelling, and ecological risk assessment. The overall framework is described in Sharp *et al.* (2011) and has been variously applied and improved in multiple iterations (Waugh *et al.* 2008, developed further by Sharp 2009, Waugh and Filippi 2009, Filippi *et al.* 2010, Richard *et al.* 2011). The method applies the “exposure-effects” approach where exposure refers to the number of fatalities arising from an activity and effect refers to the consequence of that exposure for the population. The relative encounter rate of each seabird taxon with each fishery group is estimated as a function of the spatial overlap between seabird distributions (e.g., Figure 5.12) and fishing effort distributions (e.g., see Figures 5.4–5.6), and compares these estimates with observed captures from fisheries observer data to estimate vulnerability by taxon (capture rates per encounter) to each fishery group, yielding estimates of total observable captures and population-level potential fatalities from all New Zealand commercial fisheries. Impact estimates are subsequently compared with population estimates and biological characteristics to yield estimates of population-level risk.

The current level 2 risk assessment (i.e., as described by Richard *et al.* 2011) estimated the risk posed to each of 64 seabird taxa by trawl and longline fisheries within New Zealand’s TS and EEZ. Insufficient information was available to include some other fisheries thought to pose substantial risk

to seabirds, especially setnet. For each taxon, the risk was assessed by dividing the estimated number of potential fatalities by an estimate of Potential Biological Removals (PBR, after Wade 1998). This index represents the amount of human-induced mortality a population can sustain without compromising its ability to achieve and maintain a population size above its maximum net productivity (MNPL) or to achieve rapid recovery from a depleted state. In the risk assessment, PBR was estimated from the best available information on the demography of each taxon (Figure 5.13). Because estimates of seabirds' demographic parameters and of fisheries related mortality are imprecise, the uncertainty around the demographic and mortality estimates was propagated through the analysis. This allowed uncertainty in the resulting risk to be calculated, and also allowed the identification of parameters where improved precision would reduce overly large uncertainties. However, not all sources of uncertainty could be included, and the results are best used as a guide in the setting of research and management priorities. In general, seabird demographics, the distribution of seabirds within New Zealand waters, and sources of cryptic mortality were poorly known.

Amongst the 64 studied taxa, black (Parkinson's) petrel (*Procellaria parkinsoni*) clearly stood out as at most risk from commercial fishing activities within the New Zealand Exclusive Economic Zone (estimated annual potential fishing-related fatalities almost 10 times higher than the PBR, Figures 5.14 and 5.15). Seven other taxa had estimated annual potential fatalities with 95% confidence intervals of their risk ratios completely above one. These were grey-headed albatross, Chatham albatross, Westland petrel, light-mantled albatross, Salvin's albatross, fleshfooted shearwater, and Stewart Island shag. For a further 12 taxa, the confidence interval of the risk ratio included one.

Small inshore fisheries, especially trawl fisheries targeting flatfish, and small bottom and surface fisheries, were associated with the highest estimated risk to seabirds. This was due to a combination of low observer coverage, high effort, and overlap with the distributions of many seabirds. In fisheries where there were few observations, the number of potential fatalities was estimated in a precautionary way, with the estimates being biased toward the high end of the range of values that were consistent with the observer data. In these poorly observed fisheries, the risk estimates are often primarily associated with the lack of information. Of the taxa that had a risk ratio greater than one, the risk for four of them (grey-headed albatross, Westland petrel, Chatham albatross, and light-mantled albatross) was associated with low observer coverage in inshore fisheries that overlap with the distribution of these birds. Increasing the number of observations in inshore trawl and small vessel longline fisheries, especially in FMAs 1, 2, 3, and 7, would increase the precision of the estimated fatalities. The risk was estimated independently for each fishery, and it was assumed that the vulnerabilities of seabirds to capture in different fisheries were independent. This has the consequence that birds (such as light-mantled sooty albatross) may be caught infrequently in well observed fisheries, but still have high risk associated with poorly observed fisheries.




Figure 5.12 (reproduced from Richard *et al.* 2011 supplementary material): Captures and relative density of White-capped albatross (top) and Chatham petrel (bottom) showing large differences in the extent of distributions and overlap with fishing effort (in grey), and in the number of observed captures. The

distribution base maps were obtained from NABIS (white-capped albatross) and the BirdLife single-layer range maps (Chatham petrel).




Figure 5.13 (reproduced from Richard *et al.* 2011): Diagram of the modelling approach to calculate the risk index for each taxon.  $N_{BP}$ , number of annual breeding pairs;  $N$ , total number of birds over one year old;  $N_{BPmin}$ , lower 25% of the distribution of  $N_{BP}$ ;  $Nmin$ , lower 25% of the distribution of the total number of birds over one year old;  $r_{max}$ , maximum population growth rate;  $f$ , recovery factor;  $PBR$ , Potential Biological Removal;  $P$ , proportion of adults breeding in a given year;  $A$ , age at first reproduction;  $S$ , annual adult survival rate.






Figure 5.14 (reproduced from Richard *et al.* 2011): Mean annual potential seabird fatalities in the assessed fishery groups (colour bars) and the PBR (grey bars), for each of the 64 studied taxa. The bars indicate the 95% confidence intervals of the distributions. Taxa are sorted in decreasing order of the lower confidence level of the number of fatalities.



**Figure 5.15 (reproduced from Richard *et al.* 2011): Risk ratio (total annual potential fatalities / PBR) for each of the studied taxa except black-browed albatross. The risk ratio is displayed on a logarithmic scale. The threshold where the number of potential bird fatalities equals the PBR is presented by the vertical black line. The bars indicate the 95% confidence intervals of the distributions. Taxa are sorted in decreasing order of the lower confidence level of the risk ratio.**

Many limitations were identified in the risk assessment. These may result in biased estimates (either too high or too low) of the risk of fishing to some seabirds. Moreover, some fisheries were not included in our analysis, and other sources of human-induced mortality were ignored. The conclusions of our results should therefore be interpreted with caution, as some taxa might be at risk, even if their risk ratio was estimated to be lower than one. Conversely, the fisheries-related fatalities may be overestimated in poorly observed fisheries because the method is designed to answer the question “how bad could it be?” (e.g., Figure 5.16 which shows the results of different estimation approaches and questions). The method assumed a high number of captures in the absence of data to the contrary, so the estimated potential fatalities in poorly-observed fisheries may be higher than the actual fatalities.



**Figure 5.16 (reproduced from Richard *et al.* 2011): Comparison of the number of potential annual captures (without cryptic mortality) estimated using the risk assessment method used by Richard *et al.* (2011), a simple ratio scalar, and statistical modelling, for white-chinned petrel, white-capped albatross, sooty shearwater, and all birds combined, in trawl, bottom longline, and surface longline fisheries. Each symbol represents the mean and the 95% confidence interval of an estimate.**

The method described by Richard *et al.* (2011) offers the following advantages that make it particularly suitable for assessing risk to multiple seabird populations from multiple fisheries:

- risk is assessed separately for each seabird taxon; fisheries managers must assess risk to seabirds with reference to units that are biologically meaningful;
- the method does not rely on the existence of universal or representative fisheries observer data to estimate seabird mortality (fisheries observer coverage is generally too low and/or too spatially unrepresentative to allow direct impact estimation at the species or subspecies level); the method can be applied to any fishery for which at least some observer data exists;
- the method does not rely on detailed population models (the necessary data for which are unavailable for the great majority of taxa) because risk is estimated as a function of population-level potential fatalities and biological parameters that are generally available from published sources;
- the method assigns risk to each taxon in an absolute sense, i.e. taxa are not merely ranked relative to one another; this allows the definition of biologically meaningful performance standards and ability to track changes in performance over time and in relation to risk management interventions;
- risk scores are quantitative and objectively scalable between fisheries or areas, so that risk at a population level can be disaggregated and assigned to different fisheries or areas based on their proportional contribution to total impact to inform risk management prioritisation;

- the method allows explicit statistical treatment of uncertainty, and does not conflate uncertainty with risk; numerical inputs include error distributions and it is possible to track the propagation of uncertainty from inputs to estimates of risk; and
- the method readily incorporates new information; assumptions in the assessment are transparent and testable and, as new data becomes available, the consequences for the subsequent impact and risk calculations arise logically without the need to revisit other assumptions or repeat the entire risk assessment process.

The key disadvantages of the method are that:

- fisheries for which no observer information on seabird interactions is available cannot be included in the analysis
- the assumption that the vulnerabilities of particular seabirds to capture in different fisheries are independent does not allow “sharing” of scarce observer information between fisheries within the risk assessment
- the spatial overlap method relies on appropriate spatial and temporal scales for the distributions of birds and fishing effort being used; use of inappropriate scales can lead to misleading results
- strong assumptions have to be made about the distribution and productivity of some taxa, the relative vulnerability of different taxa to capture by particular fisheries, cryptic mortality associated with different fishing methods, and the applicability of the allometric method of estimating Potential Biological Removals.

Most of these limitations are a result of the scarcity of relevant data on seabird populations and fisheries impacts and can be addressed only through the collection of more information or, in some cases, sensitivity testing. In particular, it was not possible to include some fishery groups identified by Rowe’s (2010b) level 1 analysis as posing substantial risk to seabirds. Notable among these fisheries was the commercial setnet fishery group. In the absence of quantitative information for these fishery groups, the Ministry of Fisheries combined the level 1 and level 2 results to generate a comprehensive assessment of seabird risk across all New Zealand seabirds and fisheries (Table 5.14). Apart from filling some important information gaps in the assessment, the level 1 results were also useful as a cross-check on the level 2 outputs. A number of likely misleading results were identified in this way, including those from poor input data (e.g. spatial distribution layers) or faulty structural assumptions for particular seabird taxa, and these were noted so that inappropriate conclusions were not made and to provide for better treatment in subsequent iterations of the level 2 analysis.

At the time of going to press, a major update and revision of the level 2 risk assessment published by Richard *et al.* (2011) was undergoing final review. This revision includes several substantial improvements on the 2011 version including:

- fisheries and observer data from 2006/07 onwards (i.e., post-mitigation only, allowing a better estimate of current risk)
- inclusion of set net fisheries (obviating the need to combine level 1 and level 2 analyses)
- revised bird distributions
- inclusion of seasonal stratification of bird distribution and overlap with fisheries
- an integrated approach to estimating species-specific vulnerability to particular fisheries
- correction of a bias in the estimation of productivity from age at first reproduction and annual adult survival rate
- inclusion of uncertainty in estimates of cryptic mortality

This revised risk assessment is expected to be available early in 2013.

**Table 5.14: Combined level 2 and level 1 risk assessments for seabird taxa with a risk ratio of 0.3 or greater (i.e., mean potential fatalities 30% of the estimated PBR or greater). INS, inshore trawl fisheries including for flatfish; SQU, squid trawl fisheries; SCI, scampi trawl fisheries; OFF, other offshore trawl fisheries; BLL, bottom longline fisheries; SLL, surface longline fisheries; SN, setnet (from level 1); Other, all other fisheries considered in the level 1 risk assessment. \* indicates an unreliable assessment.**

Taxon	INS	SQU	SCI	OFF	BLL	SLL	SN	Other	Risk ratio
Black (Parkinson's) petrel	4.37	0.12	0.17	0.37	5.56	0.41	0.00	0.45	11.45
Black-browed albatross *	1.07	0.02	0.04	0.64	2.46	1.37	0.00	0.00	* 5.59
New Zealand king shag	1.84	0.00	0.00	0.05	0.15	0.00	1.51	0.91	4.46
Grey-headed albatross *	2.38	0.03	0.09	0.37	0.52	0.07	0.00	0.00	* 3.46
Westland petrel	1.99	0.12	0.05	0.36	0.59	0.15	0.00	0.00	3.26
Chatham albatross	1.81	0.04	0.08	0.19	0.55	0.03	0.00	0.00	2.70
Stewart Island shag	1.59	0.00	0.00	0.00	0.01	0.00	1.01	0.00	2.62
Northern giant-petrel	1.65	0.06	0.12	0.32	0.34	0.05	0.00	0.00	2.55
Pitt Island shag	0.00	0.00	0.00	0.01	0.12	0.00	1.51	0.80	2.45
Flesh-footed shearwater	0.72	0.01	0.31	0.08	1.12	0.19	0.00	0.00	2.42
Chatham Island shag	0.00	0.00	0.00	0.02	0.17	0.00	1.51	0.60	2.31
Salvin's albatross	1.43	0.05	0.22	0.22	0.34	0.03	0.00	0.00	2.29
Light-mantled albatross *	1.46	0.02	0.05	0.22	0.34	0.04	0.00	0.00	* 2.14
Northern royal albatross	0.96	0.05	0.03	0.21	0.31	0.54	0.00	0.00	2.09
Campbell albatross	0.61	0.06	0.06	0.24	0.51	0.33	0.00	0.00	1.81
New Zealand storm-petrel	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.51	1.51
Yellow-eyed penguin	0.08	0.00	0.00	0.01	0.02	0.00	1.26	0.00	1.38
Spotted shag	0.47	0.00	0.00	0.00	0.03	0.01	0.75	0.00	1.27
Fiordland crested penguin	0.07	0.00	0.00	0.01	0.33	0.04	0.80	0.00	1.25
Wandering albatross	0.00	0.00	0.00	0.00	0.00	1.21	0.00	0.00	1.21
Southern Buller's albatross	0.36	0.17	0.03	0.38	0.05	0.20	0.00	0.00	1.19
Gibson's albatross	0.23	0.02	0.01	0.07	0.15	0.67	0.00	0.00	1.16
Antipodean albatross	0.22	0.02	0.01	0.06	0.14	0.64	0.00	0.00	1.10
Hutton's shearwaters	0.05	0.00	0.00	0.01	0.04	0.00	1.01	0.00	1.10
Pied shag	0.00	0.00	0.00	0.00	0.00	0.00	0.75	0.30	1.06
South Georgia diving-petrel	0.00	0.00	0.00	0.00	0.00	0.00	0.60	0.30	0.91
Indian yellow-nosed albatross	0.00	0.00	0.00	0.00	0.00	0.60	0.00	0.00	0.60
White-capped albatross	0.32	0.34	0.02	0.09	0.02	0.01	0.00	0.00	0.79
Sooty shearwater *	0.01	0.01	0.00	0.00	0.00	0.00	0.75	0.00	* 0.77
White-chinned petrel	0.11	0.37	0.01	0.09	0.15	0.03	0.00	0.00	0.77
Fluttering shearwater	0.00	0.00	0.00	0.00	0.00	0.00	0.75	0.00	0.75
Little black shag	0.00	0.00	0.00	0.00	0.00	0.00	0.75	0.00	0.75
Northern blue penguin	0.00	0.00	0.00	0.00	0.00	0.00	0.75	0.00	0.75
White-flipped blue penguin	0.00	0.00	0.00	0.00	0.00	0.00	0.75	0.00	0.75
Northern Buller's albatross	0.19	0.00	0.05	0.14	0.19	0.17	0.00	0.00	0.75
Southern royal albatross	0.31	0.05	0.02	0.07	0.08	0.21	0.00	0.00	0.74
Cape petrel *	0.27	0.00	0.03	0.14	0.21	0.03	0.00	0.00	* 0.69
Southern giant petrel	0.00	0.00	0.00	0.00	0.15	0.00	0.00	0.00	0.15
Chatham Island blue penguin	0.00	0.00	0.00	0.00	0.00	0.00	0.45	0.00	0.45
Grey petrel	0.13	0.00	0.00	0.03	0.08	0.12	0.00	0.00	0.37
Magenta petrel	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.30	0.33

#### 5.4.3.4. Fully quantitative modelling

Fully quantitative population modelling has been conducted only for southern Buller's albatross, black (Parkinson's) petrel, white capped albatross (mollymawk), and Gibson's (wandering) albatross. Data of similar quality and quantity are available for Antipodean (wandering) albatross, and this work should be commissioned soon, but data for other species or populations appear unlikely to be adequate for comprehensive population modelling. The poor estimates of observable and cryptic fishing-related mortality have restricted such work to comprehensive population modelling rather than formal assessment of risk.

##### 5.4.3.4.1. Quantitative models for southern Buller's albatross

Francis *et al.* (2008, see also Francis and Sagar 2012) assessed the status of the Snares Islands population of southern Buller's albatross (*Thalassarche bulleri bulleri*). They estimated (see also Sagar and Stahl 2005) that the adult population had increased about 5-fold since about 1950 (Figure 5.17) at a rate of about 2% per year, and concluded from this that the risk to the viability of this population posed by fisheries had been small. This conclusion depends critically on the reliability of the first census of nesting birds conducted in 1969, but the authors give compelling reasons to trust that information. They noted, however, that population growth had slowed by about 2005 (and perhaps reversed) and adult survival rates were falling, but could discern neither the cause nor significance of these changes because they had included survival data only up to 2007. An additional 5 years of survival and other demographic data have since been recorded (Sagar *et al.* 2010) and all monitored sites at the Snares Islands show substantial declines in the number of breeding pairs since 2006. The modelling has not yet been repeated.




Figure 5.17 (reproduced from Francis *et al.* 2008): Estimates from model SBA21 of numbers of breeders (solid line) and adults (broken line) in each year. Also shown are the census observations (after (Sagar and Stahl 2005) of numbers of breeders (crosses), with assumed 95% confidence intervals (vertical lines).

Fishery discards are an important component of the diet of chicks, but Francis *et al.* (2008) were not able to assess whether the associated positive effect on population growth (e.g., from increased breeding success) is greater or less than the negative effect of fishing-related mortality.

#### 5.4.3.4.2. Quantitative models for black petrel

Francis and Bell (2010) analysed data from the main population of black (Parkinson's) petrel (*Procellaria parkinsoni*), which breeds on Great Barrier Island. Abundance data from transect surveys were used to infer that the population was probably increasing at a rate between 1.2% and 3.1% per year. Mark-recapture data were useful in estimating demographic parameters, like survival and breeding success, but contained little information on population growth rates. Fishery bycatch data from observers were too sparse and imprecise to be useful in assessing the contribution of fishing-related mortality. Francis and Bell (2010) suggested that, because the population was probably increasing, there was no evidence that fisheries posed a risk to the population at that time. They cautioned that this did not imply that there was clear evidence that fisheries do not pose a risk.

Subsequent analysis (Bell *et al.* 2012) included an additional line transect survey in 2009/10 in which the breeding population was estimated to be ~22% lower than in 2004/05 (the latest available to Francis and Bell, 2010). Updating the model of Francis and Bell (2010) made little difference to estimates of demographic parameters such as adult survival, age at first breeding, and juvenile survival (which had 95% confidence limits of 0.67 and 0.91). The uncertainty in juvenile survival gave rise to uncertainty in the estimated population trend, with a mean rate of population growth over the modelling period ranging from -2.5% per year (if juvenile survival = 0.67) to +1.6% per year (if juvenile survival = 0.91, close to the average annual survival rate for older birds) (Figure 5.18). Bell *et al.* (2012) concluded that the mean rate of change of the population over the study period had not exceeded 2% per year, though the direction of change was uncertain.




Figure 5.18 (reproduced from Bell *et al.* 2012): Likelihood profile for annual probability of juvenile survival showing: A, the loss of fit (the horizontal dotted line shows a 95% confidence interval for this parameter); and B, population trajectories corresponding to different values of juvenile survival, together with population estimates from transect counts (crosses with vertical lines indicating 95% confidence intervals. Note that the 1988 population estimate was not used in the model.

#### 5.4.3.4.3. Quantitative models for white-capped albatross

Francis (2012) described quantitative models for white-capped albatross (*Thalassarche steadi*), New Zealand's most numerous breeding albatross, and the most frequently captured, focussing on the population breeding at the Auckland Islands. After a correction for a probable bias introduced by sampling at different times of day in one of the surveys, aerial photographic counts by Baker *et al.* (2007, 2008, 2009, and 2010, see also Table 5.15) suggest that the adult population declined at about 9.8% per year between 2006 and 2009. However, this estimate is imprecise and is not easily reconciled with the high adult survival rate (0.96) estimated from mark-recapture data. Francis (2012) also compared the trend with his estimate of the global fishing-related fatalities of white-capped albatross (slightly over 17 000 birds per year, about 30% of which is taken in New Zealand fisheries) and found that fishing-related fatalities were insufficient to account for the number of deaths implied by a decline of 9.8% per year (roughly 22 000 birds per year over the study period). The scarcity of information on cryptic mortality makes these estimates and conclusions uncertain, however.


**Table 5.15 (data from Baker *et al.* 2007, 2008, 2009, 2010): Aerial-photographic counts of breeding pairs of white-capped albatrosses on three islands in the Auckland Islands group in December 2006–2009. Confidence limits for these counts published by Baker (*op. cit.*) were based on a Poisson model and were very narrow (although uncertainty introduced by the proportion of non-nesting birds at the colonies during photography was not included).**

Year	Disappointment	SW Cape	Adams	Total
2006	110 649	6 548	–	117 197
2007	86 080	4 786	79	90 945
2008	91 694	5 264	131	97 089
2009	70 569	4 161	132	74 862


#### 5.4.3.4.4. Quantitative models for Gibson's albatross

Francis *et al.* (in press) concluded there is cause for concern about status of the population of Gibson's wandering albatross (*Diomedea gibsoni*) on the Auckland Islands. Since 2005, the adult population has been declining at 5.7%/yr (95% c.i. 4.5–6.9%) because of sudden and substantial reductions in adult survival, the proportion of adults breeding, and the proportion of breeding attempts that are successful (Figure 5.19). Forward projections showed that the most important of these to the future status of this population is adult survival (Figure 5.20).

The population in 2011 was 64% (58–73%) of its estimated size in 1991. The breeding population dropped sharply in 2005, to 59% of its 1991 level, but has been increasing since 2005 at 4.2% per year (2.3–6.1%). The 2011 breeding population is estimated to be only 54% of the average of 5831 pairs estimated by Walker & Elliott (1999) for 1991–97.



**Figure 5.19:** Estimated population trajectories for the whole Auckland Islands population of Gibson's wandering albatross. These were calculated by scaling up Francis *et al.*'s (in press) GIB5 trajectories to match the Walker & Elliott (1999) estimate for the whole population.



**Figure 5.20:** Estimated population trajectory for adults from Francis *et al.*'s (in press) model GIB5 with 20-year projections under five alternative scenarios about three demographic parameters: adult survival (adsvr); breeding success (Psuccess); and proportion of adults breeding. These scenarios differ according to whether each parameter remains at its status quo (=2011) level or recovers immediately to its 1991 level.

Francis *et al.* (in press) found it difficult to assess the effect of fisheries mortality on the viability of this population because, although some information exists about captures in New Zealand and Australian waters, the effect of fisheries in international waters is unknown. Three conclusions are possible from the available data: most fisheries mortality of Gibson's is caused by surface longlines; mortality from fishing within the New Zealand EEZ is now probably lower than it was; and there is no indication that the sudden and substantial drops in adult survival, the proportion breeding, and breeding success were caused primarily by fishing.

#### 5.4.3.4.5. Other quantitative models

This section is not intended to cover all quantitative modelling of seabird populations, rather to focus on recent studies that sought to assess the impact of fishing-related mortality.

Maunder *et al.* (2007) sought to assess the impact of commercial fisheries on the Otago Peninsula yellow-eyed penguins using mark-recapture data within a population dynamics model. They found the data available at that time inadequate to assess fisheries impacts, but evaluated the likely utility of additional information on annual survival or an estimate of bycatch for a single year. Including auxiliary information on average survival in the absence of fishing allowed estimation of the fishery impact, but with poor precision. Including an estimate of fishery-related mortality for a single year improved the precision in the estimated fishery impact. The authors concluded that there was insufficient information to determine the impact of fisheries on yellow-eyed penguins and that quantifying fishing-related mortality over several years was required to undertake such an assessment using population a modelling approach.

Fletcher *et al.* (2008) sought to assess the potential impact of fisheries on Antipodean and Gibson's wandering albatrosses (*Diomedea antipodensis antipodensis* and *D. a. gibsoni*); black petrel (*Procellaria parkinsoni*) and southern royal albatross (*Diomedea epomophora*). Because of problems with the available fisheries and biological data, they were unable to use their models to predict the impact of a change in fishing effort on the population growth rate of a given species. Instead, they used the models to estimate the impact that changes in demographic parameters like annual survival are likely to have on population growth rate. They found that: reducing breeder survival rate by k percentage points will lead to a reduction in the population growth rate of about 0.3k percentage points (0.4 for black petrel); and a reduction of k percentage points in the survival rate for each stage in the life cycle (juvenile, pre-breeder, non-breeder and breeder) will lead to a reduction in the population growth rate of approximately k percentage points. Fletcher *et al.* (2008) also made estimates of PBR for 23 New Zealand seabird taxa and summarise and tabulated non-fishing-related threats for 38 taxa.

Newman *et al.* (2009) combined survey data with demographic population models to estimate the total population of sooty shearwaters within New Zealand. They estimated the total New Zealand population between 1994 and 2005 to have been 21.3 (95% c.i. 19.0–23.6) million birds. The harvest of “muttonbirds” was estimated to be 360 000 (320 000–400 000) birds per year, equivalent to 18% of the chicks produced in the harvested areas and 13% of chicks in the New Zealand region. This directed harvest is much larger than estimates of captures in key fisheries (Table 5.4) or potential fatalities in the level 2 risk assessment (Figure 5.16). Newman *et al.* (2009) did not assess the likely impact of fishing-related mortality but concluded that the much larger directed harvest was not an adequate explanation for the observed declines in the past three decades.

#### 5.4.3.4.6. General conclusions from quantitative modelling

Fully quantitative modelling has now been conducted for four of the five seabird populations for which apparently suitable data are available. That modelling suggests very strongly that one population had been increasing steadily (southern Buller's albatross, but note this trend may have reversed) and another is declining quite rapidly (Gibson's albatross). White-capped albatross and black petrel both more likely to be declining than not but, even for these relatively data rich populations, the conclusions are uncertain. General conclusions from the modelling conducted to date, therefore, can be summarised as:

- Very few seabird populations have sufficient data for modelling
- Except for the two most complete data sets (southern Buller's and Gibson's albatross) it has been difficult to draw firm conclusions about trends in population size.

- Information from surveys or census counts is much more powerful for detecting trends in population size than data from the tagging programmes and plot monitoring implemented for New Zealand seabirds to date.
- The available information on incidental captures in fisheries have not allowed rigorous tests of the role of fishing-related mortality in driving population trends
- Although comprehensive modelling provides additional information to allow interpretation, we will have to rely on level 2 risk assessment approaches for much of our understanding of the relative risks faced by different seabird taxa and posed by different fisheries.

#### 5.4.3.5. Sources of uncertainty in risk assessments

There are several outstanding sources of uncertainty in modelling the effects of fisheries interactions on sea birds, especially for the complete assessment of risk to individual seabird populations.

##### 5.4.3.5.1. Scarcity of information on captures and biological characteristics of affected populations

These sources of uncertainty can be explored within the analytical framework of the level 2 risk assessment (Richard *et al.* 2011), noting that the results of that exploration are constrained by the structure of that analysis. Richard *et al.* (2011) provided plots of such an exploration for four example taxa (Figure 5.21). It can be concluded from this analysis that substantially more precise estimates of risk would be available for black petrel and Stewart Island shag if better estimates of potential captures were available. Conversely, substantially more precise estimates of risk would be available for Salvin's albatross and flesh-footed shearwater if better estimates of average adult survival were available. This analysis is a powerful way of assessing the priorities for collection of new information, including research.




Figure 5.21 (reproduced from Richard *et al.* 2011): Sensitivity of the risk ratio to the uncertainty in the mean number of annual potential fatalities (F, reflecting the uncertainty in vulnerability), the adult annual survival rate (S), the number of annual breeding pairs (N), the proportion of adults breeding in a given year (P), the age at first reproduction (A), and to the distribution map (D), for the taxa most at risk (lower bound of the 95% c.i. above 1). This sensitivity is expressed as the percentage reduction in the 95% confidence interval of the risk ratio when each parameter is fixed to its mean.

##### 5.4.3.5.2. Scarcity of information on cryptic mortality

Cryptic mortality is particularly poorly understood but has substantial influence on the results of the risk assessment. Richard *et al.* (2011) provided a description of the method used to incorporate cryptic

mortality into their estimates of potential fatalities in the level-2 risk assessment (their Appendix B authored by B. Sharp, MPI). This method builds on the published information from Brothers *et al.* (2010) for longline fisheries and Watkins *et al.* (2008) and Abraham (2010) for trawl fisheries. Brothers *et al.* (2010) observed almost 6 000 seabirds attempting to take longline baits during line setting, of which 176 (3% of attempts) were seen to be caught. Of these, only 85 (48%) were retrieved during line hauling. They concluded that using only observed captures to estimate seabird fatalities grossly underestimates actual levels in pelagic longline fishing. Similarly, Watkins *et al.* (2008) observed 2454 interactions between seabirds and trawl warps in the South African hake fishery over 189.8 hours of observation. About 11% of those interactions (263) involved birds, mostly albatrosses, being dragged under the water by the warps, and 30 of those submersions were observed to be fatal. Of the 30 birds observed killed on the warps, only two (both albatrosses) were hauled aboard and would have been counted as captures by an observer in New Zealand. Aerial collisions with the warps were about 8 times more common but appeared mostly to have little effect (although one white-chinned petrel suffered a broken wing which would almost certainly have fatal consequences).

Given the relatively small sample sizes in both of these trials, there is substantial (estimatable) uncertainty in the estimates from the trials themselves and additional (non-estimatable) uncertainty related to the extent to which these trials are representative of all fishing of a given type, particularly as both trials were undertaken overseas. The binomial 95% confidence range (calculated using the Clopper-Pearson “exact” method) for the ratio of total fatalities to observed captures in Brothers *et al.*’s (2010) longline trial is 1.8–2.5 (mean 2.1), and that for Watkins *et al.*’s trawl warp trial is 5–122 (mean 15.0 fatalities per observed capture). Abraham (2010) estimated that there were 244 (95% c.i. 190–330) warp strikes by large birds for every one observed captured, and 6 440 (3400–20 000) warp strikes by small birds for every one observed captured (although small birds tend to be caught in the net rather than by warps). There is also uncertainty in the relative frequencies and consequences of different types of encounters with trawl warps in New Zealand fisheries (Abraham 2010, Richard *et al.* 2011 Appendix B).

#### 5.4.3.5.3. Mortalities in non-commercial fisheries.


Little is known about the nature and extent of incidental captures of seabirds in non-commercial fisheries, either in New Zealand or globally (Abraham *et al.* 2010). In New Zealand, participation in recreational fishing is high and 2.5% of the adult population are likely to be fishing in a given week (mostly using rod and line). Because of this high participation rate, even a low rate of interactions between individual fishers and seabirds could have population-level impacts. A boat ramp survey of 765 interviews at two locations during the summer of 2007–08 revealed that 47% of fishers recalled witnessing a bird being caught some time in the past. Twenty-one birds were reported caught on the day of the interview at a capture rate of 0.22 (95% c.i.: 0.13–0.34) birds per 100 hours of fishing. Observers on 57 charter trips recorded seabird captures at rate of 0.36 (0.09–0.66) birds per 100 fisher hours. The most frequently reported type of bird caught in rod and line fisheries were petrels and gulls. Captures of albatrosses, shags, gannets, penguins, and terns were also recalled.

The ramp surveys reported by Abraham *et al.* (2010) were limited and covered only two widely-separated parts of the New Zealand coastline. However, they also report two other pieces of information that suggest non-commercial captures are likely to be very widespread. First, the Ornithological Society of New Zealand’s beach patrol scheme records seabird hookings and entanglements as a common occurrence throughout New Zealand. Second, returns of banded birds caught in fisheries (separating commercial and non-commercial fisheries is very difficult) are very widely distributed around the coast (Figure 5.22).

Noting that our understanding of seabird capture rates in amateur fisheries is very sketchy, it is possible to make first-order estimates of total captures using information on fishing effort. For example, in the north-eastern region where most of Abraham *et al.*’s (2010) interviews were conducted, there were an estimated 4.8 (4.4–5.2) million fisher hours rod and line fishing from trailer

boats in 2004–05 (Hartill *et al.* 2007). Applying Abraham *et al.*'s (2010) capture rate leads to an estimate of 11 500 (6600–17 200) captures per year in this area. Based on estimates of nationwide recreational fishing effort, this could increase to as many as 40 000 bird captures annually. Most birds captured by amateur fishers were reported to have been released unharmed (77% of the incidents recalled) and only three people reported incidents where the bird died. Because of likely recall biases and the qualitative nature of the survey, the fate of birds that are captured by amateur fishers remains unclear.

Non-commercial fishers are allowed to use setnets in New Zealand and two studies suggest that these have an appreciable bycatch of seabirds. A study of captures in non-commercial setnets in Portobello Bay, Otago Harbour, between 1977 and 1985 (Lalas 1991) suggested spotted shags were the most frequently caught taxa (82 recorded, compared with 14 Stewart Island shags and two little shags). Lalas (1991) suggested that up to 800 spotted shags (20% of the local population) may have been caught in the summer of 1981/82. A broader-scale study of yellow-eyed penguin mortality in setnets in southern New Zealand (Darby and Dawson 2000) suggested non-negligible captures of this species by non-commercial fishers, also reporting other seabirds like spotted shags and little blue penguin.



**Figure 5.22 (reproduced from Abraham *et al.* 2010): Distribution of the reported capture locations for banded seabirds reported as being captured in fishing gear, 1952–2007. Note, band recovery locations are reported with low spatial precision and some of the inland locations may be correct.**

#### 5.4.3.5.4. Out of zone mortality.

Robertson *et al.* (2003) mapped the distribution of the 25 breeding (mainly endemic) New Zealand seabird taxa they considered most at risk outside New Zealand waters. These ranged widely: 4 used the South Atlantic; 4 the Indian Ocean; 22 Australian waters and the Tasman Sea; 15 used the South Pacific Ocean as far afield as Chile and Peru; and 6 used the North Pacific Ocean as far north as the Bering Sea. These taxa therefore use the national waters of at least 18 countries. For example, the

level-2 risk assessment described by Richard *et al.* (2011) includes only that part of the range of each taxon contained within New Zealand waters, but many including commonly-caught seabirds like white-capped albatross and white-chinned petrel range much further and are vulnerable to fisheries in other parts of the world. For instance, fatalities of white-capped albatross outside the New Zealand EEZ greatly exceed fatalities within the zone (Baker 2007, Francis 2012, Table 5.16), and more than 10 000 white-chinned petrel are killed off South America each year (Phillips *et al.* 2006), noting that reliable records are not available for most of the fisheries involved. Based on similar analyses, Moore and Zydels (2008) concluded that a population-based, multi-gear and multi-national framework is required to identify the most significant threats to wide-ranging seabird populations and to prioritize mitigation efforts in the most problematic areas. To that end, the Agreement for the Conservation of Albatrosses and Petrels (ACAP) adopted a global prioritisation framework at the Fourth Session of the Meeting of the Parties (MoP4) in April 2012 (ACAP 2012).

**Table 5.16 (after Francis 2012): Estimates of the number of white-capped albatrosses killed annually, by fishery. The first two columns are from Baker *et al.* (2007) (mid-point where a range was presented), including their assessment of reliability (L = low, M-H = medium-high, H = high). Updated estimates are from Watkins *et al.* (2008, \*) and Petersen *et al.* (2009, \*\*). Estimates not already corrected for cryptic mortality are either doubled to allow for this (\*\*\*) or replaced by estimates of potential fatalities from Richard *et al.* (2011, \*\*\*), noting that potential fatalities may considerably overestimate actual fatalities.**

Fishery	From Baker <i>et al.</i> 2007		Updated	Incl. Cryptic mortality
South African demersal trawl	4 750	(L)	* 6650	6 650
Asian distant-water longline	1 255	(L)	–	*** 2 510
Namibian demersal trawl	910	(L)	* 1270	1 270
Namibian pelagic longline	180	(L)	** 195	*** 390
NZ hoki and squid trawl	513	(MH)	–	**** 4 920
NZ longline	60	(MH)	–	**** 199
Australian (line fisheries)	15	(MH)	–	*** 30
South African pelagic longline	570	(H)	** 570	*** 1 140
Total	8 210	–	–	17 110

#### 5.4.3.5.5. Other sources of anthropogenic mortality.


Taylor (2000) listed a wide range of threats to New Zealand seabirds including introduced mammals, avian predators (weka), disease, fire, weeds, loss of nesting habitat, competition for nest sites, coastal development, human disturbance, commercial and cultural harvesting, volcanic eruptions, pollution, plastics and marine debris, oil spills and exploration, heavy metals or chemical contaminants, global sea temperature changes, marine biotoxins, and fisheries interactions. Relatively little is known about most of these factors, but the parties to ACAP have agreed a formal prioritisation process to address and prioritise major threats (ACAP 2012). Croxall *et al.* (2012) identified the main priorities as: protection of Important Bird Area (IBA) breeding, feeding, and aggregation sites; removal of invasive, especially predatory, alien species as part of habitat and species recovery initiatives. Lewison *et al.* (2012) identified similar research priorities (in addition to direct fishing-related mortality), including: understanding spatial ecology; tropho-dynamics; response to global change; and management of anthropogenic impacts such as invasive species, contaminants, and protected areas. Non fishing-related threats to seabirds in New Zealand are largely the mandate of the Department of Conservation and a detailed description is beyond the scope of this document (although causes of mortality other than fishing are clearly relevant to the interpretation of risk assessment restricted to the direct effects of fishing). These threats are identified in DOC's Action Plan for Seabird Conservation in New Zealand (Taylor 2000).

## 5.5. Indicators and trends

Population size	Multiple species and populations: see Taylor (2000)																																			
Population trend	Multiple species and populations: see Taylor (2000)																																			
Threat status	Multiple species and populations: see Miskelly <i>et al.</i> (2008) and updates																																			
Number of interactions	<p>In the 2010/11 October fishing year, there were an estimated 4931 seabird captures (excluding cryptic mortalities) across all trawl and longline fisheries (excluding about 14% of bottom longline effort that could not be included in the models) (Data version v20121101). About 57% of the captures were in trawl fisheries, 15% in surface longline fisheries, and 28% in bottom longline fisheries:</p> <table><tr><td>Bird group</td><td>Trawl</td><td>Surface longline</td><td>Bottom longline</td><td>All these methods</td></tr><tr><td>White-capped albatross</td><td>356</td><td>84</td><td>2</td><td>442</td></tr><tr><td>Other albatrosses</td><td>808</td><td>287</td><td>257</td><td>1 352</td></tr><tr><td>White-chinned petrel</td><td>540</td><td>34</td><td>422</td><td>996</td></tr><tr><td>Sooty shearwater</td><td>488</td><td>2</td><td>69</td><td>559</td></tr><tr><td>Other birds</td><td>596</td><td>333</td><td>652</td><td>1 581</td></tr><tr><td>All birds combined</td><td>2 788</td><td>740</td><td>1 403</td><td>4 931</td></tr></table>	Bird group	Trawl	Surface longline	Bottom longline	All these methods	White-capped albatross	356	84	2	442	Other albatrosses	808	287	257	1 352	White-chinned petrel	540	34	422	996	Sooty shearwater	488	2	69	559	Other birds	596	333	652	1 581	All birds combined	2 788	740	1 403	4 931
Bird group	Trawl	Surface longline	Bottom longline	All these methods																																
White-capped albatross	356	84	2	442																																
Other albatrosses	808	287	257	1 352																																
White-chinned petrel	540	34	422	996																																
Sooty shearwater	488	2	69	559																																
Other birds	596	333	652	1 581																																
All birds combined	2 788	740	1 403	4 931																																
Trend in interactions	<p>Captures of all birds combined show a decreasing trend between 2002/03 and 2010/11 (Data version v20121101) but there are substantial differences in trends between species and fisheries. Captures of white-capped albatross have decreased, especially in offshore trawl fisheries, whereas captures of white-chinned petrel have increased:</p> <div><p>The figure consists of six stacked bar charts arranged in a 3x2 grid. Each chart shows the estimated captures of a specific bird species or group from 2003 to 2011. The y-axis for each chart is labeled 'Estimated captures'. The x-axis for each chart shows the years 2003, 2005, 2007, 2009, and 2011. The legend for all charts indicates three categories: Trawl (blue), SLL (red), and BLL (green). The species and groups are: White-capped albatross, Other albatrosses, Sooty shearwater, White-chinned petrel, Other birds, and All birds combined. The charts show varying trends over time, with some species showing a general decrease in captures and others showing an increase or more stable levels.</p></div>																																			
<p>Capture rate trends (excluding cryptic mortalities) are described for the four fisheries estimated to account for most of captures of a species (accounting for 80% or more of the total). Capture rates of white-capped albatross have fallen in trawl fisheries for</p>																																				

*Trend in interactions contd.*

hoki and squid but have remained steady in inshore trawl fisheries and increased in the southern bluefin tuna fishery. Capture rates for white-chinned petrel have increased in trawl fisheries for squid and scampi but have remained steady in longline fisheries. Capture rates of sooty shearwater have declined in the ling longline fishery but have fluctuated without trend in other key fisheries.



## 5.6. References

- Abraham ER (2007). Summary of data collected during the southern squid fishery mincing trial. Research report prepared by Dragonfly for the Department of Conservation, Wellington, New Zealand. 39 p.
- Abraham ER (2009). Batching of waste to reduce seabird numbers behind trawl vessels. Unpublished report prepared for the Department of Conservation, Wellington, New Zealand.
- Abraham ER (2010). Warp strike in New Zealand trawl fisheries, 2004–05 to 2008–00. *New Zealand Aquatic Environment and Biodiversity Report No. 60*. 29 p.
- Abraham ER (2010). Mincing offal to reduce the attendance of seabirds at trawlers. Report prepared by Dragonfly for Department of Conservation, Wellington, New Zealand. 28 p.
- Abraham ER; Berkenbusch KN; Richard Y (2010). The capture of seabirds and marine mammals in New Zealand non-commercial fisheries *New Zealand Aquatic Environment and Biodiversity Report No. 64*. 52 p.
- Abraham ER; Pierre JP; Middleton DAJ; Cleal J; Walker NA; Waugh SM (2009). Effectiveness of fish waste management strategies in reducing seabird attendance at a trawl vessel. *Fisheries Research* 95: 210–219
- Abraham ER; Thompson FN (2009a). Capture of protected species in New Zealand trawl and longline fisheries, 1998–99 to 2006–07. *New Zealand Aquatic Environment and Biodiversity Report No. 32*.
- Abraham ER; Thompson FN (2009b). Warp strike in New Zealand trawl fisheries, 2004–05 to 2006–07. *New Zealand Aquatic Environment and Biodiversity Report No. 33*. 21 p.
- Abraham ER; Thompson FN (2010). Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2006–07. Final Research Report for research project PRO2007-01 (Unpublished report held by Ministry of Fisheries, Wellington).
- Abraham ER; Thompson FN (2011a). Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 79*. 74 p.
- Abraham ER; Thompson FN (2011b). Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 80*. 170 p.
- Abraham ER; Thompson FN; Oliver MD (2010). Summary of the capture of seabirds, marine mammals and turtles in New Zealand commercial fisheries, 1998–99 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 45*.
- ACAP (2011). Report of the Fourth Meeting of the Seabird Bycatch Working Group, Guayaquil, Ecuador, 22–24 August 2011. 94 p. AC6 Doc 14 Rev4.
- ACAP (2012) ACAP Conservation Priorities. Fourth Meeting of the Parties, Lima, Peru, 23 – 27 April 2012, available at <http://www.acap.aq/docman/english/meeting-of-the-parties/mop4/mop4-final-report>. 13 p.
- Arcos JM; Oro D (1996). Changes in foraging range of Audouin's gulls *Larus audouinii* in relation to a trawler moratorium in the western Mediterranean. *Waterbirds* 19:128–131.
- Astles KL; Holloway MG; Steffe A; Green M; Ganassin C; Gibbs PJ (2006). An ecological method for qualitative risk assessment and its use in the management of fisheries in New South Wales, Australia. *Fisheries Research* 82: 290–303.
- Ayers, D.; Francis, M.P.; Griggs, L.H.; Baird, S.J. (2004). Fish bycatch in New Zealand tuna longline fisheries, 2000/01 and 2001/02. *New Zealand Fisheries Assessment Report 2004/46*. 47 p.
- Baird SJ 1994: Nonfish Species and Fisheries Interactions Working Group Report. N.Z. Fisheries Assessment Working Group Report 94/1. MAF Fisheries, N.Z. Ministry of Agriculture and Fisheries, Wellington. 54 p.
- Baird SJ 1995: Nonfish Species and Fisheries Interactions Working Group Report — April 1995. N.Z. Fisheries Assessment Working Group Report 95/1. MAF Fisheries, N.Z. Ministry of Agriculture and Fisheries, Wellington. 24 p.
- Baird SJ 1996: Nonfish Species and Fisheries Interactions Working Group Report — May 1996. N.Z. Fisheries Assessment Working Group Report 96/1. Ministry of Fisheries, Wellington. 34 p.
- Baird SJ 1997: Report on the incidental capture of nonfish species during fishing operations in New Zealand waters. (Unpublished report completed as part of the Ministry of Fisheries SANF01 contract.) 15 p plus appendices on New Zealand fur seal-trawl fishery interaction (54 p) and seabird-tuna longline fishery interaction (34 p).
- Baird SJ (1998). Estimation of nonfish bycatch in commercial fisheries in New Zealand waters, 1990–91 to 1993–94. Final Research Report for Ministry of Fisheries Research Project ENV9701.
- Baird SJ 1999: Estimation of nonfish bycatch in commercial fisheries in New Zealand waters, 1997–98. Unpublished report completed for Objective 1 of Ministry of Fisheries Project ENV9801. 57 p.
- Baird SJ (2000a). Estimation of the incidental capture of seabird and marine mammal species in commercial fisheries in New Zealand waters, 1998–99. *New Zealand Fisheries Assessment Report 2001/14* 43 p.
- Baird SJ (2000b). Incidental capture of seabirds in New Zealand tuna longline fisheries. In Flint, E., & Swift, K. (Eds.), p. 126, Second Albatross Conference on the Biology and Conservation of Albatrosses and other Petrels, Honolulu, Hawaii, USA, 8–12 May 2000. Abstracts and Oral Presentations. *Marine Ornithology* 28: 125–152.
- Baird SJ (2001a). Estimation of the incidental capture of seabird and marine mammal species in commercial fisheries in New Zealand waters, 1998–99. *New Zealand Fisheries Assessment Report 2001/14*. 43 p.
- Baird SJ (Comp. & Ed.) (2001b): Report on the International Fishers' Forum on Solving the incidental Capture of Seabirds in Longline Fisheries, Auckland, New Zealand, 6–9 November 2000. Department of Conservation, Wellington, New Zealand. 63 p.
- Baird SJ (2003). New Zealand breeding seabirds: human-induced mortality — a review. Draft report prepared for the Ministry of Fisheries Project ENV2000/09. 74 p.

- Baird SJ (2004a). Estimation of the incidental capture of seabird and marine mammal species in commercial fisheries in New Zealand waters, 1999–2000. New Zealand Fisheries Assessment Report 2004/41. 56 p.
- Baird SJ (2004b). Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2000–01. New Zealand Fisheries Assessment Report 2004/58. 63 p.
- Baird SJ (2004c). Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2001–02. New Zealand Fisheries Assessment Report 2004/60. 51 p.
- Baird SJ (2005). Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2002–03. New Zealand Fisheries Assessment Report 2005/2. 50 p.
- Baird SJ; Francis M; Griggs LH; Dean HA (1998). Annual review of bycatch in southern bluefin and related tuna longline fisheries in the New Zealand 200 n. mile Exclusive Economic Zone. CCSBT-ERS/9806/31. (Third Meeting of the Ecologically Related Species Working Group, Tokyo, 9–13 June 1998.)
- Baird SJ; Gilbert DJ (2010). Initial assessment of risk posed by commercial fisheries to seabirds breeding in New Zealand waters. *Aquatic Environment and Biodiversity Report No. 50*. 99 p.
- Baird SJ; Gilbert DJ; Smith MH (2006) Review of environmental risk assessment methodologies with relevance to seabirds and fisheries within New Zealand waters. Final Research Report for Ministry of Fisheries project ENV2005/01 Objective 3.
- Baird SJ; Griggs LH (2004). Estimation of within-season chartered southern bluefin tuna (*Thunnus maccoyii*) longline seabird incidental captures, 2002. New Zealand Fisheries Assessment Report 2004/42. 15 p.
- Baird SJ; Sanders BM; Dean HA; Griggs LH (1999). Estimation of nonfish bycatch in commercial fisheries in New Zealand waters, 1990/91 to 1993/94. Final Research Report for Ministry of Fisheries Project ENV9701 Objective 1. 63 p.
- Baird SJ; Smith MH (2008). Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2005–06. *New Zealand Aquatic Environment and Biodiversity Report No. 18*. 124 p.
- Baker GB; Double MC; Gales R; Tuck GN; Abbott CL; Ryan PG; Petersen SL; Robertson CJR; Alderman R (2007). A global assessment of the impact of fisheries related mortality on shy and white-capped albatrosses: conservation implications. *Biological Conservation* 137: 319–333.
- Baker GB; Jensz K; Cunningham R (2009). Data collection of demographic, distributional and trophic information on the white-capped albatross to allow estimation of effects of fishing on population viability - 2008 field season. Final Research Report for Ministry of Fisheries project PRO2006-01H. 13 p.
- Baker GB; Jensz K; Cunningham R (2010). Data collection of demographic, distributional and trophic information on the white-capped albatross to allow estimation of effects of fishing on population viability - 2009 field season. Final Research Report for Ministry of Fisheries project PRO2006-01I. 13 p.
- Baker GB; Jensz K; Double M; Cunningham R (2007). Data collection of demographic, distributional and trophic information on selected seabird species to allow estimation of effects of fishing on population viability. Final Research Report for Ministry of Fisheries project PRO2006-01F. 12 p.
- Baker GB; Jensz K; Double M; Cunningham R (2008). Data collection of demographic, distributional and trophic information on selected seabird species to allow estimation of effects of fishing on population viability. Final Research Report for Ministry of Fisheries project PRO2006-01G. 11 p.
- Bartle JA (1991). Incidental capture of seabirds in the New Zealand subantarctic squid trawl fishery, 1990. *Bird Conservation International* 1: 351–359.
- Bartumeus F; Giuggioli L; Louzao M; Bretagnolle V; Oro D; Levin SA (2010). Fishery discards impact on seabird movement patterns at regional scales. *Current Biology* 20: 215–222.
- Bell EA; Sim JL; Scofield P; Francis RICC (2012). Population parameters of the black petrels (*Procellaria parkinsoni*) on Great Barrier Island (Aotea Island), 2009/10. Report for Conservation Services Programme Project POP2009-01, near-final (post-review) draft. 82 p. Department of Conservation, Wellington.
- Brothers NP (1991). Albatross mortality and associated bait loss in the Japanese longline fishery in the Southern Ocean. *Biological Conservation* 55: 255–268.
- Brothers NP; Cooper J; Lokkeborg S (1999). The incidental catch of seabirds by longline fisheries: worldwide review and technical guidelines for mitigation. FAO Fisheries Circular No. 937. 107 p.
- Brothers NP; Duckworth AR; Safina C; Gilman EL (2010). Seabird bycatch in pelagic longline fisheries is grossly underestimated when using only haul data. *PLoS ONE* 5: e12491.
- Bull LS (2007). A review of methodologies for mitigating incidental catch of seabirds in New Zealand Fisheries. *DOC Research & Development Series* 263. Department of Conservation, Wellington. 57p.
- Bull LS (2009). New mitigation measures reducing seabird by-catch in trawl fisheries. *Fish and Fisheries* 8: 31–56.
- Campbell JL; Gallagher C (2007). Assessing the relative effects of fishing on the New Zealand marine environment through risk analysis. *ICES Journal of Marine Science*, 64: 256–270.
- Carey, C. (2005). Carefree's cunning contraption. *Seafood New Zealand* 13(6): 44–45.
- Croxall JP (2008). The role of science and advocacy in the conservation of Southern Ocean albatrosses at sea. *Bird Conservation International* 18: S13–29.
- Croxall JP; Butchart SHM; Lascelles B; Stattersfield AJ; Sullivan B; Symes A; Taylor P (2012). Seabird conservation status, threats and priority actions: a global assessment. *Bird Conservation International* 22: 1–34
- Croxall JP; Prince PA; Rothery P; Wood AG (1998). Population changes in albatrosses at South Georgia. Pages 68–83 in G Robertson; R Gales, eds. *Albatross biology and conservation*: Chipping Norton, Australia: Surrey Beatty & Sons.
- Crysel, S. (2002). Baffling birds brings benefits. *Seafood New Zealand* 10(10): 60–61.
- Darby JT; Dawson SM (2000). Bycatch of yellow-eyed penguin (*Megadyptes antipodes*) in gillnets in New Zealand waters 1979–1997. *Biological Conservation* 93: 327–332.
- Dillingham PW; Fletcher D (2011). Potential biological removal of albatrosses and petrels with minimal demographic information. *Biological Conservation* 144: 1885–1894

- Favero M; Blanco G; Garcia G; Copello S; Seco Pon JP; Frere E; Quintana F; Yorio P; Rabuffetti F; Canete G; Gandini P (2010). Seabird mortality associated with ice trawlers in the Patagonian shelf: effect of discards on the occurrence of interactions with fishing gear. *Animal Conservation* 14:131–139.
- Filippi D; Waugh SM; Nicol S (2010). Revised spatial risk indicators for seabird interactions with longline fisheries in the western and central Pacific. Western and Central Pacific Fisheries Commission, Scientific Committee (2010) Paper EB-IP-01.
- Fletcher D; MacKenzie D; Dillingham P; (2008). Modelling of impacts of fishing-related mortality on New Zealand seabird populations. Final Research Report for Ministry of Fisheries project, 17 December 2008.
- Fletcher WJ (2005). The application of qualitative risk assessment methodology to prioritize issues for fisheries management. *ICES Journal of Marine Science* 62:1576–1587.
- FAO (2009). Fishing operations. 2. Best practices to reduce incidental catch of seabirds in capture fisheries. *FAO Technical Guidelines for Responsible Fisheries. No. 1, Suppl. 2*. Rome, FAO. 2009. 49p.
- FAO (2010). Report of the Technical Consultation to Develop International Guidelines on Bycatch Management and Reduction of Discards, Rome, 6–10 December 2010. *FAO Fisheries and Aquaculture Report No. 957*. 41 p.
- Francis RICC (2012). Fisheries risks to the population viability of white-capped albatross *Thalassarche steadi*. *New Zealand Aquatic Environment and Biodiversity Report. No. 104*. 24 p.
- Francis RICC; Bell EA (2010). Fisheries risks to population viability of black petrel (*Procellaria parkinsoni*). *New Zealand Aquatic Environment and Biodiversity Report No. 51*. Ministry of Fisheries, Wellington
- Francis RICC; Elliott G; Walker K (in press). Fisheries risks to the viability of Gibson's wandering albatross *Diomedea gibsoni*. *New Zealand Aquatic Environment and Biodiversity Report*.
- Francis RICC; Sagar PM (2012): Modelling the effect of fishing on southern Buller's albatross using a 60-year dataset, *New Zealand Journal of Zoology* 39:1, 3–17.
- Francis RICC; Sagar PM; Fu D (2008). Status of the Snares Islands population of southern Buller's albatross. Final Research Report for Year 2 of Ministry of Fisheries Research Project PRO200602. 138 p.
- Furness RW (2003). Impacts of fisheries on seabird communities. *Sci. Mar.* 67:33–45.
- Furness RW; Ensor K; Hudson AV (1992). The use of fishery waste by gull populations around the British Isles. *Ardea* 80: 105–113.
- Hartill B; Bian R; Armiger H; Vaughan M; Rush N (2007). Recreational marine harvest estimates of snapper, kahawai, and kingfish in QMA 1 in 2004–05. *New Zealand Fisheries Assessment Report 2007/26*. 44 p.
- Hobday AJ; Smith A; Webb H; Daley R; Wayte S; Bulman C; Dowdney J; Williams A; Sporic M; Dambacher J; Fuller M; Walker T (2007). Ecological risk assessment for the effects of fishing: methodology. Report R04/1072 for the Australian Fisheries Management Authority, Canberra.
- Kirby J; Hobday A (2007) Ecological risk assessment for the effects of fishing the western and central Pacific Ocean: productivity-susceptibility analysis. Paper presented to the Scientific Committee of the Third Regular Session of the Western and Central Pacific Fisheries Commission, WCPFC-SC3-EB SWG/WP-1
- Lalas C (1991). Assessment of bird kills in set nets in Otago Harbour over a period of eight years (1977–1985). Unpublished report held by Department of Conservation, Dunedin.
- Lewison RL; Crowder LB; Read AJ; Freeman SA (2004). Understanding impacts of fisheries bycatch on marine megafauna. *Trends in Ecology and Evolution* 19: 598–604.
- Lewison RL; Oro D; Godley BJ; Underhill L; Bearhop S; Wilson RP; Ainley D; Arcos JM; Boersma PD; Borboroglu PG; Boulenger T; Frederiksen M; Genovart M; González-Solís J; Green JA; Grémillet D; Hamer KC; Hilton GM; Hyrenbach KD; Martínez-Abraín A; Montevecchi WA; Phillips RA; Ryan PG; Sagar PM; Sydeman WJ; Wanless S; Watanuki Y; Weimerskirch H; Yorio P (2012). Review: Research priorities for seabirds: improving conservation and management in the 21st century. *Endangered Species Research* 17: 93–121.
- Louzao M; Arcos JM; Guijarro B; Valls M; Oro D (2011). Seabird-trawling interactions: factors affecting species-specific to regional community utilisation of fisheries waste. *Fisheries Oceanography* 20: 263–277.
- Maunder MN; Houston DM; Dunn A; Seddon PJ; Kendrick TH (2007). Assessment to risk of yellow-eyed penguins *Megadyptes antipodes* from fisheries incidental mortality in New Zealand fisheries and definition of information requirements for managing fisheries related risk. Final Research Report for Ministry of Fisheries project ENV2005-13. 28 p.
- Melvin EF; Dietrich KS; Fitzgerald S; Cordoza T (2010). Reducing seabird strikes with trawl cables in the pollock catcher-processor fleet in the eastern Bering Sea. Agreement on the Conservation of Albatrosses and Petrels, SBWG-3 Doc 14 Rev1, Hobart, Australia, 18 p
- MFish-DOC (2004). National plan of action to reduce the incidental catch of seabirds in New Zealand fisheries. NZ Ministry of Fisheries and Department of Conservation, Wellington. 62 p.
- Middleton DAJ; Abraham ER (2007). The efficacy of warp strike mitigation devices: trials in the 2006 squid fishery. Final Research Report for Ministry of Fisheries research project IPA2006/02.
- Miskelly CM; Dowding JE; Elliott GP; Hitchmough RA; Powlesland RG; Robertson HA; Sagar PM; Scofield RP; Taylor GA (2008). Conservation status of New Zealand birds, 2008. *Notornis* 55: 117–135
- Moore JE; Zydels R (2008). Quantifying seabird bycatch: where do we go from here? *Animal Conservation* 11: 257–259
- Newman J; Scott D; Bragg C; McKechnie S; Moller H; Fletcher D (2009). Estimating regional population size and annual harvest intensity of the sooty shearwater in New Zealand. *New Zealand Journal of Zoology* 36: 307–323.
- Oro D; Cam E; Pradel R; Martínez-Abraín A (2004). Influence of food availability on demography and local population dynamics in a long-live seabird. *Proc. R. Soc. Lond. B Biol. Sci.* 271:387–396.
- Oro D; Furness RW (2002). Influences of food availability and predation on survival of kittiwakes. *Ecology* 83:2516–2528.
- Oro D; Pradel R; Lebreton JD (1999). Food availability and nest predation influence life history traits in Audouin's gull, *Larus audouinii*. *Oecologia* 118:438–445.

- Oro D; Ruiz X; Jover L; Pedrocchi V; Gonzalez-Solis J (1997). Audouin's gull diet and adult time budget responses on changes in food availability induced by commercial fisheries. *Ibis* 139:631–637.
- OSNZ (2010). Checklist of the birds of New Zealand and the Ross Dependency, Antarctica. Fourth Edition. Te Papa Press in association with the Ornithological Society of New Zealand Inc. Wellington.
- Petersen SL; Honig MB; Ryan PG; Underhill LG (2009). Seabird bycatch in the pelagic longline fishery off southern Africa. *African Journal of Marine Science* 31: 191–204.
- Phillips RA; Silk JRD; Croxall JP; Afanasyev V (2006). Year-round distribution of white-chinned petrels from South Georgia: relationships with oceanography and fisheries. *Biological Conservation* 129:336–347.
- Pierre JP; Abraham ER; Middleton DAJ; Cleal J; Bird R; Walker NA; Waugh SM (2010). Reducing interactions between seabirds and trawl fisheries: Responses to foraging patches provided by fish waste batches. *Biological Conservation* 143: 2779–2788.
- Pierre JP; Abraham ER; Cleal J; Middleton DAJ (2012). Reducing effects of trawl fishing on seabirds by limiting foraging opportunities provided by fishery waste. *Emu* 112: 244–254. CSIRO.
- Pierre JP; Abraham ER; Middleton DAJ; Cleal J; Bird R; Walker NA; Waugh SM (2010). Reducing interactions between trawl fisheries and seabirds: responses to foraging patches provided by fish waste batches. *Biological Conservation* 143: 2779–2788.
- Pierre JP; Abraham ER; Richard Y (2012). Concluding six years of research on seabird bycatch reduction through modified discharge management regimes: Is batch discharge better than ad-hoc discharge from trawl vessels? SBWG-4 Doc 14, Progress report on ACAP Project Application 2010-04.
- Poncet S; Robertson G; Phillips RA; Lawton K; Phalan B; Trathan PN; Croxall JP (2006). Status and distribution of wandering, black-browed and grey-headed albatrosses breeding at South Georgia. *Polar Biology* 29: 772–781.
- Ramm KC (2011). Conservation Services Programme Observer Report for the period 1 July 2008 to 30 June 2009. Available at [www.doc.govt.nz/documents/science-and-technical/2008-09-csp-observer-report.pdf](http://www.doc.govt.nz/documents/science-and-technical/2008-09-csp-observer-report.pdf)
- Ramm KC (2012). Conservation Services Programme Observer Report for 1 July 2009 to 30 June 2010. Available: [www.doc.govt.nz/documents/conservation/marine-and-coastal/marine-conservation-services/csp-observer-report-2009-10.pdf](http://www.doc.govt.nz/documents/conservation/marine-and-coastal/marine-conservation-services/csp-observer-report-2009-10.pdf)
- Rice J. 2006. Impacts of Mobile Bottom Gears on Seafloor Habitats, Species, and Communities: A Review and Synthesis of Selected International Reviews. *Canadian Science Advisory Secretariat Research Document* 2006/057. 35 p. (available from: [http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006\\_057\\_e.pdf](http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006_057_e.pdf)).
- Richard Y; Abraham EA; Filippi D (2011). Assessment of the risk to seabird populations from New Zealand commercial fisheries. Final Research Report for Ministry of Fisheries projects IPA2009-19 and IPA 2009-20.
- Robertson CJR; Bell EA; Sinclair N; Bell BD (2003). Distribution of seabirds from New Zealand that overlap with fisheries worldwide. *Science for Conservation* 233. Department of Conservation, Wellington. 102 p.
- Rowe SJ (2009). Conservation Services Programme Observer Report for the period 1 July 2004 to 30 June 2007. *DOC Marine Conservation Services Series 1*. Department of Conservation, Wellington. 97p
- Rowe SJ (2010). Conservation Services Programme Observer Report for the period 1 July 2007 to 30 June 2008. *DOC Marine Conservation Services Series 4*. Department of Conservation, Wellington. 97p
- Rowe SJ (2010). Level 1 risk assessment for incidental seabird mortality associated with New Zealand Fisheries in the NZ-EEZ. Department of Conservation. 75 pp.
- Sagar PM; Torres L; Thompson D (2010). Demography and distribution of Buller's Albatrosses *Thalassarche bulleri bulleri*: Final research report of the 2010 field season. NIWA Client Report: CHC2010-156, December 2010, prepared for Department of Conservation project
- Sagar PM; Stahl JC (2005). Increases in the numbers of breeding pairs in two populations of Buller's albatross (*Thalassarche bulleri bulleri*). *Emu* 105: 49–55.
- Sharp B (2009). A risk assessment framework for incidental seabird mortality associated with New Zealand fishing in the New Zealand EEZ. Unpublished report to the Ministry of Fisheries May 2009.
- Sharp B; Parker S; Smith N (2009). An impact assessment framework for bottom fishing methods in the CCAMLR area. *CAMMLR Science*, 15 June 2009.
- Soykan CU; Moore JE; Zydels R; Crowder LB; Safina C; Lewison RL (2008). Why study bycatch? An introduction to the Theme Section on fisheries bycatch. *Endangered Species Research* 5: 91–102.
- Sullivan BJ; Brickle P; Reid TA; Bone DG; Middleton DAJ (2006). Mitigation of seabird mortality on factory trawlers: Trials of three devices to reduce warp cable strikes. *Polar Biology* 29: 745–753.
- Sullivan, BJ; Reid TA; Bugoni L (2006). Seabird mortality on factory trawlers in the Falkland Islands and beyond. *Biological Conservation* 131: 495–504.
- Taylor GA (2000). Action plan for seabird conservation in New Zealand. Part A, threatened seabirds. *Threatened Species Occasional Publication No. 16*. 234 p. Dept. of Conservation, Wellington.
- Thompson FN; Abraham ER (2009). Six monthly summary of the capture of protected species in New Zealand commercial fisheries, summer 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 35*.
- Tuck GN; Polacheck T; Croxall JP; Weimerskirch H; Ryan P; Nel D; Wayte S; Bulman CM (2004). Modelling the impact of fishery incidental mortality on three populations of wandering albatross. Pages 149–241 in GN Tuck (ed.) A comprehensive study of the ecological impacts of the worldwide pelagic longline industry: Southern Hemisphere studies. Melbourne, Australia: CSIRO.
- Tuck GN; Phillips RA; Small C; Thomson RB; Klaer N; Taylor F; Wanless R; Arrizabalaga H (2011) An assessment of seabird–fishery interactions in the Atlantic Ocean. *ICES Journal of Marine Science* 68: 1628–1637.
- Vié J-C; Hilton-Taylor C; Stuart SN (eds.) (2009). Wildlife in a changing world – an analysis of the 2008 IUCN Red List of Threatened Species. Gland, Switzerland: IUCN. 180 pp.

- Wagner EL; Boersma PD (2011). Effects of fisheries on seabird community ecology. *Reviews in Fisheries Science* 19: 157–167.
- Walker K; Elliott G (2009). Gibson's wandering albatross research Adams Island 2009. 11 p. Unpublished report produced for the Department of Conservation.
- Watkins BP; Petersen SL; Ryan PG (2008). Interactions between seabirds and deep-water hake trawl gear: an assessment of impacts in South African waters. *Animal Conservation* 11: 247–254.
- Waugh SM; Baker GB; Gales R; Croxall JP (2008). CCAMLR process of risk assessment to minimise the effects of longline fishing mortality on seabirds. *Marine Policy* 32: 442–454.
- Waugh SM; Filippi D; Walker NA; Kirby DS (2008). Updated preliminary results of an ecological risk assessment for seabirds and marine mammals with risk of fisheries interactions. Western Central Pacific Fisheries Commission Scientific Committee 4th Regular Session, August 2008, Port Moresby, Papua New Guinea. WCPFC-SC4-2008/EB-WP-2.
- Waugh SM; Filippi D (2009). Ecological risk assessment for seabirds in New Zealand fisheries: primary components of ecological risk assessment methodology and calculating vulnerability and productivity for species. Research Progress Report for Ministry of Fisheries Research Project IPA2009-19.
- WGSE (2011). Report of the International Council for the Exploration of the Sea (ICES) Working Group on Seabird Ecology (WGSE), 1–4 November 2011, Madeira, Portugal. ICES CM 2011/SSGEF:07. 73 p.
- Wienecke B; Robertson G (2002). Seabird and seal-fisheries interactions in the Australian Patagonian toothfish *Dissostichus eleginoides* trawl fishery. *Fisheries Research* 54:253–265.

## **THEME 2: NON-PROTECTED BYCATCH**

## 6. Non-protected species (fish and invertebrates) bycatch

<i>Scope of chapter</i>	<p>This chapter outlines the main non-protected bycatch species (fish and invertebrates) and annual levels and trends in bycatch and discards in New Zealand’s major offshore fisheries. Research in this field is conducted fishery by fishery and this first summary of current knowledge reflects that strategy. It is expected that future summaries will be aligned to the general format used in other sections of this report, and be based on fishing method, habitat type, region, or a combination of these.</p> <p>The fisheries summarised are as follows:</p> <table><tr><td><i>Trawl fisheries:</i></td><td><i>Longline fisheries:</i></td><td><i>Other fisheries</i></td></tr><tr><td>Arrow squid</td><td>Ling</td><td>Albacore troll</td></tr><tr><td>Hoki/hake/ling</td><td>Tuna</td><td>Skipjack purse seine</td></tr><tr><td>Jack mackerel</td><td></td><td></td></tr><tr><td>Southern blue whiting</td><td></td><td></td></tr><tr><td>Orange roughy</td><td></td><td></td></tr><tr><td>Oreo</td><td></td><td></td></tr><tr><td>Scampi</td><td></td><td></td></tr></table>	<i>Trawl fisheries:</i>	<i>Longline fisheries:</i>	<i>Other fisheries</i>	Arrow squid	Ling	Albacore troll	Hoki/hake/ling	Tuna	Skipjack purse seine	Jack mackerel			Southern blue whiting			Orange roughy			Oreo			Scampi		
<i>Trawl fisheries:</i>	<i>Longline fisheries:</i>	<i>Other fisheries</i>																							
Arrow squid	Ling	Albacore troll																							
Hoki/hake/ling	Tuna	Skipjack purse seine																							
Jack mackerel																									
Southern blue whiting																									
Orange roughy																									
Oreo																									
Scampi																									
<i>Area</i>	All areas and fisheries																								
<i>Focal localities</i>	<p><i>Arrow squid:</i> Auckland Islands and Stewart/Snares Shelf (80–300 m).</p> <p><i>Hoki/hake/ling:</i> Chatham Rise, West Coast South Island, Campbell Plateau, Puysegur Bank, and Cook Strait (200–800 m).</p> <p><i>Jack mackerel:</i> West Coast of the North and South Islands, Chatham Rise, and Stewart-Snares Shelf (0–300 m).</p> <p><i>Southern blue whiting:</i> Campbell Plateau and Bounty Plateau (250–600 m).</p> <p><i>Orange roughy:</i> The entire New Zealand region (700–1200 m).</p> <p><i>Oreos:</i> South Chatham Rise, Pukaki Rise, Bounty Plateau, and Southland (700–1200 m).</p> <p><i>Scampi:</i> East coasts of the North and South Islands, Chatham Rise, and Auckland Islands (300–450 m).</p> <p><i>Ling longline:</i> Chatham Rise, Bounty Plateau, and Campbell Plateau (150–600 m).</p> <p><i>Tuna longline:</i> Surface waters off the east coast of the North Island and west coast of the South Island.</p> <p><i>Albacore troll fishery:</i> Surface waters off the west coasts of the North and South Islands.</p>																								

	<i>Skipjack purse seine fishery</i> : Northern North Island
<i>Key issues</i>	<p>Under-utilisation (including shark finning) of high volume, low value bycatch species, especially rattails, spiny dogfish, deepsea sharks, blue sharks, porbeagle sharks, and swimming crabs.</p> <p>Potential for considerable reduction of discards by discretionary fishing practices such as the use of mid-water nets, where practicable, and meal plants.</p> <p>Unseen mortality in longline fisheries due to predation by large fish and sharks, marine mammals, seabirds, and sea lice.</p>
<i>Emerging issues</i>	<p>Trends of increasing rates and levels of bycatch and discarding in several categories of catch, especially non-QMS fish species and invertebrates.</p> <p>The effect on bycatch rates in the ling longline fishery of a change to heavier fishing gear (including integrated weights) as used in the Antarctic toothfish fishery.</p> <p>Increasing trawl lengths in the squid, scampi, and orange roughy fisheries due to changes in fishing gear or reduction of target species catch rates—leading to greater bycatch levels in some categories.</p>
<i>MPI Research (current)</i>	<p>DAE201002 (bycatch and discards in deepwater fisheries)</p> <p>DEE201004 (ecological risk assessment in deepwater fisheries)</p> <p>DEE201005A (environmental indicators in deepwater fisheries)</p> <p>HMS200901 (bycatch in tuna longline fisheries)</p>
<i>Other Govt Research (current)</i>	None
<i>Links to 2030 objectives</i>	Objective 6: Manage impacts of fishing and aquaculture.
<i>Related chapters/issues</i>	NPOA sharks

## 6.1. Context

Management of non-protected species bycatch aligns with Fisheries 2030 Objective 6: *Manage impacts of fishing and aquaculture*.

The management of non-protected species bycatch in the deepwater and middle-depth fisheries is described in the National Fisheries Plan for Deepwater and Middle-depth Fisheries (the National Deepwater Plan). Under the National Deepwater Plan, the objective most relevant for management of non-protected species bycatch is Management Objective 2.4: *Identify and avoid or minimise adverse effects of deepwater and middle-depth fisheries on incidental bycatch species*. Specific objectives for the management of non-protected species bycatch will be outlined in the fishery-specific chapters of the National Deepwater Plan. Estimation of non-protected species bycatch is carried out for each of the Tier-1 deepwater fisheries on an annual rotational basis, with each of the following fisheries updated about every 4–5 years:

- Arrow squid
- ling bottom longline
- hoki/hake/ling trawl
- Jack mackerel trawl
- southern blue whiting trawl
- orange roughy/oreo trawl
- scampi trawl

Non-protected fish species bycatch in the Highly Migratory Species (HMS) is addressed in the HMS fish plan. Tuna fisheries incidental bycatch has been regularly examined, with updates every 2–3 years. Some data on bycatch in the Albacore troll fishery and the skipjack tuna purse seine fishery are also available.

The three National Fisheries Plans for Inshore species (finfish, shellfish and freshwater fisheries) also include objectives which address non-protected species bycatch, but research on these objectives has yet to be conducted. However, summaries of the main bycatch species are occasionally included in reports from fisheries characterisation projects, for example school shark, red gurnard, and elephantfish (Starr In Prep; Starr *et al.* 2010a, b, c, Starr & Kendrick 2012).

## **6.2. Global understanding**

Bycatch of unwanted, low value species and discarding of these and of target species that are damaged or too small to process are significant issues in many fisheries worldwide. Few, if any, fisheries are completely without bycatch and this issue has been the subject of innumerable studies and international meetings. Saila (1983) made the first comprehensive global assessment and estimated, albeit with very poor information, that at least 6.7 million tonnes was discarded each year. Alverson *et al.* (1994) extended that work and estimated the global bycatch at 27.0 (range 17.9–39.5) million tonnes each year. An update by Kelleher (2005) suggested global bycatch of about 8% of the global catch, or 7.3 million tonnes, in 1999–2001.

Tropical shrimp trawl fisheries typically have the highest levels of unwanted bycatch, with an average discard rate of 62% (Kelleher 2005), accounting for about one-quarter to one-third of global bycatch. Discard rates in demersal trawl fisheries targeting finfish are typically much lower but, because they are so widespread, their contribution to global discards is considerable. Tuna longline fisheries have the next largest contribution and tend to have greater unwanted bycatch than other line fisheries (Kelleher 2005).

The estimated global level of discards has reduced considerably since the first estimates were made, but differences in the methodology and definition of bycatch used (Kelleher 2005, Davies *et al.* 2009) make it difficult to quantify the decline. The main reasons for the decline in bycatch are thought to have been a combination of higher retention rates, better fisheries management, and improved fishing methods.

Bycatch and discard estimation is frequently very coarse, and estimates of rates based on occasional surveys are often scaled up to represent entire fisheries and applied across years, or even to other fisheries (e.g., Bellido *et al.* 2011). Data from dedicated fisheries observers are also frequently used for individual fisheries, and these are considered to provide the most accurate results, providing that discarding is not illegal (leading to bias due to “observer effects”, Fernandes 2011). Ratio estimators similar to those applied in New Zealand fisheries are frequently used to raise observed bycatch and discard rates to the wider fishery, and the methods used in New Zealand fisheries are broadly similar to those used elsewhere (e.g., Fernandes 2011, Borges *et al.* 2005).

Discard data are increasingly incorporated into fisheries stock assessments and management decision-making, especially with the move towards an Ecosystem Approach to Fisheries (EAF) (Bellido *et al.* 2011), and as third party fishery certification schemes examine more closely the effects of fishing on the ecosystem. They can also be used to assess impacts on non-target species (e.g., Pope *et al.* 2000, Casini *et al.* 2003, Piet *et al.* 2009).

### 6.3. State of knowledge in New Zealand

Estimation of annual bycatch and discard levels of non-protected species in selected New Zealand fisheries have been undertaken at regular intervals since 1998 (Table 6.1).

**Table 6.1: Summary of research into bycatch in New Zealand fisheries**

Fishery	Report
Arrow squid	Anderson <i>et al.</i> (2000) Anderson (2004b) Ballara and Anderson (2009) Anderson (In Press)
Ling bottom longline	Anderson <i>et al.</i> (2000) Anderson (2008)
Hoki trawl	Clark <i>et al.</i> (2000) Anderson <i>et al.</i> (2001) Anderson and Smith (2005) Ballara <i>et al.</i> (2010)
Hake trawl	Ballara <i>et al.</i> (2010)
Ling trawl	Ballara <i>et al.</i> (2010)
Jack mackerel trawl	Anderson <i>et al.</i> (2000) Anderson (2004b) Anderson (2007)
Southern blue whiting trawl	Clark <i>et al.</i> (2000) Anderson (2004a) Anderson (2009b)
Orange roughy	Clark <i>et al.</i> (2000) Anderson <i>et al.</i> (2001) Anderson (2009a) Anderson (2011)
Oreo trawl	Clark <i>et al.</i> (2000) Anderson (2004a) Anderson (2011)
Scampi trawl	Anderson (2004b) Ballara and Anderson (2009)
Tuna longline	Francis <i>et al.</i> (1999a, 1999b) Ayers <i>et al.</i> (2004) Francis <i>et al.</i> (2004) Griggs <i>et al.</i> (2007) Griggs <i>et al.</i> (2008) Griggs & Baird (In Press)
Albacore troll fishery	Griggs <i>et al.</i> (In Press)
Skipjack purse seine fishery	Griggs (unpublished data)

The estimation process uses rates of bycatch and discards in various categories (in most cases “all QMS species combined”, “all non-QMS species combined”, “all invertebrate species combined”) and fishery strata in the observed fraction of the fishery, and effort statistics from the wider fishery, to calculate annual bycatch and discard levels. This ratio-based approach calculates precision by incorporating a multi-step bootstrap algorithm which takes into account the effect of correlation between trawls in the same observed trip and stratum. Estimates of the annual bycatch of a wide range of individual species were also made in the most recent analysis of the arrow squid fishery (Anderson In Press).

The approach used in these analyses relies heavily on an appropriate level and spread of observer effort being achieved, and this is examined in detail in each report. Although details of bycatch and discards are also recorded directly by vessel skippers for the entire fishery, through catch effort forms, these data are often incomplete as the forms list only the top 5 catch species, discards are not well recorded, and they generally lack the accuracy and precision of observer data. Nevertheless, annual bycatch totals are also derived from these data, but only as secondary estimates.

### **6.3.1. Arrow squid trawl fishery**

Since 1990–91 the level of observer coverage in this fishery has ranged from 6% to 53% of the total annual catch, and has been higher in more recent years due to the management measures imposed for the protection of New Zealand sealions (*Phocarctos hookeri*). This coverage has been spread across the fleet and annually 10–68% of all vessels targeting arrow squid have been observed, with this fraction increasing over time. Observers have covered the full size range of vessels operating in the fishery, although the smallest vessels have been slightly undersampled and the largest oversampled.

The observer effort was mostly focussed on the main arrow squid fisheries around the Auckland Islands and Stewart-Snares Shelf, but the smaller fisheries on the Puysegur Bank and off Banks Peninsula were also covered, although less consistently. Observer coverage was more focussed on the central period of the arrow squid season, February to April, than the fleet was in general – with fishing in January and May slightly undersampled.


Appropriate stratification for the analyses was determined using linear mixed-effect models (LMEs) to identify key factors influencing variability in the observed rates of bycatch and discarding. This approach addresses the significant vessel-to-vessel and trip-to-trip differences in bycatch and discard rates in this fishery by treating the trip variable as a random effect (whereby the trip associated with each record is assumed to be randomly selected from a population of trips) and treating other variables as fixed effects. This process consistently identified the separate fishery areas (Auckland Islands, Stewart-Snares Shelf, Puysegur Bank, Banks Peninsula) as having the greatest influence on bycatch and discard rates (with trawl duration of secondary importance) and so area was used in all cases to stratify the calculation of annual levels.

Since 1990–91, over 470 bycatch species or species groups have been identified by observers in this fishery, most being non-commercial species (including invertebrate species) caught in low numbers. Arrow squid have accounted for about 80% of the total estimated catch recorded by observers. The main bycatch species or species groups were the QMS species barracouta (8.5%), silver warehou (2.5%), spiny dogfish (1.7%), and jack mackerel (1.1%); of these only spiny dogfish were mostly discarded (Figure 6.1).

Of the other invertebrate groups crabs (0.8%), in particular smooth red swimming crabs (*Nectocarcinus bennetti*) (0.5%), were caught in the greatest amounts and these were mostly discarded. Smaller amounts of octopus and squid, sponges, cnidarians, and echinoderms were also often caught and discarded.

When combined into broader taxonomic groups, bony fish (excluding rattails, tuna, flatfish, and eels) contributed the most bycatch (16.5% of the total catch), followed by sharks and dogfish (1.9%), crustaceans (0.8%), and rattails (0.2%). The combined bycatch of all other fish (tuna, rays & skates, chimaeras, flatfish, and eels) accounted for a further 0.5% of the total catch.

More than 75% of the sharks & dogfish, rattails, and eels were discarded, whereas about half the flatfish were retained, as were most of the tuna, rays & skates, chimaeras, and other fish not in any of these groups. The fish species discarded in the greatest amounts were spiny dogfish, redbait, rattails, and silver dory. Of the invertebrates, virtually all the echinoderms, other squid, sponges, cnidarians, and polychaetes were discarded, but crustaceans, octopuses, and other molluscs were often retained.



**Figure 6.1: Percentage of the total catch contributed by the main bycatch species (those representing 0.05% or more of the total catch) in the observed portion of the arrow squid fishery, and the percentage discarded. The “Other” category is the sum of all bycatch species representing less than 0.05% of the total catch.**

Total annual bycatch in the arrow squid fishery ranged from about 4500 t to 25 000 t, with low levels in the early 1990s and after 2007–08, and a peak in the early 2000s (Figure 6.2). The large majority of the bycatch comprised QMS species, with less than 1000 t of non-QMS species and invertebrate species bycatch in most years.

Estimated total annual discards ranged from just over 200 t in 1995–96 to about 5500 in 2001–02 and, like bycatch, peaked in the early 1990s and were at relatively low levels after 2006–07 (Figure 6.3). The majority of discards were QMS species (about 62% over all years), followed by non-QMS species (19%), invertebrate species (11%), and arrow squid (7%).





Figure 6.2: Annual estimates of bycatch in the arrow squid trawl fishery, for QMS species, non-QMS species, invertebrates (INV), and overall for 1990–91 to 2010–11. Also shown (in grey) are estimates of bycatch in each category (excluding INV) calculated for 1999–2000 to 2005–06 (Ballara & Anderson 2009). Error bars indicate 95% confidence intervals. The red lines show the fit of a locally-weighted polynomial regression to annual bycatch. In the bottom panel the solid black line shows the total annual reported trawl-caught landings of arrow squid (Ministry of Fisheries 2011), with circles indicating years in which the fishery closed early after reaching the sea lion FRML; and the dashed line shows annual effort (scaled to have mean equal to that of total bycatch).



**Figure 6.3:** Annual estimates of discards in the arrow squid trawl fishery, for arrow squid (SQU), QMS species, non-QMS species, invertebrates (INV), and overall for 1990–91 to 2010–11. Also shown (in grey) are estimates of discards in each category (excluding INV) calculated for 1999–2000 to 2005–06 (Ballara & Anderson 2009). Error bars indicate 95% confidence intervals. The red lines show the fit of a locally-weighted polynomial regression to annual discards.

### 6.3.2. Ling longline fishery

The first analysis of bycatch and discards in this fishery covered the period from 1990–91 to 1997–98, and the second (and latest) analysis covered the following years up to 2005–06. To enable a comparison of estimates between studies, which used slightly different methodologies, the 1994–95 fishing year was re-assessed in the recent analysis. In addition to estimating the bycatch of all quota species combined, and all non-quota species combined, in the recent analysis annual bycatch was estimated separately for three commonly caught individual species, spiny dogfish, red cod, and ribaldo. Comparative estimates of only total annual bycatch are available from the first analysis for 1990–91 to 1997–98.

The ratio estimator used in these analyses to calculate bycatch and discard rates was based on the number of hooks set. The ratios were applied to hook number totals calculated from commercial catch-effort data to make annual estimates for the target fishery as a whole.

Regression tree methods were used to minimise the number of levels of season and area variables used to stratify data for the calculation of annual discard bycatch totals in all categories with minimal loss of explanatory power. This reduced the number of areas in each category from eight down to between two and four, and split the year into three or four periods. The area variables created in this way tended to have more explanatory power. .

Between 1998–99 and 2005–06 only 9% of the vessels operating in this fishery were observed (14 vessels in all) but these tended to be the main operators (including most of the larger autoliners) and accounted for between 7.7% and 52.5% of the annual target ling catch and 7.8% to 61% of the annual number of longlines set during these years. The annual number of observed sets ranged from 324 to 1605 compared with the total target fishery effort of about 2500 to 4150 sets. Observer coverage before 1998–99 was very low, exceeding 5% of the annual target ling catch only in 1994–95 and 1996–97.

Ling accounted for 68% of the total estimated catch from all observed sets targeting ling between 1998–99 and 2005–06, and spiny dogfish accounted for about a further 14%. About half of the remaining 18% of the catch comprised other commercial species; especially red cod (*Pseudophycis bachus*), (2.3%), ribaldo (*Mora moro*) (2.2%), rough skates (*Zearaja nasuta*, 1.9%), smooth skates (*Dipturus innominatus*) (1.8%), and sea perch (*Helicolenus* spp.) (1.2%). Altogether, 93% of the observed catch was comprised of QMS species, representing 40 of the 96 species in the QMS prior to 1 October 2007. Over 130 species or species groups were identified by observers, the majority being non-commercial species caught in low numbers, especially black cod (*Paranotothenia magellanicus*) and Chondrichthyans, often unspecified but including shovelnose spiny dogfish (*Deania calcea*), *Etmopterus* species, and seal sharks (*Dalatias licha*). A surprising number of echinoderms, especially starfish (of which almost 200 000 were observed caught during the period), anemones, crustaceans, and other invertebrates were also recorded by observers.

Total annual bycatch estimates for 1998–99 to 2005–06 ranged from about 2200 t to 3700 t, compared with approximate target species catches in the same period of between about 3500 and 8700 t. A large part of this bycatch (40–50%) comprised a single species, spiny dogfish, and 80% of the bycatch were quota species (Figures 6.4 & 6.5). Bycatch levels decreased during the period, in line with decreasing effort in the fishery. Total bycatch estimates for the years before 1998–99 ranged from about 880 t to 3900 t. Differences in methodology between the two studies, coupled with generally low observer coverage, resulted in significantly different estimates of total bycatch for 1994–95.





Figure 6.4: Annual estimates of fish bycatch in the target ling longline fishery, calculated for commercial (QMS) species (COM), non-commercial (non-QMS) species (OTH), and overall (TOT) for the years 1994-95 and 1998-99 to 2005-06 (in black). Also shown (in grey) are estimates of total bycatch calculated for the period 1990-91 to 1997-98 by Anderson *et al.* (2000). Error bars show the 95% confidence intervals.



**Figure 6.5: Annual estimates of the bycatch of spiny dogfish (SPD), red cod (RCO), and ribaldo (RIB) in the target ling longline fishery for the years 1994–95 and 1998–99 to 2005–06. Error bars show the 95% confidence intervals.**

Total annual discard estimates for 1998–99 to 2005–06 ranged from about 1400 t to 2400 t, and generally decreased during the period (Figure 6.6). About 70–75% of these discarded fish were quota species, and 60–70% spiny dogfish, the remainder being non-quota, generally non-commercial, species. Ling were discarded in small amounts (40–90 t per year), these discards generally being attributable to fish being lost on retrieval or predated by marine mammals and birds. Estimated annual discards were generally lower for the earlier period (1990–91 to 1997–98) and between about 350 t and 1600 t. Total discard estimates for 1994–95 were similar for the two studies.




Figure 6.6: Annual estimates of fish discards in the target ling longline fishery, calculated for ling (LIN), commercial (QMS) species (COM), non-commercial (non-QMS) species (OTH), and overall (TOT) for the years 1994–95 and 1998–99 to 2005–06 (in black). Also shown (in grey) are estimates of the ling and total discards calculated for 1990–91 to 1997–98 by Anderson *et al.* (2000). Error bars show the 95% confidence intervals.

### 6.3.3. Hoki/hake/ling trawl fishery


Earlier reports were limited to the hoki target fishery and only the most recent report considers bycatch and discards for the fishery as defined by the three target species combined—but hoki is dominant in this fishery, accounting for over 90% of the catch.

Observer coverage in the hoki, hake, and ling trawl fishery between 2000–01 and 2006–07 ranged from 11% to 21% of the annual target fishery catch, and 78 separate vessels were observed, covering

the full range of vessel sizes. The annual number of observed tows decreased from 3580 in 2000–01 to 1999 in 2006–07. Coverage has been spread over the geographical range of this fishery, with high sampling throughout the west coast South Island (WCSI) and Chatham Rise fishing grounds and, less frequently, in the Sub-Antarctic. Lower levels of sampling have been achieved in the Cook Strait and Puysegur fisheries, and coverage was lower still around the North Island although this area accounts for very little of the overall catch. Good observer coverage was achieved during the hoki spawning season (July to early September), but coverage outside of this period was variable and under-representative in some months in some years, especially in the Sub-Antarctic, Chatham Rise and Puysegur fisheries.

Hoki, hake, and ling accounted for 87% (77%, 6%, and 4% respectively) of the total observed catch from trawls targeting hoki, hake, and ling between 2000–01 and 2006–07. The remaining 13% comprised a large range of species, especially javelinfish (2.1%), silver warehou (1.7%), rattails (1.4%), and spiny dogfish (1.1%). In total, over 470 species or species groups have been identified by observers, the majority of which are non-commercial species caught in low numbers. Chondrichthyans in general, often unspecified but including spiny dogfish and basking shark, have accounted for much of the non-commercial catch. Echinoderms, squids, crustaceans, and other unidentified invertebrates were also well represented in the bycatch of this fishery.

Total bycatch in the hoki, hake, and ling fishery between 2000–01 and 2006–07 ranged from about 36 000 to 58 000 t per year (compared to the combined total landed catch of hoki, hake, and ling of 130 000 to 238 000 t). Estimates of total bycatch for 1990–91 to 1998–99 from earlier projects (for the hoki target fishery alone), ranged from about 15 000 t to 60 000 t (Figure 6.7). Overall, total bycatch increased during the 1990s to a peak in the early 2000s, and has since declined slowly. Annual bycatch for the 1990–01 to 2006–07 period was also estimated for commercial species (QMS species and species which were generally retained (>75%) and comprised 0.1% or more of the total observed catch) and non-commercial species, rather than QMS and non-QMS species. Roughly similar amounts of these two categories were caught overall, and each showed a similar pattern over time to total bycatch.



**Figure 6.7: Annual estimates of fish bycatch in the target hoki, hake and ling trawl fishery, calculated for commercial species, non-commercial species, and overall for 2000–01 to 2006–07 (black). Also shown (in light grey) are the equivalent bycatch estimates calculated for 1990–91 to 1998–99 by Anderson *et al.* (2001), and for the years 1990–91, 1994–95, 1998–99 and 1999–2000 to 2002–03 by Anderson and Smith (2004), (in dark grey). Error bars show the 95% confidence intervals.**

Total annual discard estimates for 2000–01 to 2006–07 ranged from about 5500 to 29 000 t per year with the main species being discarded including spiny dogfish, rattails, javelinfish, hoki, and shovelnose dogfish. Total annual discards for 1990–91 to 1998–99 were between 6600 t and 17 900 t, and overall there has been no obvious trend in total discards (Figure 6.8). The target species (hoki, hake, and ling) made up 9.7% of total observed discards. Discard rates were strongly influenced by the use of meal plants on fishing vessels; discards of non-commercial species on factory vessels without meal plants was up to twice the level of discards for vessels with meal plants. The use of meal plants, especially for species such as javelinfish and other rattails, has become more prevalent in recent years.





Figure 6.8: Annual estimates of fish discards in the target hoki, hake, and ling trawl fishery, calculated for commercial species, non-commercial species, hoki, and overall for the period 2000–01 to 2006–07 (black). Also shown (in light grey) are the equivalent discard estimates calculated for the period 1990–91 to 1998–99 by Anderson *et al.* (2001), and for 1990–91, 1994–95, 1998–99 and 1999–2000 to 2002–03 by Anderson and Smith (2004), (in dark grey). Error bars show the 95% confidence intervals.

#### **6.3.4. Jack mackerel trawl fishery**

Estimates of annual bycatch in this fishery are available for 1990–91 to 2004–05, with this fishery due to for reassessment in 2013. The annual level of observer coverage in this fishery has varied between 8% and 27% of the target fishery catch but was usually between 15% and 20%. For the most recent period examined, 2001–02 to 2004–05, the majority of the observer effort has focussed on the main fishery, off the west coasts of the North and South Islands, with some additional coverage on the Stewart/Snares Shelf and Chatham Rise fisheries. However, in 2003–04 and 2004–05, there was a total of only 12 trawls observed outside of the western fishery. During this time the fishery was dominated by seven large trawlers and observers were able to complete a trip on each vessel in most years. The fishery runs year round, and although there were significant periods in each year when commercial fishing effort was not observed, coverage encompassed all seasons for the four years combined.

Jack mackerel species accounted for 70% of the total estimated catch from all trawls targeting jack mackerel between 2001–02 and 2004–05. The remaining 30% mostly comprised other commercial species; especially barracouta (15.6%), blue mackerel (4.8%), frostfish (3.1%), and redbait (2.7%). Overall about 130 species or species groups were identified by observers, and about half of these were non-commercial, non-QMS species caught in low numbers. The species most discarded was the spiny dogfish, which comprised about 0.5% of the total catch. The bycatch of non-QMS invertebrate species has yet to be closely studied in this fishery, but species of squid, salps, jellyfish were the most commonly recorded by observers during this period.

Total bycatch in the jack mackerel trawl fishery between 2001–02 and 2004–05 ranged from about 7700 t to 11 900 t. Estimates of total bycatch for 1990–91 to 2003–04 from earlier projects ranged from about 5400 t to 15 500 t (Figure 6.9). After an abrupt increase in the late 1990s, annual bycatch steadily decreased to a level comparable to that of the 1990–91 to 1996–97 period. This bycatch almost entirely comprised commercial (mainly QMS) species.



**Figure 6.9: Annual estimates of fish bycatch in the target jack mackerel trawl fishery for the 2001-02 to 2004-05 fishing years (in black), calculated for commercial species (COM), non-commercial species (OTH), and overall (TOT). Also shown (in grey) are estimates of overall bycatch calculated for 1990-91 to 2000-01 by Anderson *et al.* (2000) and Anderson (2004a). Error bars show the 95% confidence intervals.**

Total annual discards decreased between 2001-02 and 2004-05, continuing a trend that began in 1998-99, to a level of only 90-100 t per year. This is about 5% of the level of 1997-98 (1850 t), when annual discards were at their greatest, and is lower than in any year since 1990-91 (Figure 4.10). Discards of the target species were about 200-400 t per year prior to 1998-99 but thereafter decreased to only about 10 t per year, mainly due to the absence of recorded losses of large quantities of fish through rips in the net or intentional releases of fish during landing. Discards comprised a roughly equal amount of commercial and non-commercial species in the recent study, although commercial species discards were substantially greater in 2001-02.




Figure 6.10: Annual estimates of fish discards in the target jack mackerel trawl fishery for the 2001-02 to 2004-05 fishing years (in black), calculated for jack mackerel (JMA), commercial species (COM), non-commercial species (OTH), and overall (TOT). Also shown (in grey) are estimates of jack mackerel and overall discards calculated for 1990-91 to 2000-01 by Anderson *et al.* (2000) and Anderson (2004a). Error bars show the 95% confidence intervals.

### 6.3.5. Southern blue whiting trawl fishery


In the most recent study, covering the period 2002-03 to 2006-07, the ratio estimator used to calculate bycatch and discard rates in this fishery was based on trawl duration. Linear mixed-effect models (LMEs) identified fishing depth as the key variable influencing bycatch rates and discard rates in this fishery, and regression tree methods were used to optimise the number of levels of this variable in order to stratify the calculation of annual bycatch and discard totals in each catch category.

The key categories of catch/discards examined were; southern blue whiting, other QMS species combined, commercial species combined (as defined above for hoki/hake/ling), non-commercial species combined, and three commonly caught individual species, hake, hoki, and ling.

The level of observer coverage represented between about 22% and 53% of the target fishery catch between 2002–03 and 2006–07 and similar levels were reported from earlier reports, for 1990–91 to 2001–02. The spread of observer data, across a range of variables, has shown no significant shortcomings, due to a combination of the highly restricted distribution of the southern blue whiting fishery over space and time of year, a stable and uniform fleet composition, and a high level of observer effort.

Southern blue whiting accounted for more than 99% of the total estimated catch from all observed trawls targeting southern blue whiting between 2002–03 and 2006–07. About half the remaining total catch was made up of ling (0.2%), hake (0.1%), and hoki (0.1%). These three species, along with other QMS species, comprised over 80% of the total bycatch. In all, over 120 species or species groups were identified by observers, most being non-commercial species caught in low numbers. Porbeagle sharks (introduced into the QMS in 2004), javelinfish and other rattails, and silverside, accounted for much of remaining bycatch. Invertebrate species (mainly sponges, crabs, and echinoderms) were also recorded by observers, but no taxon accounted for more than 0.01% of the total observed catch.

Total annual bycatch estimates for 2002–03 and 2006–07 ranged from about 40 t to 390 t, compared with approximate target species catches in the same period of about 22 000 to 42 000 t. This bycatch was fairly evenly split between commercial species (55%) and non-commercial species (45%), although QMS species accounted for about 80% of the total bycatch during this period. Total annual bycatch decreased during the period, to an all-time low of 40 t in 2006–07. Total annual bycatch estimates for 1990–91 to 2001–02, from earlier reports, were mostly between about 60 t and 500 t but reached nearly 1500 t in 1991–92 (Figure 6.11). This year immediately preceded the introduction of southern blue whiting into the QMS, and effort and catch were exceptionally high.



**Figure 6.11: Annual estimates of fish bycatch in the southern blue whiting trawl fishery, calculated for QMS species, non-commercial species (OTH), and overall (TOT) for 2002–03 to 2006–07 (in black). Also shown (in grey) are estimates of bycatch in each category (excluding QMS) for 1990–91 to 2001–02 (Anderson 2004a). Error bars show the 95% confidence intervals. Note: the 98–00 fishing year encompasses the 18 months between September 1998 and March 2000, the transitional period between a change from an Oct–Sep to Apr–Mar fishing year. The dark line in the bottom panel shows the total annual estimated landings of SBW (Ministry of Fisheries 2009).**

Total annual discard estimates between 2002–03 and 2006–07 ranged from about 90 t to 250 t per year. Discard amounts sometimes exceeded bycatch due to the large contribution of the target species (50–230 t per year) to total discards – the result usually of fish losses during recovery of the trawl. Discarding of commercial species was virtually non-existent in most years and discards of non-commercial species amounted to only 10–50 t per year. The main species discarded were southern blue whiting, rattails and porbeagle sharks. Total annual discard estimates for 1990–91 to 2001–02, from earlier reports, were mostly between about 140 t and 750 t but were about 1200 t in 1991–92 (Figure 6.12). Discards of southern blue whiting (and therefore total discards) decreased substantially at the end of the 1990s and have remained at low levels, below 250 t per year, at least up until 2006–0




Figure 6.12: Annual estimates of fish discards in the southern blue whiting trawl fishery, calculated for the target species (SBW), QMS species, non-commercial species (OTH), and overall (TOT) for 2002–03 to 2006–07 (in black). Also shown (in grey) are estimates of discards in each category (excluding QMS) calculated for 1990–91 to 2001–02 by Anderson (2004a). Error bars show the 95% confidence intervals. The dark line shows the total annual estimated landings of SBW (Ministry of Fisheries 2009).

### 6.3.6. Orange roughy trawl fishery

In the most recent study, covering the period 1990–91 to 2008–09, the ratio estimator used to calculate bycatch and discard rates in the orange roughy fishery was based on the number of trawls. Linear mixed-effect models (LMEs) identified trawl duration as the key variable influencing bycatch rates and discard rates in this fishery, and regression tree methods were used to optimise the number of levels of this variable in order to stratify the calculation of annual bycatch and discard totals in each catch category.

The key categories of catch/discards examined were; orange roughy, other QMS species (excluding oreos) combined, commercial species combined (as defined above for hoki/hake/ling), and non-commercial species combined.

The level of observer coverage in this fishery has been relatively high over the entire period of the fishery—more than 10% (in terms of the total fishery catch) in all but one year, and over 50% in some years. Observer coverage was not evenly spread across all parameters of the orange roughy fishery, the most widespread of any New Zealand fishery, with notable undersampling of smaller vessels, the east coast fisheries in QMAs ORH 2A, ORH 2B, and ORH 3A, and some of the earlier years of the period.

For the recent orange roughy fishery (since 2005–06), orange roughy accounted for about 84% of the total observed catch. Much of the remainder of the total catch (about 10%) comprised oreo species: mainly smooth oreo (8%), and black oreo (2.1%). Rattails (various species, 0.8%) and shovelnose spiny dogfish (*Deania calcea*, 0.6%) were the species most adversely affected by this fishery, with over 90% discarded. Other fish species frequently caught and usually discarded included deepwater dogfishes (family Squalidae), especially *Etmopterus* species, the most common of which is likely to have been Baxter's dogfish (*E. baxteri*), slickheads, and morid cods, especially Johnson's cod (*Halargyreus johnsonii*) and ribaldo. In total, over 250 bycatch species or species groups were observed, most being non-commercial species, including invertebrate species, caught in low numbers. Squid (mostly warty squid, *Onykia* spp.) were the largest component of invertebrate catch, followed by various groups of coral, echinoderms (mainly starfish), and crustaceans (mainly king crabs, family Lithodidae).

Total annual bycatch in the orange roughy fishery since 1990–91 ranged from about 2300 t to 27 000 t, and declined over time alongside the decline in catch and effort in this fishery to be less than 4000 t in each of the last four years estimated (Figure 6.13). Bycatch mostly comprised commercial species, with non-commercial species accounting for only 5–10% of the total bycatch in the recent period.

Estimated total annual discards also decreased over time, from about 3400 t in 1990–91 to about 300 t in 2007–08 (Figure 6.14), and since about 2000 were almost entirely non-commercial, non-QMS species. Large discards of orange roughy and other commercial species, more prevalent early in the fishery, were often due to fish lost from torn nets during hauling.




Figure 6.13: Annual estimates of fish bycatch in the orange roughy trawl fishery, calculated for commercial species (COM), non-commercial species (OTH), QMS species, and overall for 1990–91 to 2008–09 (black points). Also shown (grey points) are earlier estimates of bycatch in each category (excluding QMS) calculated for 1990–91 to 2004–05 (Anderson *et al.* 2001, Anderson 2009a). Error bars show the 95% confidence intervals. The black line in the bottom panel shows the total annual estimated landings of orange roughy (O. Anderson & M. Dunn (NIWA), unpublished data).




Figure 6.14: Annual estimates of fish discards in the orange roughy trawl fishery, calculated for the target species (ORH), commercial species (COM), non-commercial species (OTH), QMS species, and overall for 1990–91 to 2008–09 (black points). Also shown (grey points) are estimates of discards in each category (excluding QMS) calculated for 1990–91 to 2004–05 (Anderson *et al.* 2001, Anderson 2009a). Error bars show the 95% confidence intervals. The black line in the bottom panel shows the total annual estimated landings of orange roughy (O. Anderson & M. Dunn (NIWA), unpublished data).

### 6.3.7. Oreo trawl fishery

In the most recent study, covering the period 1990–91 to 2008–09, the ratio estimator used to calculate bycatch and discard rates in the oreo fishery was based on the number of trawls. Linear mixed-effect models (LMEs) identified trawl duration as the key variable influencing bycatch rates and discard rates in this fishery, and regression tree methods were used to optimise the number of levels of this variable in order to stratify the calculation of annual bycatch and discard totals in each catch category.

The key categories of catch/discards examined were; oreos, other QMS species (excluding orange roughy) combined, commercial species combined (as defined above for hoki/hake/ling), and non-commercial species combined.

The oreo fishery is strongly linked to the orange roughy fishery, and only about 15% of the observed trips examined in the study predominantly targeted oreos, and nearly 30% of the observed trawls targeting oreos were from trips which predominantly targeted orange roughy. The coverage of the oreo fishery is therefore partly determined by the operations of the orange roughy fishery.

The annual number of observed trawls in the oreo fishery ranged from 30 in 1991–92 to 1006 in 2006–07 and the number of vessels observed ranged from 2 to 12. The level of coverage remained at a relatively consistent level after the mid-1990s, despite a decrease in the total catch and effort. Observer coverage was mostly restricted to the main fisheries on the South Chatham Rise and further south. Within this region, few locations were not covered by observers during the 19 years examined, but in the smaller fisheries, on the North Chatham Rise, Louisville Ridge, and the east coast from Kaikoura to East Cape, coverage was minimal. The match of observer coverage to commercial effort was relatively good, especially compared with the orange roughy fishery. Some oversampling on the south Chatham Rise occurred in some periods, e.g., 2001–2005 and 2008–09, and undersampling in the Pukaki/Bounty fisheries in 2005–06 and 2008–09, but elsewhere, and at other times, the spread of coverage was nearly ideal. The full range of vessel sizes (mainly between 300 t and 3000 t) was covered by observers, although small vessels were somewhat underrepresented and large vessels overrepresented. The fleet has shrunk in recent years and the remaining vessels are observed more regularly, with 30–60% of the fleet hosting observers annually since 2002–03.

Oreo species accounted for about 92% of the total estimated catch from all observed trawls targeting oreos after 1 October 2002. Orange roughy (3.5%) was the main bycatch species, with no other species or group of species accounting for more than 0.6% of the total catch. Hoki were the next most common bycatch species, followed by rattails, deepwater dogfish (especially Baxter's dogfish and seal shark (*Dalatias licha*)), slickheads, and basketwork eel (*Diastobranchus capensis*), all of which were usually discarded. Ling were also frequently caught, but only comprised about 0.25% of the total catch. In total, over 250 species or species groups were identified by observers in the target fishery, including numerous invertebrates. As in the orange roughy fishery, corals, squids and octopuses, king crabs, and echinoderms were the main groups caught. Coral, in particular, was a substantial part of the bycatch, accounting for almost 0.4% of the total catch.

Total annual bycatch in the oreo fishery since 1990–91 has ranged from about 270 t to 2200 t and, apart from some higher levels in the late 1990s, not shown any obvious trends (Figure 6.15). Bycatch has been split almost evenly between commercial and non-commercial species overall, although after 2002 about 60% of the bycatch comprised commercial species.

Discards in the oreo fishery remained relatively stable over time, ranging from about 260 t to 750 t per year, with higher levels in the late 1990s than in the early 1990s or 2000s (Figure 6.16). Discards mainly comprised non-commercial, non-QMS species, but also included a significant component of the target species in most years.




Figure 6.15: Annual estimates of fish bycatch in the oreo trawl fishery, calculated for commercial species (COM), non-commercial species (OTH), QMS species, and overall for 1990-91 to 2008-09 (black points). Also shown (grey points) are estimates of bycatch in each category (excluding QMS) calculated for 1990-91 to 2001-02 (Anderson 2004a). Error bars show the 95% confidence intervals. The black line in the bottom panel shows the total annual estimated landings of oreos (Ministry of Fisheries 2010).




Figure 6.16: Annual estimates of fish discards in the oreo trawl fishery, calculated for the target species (OEO), commercial species (COM), non-commercial species (OTH), QMS species, and overall for 1990–91 to 2008–09 (black points). Also shown (grey points) are estimates of discards in each category (excluding QMS) calculated for 1990–91 to 2001–02 (Anderson 2004a). Error bars show the 95% confidence intervals. The black line in the bottom panel shows the total annual estimated landings of oreos (Ministry of Fisheries 2010).

### 6.3.8. Scampi trawl fishery


In the most recent study, covering the period 1990–91 to 2009–10, the ratio estimator used to calculate bycatch and discard rates in the scampi fishery was based on the number of trawls. Linear mixed-effect models (LMEs) identified fishery area as the key variable influencing bycatch rates and discard rates.

The key categories of catch/discards examined were; all QMS species combined, all non-QMS species combined, all invertebrate species combined, javelinfish, and all other rattail species combined.

Observer coverage in the scampi fishery has been relatively low compared with most of the other fisheries assessed. The long-term level of observer coverage in the orange roughy, oreo, arrow squid, southern blue whiting, and ling longline fisheries is greater than 18% of the target fishery catch (and over 40% for southern blue whiting) whereas in the scampi fishery (and also in the jack mackerel fishery) long-term coverage has only been about 11–12%. However, annual coverage in the scampi fishery was greater than 10% in most years and fell below 5% only once (in 2000–01).

The annual number of observed trawls in the fishery ranged from 142 to 797, but has been over 300 trawls in most years. The number of vessels observed in each year ranged from 3 to 8 (equivalent to 33–66% of the fleet) and was very constant—5 or 6 vessels in most years. Analysis of the spread of observer effort compared with that of the scampi fishery as a whole, across a range of variables, indicated that this coverage was reasonably well spread. Although some less important regions of the fishery received little or no coverage (e.g. the central Chatham Rise, where commercial scampi fishing has only recently developed, and west coast South Island), the main scampi fisheries were consistently sampled throughout the period examined. Vessels were mostly of a similar size, and the small amount of effort by larger vessels was adequately covered, as was the full depth range of the fishery and (despite highly intermittent sampling in several years) all periods of the year.

Over 450 bycatch species or species groups were observed in the scampi target fishery catch, most being non-commercial species, including invertebrate species, caught in low numbers. Scampi accounted for only about 17% of the total estimated catch from all observed trawls targeting scampi since 1 October 1990. The main bycatch species or species groups were javelinfish (16%), other (unidentified) rattails (13%), sea perch (*Helicolenus* spp., 8.4%), ling (7.5%), and hoki (6.1%). The first three of these bycatch groups were mostly discarded (Figure 6.17). Of the other invertebrate groups, unidentified crabs (1.1%) and unidentified starfish (0.8%) were caught in the greatest amounts. When combined into broader taxonomic groups, bony fish (excluding rattails) contributed the most to total bycatch (40%), followed by rattails (29%), rays and skates (3.5%), sharks and dogfish (2.3%), crustaceans (2.2%), chimaeras (2.0%), echinoderms (1.6%), and cnidarians (0.6%). A large percentage of the bycatch in these groups was discarded, and was less than 85% only for bony fish (excluding rattails) (33%), rays and skates (67%), and chimaeras (28%).



**Figure 6.17: Percentage of the total catch contributed by the main bycatch species (those representing 1% or more of the total catch) in the observed portion of the scampi fishery, and the percentage discarded. The “Other” category is the sum of all other bycatch species (fish and invertebrates) representing less than 1% of the total catch.**

Total annual bycatch since 1990–91 ranged from about 2100 t to 9200 t and, although highly variable, showed a significant decline over the past 20 years – driven mainly by a decline in the bycatch of QMS species (Figure 6.18). Annual bycatch has generally been an even mixture of QMS and non-QMS species, with invertebrate species (although showing a significant increase over time) accounting for only about 7% of the total bycatch for the whole period. Rattails (split evenly between javelinfish and all other species combined) accounted for 30–80% of the annual non-QMS bycatch. Comparison of bycatch rates with relative biomass estimates from trawl surveys to test for similarity of trends over time was possible for the Chatham Rise and Auckland Islands fishery areas, but these were inconclusive.

Total annual discards ranged from 6790 t in 1995–96 to 1430 t in 2005–06 and, although showing a general decrease since 2001–02, there was no significant trend in overall discard levels since 1990–91 (Figure 6.19). Discards were dominated by non-QMS species (overall about 75%) followed by QMS species (16%) and invertebrates (9%). Rattail species accounted for nearly 60% of the non-QMS discards and about 45% of all discards.






Figure 6.18: Annual estimates of bycatch in the scampi trawl fishery, for QMS species, non-QMS species, invertebrates (INV), and overall for 1990–91 to 2009–10. Also shown (in grey) are estimates of bycatch in each category (excluding INV) calculated for 1999–2000 to 2005–06 (Ballara & Anderson 2009). Error bars indicate 95% confidence intervals. The straight lines show the fit of a weighted regression to annual bycatch. In the bottom panel the solid black line shows the total annual reported landings of scampi (Ministry of Fisheries 2011) and the dashed line shows annual effort (scaled to have mean equal to that of total bycatch).



**Figure 6.19:** Annual estimates of discards in the scampi trawl fishery, for QMS species, non-QMS species, invertebrates (INV), and overall for 1990–91 to 2009–10. Also shown (in grey) are estimates of discards in each category (excluding INV) calculated for 1999–2000 to 2005–06 (Ballara & Anderson 2009). Error bars indicate 95% confidence intervals. The straight lines show the fit of the weighted regression to annual discards.

### 6.3.9. Tuna longline fishery

The New Zealand tuna longline fishery was dominated by the foreign licensed vessels during the 1980s, but is now comprised of chartered Japanese vessels and New Zealand domestic vessels. The domestic fishing fleet has been the dominant fleet in the fishery since 1993–94 (Figure 6.20).



**Figure 6.20: Effort (hooks set) in the tuna longline fishery. Black bars are Foreign and Charter vessels, white bars are NZ domestic vessels.**

The Japanese charter fleet mainly target southern bluefin tuna off the west coast South island (WCSI), and domestic vessels target mainly southern bluefin tuna and bigeye tuna and the fishery is concentrated on the east coast of the North Island (ECNI) with some fishing for southern Bluefin tuna on the WCSI.

The most recent analysis of fish bycatch in tuna longline fisheries was the 2006–07 to 2009–10 fishing years (Griggs & Baird 1012)

Observer effort has mainly focused on the Japanese charter vessels (all vessels covered and usually about 80% of hooks observed), with lower coverage of the domestic fishery (approximately 7-8% during 2006–07 to 2009–10). Most of the fishing effort is carried out by the domestic fleet so this fleet is under-observed.

During 2006–07 to 2009–10, 111 074 fish and invertebrates from at least 62 species or species groups were observed. Most species were rarely observed, with only 37 species (or species groups) exceeding 100 observations between 1988–89 and 2009–10. The most commonly observed species over all years were blue shark, albacore tuna, and Ray's bream, these three making up nearly 70% of the catch by numbers. Blue shark and Ray's bream were the most abundant and second most abundant species in each of the four fishing years 2006–07 to 2009–10 (Table 6.2). Other important non-target species were albacore, lancetfish, bigscale pomfret, dealfish, porbeagle shark, swordfish, moonfish, mako shark, deepwater dogfish, sunfish, and oilfish. The catch composition varied with fleet and area fished.

QMS bycatch species are blue sharks, mako sharks, porbeagle sharks, school shark, moonfish, Ray's bream, and swordfish. Swordfish is also sometimes targeted.

**Table 6.2: Species composition of observed tuna longline catches. Number of fish observed are shown for 2006-07 to 2009-10 and all fish observed since 1988-89. Top 30 species.**

	Species	Scientific Name	2006–07 2009–10	to Total number
1	Blue shark	<i>Prionace glauca</i>	38162	182628
2	Albacore tuna	<i>Thunnus alalunga</i>	9854	101316
3	Rays bream	<i>Brama brama</i>	25277	98205
4	Southern bluefin tuna	<i>Thunnus maccoyii</i>	10373	43291
5	Porbeagle shark	<i>Lamna nasus</i>	2235	19011
6	Dealfish	<i>Trachipterus trachipterus</i>	2304	17185
7	Lancetfish	<i>Alepisaurus ferox</i> & <i>A. brevirostris</i>	5661	14383
8	Moonfish	<i>Lampris guttatus</i>	1683	9134
9	Deepwater dogfish	Squaliformes	1600	9112
10	Swordfish	<i>Xiphias gladius</i>	2213	8286
11	Big scale pomfret	<i>Taractichthys longipinnis</i>	2954	7818
12	Oilfish	<i>Ruvettus pretiosus</i>	711	7542
13	Mako shark	<i>Isurus oxyrinchus</i>	1676	6162
14	Rudderfish	<i>Centrolophus niger</i>	373	4907
15	Butterfly tuna	<i>Gasterochisma melampus</i>	617	4469
16	Escolar	<i>Lepidocybium flavobrunneum</i>	643	4422
17	Bigeye tuna	<i>Thunnus obesus</i>	1240	4390
18	School shark	<i>Galeorhinus galeus</i>	419	3620
19	Yellowfin tuna	<i>Thunnus albacares</i>	97	3342
20	Sunfish	<i>Mola mola</i>	1000	2755
21	Pelagic stingray	<i>Pteroplatytrygon violacea</i>	585	2398
22	Hoki	<i>Macruronus novaezelandiae</i>	265	2021
23	Thresher shark	<i>Alopias vulpinus</i>	169	1400
24	Skipjack tuna	<i>Katsuwonus pelamis</i>	38	1151
25	Dolphinfish	<i>Coryphaena hippurus</i>	134	608
26	Flathead pomfret	<i>Taractes asper</i>	158	516
27	Striped marlin	<i>Tetrapturus audax</i>	59	468
28	Black barracouta	<i>Nesiarchus nasutus</i>	51	386
29	Barracouta	<i>Thyrsites atun</i>	10	357
30	Pacific bluefin tuna	<i>Thunnus orientalis</i>	34	222

Most blue, porbeagle, mako, and school sharks were processed in some way, either being finned or retained for their flesh, but there were significant fleet differences. Blue sharks were mainly just finned. Most albacore, swordfish, yellowfin tuna, moonfish and Ray's bream were retained. Most bigscale pomfret, escolar, oilfish and rudderfish were discarded, with some year and fleet differences. Almost all deepwater dogfish, dealfish, and lancetfish were discarded.

### 6.3.10. Albacore troll fishery

This fishery is comprised entirely of small domestic vessels fishing over the summer months mainly on the west coast of the North and South Island, especially WCSI.

Observers began to go to sea on troll vessels in 2007. The first 2 years were a trial period with one trip observed and targets were set in 2009. Coverage has ranged from 0.5-1.5% of days fished over the 2009–10 to 2010–12 fishing years.

Albacore has made up 93.5% of the observed catch over the past six years, followed by Ray's bream (3.1%) and Skipjack tuna (2.1%) and small numbers (<1%) of a few other species (Table 6.3).

**Table 6.3: Species composition of observed albacore troll catches for 2006–07 to 2011–12.**

Species	Scientific name	Number of fish caught						Total of 6 years
		2006–07	2007–08	2008–09	2009–10	2010–11	2011–12	
Albacore tuna	<i>Thunnus alalunga</i>	1684	1776	1755	5403	4905	2772	18295
Rays bream	<i>Brama brama</i>		18	12	537	35	7	609
Skipjack tuna	<i>Katsuwonus pelamis</i>	1	2	26	20	359	2	410
Barracouta	<i>Thyrsites atun</i>			1		24	13	38
Kahawai	<i>Arripis trutta</i>			6		3	14	23
Kingfish	<i>Seriola lalandi</i>			2	4	4		10
Dolphinfish	<i>Coryphaena hippurus</i>				1			1
Mako shark	<i>Isurus oxyrinchus</i>						1	1
Unidentified		2			174			176

### 6.3.11. Skipjack purse seine fishery

Skipjack tuna makes up 98.9% of the catch observed on purse seine vessels in NZ waters.

Catch composition from eight observed purse seine trips operating within New Zealand fisheries waters in 2010 and 2011 can be seen in Table 6.4.

**Table 6.4: Species composition of observed skipjack purse seine catches in 2010 and 2011.**

Common name	Scientific name	2010–2011 Observed catch weight (kg)	% Catch
Skipjack tuna	<i>Katsuwonus pelamis</i>	3 600 988	98.92
Jack mackerel	<i>Trachurus</i> spp.	22 090	0.61
Jellyfish	Scyphozoa	6 740	0.19
Blue mackerel	<i>Scomber australasicus</i>	4 040	0.11
Manta ray	<i>Mobula japanica</i>	2 122	0.06
Sunfish	<i>Mola mola</i>	1 456	0.04
Striped marlin	<i>Tetrapturus audax</i>	820	0.02
Mako shark	<i>Isurus oxyrinchus</i>	517	0.01
Albacore tuna	<i>Thunnus alalunga</i>	422	0.01
Porcupine fish	<i>Tragulichthys jaculiferus</i>	343	0.01
Flying fish	Exocoetidae	174	<0.01
Frigate tuna	<i>Auxis thazard</i>	100	<0.01
Hammerhead shark	<i>Sphyrna zygaena</i>	80	<0.01
Frostfish	<i>Lepidopus caudatus</i>	79	<0.01
Thresher shark	<i>Alopias vulpinus</i>	75	<0.01
Salps	Thaliacea	57	<0.01
Barracouta	<i>Thyrsites atun</i>	42	<0.01
Moonfish	<i>Lampris guttatus</i>	40	<0.01
Discfish	<i>Dirtemus argenteus</i>	25	<0.01
Electric ray	<i>Torpedo fairchildi</i>	21	<0.01
Slender tuna	<i>Allothunnus fallai</i>	20	<0.01
Blue shark	<i>Prionace glauca</i>	10	<0.01
Garfish	<i>Hyporhamphus ihi</i>	5	<0.01
Pilot fish	<i>Naucrates ductor</i>	5	<0.01
Porbeagle shark	<i>Lamna nasus</i>	5	<0.01
Smooth skate	<i>Dipturus innominatus</i>	5	<0.01
Pilchard	<i>Sardinops neopilchardus</i>	3	<0.01
Starfish	Asteroidea & ophiuroidea	3	<0.01
Dealfish	<i>Trachipterus trachipterus</i>	2	<0.01
	<i>Nototodarus sloanii</i> & <i>n</i>		
Arrow Squid	<i>gouldi</i>	2	<0.01
Dolphinfish	<i>Coryphaena hippurus</i>	1	<0.01
Gurnard	<i>Chelidonichthys kumu</i>	1	<0.01
John dory	<i>Zeus faber</i>	1	<0.01
Decapod	Crustacea	1	<0.01

## 6.4. Indicators and trends

A standard measure that can be used to indicate the degree of wastefulness in a fishery is the level of annual discards as a fraction of the catch of the target species. The most recent estimates of this measure are provided in Table 6.5 for those fisheries where the necessary data were available.

**Table 6.5: Fishery efficiency. Kilograms of discards per kilogram of target species catch. The figures represent the most recent estimate, from published reports.**

Fishery	Discards/target species catch (kg)
Arrow squid trawl	0.02–0.07
Ling longline	0.35
Hoki/hake/ling trawl	0.03
Jack mackerel trawl	0.011
Southern blue whiting trawl	0.005
Orange roughy trawl	0.03–0.06
Oreo trawl	0.02–0.03
Scampi trawl	3.5

Some general trends have been identified in some fisheries, especially those examined within more recent MPI projects where the determination of trends in the rates and levels of bycatch over time has been an explicit objective (Table 6.6).

**Table 6.6: Trends in non-protected species bycatch from recent MPI projects where trend determination has been an objective.**

Fishery	Trends
Arrow squid trawl	<p>Linear regression modelling of observer catch data indicated increasing bycatch rates over time (positive slopes) in all species categories and areas except for QMS species in the Stewart-Snares Shelf and Banks Peninsula fisheries. These trends were statistically significant (<math>p &lt; 0.05</math>) for non-QMS species in the Stewart-Snares Shelf fishery and for invertebrate species in all areas. Bycatch levels for the fishery as a whole also increased over time in each species category, and this increase was significant (<math>p &lt; 0.05</math>) for invertebrates</p> <p>Discard rates increased over time in all species categories and areas except for arrow squid in the Banks Peninsula fishery. These trends were statistically significant (<math>p &lt; 0.05</math>) for QMS species in the Auckland Islands fishery, non-QMS species in the Stewart-Snares Shelf fishery, and for invertebrate species in the Auckland Islands and Banks Peninsula fisheries. Discard levels for the fishery as a whole increased over time in all species categories, and this increase was significant (<math>p &lt; 0.05</math>) for non-QMS species discards and total discards.</p>
Orange roughy trawl	<p>Increased non-commercial species bycatch between the mid-1990s and mid-2000s was shown to strongly correlate with an overall increase in mean trawl length in the fishery resulting from increasing effort away from undersea features.</p>
Scampi trawl	<p>Linear regression modelling of observer catch data indicated significant trends of decreasing bycatch over time for QMS species and total species bycatch and a significant trend of increasing bycatch for invertebrates.</p> <p>A significant trend of increasing discards over time was shown for invertebrates, both rattail categories, and for rattails overall.</p>

	Recent fleet-wide alterations to the nets providing escape gaps for larger unwanted fish species (e.g., skates) may be responsible for the above trends. These escape gaps allow for longer tows, as the nets fill up less rapidly, and may lead to greater catches of benthic invertebrates and smaller fish species.
--	--

## 6.5. References

- Alverson DL; Freeberg MH; Murawski SA; Pope JG (1994). A global assessment of fisheries bycatch and discards. FAO Technical Paper No. 339. Rome. 233 p.
- Anderson, O.F. (2004a). Fish discards and non-target fish catch in the fisheries for southern blue whiting and oreos. New Zealand Fisheries Assessment Report 2004/9. 40 p.
- Anderson, O.F. (2004b). Fish discards and non-target fish catch in the trawl fisheries for arrow squid, jack mackerel, and scampi in New Zealand waters. New Zealand Fisheries Assessment Report 2004/10. 61 p.
- Anderson, O.F. (2007). Fish discards and non-target fish catch in the New Zealand jack mackerel trawl fishery, 2001–02 to 2004–05. New Zealand Aquatic Environment and Biodiversity Report 8. 36 p.
- Anderson, O.F. (2008). Fish and invertebrate bycatch and discards in ling longline fisheries, 1998–2006. New Zealand Aquatic Environment and Biodiversity Report 23. 43 p.
- Anderson, O.F. (2009a). Fish discards and non-target fish catch in the New Zealand orange roughy trawl fishery, 1999–2000 to 2004–05. New Zealand Aquatic Environment and Biodiversity Report 39. 40 p.
- Anderson, O.F. (2009b). Fish and invertebrate bycatch and discards in southern blue whiting fisheries, 2002–07. New Zealand Aquatic Environment and Biodiversity Report 43. 42 p.
- Anderson, O.F.; (2011). Fish and invertebrate bycatch and discards in orange roughy and oreo fisheries from 1990–91 until 2008–09. New Zealand Aquatic Environment and Biodiversity Report 67.
- Anderson, O.F. (In Press). Fish and invertebrate bycatch and discards in New Zealand arrow squid fisheries from 1990–91 until 2010–11. New Zealand Aquatic Environment and Biodiversity Report.
- Anderson, O.F., Clark, M.R., & Gilbert, D.J. (2000). Bycatch and discards in trawl fisheries for jack mackerel and arrow squid, and in the longline fishery for ling, in New Zealand waters. NIWA Technical Report 74. 44 p.
- Anderson, O.F., Gilbert, D.J., & Clark, M.R. (2001). Fish discards and non-target catch in the trawl fisheries for orange roughy and hoki in New Zealand waters for the fishing years 1990–91 to 1998–99. New Zealand Fisheries Assessment Report 2001/16. 57 p.
- Anderson, O.F.; Clark, M.R. (2003). Analysis of bycatch in the fishery for orange roughy, *Hoplostethus atlanticus*, on the South Tasman Rise. Marine and Freshwater Research 54: 643–652.
- Anderson, O.F.; Smith, M.H. (2005). Fish discards and non-target fish catch in the New Zealand hoki trawl fishery, 1999–2000 to 2002–03. New Zealand Fisheries Assessment Report 2005/3. 37 p.
- Ayers, D.; Francis, M.P.; Griggs, L.H.; Baird, S.J. (2004). Fish bycatch in New Zealand tuna longline fisheries, 2000–01 and 2001–02. New Zealand Fisheries Assessment Report 2004/46. 47 p.
- Ballara, S.L.; Anderson, O.F. (2009). Fish discards and non-target fish catch in the trawl fisheries for arrow squid and scampi in New Zealand waters. New Zealand Aquatic Environment and Biodiversity Report 38. 102 p.
- Ballara, S.L.; O'Driscoll, R.L.; Anderson, O.F. (2010). Fish discards and non-target fish catch in the trawl fishery for hoki, hake, and ling in New Zealand waters. New Zealand Aquatic Environment and Biodiversity Report 48.
- Bellido, J.M.; Santos, B. M.; Pennino, G. M.; Valeiras, X.; Pierce, G.J. (2011). Fishery discards and bycatch: solutions for an ecosystem approach to fisheries management? Hydrobiologia, 670. 317–333
- Borges, L.; Zuur, A. F.; Rogana, E.; Officer, R. 2005. Choosing the best sampling unit and auxiliary variable for discards estimations. Fisheries Research, 75: 29–39.
- Casini M; Vitale F; Cardinale M (2003). Trends in biomass and changes in spatial distribution of demersal fish species in Kattegatt and Skagerrak between 1981 and 2003. ICES CM 2003/Q:14
- Clark, M.R., Anderson, O.F., & Gilbert, D.J. (2000). Discards in trawl fisheries for southern blue whiting, orange roughy, hoki, and oreos in New Zealand waters. NIWA Technical Report 71. 73 p.
- Davies, R. W. D.; Cripps, S. J.; Nickson, A; Porter, G. (2009). Defining and estimating global marine fisheries bycatch. Marine Policy 33: 661–672.
- Fernandes, P. G.; Coull, K.; Davis, C.; Clark, P.; Catarino, R.; Bailey, N.; Fryer, R.; Pout, A. (2011). Observations of discards in the Scottish mixed demersal trawl fishery. ICES Journal of Marine Science, 68: 1734–1742.
- Francis, M. P.; Griggs, L. H.; Baird, S. J.; Murray, T. E.; Dean, H. A. (1999a). Fish bycatch in New Zealand tuna longline fisheries. NIWA Technical Report 55. 70 p.
- Francis, M. P.; Griggs, L. H.; Baird, S. J.; Murray, T. E.; Dean, H. A. (1999b). Fish bycatch in New Zealand tuna longline fisheries, 1988–89 to 1997–98. NIWA Technical Report 76. 79 p.
- Francis, M.P.; Griggs, L.H.; Baird, S.J. (2004). Fish bycatch in New Zealand tuna longline fisheries, 1998–99 to 1999–2000. New Zealand Fisheries Assessment Report 2004/22. 62 p.
- Griggs, L.; Doonan, I. McKenzie, A.; Fu, D. (In Press). Monitoring the length structure of commercial landings of albacore (*Thunnus alalunga*) during the 2009–10 to 2011–12 fishing years. New Zealand Fisheries Assessment Report.
- Griggs, L.H.; Baird, S.J.; Francis, M.P. (2007). Fish bycatch in New Zealand tuna longline fisheries, 2002–03 to 2004–05. New Zealand Fisheries Assessment Report 2007/18. 58 p.
- Griggs, L.H.; Baird, S.J.; Francis, M.P. (2008). Fish bycatch in New Zealand tuna longline fisheries in 2005–06. New Zealand Fisheries Assessment Report 2008/27. 47 p.
- Griggs, L.H.; Baird, S.J.; (In Press). Fish bycatch in New Zealand tuna longline fisheries in 2006–07 to 2009–10. New Zealand fisheries Assessment Report.
- Kelleher, K.: (2005). Discards in the world's marine fisheries. An update. FAO Fisheries Technical Paper No. 470. FAO, Rome: 131 pp.
- Ministry of Fisheries (2009). Report from the Fisheries Assessment Plenary, May 2010: stock assessments and yield estimates. Wellington, New Zealand Ministry of Fisheries. 1040p.

- Ministry of Fisheries (2010). Report from the Fisheries Assessment Plenary, May 2010: stock assessments and yield estimates. Wellington, New Zealand Ministry of Fisheries. 1158p.
- Ministry of Fisheries. (2011). Report from the Fisheries Assessment Plenary, May 2011: stock assessments and yield estimates. Ministry of Fisheries, Wellington, New Zealand. 1178p.
- Pope JG; MacDonald DS; Daan N; Reynolds JD; Jennings S (2000). Gauging the impact of fishing mortality on non-target species. ICES Journal of Marine Science 57: 689–696.
- Saila S (1983). Importance and assessment of discards in commercial fisheries. FAO Circular No. 765. Rome. 62 p.
- Starr P.J. (In prep). Stock assessment of east coast South Island elephantfish (ELE 3). New Zealand Fisheries Assessment Report xxxx/xx: 32p.
- Starr P.J., Kendrick T.H. (2012). GUR 3 Fishery Characterisation and CPUE Report. SINS-WG-2012-14v2. 72 pp.
- Starr P.J., Kendrick T.H., Bentley, N. (2010a.) Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation, CPUE analysis and logbook data for SCH 3. Document 2010/07-v2, 62 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3874.aspx>)
- Starr P.J., Kendrick T.H., Bentley, N. (2010b). Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation, CPUE analysis and logbook data for SCH 5. Document 2010/08-v2, 65 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3875.aspx>)
- Starr P.J., Kendrick T.H., Bentley, N. (2010c). Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation, CPUE analysis and logbook data for SCH 7 and SCH 8. Document 2010/09-v2, 149 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3876.aspx>)

## **THEME 3: BENTHIC IMPACTS**

## 7. Benthic (seabed) impacts

<i>Scope of chapter</i>	This chapter outlines the main effects of mobile bottom (or demersal) fishing gear on seabed habitats and communities. All trawl gears contacting the seabed and shellfish dredges are included. Danish seines and more or less static methods like bottom longline and potting are excluded in this first version, as are fisheries outside the EEZ.
<i>Area</i>	All of the New Zealand Territorial Sea (TS) and Exclusive Economic Zone (EEZ). There will be some relevance for out-of-zone bottom trawl fisheries.
<i>Focal localities</i>	Areas that are fished more frequently and habitats that are more sensitive to disturbance are likely to be most affected; areas that are closed to bottom impacting methods will not be directly affected. Bottom trawling in the EEZ is most intense on the western flanks and to the southwest of the Chatham Rise, the edge of the Stewart-Snares Shelf, south of the Auckland Islands, and off the northwest coast of the South Island. Because of the low spatial resolution of reporting up to 2006/07, the spatial distribution of trawling within the TS is less well understood. Shellfish dredges probably have the greatest effect but their footprint is much smaller than that of bottom trawl fisheries and in generally shallow waters.
<i>Key issues</i>	Habitat modification, potential loss of biodiversity, potential loss of benthic productivity, potential modification of important breeding or juvenile fish habitat leading to reduced fish recruitment.
<i>Emerging issues</i>	Potential for effects on habitats of particular significance to fisheries management (HPSFM). The need for (and opportunities presented by) better spatial information on inshore fisheries from finer scale reporting of fishing locations (including logbooks). Cumulative effects and interactions with other stressors (including existing effects, especially in the coastal zone, and climate change).
<i>MFish Research (current)</i>	BEN2007/01, <i>Assessing the effects of fishing on soft sediment habitat, fauna, and processes</i> ; DAE2010/04, <i>Monitoring the trawl footprint for deepwater fisheries</i> ; DAE2010/01, <i>Taxonomic identification of benthic samples</i> ; DEE2010/05, <i>Development of a suite of environmental indicators for deepwater fisheries</i> ; DEE2010/06, <i>Design a programme to monitor trends in deepwater benthic communities</i> ; BEN2012-01, <i>Spatial overlap of mobile bottom fishing methods and coastal benthic habitats</i> .
<i>NZ Government Research (current)</i>	MSI (ex-FRST) programmes: C01X0907, <i>Coastal Conservation Management</i> ; C01X0906, <i>Impacts of resource use on vulnerable deep-sea communities</i> ; C01X0808, <i>Deepsea mining of the Kermadec Ridge</i> . Previous OBI programmes <i>Coasts &amp; Oceans</i> C01X0501 and <i>Marine Biodiversity &amp; Biosecurity</i> C01X0502 are now part of NIWA core funding.
<i>Links to 2030 objectives</i>	Objective 6: Manage impacts of fishing and aquaculture
<i>Related chapter/ issues</i>	Habitats of particular significance for fisheries management (HPSFM), marine environmental monitoring, marine mining/sand extraction, land-based effects.

## 7.1. Context

For the purpose of this document, mobile bottom fishing methods include all types of trawl gear that are used in contact with the seabed, Danish seines, and various designs of shellfish dredges. The information available on the distribution and effects of Danish seining is poor relative to that on trawls and dredges, so that method is not considered here in detail. The benthic effects of other methods of catching fish on or near the seabed that do not involve deliberately towing or dragging fishing gear across the seabed are thought to be considerably less than those of the mobile methods (although not always negligible) and these methods are not considered in this version.

Trawls and dredges are used to catch a relatively high proportion of commercial landings in New Zealand and such methods can represent the only effective and economic way of catching some species. However, the resulting disturbance to seabed habitats and communities may have consequences for biodiversity and ecosystem services, including fisheries and other secondary production. The guiding sections of the Fisheries Act 1996 for managing the effects fishing, including benthic effects, are s.8(2)(b) which specifies that “ensuring sustainability” (s.8(1)) includes “*avoiding, remedying, or mitigating any adverse effects of fishing on the aquatic environment*” and s.9 which specifies a principle that “*biological diversity of the aquatic environment should be maintained*”. Also potentially relevant is the principle in s.9 that “*habitat of particular significance for fisheries management should be protected*” (see the chapter on Habitats of Particular Significance for Fisheries Management for more details).

One approach to managing the effects of mobile bottom fishing methods is through the use of spatial controls. A wide variety of such controls apply in New Zealand waters (Figure 7.1). Some of these controls were introduced specifically to manage the effects of trawling, shellfish dredging, and Danish seining in areas or habitats considered sensitive to such disturbance (e.g., the bryozoans beds off Separation Point, between Golden and Tasman Bays, and the sponge-dominated fauna to the north of Spirits and Tom Bowling Bays in the far north). Other closures exist for other reasons but have the effect of protecting certain areas of seabed from disturbance by mobile bottom fishing methods. These include no-take marine reserves, pipeline and power cable exclusion zones, and areas set aside to protect marine mammals (e.g., see Figure 7.2 for trawl closures introduced in 2008 to protect Hector’s and Maui’s dolphins). Marine reserves provide marine protection in a range of habitats within the Territorial Sea. Although marine reserves provide a higher level of protection by prohibiting all extractive activities, most tend to be small. New Zealand’s 34 marine reserves protect about 7.6% of New Zealand’s Territorial Sea; however, 99% of this is in two marine reserves in the territorial seas around offshore island groups in the far north and far south of New Zealand’s EEZ (Helson *et al.* 2009). Until 2000, most closures that had the effect of protecting areas of seabed from disturbance by trawling and dredging were in the Territorial Sea.

In the Exclusive Economic Zone, 18 seamount closures were established in 2000 to protect representative underwater topographic features from bottom trawling and dredging (Brodie and Clark 2003, see Figure 7.1). These areas include 25 features, including 12 large seamounts >1000 m high, covering 2% (81, 000 km<sup>2</sup>) of the EEZ. The seamount areas are closed to all types of trawling and dredging. In 2006, members of the fishing industry proposed the closure of about 31% of the EEZ to bottom trawling and dredging in Benthic Protection Areas (BPAs), including the existing seamount closures. The design criteria for the BPAs were they should be large, relatively unfished, have simple boundaries, and be broadly representative of the marine environment. After a consultation process, a substantially revised package of BPAs (including three additional areas totalling 13,887 km<sup>2</sup>, 10 additional active hydrothermal vents, and 35 topographic features) that complemented the existing seamount closures was implemented by regulation in 2007 (Helson *et al.* 2009, Figure 7.3). BPAs cover about 1.1 million km<sup>2</sup> (30%) of New Zealand’s EEZ and are closed to trawling on or close to the bottom. Midwater trawling well off the bottom is permitted in the BPAs if two observers are on

board and an approved net monitoring system is used. Much of the seabed within BPAs is below trawlable depth (maximum trawlable depth is about 1600 m) and all are outside the Territorial Sea. In combination, the seamount closures and the BPAs include: 28% of topographic features (a term that includes underwater hills, knolls, and seamounts); 52% of seamounts over 1000 m high; and 88% of known active hydrothermal vents.




Figure 7.1: Map, from Baird and Wood 2010, of the major spatial restrictions to trawling present at some stage during 1989–90 to 2004–05 and the Ministry for Primary Industries Fishery Management Areas (FMAs) within the outer boundary of the New Zealand EEZ. Vessels longer than 28 m may not trawl within the TS and additional restrictions are specified in the Fisheries (Auckland Kermadecs Commercial Fishing) Regulations 1986, the Fisheries (Central Area Commercial Fishing) Regulations 1986, the Fisheries (Challenger Area Commercial Fishing) Regulations 1986 the Fisheries (South East Area Commercial Fishing) Regulations 1986, and the Fisheries (Southland and Sub-Antarctic Areas Commercial Fishing) Regulations 1991.





Figure 7.2: Maps from Ministry of Fisheries website showing the general locations of areas closed to trawling to protect Hector's and Maui's dolphins. Note scales differ. ([http://www.fish.govt.nz/en/nz/Consultations/Archive/2008/Hectors+dolphins/Decisions.htm?wbc\\_purpose=Basic&WBCMODE=PresentationUnpublished%2525252525252cPresentationUnpublished](http://www.fish.govt.nz/en/nz/Consultations/Archive/2008/Hectors+dolphins/Decisions.htm?wbc_purpose=Basic&WBCMODE=PresentationUnpublished%2525252525252cPresentationUnpublished))



**Figure 7.3:** Map from Ministry of Fisheries website showing the general locations of Benthic Protection Areas (BPAs) ([http://www.fish.govt.nz/en-nz/Environmental/Seabed+Protection+and+Research/Benthic+Protection+Areas.htm?wbc\\_purpose=basic&WBCMODE=presentationunpublished&MSHiC=65001&L=10&W=BPA%20&Pre=%3Cspan%20class%3d'SearchHighlight'%3E&Post=%3C/span%3E](http://www.fish.govt.nz/en-nz/Environmental/Seabed+Protection+and+Research/Benthic+Protection+Areas.htm?wbc_purpose=basic&WBCMODE=presentationunpublished&MSHiC=65001&L=10&W=BPA%20&Pre=%3Cspan%20class%3d'SearchHighlight'%3E&Post=%3C/span%3E)). See also Helson *et al.* 2009.

## 7.2. Global understanding

Concerns about the use of towed fishing gear on benthic habitats were first raised by fishermen in the fourteenth century in the UK (Lokkeborg 2005). They were worried about the capture of juvenile fish and the detrimental effects on food sources for harvestable fish. Despite this long history of concern, it is really only in the last 20 years that research efforts have focused strongly on the effects of mobile bottom fishing methods on benthic (seabed) communities, biodiversity, and production. This activity, combined with controversy around fishing effects, has spawned numerous reviews in the past 10 years that seek to summarise or synthesise the information (Jones 1992, Dayton *et al.* 1995; Jennings and Kaiser 1998; Watling and Norse 1998; Lindeboom and deGroot 1998, Auster and Langton 1999; Hall 1999; ICES 2000a and b, Kaiser and de Groot 2000; NMFS 2002, NRC 2002, Dayton *et al.* 2002; Thrush and Dayton 2002; Lokkeborg 2005, Barnes and Thomas 2005, Clark and Koslow 2007).

Benthic habitats provide shelter and refuge for juvenile fish and the associated fauna can be the prey of demersal fish species. Towed fishing gears (particularly trawl doors), affect benthic habitats and organisms and the level of effect will depend on the type of trawl doors and ground gear used, and the physical and biological characteristics of seabed habitats in the fishing grounds. The effects are

difficult to assess because of the complexity of benthic communities and their temporal and spatial variations, and interpretation can be complicated by environmental gradients or change. For reasons of accessibility, cost, and tractability, most research on seabed disturbance caused by human activities worldwide has been carried out in coastal systems, and our understanding of the effects of physical disturbance in the sparse but highly diverse communities of the deep sea has developed only recently. The reviews above broadly indicate that numerical abundance of many invertebrates declines (sometimes substantially) after mining, trawling, or other major disturbance. Trawling and dredging can re-suspend sediment and can, depending on sediment and local currents, alter sediment characteristics. Physical effects include furrows and berms from trawl doors, furrows from the bobbins and rock hoppers, and sediment resorting, but the magnitude of these depends on sediment type, currents, and wave action (if any). Bottom trawling can also alter natural sediment fluxes and reduce the depth of the oxic layer in sediments (Churchill 1989, Warnken *et al.* 2003, Bradshaw *et al.* 2012), and trawling can modify the shape of the upper continental slope (Puig *et al.* 2012), reducing morphological complexity and benthic habitat heterogeneity. The mixing of sediments and overlying water can alter the chemical makeup of the sediment and have considerable effects in deep, stable waters (Rumohr, 1998). Chemical release from the sediment can also be changed, as shown for phosphate in the North Sea (ICES 1992, noting lower fluxes were observed after trawling events). Trawling can alter benthic communities, reduce total biomass of benthic species, and increase predation by scavengers. Sites subject to greater natural disturbance are generally thought less susceptible to change from bottom contact fishing (but see Schratzberger *et al.* 2009 who concluded that common anthropogenic disturbances differ fundamentally from natural disturbance).

There has been less work on the effects of other methods of catching demersal fish or crustaceans that do not involve deliberately towing or dragging fishing gear across the seabed, but some such methods can have non-negligible effects (e.g., Sharp *et al.* 2009, Williams *et al.* 2011). Studies of recovery dynamics are rarer still, but a return to pre-disturbance levels after such changes can take up to several years, even in some sites subject to considerable natural disturbance (see Kaiser *et al.* 2006 for a summary). In shallow regions with mobile sediments, the effects are generally difficult to detect and recovery can be rapid (e.g., Jennings *et al.* 2005). Hard-bottom fauna is predicted to recover most slowly and Williams *et al.* (2010) concluded that hard-bottom fauna on seamounts did not show signs of recovery within 5–10 years on Australasian seamounts. Recovery rate is typically correlated with the spatial extent of a disturbance event (e.g., Hall 1994, Kaiser *et al.* 2003, see also Figure 7.4) and the effects of some “catastrophic” natural disturbance events, such as large-scale marine mudslides, can be detected for hundreds of years, even for taxa thought to be robust to physical disturbance such as nematodes (Hinz *et al.* 2008).




Figure 7.4: General relation between the spatial extent of disturbance events and the time taken to recover from such events in marine systems (after Kaiser *et al.* 2003). Blue dots signal human impacts, including fishing in habitats of different abilities to recover, and black dots signal natural disturbance.

Rice (2006) summarised the findings of five major reviews of the effects of mobile bottom-contacting fishing gears on benthic species, communities, and habitats (available at: [http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006\\_057\\_e.pdf](http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006_057_e.pdf)). In this “review of reviews” Rice (2006) summarised the findings of the multiple working groups that contributed to the reviews as follows:

**Rice’s (2006) conclusions about the effects on habitats of mobile bottom fishing gears were that they:**

- can damage or reduce structural biota (*All reviews, strong evidence or support*).
- can damage or reduce habitat complexity (*All reviews, variable evidence or support*).
- can reduce or remove major habitat features such as boulders (*Some reviews, strong evidence or support*).
- can alter seafloor structure (*Some reviews, conflicting evidence for benefits or harm*).

**Other emergent conclusions on habitat effects included:**

- There is a gradient of effects, with greatest effects on hard, complex bottoms and least effect on sandy bottoms (*All reviews, strong support, with qualifications*).
- There is a gradient of effects, with greatest effects on low energy environments and least (often negligible) effect on high-energy environments (*All reviews, strong support*).
- Trawls and mobile dredges are the most damaging of the gears considered (*Three of the reviews considered other gears; all drew this conclusion, often with qualifications*).

**Mobile bottom gears affect benthic species and communities in that they:**

- can change the relative abundance of species (*All reviews, strong evidence or support*).
- can decrease the abundance of long-lived species with low turnover rates (*All reviews, moderate to strong evidence or support*).
- can increase the abundance of short-lived species with high turnover rates (*All reviews, moderate to occasionally strong evidence or support*).
- affect populations of surface-living species more often and to greater extents than populations of burrowing species (*All reviews, weak to occasionally strong evidence or support*).
- have lesser effects in high-energy or frequent natural disturbance environments than in low energy environments where natural disturbances are uncommon (*Four reviews (the other did not address the factor), strong evidence or support*).
- affect populations of structurally fragile species more often and to greater extents than populations of “robust” species (*All reviews, variable evidence and support*).
- Abundance of scavengers increases temporarily in areas where bottom trawls have been used (*Three reviews, variable support or evidence, all argue for the effects being transient*).
- Rates of nutrient cycling or sedimentation are increased in areas where bottom trawls have been used (*Two reviews, mixed views on magnitude of effects and conditions under which they occur*).

**Considerations in the application or adoption of mitigation measures:**

- The effect of mobile fishing gears on benthic habitats and communities is not uniform. It depends on:
  - the features of the seafloor habitats, including the natural disturbance regime (*All reviews, strong evidence or support*);
  - the species present (*All reviews, strong evidence or support, though not mentioned by NMFS panel*);
  - the type of gear used and methods of deployment (*All reviews, moderate to strong evidence support*);
  - the history of human activities, particularly past fishing, in the area of concern (*All reviews, strong evidence or support*).

- Recovery time from trawl-induced disturbance can take from days to centuries, and depends on the same factors as listed above. (*All reviews, strong evidence or support*).
- Given the above considerations, the effect of mobile bottom gears has a monotonic relationship with fishing effort, and the greatest effects are caused by the first few fishing events (*All reviews, moderate to strong evidence or support*).
- Application of mitigation measures requires case specific analyses and planning; there are no universally appropriate fixes (*Three reviews, moderate to strong evidence or support. The issue of implementing mitigation was not addressed in the FAO review. It was also stressed in the US National Academy of Sciences review and discussed in the ICES review that extensive local data are not necessary for such case-specific planning. The effects of mobile bottom gears on seafloor habitats and communities are consistent enough with well-established ecological theory, and across studies, that cautious extrapolation of information across sites is legitimate*).

Rice (2006) concluded “*These overall conclusions on impacts and mitigation measures, and recommendations for management action form a coherent and consistent whole. They are relevant to the general circumstances likely to be encountered in temperate, sub-boreal, and boreal seas on coastal shelves and slopes, and probably areas ... beyond the continental shelves. They allow use of all relevant information that can be made available on a case by case basis, but also guide approaches to management in areas where there is little site-specific information.*”

Since Rice’s (2006) paper, Kaiser *et al.* (2006) published a meta-analysis of 101 separate manipulative experiments that confirms many of Rice’s findings. Shellfish dredges have the greatest effect of the various mobile bottom fishing gears, biogenic habitats are the most sensitive to such disturbance (especially for attached fauna on hard substrates) and unconsolidated, coarse sediments (e.g., sands) are the least sensitive. Kaiser *et al.* (2006) concluded that recovery from disturbance events can take months to years, depending on the combination of fishing method and benthic habitat type. This meta-analysis of manipulative experiments was an important development, reinforcing the inferences drawn from multiple mensurative observations at much larger scale (“fisheries scale”) in New Zealand (e.g., Thrush *et al.* 1998, Cryer *et al.* 2002) and overseas (e.g., Craeymeersch *et al.* 2000, McConnaughey *et al.* 2000, Bradshaw *et al.* 2002, Blyth *et al.* 2004, Tillin *et al.* 2006, Hiddink *et al.* 2006). This is a powerful combination that implies substantial generality of the findings.

The international literature is, therefore, clear that bottom (demersal) trawling and shellfish dredging are likely to have largely predictable and sometimes substantial effects on benthic community structure and function. However, the positive or negative consequences for ecosystem processes such as production had not been addressed until more recently (e.g., Jennings *et al.* 2001, Reiss *et al.* 2009, Hiddink *et al.* 2011). It has been mooted that frequent disturbance should lead to the dominance of smaller species with faster life histories and that, because smaller species are more productive than larger ones, system productivity and production should increase under trawling disturbance. However, when this proposition has been tested, it has not been supported by data in real fishing situations (e.g., Jennings 2002, Hermesen *et al.* 2003, Reiss *et al.* 2009) and where overall productivity has been assessed, it decreases with increasing trawling disturbance.

For example, Veale *et al.* (2000) examined spatial patterns in the scallop fishing grounds in the Irish Sea and found that total abundance, biomass, and secondary production (including that of most individual taxa examined) decreased significantly with increasing fishing effort. Echinoids, cnidarians, prosobranch molluscs, and crustaceans contributed most to the differences. Jennings *et al.* (2001) showed that, in the North Sea, trawling led to significant decreases in infaunal biomass and production in some areas even though production per unit biomass rose with increased trawling disturbance. The expected increase in relative production did not compensate for the loss of total production that resulted from the depletion of large-bodied species and individuals. Hermesen *et al.* (2003) found that mobile fishing gear disturbance had a conspicuous effect on benthic megafaunal production on Georges Bank, and cessation of such fishing led to a marked increase in benthic

megafaunal production, dominated by scallops and urchins. Hiddink *et al.* (2006) estimated that more than half of the southern North Sea was trawled sufficiently frequently to depress benthic biomass by 10% or more, and that 27% was in a state where benthic production was depressed by 10% or more. They estimated that recovery from this situation would take 2.5–6 years or more once fishing effort had been eliminated. They further estimated that fishing reduced benthic biomass and production by 56% and 21%, respectively, compared with an unfished situation. Reiss *et al.* (2009) found that, although sediment composition was the most important driver of benthic community structure in their North Sea study area, the intensity of fishing effort was also important and reductions in the secondary production of the infaunal community could be detected even within this heavily fished region.

The types of models developed by Hiddink *et al.* (2006, 2011, but see also Ellis and Pantus 2001 and Dichmont *et al.* (2008) can be used to assess the likely performance of different management approaches or levels of fishing intensity. Such management-strategy-evaluation (MSE) methods involve specifying management objectives, performance measures, a suite of alternative management strategies, and evaluating these alternatives using simulation (Sainsbury *et al.* 2000). For instance, the early study by Ellis and Pantus (2001) assessed the effect of trawling on marine benthic communities by combining an implementation of the spatial and temporal behaviour of the local fishing fleet with realistic ranges for the removal and recovery of benthic organisms. The model was used to compare the outcomes of two radically different management approaches, spatial closures and reductions in fishing effort. Lundquist *et al.* (2007, 2010) used a more sophisticated spatially explicit landscape mosaic model with variable connectivity between patches to assess the implications of different spatial and temporal patterns of disturbance in the model landscape. They found that the scale of the disturbance regime (which could be trawling or any other physical disturbance) and the dispersal processes interact, and that the scales of these processes greatly influenced changes in the structure and diversity of the model community, and that recovery across the mosaic depended strongly on dispersal. System stability also decreased as dispersal distance decreased.

### **7.3. State of knowledge in New Zealand**

To understand the effects of mobile bottom fishing methods on benthic habitats, it is necessary to have knowledge on

- the distribution of such habitats,
- the extent to which mobile bottom fishing methods are used in each habitat (the overlap),
- the consequences of any such disturbance (potentially in conjunction with other disturbances or stressors), and
- the nature and speed of recovery from the disturbance.


These components will be dealt with in turn.

#### **7.3.1. Distribution of Habitats**

Mapping of benthic habitats at the large scales inherent in fisheries management is expensive and time-consuming so the New Zealand government commissioned an environmental classification to provide a spatial framework that subdivided the TS and EEZ into areas having similar environmental and biological character. This Marine Environment Classification (MEC) was launched in 2005 (Snelder *et al.* 2004, 2005, 2006) using available physical and chemical predictors, and because environmental pattern was thought a reasonable surrogate for biological pattern. The authors suggested that the MEC provided managers with a useful spatial framework for broad scale management, but cautioned that the full utility and limitations would become clear only as the MEC was applied to real issues. They described the MEC as a tool to organise data, analyses and ideas, and as only one component of the information that would be employed in any analysis. The 20-class version (Figure 7.5, Table 7.1) has been the most widely cited, although additional classification

levels provide more detail that is significantly correlated with biological layers. The 2005 MEC was not optimised for any specific ecosystem component but was “tuned” against data for demersal fish, phytoplankton, and benthic invertebrates. It performed least well as a classification of benthic invertebrates and, at the 20-class level, grouped most of the Chatham Rise and Challenger Plateau into a single class. Although separation of these two areas was evident as the MEC was driven to larger numbers of classes, their inclusion in a single class in the 20-class classification was considered counter-intuitive because their productivity and fisheries are known to be very different.

This disquiet with the predictions of the original MEC for benthic habitat classes led to the development of alternatives that might perform better for benthic systems. First of these was a classification optimised for demersal fish (Leathwick *et al.* 2006). Several variants of this classification all out-performed the original MEC for demersal fish, particularly at lower levels of classification detail and was adopted by the Ministry for the Environment for their indicators related to bottom trawling and their 2010 Environmental Snapshot where the trawl footprint is compared with putative habitats (Ministry for the Environment 2010, see also: <http://www.mfe.govt.nz/environmental-reporting/report-cards/seabed-trawling/2010/index.html>).



**Figure 7.5:** The 20-class version of the 2005 general purpose Marine Environment Classification (MEC, from Snelder *et al.* 2005). The class numbers are nominal; for attributes of each class at this level, see Table 7.1.


Based partly on this experience, the Ministry of Fisheries commissioned a Benthic-Optimised Marine Environment Classification, BOMECE. Many more physical, chemical, and biological data layers were available for the development and tuning of this classification than for the 2005 MEC. Especially relevant for benthic invertebrates was the inclusion of a layer for sediment grain size (notably absent from the MEC). Generalised Dissimilarity Modelling (GDM, Ferrier *et al.* 2002, 2006, Leathwick *et al.* 2011) was used to define the classification because this approach is well suited to the sparse and unevenly distributed biological data available. The BOMECE classes (Figure 7.6) were strongly driven by depth, temperature, and salinity into five major groups: inshore and shelf; upper slope; northern mid-depths; southern mid-depths; and deeper waters (generally beyond the fishing footprint, down to 3000 m, the limit of the analysis). Waters deeper than 3000 m could be considered an additional class.

Recent testing (Bowden *et al.* 2011) has indicated that the BOMECE out-performs the original MEC at predicting benthic habitat classes on and around the Chatham Rise, but that none of the available classifications is very good at predicting the abundance and composition of benthic invertebrates at the fine scale of the sampling undertaken (10s of metres to kilometres). This, in conjunction with the findings of Leathwick *et al.* (2006), reinforces the role of environmental classifications as broad-scale predictors of general patterns at broad scale (tens to hundreds of kilometres) when more specific biological information is not available.

Where broad scale classification methods are not applicable, other approaches have been taken. The trawl fisheries for orange roughy, oreos, and cardinalfish take place to a large extent on seamounts or other features (Clark and O'Driscoll 2003, O'Driscoll and Clark 2005). These features are often geographically small and, in common with other, localised habitats like vents, seeps, and sponge beds, do not appear on broad-scale habitat maps (e.g., at EEZ scale) and cannot realistically be predicted by broad-scale environmental classifications. Many features have been extensively mapped in recent years (e.g., Rowden *et al.* 2008), and seamount classifications based on biologically-referenced physical and environmental "proxies" have also been developed, in New Zealand waters by Rowden *et al.* (2005), and globally by Clark *et al.* (2010a&b), and Davies and Guinotte (2011) developed a method of predicting the framework-forming (i.e, physically structuring) coldwater corals that are a focus for benthic biodiversity in deepwater systems. Work continues worldwide, including in New Zealand, on the development of sampling, analytical, and modelling techniques to provide cost-effective assessments of the distribution of marine habitats at a range of scales. MPI project DEE2010/06 has been commissioned to design a monitoring programme to assess trends in deepwater benthic communities using information from trawl surveys, observers, and directed sampling. MPI project DEE2010/05 has been commissioned to develop a suite of environmental indicators for deepwater fisheries using, to the extent possible, existing data collection processes. This is an area of rapid change in the science and better techniques and data sets for predicting and mapping marine benthic habitats are likely to become available in the short to medium term. MPI project BEN2012/01 will use existing information and classifications to describe the distribution of benthic habitats in the coastal zone. NIWA has a MBIE-funded project "*Predicting the occurrence of vulnerable marine ecosystems for planning spatial management in the South Pacific region*" in collaboration with Victoria University of Wellington and the Marine Conservation Institute (USA). The research will develop a model to predict the locations of VMEs to inform New Zealand and South Pacific Regional Fisheries Management Organisation (SPRFMO) initiatives on spatial management in the South Pacific region. There may be applications within the New Zealand EEZ.

**Table 7.1: Average values for each of the eight defining environmental variables in each class of the 20-class level of the MEC classification. After Snelder *et al.* 2005.**

Class	Area (km <sup>2</sup> )	Depth	Slope	Orbital velocity	Radiation mean	SST amplitude	SST gradient	SST winter	Tidal current	2-class level	4-class level	9-class level
1	88,503	-3001	1.4	0	17.5	2.3	0.01	19.5	0.06	Oceanic	Subtropical	Deep
22	53,368	-1879	1.5	0	15.4	2.4	0.01	16.3	0.11			
9	64,306	-5345	1.4	0	14.8	2.6	0.01	16.1	0.03			Abyssal
47	60,053	-2998	1.0	0	12.1	2.4	0.01	11.6	0.07		Shelf and subtropical front	Central
55	2,213	-334	1.6	0	15.5	2.4	0.02	15.1	0.20			
63	26,626	-754	0.9	0	12.8	2.4	0.02	12.1	0.18			
178	39,360	-750	0.4	0	9.5	1.3	0.01	7.6	0.15		Sub-Antarctic	Southern
127	60,884	-4830	0.5	0	10.7	1.7	0.01	10.0	0.05			
204	18,277	-2044	3.0	0	9.2	0.9	0.01	8.0	0.08			
273	805	-2550	9.1	0	8.4	1.4	0.03	4.4	0.05			
219	93,982	-4779	0.6	0	8.9	1.0	0.01	6.7	0.04			
12	149	-94	0.9	113	17.8	2.3	0.01	19.3	0.30	Coastal	Northern	Northern
58	394	-117	0.7	57	14.7	2.2	0.03	13.0	1.09			Central
60	4,084	-112	0.3	21	14.4	2.5	0.02	13.2	0.26			
64	2,689	-38	0.3	272	14.2	2.9	0.02	12.6	0.19			
124	68	-8	0.4	836	13.4	2.3	0.02	12.7	0.00			
130	14	-10	0.4	353	14.1	2.4	0.09	11.9	0.21			
169	932	-66	0.2	113	12.4	2.7	0.04	9.9	0.21			
190	339	-321	1.9	3	12.3	2.3	0.06	9.4	0.10			
170	5,208	-129	0.3	99	10.2	1.3	0.02	9.3	0.55			Southern



**Figure 7.6: Map of the distribution of Benthic Optimised Marine Environment Classification (BOME) classes defined by multivariate classification of environmental data transformed using results from GDM analyses of relationships between environment and species turnover averaged across eight taxonomic groups of benthic species. From Leathwick *et al.* 2010.**

### 7.3.2. Distribution of Fishing

Since 1989/90, mobile bottom fishing has been reported on one of three standardised reporting forms (Table 7.2). Trawl Catch Effort and Processing Returns (TCEPRs) contain detailed spatial and other information for each trawl tow, whereas Catch Effort and Landing Returns (CELRs) include only summarised information for each day's fishing, with very limited spatial resolution. Since 2007/08, Trawl Catch and Effort Returns (TCERs) have been available for smaller, predominantly inshore trawlers. These include spatial and other information for each trawl tow but in less detail than on TCEPRs. Between 1989/90 and 2004/05, only about 25% of all mobile bottom fishing events were reported on TCEPRs. Another 25% were bottom trawls reported on CELRs, and the remaining 50% were dredge tows for shellfish reported on CELRs. The distribution of trawling reported on CELRs is not the same as that reported on TCEPRs (Figure 7.8); the smaller trawlers using CELRs are much more likely than the larger boats to fish close to the coast and target inshore species such as flatfish, red cod, tarakihi, and red gurnard (collectively 73% of all trawl tows reported on CELRs). MPI project BEN2012/01 will update the work in BEN2006/01 producing maps of swept area and footprint for more recent years.

**Table 7.2: Attributes, usage, and resolution of spatial reporting required on Trawl Catch Effort and Processing Returns (TCEPRs) Trawl Catch and Effort Returns (TCERs) and Catch Effort and Landing Returns (CELRs).**

	Trawl catch and effort reporting forms		
	TCEPR	TCER	CELR
Year of introduction	1988/89	2007/08	1988/89
Vessels using	All trawlers >28 m Other vessels as directed Other vessels optional	All trawlers 6–28 m unless exempted	Trawlers not using TCER or TCEPR Shellfish dredgers
Trawl tow reporting	Tow by tow, start and finish locations, speed, depth, gear	Tow by tow, start location, speed, depth, gear	Daily summary, number of tows, effort, gear
Spatial resolution	1 minute (lat/long)	1 minute (lat/long)	Statistical reporting area (optionally lat/long)

Baird *et al.* (2002) and Baird *et al.* (2001) described the distribution and frequency of reported fishing by mobile bottom fishing gear (dredge, Danish seine, bottom trawl, bottom pair trawl, and mid-water trawl in contact with the bottom) in New Zealand's TS and EEZ during the 1990s and up to 2004/05, respectively. They showed that fishing was highly heterogeneous (spatially), but had considerable consistency among years; sites that were fished heavily in one year were likely to be fished heavily in other years and *vice versa*. A similar but more detailed analysis was conducted for the Chatham Rise and SubAntarctic areas by Baird *et al.* (2006). Tows reported on TCEPRs were included in the main spatial analysis but some additional analysis was possible using tows reported on CELRs. Until 2006/07, many inshore vessels used CELRs and these comprised a substantial proportion of reported trawling, even for some “deepwater” species. For instance, Cryer and Hartill (2002) estimated that, in the Bay of Plenty, 78%, 75%, and 39% of trawl tows for tarakihi, gemfish, and hoki, respectively, were reported on CELR forms in the 1990s. Since 2007/08, almost all trawling effort has been reported on TCEPR or TCER forms.

Baird *et al.* (2011) calculated three annual measures of fishing effort: the number of tows, the aggregate swept area (using assumed door spreads, see Figure 7.7), and the coverage (“footprint”) of the total trawl contact. Trawls were represented spatially as tracklines between the reported start and

finish positions buffered by the assumed door spread to generate trawl polygons. The aggregate swept area for a year is the sum of the areas of the polygons and the “footprint” is the estimated area of the seabed that is covered by the polygons overlaid. The estimated swept areas and footprint do not account for any modification that might occur alongside the trawl path as represented by the swept area polygon (e.g., by suspended sediments transported by currents away from the trawl track). Baird *et al.* (2011) produced maps of the aggregate swept area by year for each of the 22 main target species or species groups, and various tables and figures describing trends. The annual number of trawls peaked in 1997–98 at 78 610 tows (swept area ~ 180 450 km<sup>2</sup>). In 2007/08, fewer than 55 000 tows were reported on TCEPRs (~ 130 800 km<sup>2</sup>)




Figure 7.7: Map from Baird *et al.* 2011 showing the intensity of bottom-contacting trawling effort reported on TCEPR forms 1989–90 to 2004–05. The colour scale indicates the aggregate swept area estimated by Baird *et al.* for each 5 x 5 km cell, all target species combined (e.g., the 36 most intensively fished 25 km<sup>2</sup> cells all had an aggregate swept area of over 2290 km<sup>2</sup> over 16 years, which translates to the seabed in those cells being swept by some part of a trawl 92 times in 16 years, or an average of 5.8 or more times each year). Updates for deepwater trawl fisheries are expected in 2013.

Baird *et al.* (2011) used reported tows on small topographic features that are a focus for orange roughy and cardinalfish fisheries by defining polygons for these tows as radii around the reported start position with the area swept estimated from the reported duration and speed of the tow. These short tows do not appear to contribute substantially to broad-scale plots like Figure 7.7, yet can represent intense fishing effort on particular, small seamount features (e.g. Rowden *et al.* 2005, O'Driscoll and Clark 2005).





Figure 7.8: Broad-scale distribution from Baird *et al.* 2011 of bottom trawl effort reported on CELRs (left) and on CELRs and TCEPRs combined (right), for all fishing years 1989–90 to 2004–05. Updates for deepwater trawl fisheries (but not inshore fisheries reporting on TCER) are expected in 2013.

After the peak of over 140 000 reported trawl tows in 1996/97 and 1997/98 (Figure 7.9) when slightly over half of all tows were reported on TCEPRs, overall trawling effort declined to less than 100 000 tows per year by 2006/07. The reported number of trawl tows has remained relatively stable at about


85–90 000 tows per year, only about 44% of which is reported on TCEPRs (virtually all other tows are reported on TCERs)

Dredging for shellfish (oysters and scallops) is conducted in a number of specific areas that have separate, smaller statistical reporting areas (Figure 7.10). Over the 16-year dataset, there were almost 1.5 million scallop dredge tows in the four main scallop fisheries and over 0.6 million oyster dredge tows in the two dredge oyster fisheries. These data are collected on CELRs, usually at the spatial scale of a scallop or oyster fishery area and the data have been summarised as the number of dredge tows. No estimates of the area swept by these dredges have been made, but the number of reported tows has declined markedly since the early 1990s (Figure 7.11).



**Figure 7.9:** The number of trawl tows reported on Trawl Catch Effort and Processing Returns (TCEPR), Catch Effort and Landing Returns (CELR) and Trawl Catch and Effort Return (TCER) between the 1989/90 and 2007/08 fishing years. Data for the 2011/12 year may be incomplete.

Our knowledge of the distribution of mobile bottom fishing effort within our TS and EEZ is, by international standards, very good; since 2007/08 we have had tow-by-tow reporting of almost all trawling with a spatial precision of about 1 nautical mile. The distribution of dredge tows for shellfish is not reported with such high precision, but records kept by fishers in industry logbooks are often much more detailed than the Ministry for Primary Industries standard returns, and have sometimes been used to support spatial analyses that would not have been possible using the standard returns (e.g., Tuck *et al.* 2006 for project ZBD2005/15 on the Coromandel scallop fishery and Michael *et al.* 2006 for project ZBD2005/04 on the Foveaux Strait oyster fishery). These studies indicate the value of records with higher spatial precision.



**Figure 7.10: Maps taken from Baird *et al.* 2011 of statistical reporting areas for the main oyster and scallop dredge fisheries (scales differ). Note that these reporting areas are generally much smaller than the standard statistical reporting areas used for most finfish reporting.**




Figure 7.11: The number of dredge tows for scallop or oysters reported on Catch Effort and Landing Returns (CELR) between the 1989/90 and 2007/08 fishing years (data from Baird *et al.* 2011 and MPI databases). Data for the 2011/12 year may be incomplete.

### 7.3.3. Overlap of Fishing and Predicted Habitat Classes

Baird and Wood (2010, project BEN200601) overlaid the 16-year trawl footprint up to 2004-05 on the 15-class BOMECS to estimate the proportion of each class that had been trawled (and reported on TCEPRs). They found that the size of the footprint and the proportion of each class trawled varied substantially between habitat classes (Figure 7.12, Table 7.3). Class O is the largest BOMECS class but has almost no reported fishing effort. Conversely, class I is one of the smaller classes but has a larger trawl footprint that overlaps about 70% of the total class area. Two contrasting classes, together with their trawl footprints, are shown in Figure 7.13. The cumulative trawl footprint from Baird and Wood's analysis overlaps about 8% of the 4.1 million km<sup>2</sup> of seafloor within the New Zealand EEZ boundary (i.e., including the Territorial Sea). However, this overlap and that for some individual BOMECS classes (particularly coastal classes A–E) will be underestimated because of the omission of CELR data from these analyses. This analysis is being updated for offshore (middle depth and deepwater) trawl fisheries under project DAE2010/04, *Monitoring the trawl footprint for deepwater fisheries*, and the results are expected to be available in early 2013. MPI project BEN2012/01, *Spatial overlap of mobile bottom fishing methods and coastal benthic habitats*, will update the work in BEN2006/01, particularly focussing on the overlap between fishing and habitats in the coastal zone.




Figure 7.12: Plots from Baird and Wood (2010) of the areas of each BOMECE Class (top), the fishing footprint up to 2004/05 shown in Figure 7.8 (centre), and percentage of each BOMECE Class area covered by the fishing footprint (bottom).

Table 7.3: Estimated area of each BOMECE class (within the outer boundary of the EEZ), the minimum and maximum values for the trawl footprint in each, and cumulative footprint over the 16 years studied by Baird and Wood (2010).

BOMECE class	Area (km <sup>2</sup> )	Min. annual footprint area (km <sup>2</sup> )	Max. annual footprint area (km <sup>2</sup> )	Cumulative (16 yr) proportion overlapped
A*	27 557	121	4 026	0.42
B*	12 420	40	484	0.19
C*	89 710	4 271	11 374	0.58
D*	27 268	377	1 602	0.30
E*	60 990	4 046	7 108	0.40
F	38 608	517	1 391	0.13
G	6 342	132	833	0.34
H	138 550	9 583	20 344	0.45
I	52 224	5 511	18 016	0.70
J	311 361	10 469	15 975	0.18
K	1 290	-	2	0.01
L	198 577	4 238	13 599	0.24
M	233 825	895	4 390	0.06
N	493 034	601	1 054	0.02
O	935 315	2	28	0.00
TS & EEZ	4 115 806	46 300	90 940	0.08




Figure 7.13: Maps from Baird and Wood (2010) showing BOMECEC classes I (left) and M (right) overlaid with the footprint of trawls on or near the seafloor reported on TCEPR forms to 2004–05 for each 25-km<sup>2</sup> cell.

#### 7.3.4. Studies of the Effects of Mobile Bottom Fishing Methods in New Zealand

The widespread nature of bottom trawling suggests that fishing is the main anthropogenic disturbance agent to the seabed throughout most of New Zealand's EEZ. Wind waves are certainly very widespread, but both field studies and modelling (Green *et al.* 1995) suggest that erosion of the seabed deeper than 50 m by waves occurs only very rarely in the New Zealand EEZ. Despite their widespread distribution at the surface, therefore, wind-waves are not a dominant feature of the long-term disturbance regime throughout most of the EEZ. In some places, especially in the coastal zone and in areas close to headlands, straits, or islands, currents and tides may dominate the natural disturbance regime and a community adapted to this type of disturbance will have developed. However, over most of the EEZ between about 100 and 1000 m depth, especially in areas where there are few strong currents, fishing is probably the major broad-scale disturbance agent.

Several studies have been conducted since 1995 in New Zealand, focussing on the effects of various dredge and trawl fishing methods on a variety of different habitats in several geographical locations (Table 7.4). Despite the diversity of these studies, and their different depths, locations, and habitat types, the results are consistent with the global literature on the effects of mobile bottom fishing gear on benthic communities. Generally, there are decreases in the density and diversity of benthic communities and, especially, the density of large, structure-forming epifauna, and long-lived organisms along gradients of increasing fishing intensity. Large, emergent epifauna like sponges and framework-forming corals that provide structured habitat for other fauna are particularly noted as being susceptible to disturbance by mobile bottom fishing methods (Cranfield *et al.* 1999, 2001, 2003, Cryer *et al.* 2000), especially on hard (non sedimentary) seabeds (Clark & Rowden 2009, Clark *et al.* 2010a&b, Williams *et al.* 2011). Even though large emergent fauna seem most susceptible, however, effects have also been shown in the sandy or silty sedimentary systems usually considered to be most resistant to disturbance (Thrush *et al.* 1995, 1998, Cryer *et al.* 2002). Also typical of the international literature is a substantial variation in the extent to which individual New Zealand studies have shown clear effects. For instance, in Foveaux Strait, Cranfield *et al.* (1999, 2001, 2003) inferred substantial

changes in the benthic system caused by over 130 years of oyster dredging, but Michael *et al.* (2006) did not support such conclusions in the same system. Subsequent review of these studies found much common ground but no overall consensus on the long-term effects of dredging on the benthic community of the strait.

These studies have focussed predominantly on changes in patterns in biodiversity associated with trawling and/or dredging and less work has been done to assess changes in ecological process or to estimate the rate of recovery from fishing. Projects that have started on recovery rates are focussed on relatively few habitats and primarily those that are known to be sensitive to physical disturbance, including by trawling or dredging (e.g., seamounts, project ENV2005/16, and areas of high current and natural biogenic structure, projects ENV9805, ENV2005/23 and BEN2009/02). Thus, the understanding of the consequences of fishing (or ceasing fishing) for sustainability, biodiversity, ecological integrity and resilience, and fish stock productivity in the wide variety of New Zealand's benthic habitats remains incomplete. Reducing this uncertainty would allow the testing of the utility and likely long-term productivity of a variety of management strategies, and enable a move towards a regime that maximises value to the nation consistent with Fisheries 2030.

**Table 7.4: Summary of studies of the effects of bottom trawling and dredging in New Zealand waters.**

Location	Approach	Key findings	References
Mercury Islands sandy sediments. Scallop dredge	Experimental	Density of common macrofauna at both sites decreased as a result of dredging at two contrasting sites; some populations were still significantly different from reference plots after 3 months.	Thrush <i>et al.</i> 1995
Hauraki Gulf various soft sediments. Bottom trawl & scallop dredge.	Observational, gradient analysis	Decreases in the density of echinoderms, longlived taxa, epifauna, especially large species, the total number of species and individuals, and the Shannon-Weiner diversity index with increasing fishing pressure (including trawl and scallop dredge). Increases in the density of deposit feeders, small opportunists, and the ratio of small to large heart urchins.	Thrush <i>et al.</i> 1998
Bay of Plenty continental slope. Scampi and other bottom trawls.	Observational, multiple gradient analyses	Depth and historical fishing activity (especially for scampi) at a site were the key drivers of community structure for large epifauna. The Shannon-Weiner diversity index generally decreased with increasing fishing activity and increased with depth. Many species were negatively correlated with fishing activity; fewer were positively correlated (including the target species, scampi).	Cryer <i>et al.</i> 1999 Cryer <i>et al.</i> 2002
Foveaux Strait, sedimentary & biogenic reef. Oyster dredge.	Observational, various	Interpretations of the authors differ. Cranfield <i>et al.</i> 's papers concluded that dredging biogenic reefs for their oysters damages their structure, removes epifauna, and exposes associated sediments to resuspension such that, by 1998, none of the original bryozoan reefs remained. Michael <i>et al.</i> concluded that there are no experimental estimates of the effect of dredging in the strait or on the cumulative effects of fishing or regeneration, that environmental drivers should be included in any assessment, and that the previous conclusions cannot be supported. The authors agree that biogenic bycatch in the fishery has declined over time in regularly-fished areas, that there may have been a reduction in biogenic reefs in the strait since the 1970s, and that simple biogenic reefs appear able to regenerate in areas that are no longer fished (dominated by byssally attached mussels or reef-building bryozoans). There is no consensus that reefs in Foveaux Strait were (or were not) extensive or dominated by the bryozoan <i>Cinctopora</i> .	Cranfield <i>et al.</i> 1999, 2001, 2003 Michael <i>et al.</i> 2006
Spirits Bay, sedimentary &	Observational, gradient	In 1999, depth was found to be the most important explanatory variable for benthic community composition but a coarse index of	Cryer <i>et al.</i> 2000

*AEBAR 2012: Benthic impacts*

biogenic areas. Scallop dredge.	analysis	<p>dredge fishing intensity was more important than substrate type for many taxonomic groups. Sponges seemed most affected by scallop dredging, and samples taken in an area once rich in sponges had few species in 1999. This area had probably been intensively dredged for scallops. Analysis of historical samples of scallop survey bycatch showed a marked decline in sponge species richness between 1996 and 1998.</p> <p>In 2006, significant differences were identified among areas within which fishing was or was not allowed. Species contributing to these differences included those identified as being most vulnerable to the effects of fishing. These differences could not be attributed specifically to fishing because of interactions with environmental gradients and uncertainty over the history of fishing. No significant change between 1999 and 2006 was identified.</p> <p>In 2010, analysis of both epifaunal and infaunal community data identified change since 2006, and significant depth, habitat and fishing effects. The combined fishing effects accounted for 15 – 30% of the total variance (about half of the explained variance). Individual species responses to fishing were examined, and those identified as most sensitive to fishing in this analysis had previously been categorised as sensitive on the basis of life history characteristics within the 2006 study.</p>	Tuck <i>et al.</i> 2009 Tuck & Hewitt 2012
Tasman & Golden Bays. Bottom trawl, scallop & oyster dredge	Observational, gradient analysis	A gradient analysis was adopted to investigate the importance of the different factors affecting epifaunal and infaunal communities in Tasman and Golden Bays. Fishing was consistently identified as an important factor in explaining variance in community structure, with recent trawl and scallop effort being more important than other fishing terms. Important environmental variables included maximum current speed, maximum wave height, depth, % mud, and salinity. Fishing accounted for 31–50% of the explained variance in epifaunal and infaunal community composition, species richness, and Shannon-Weiner diversity. Overall, models explained 30–54% of variance, and additional spatial patterns identified in the analysis explained a further 5–16% of variance.	Tuck <i>et al.</i> 2011
Graveyard complex “seamounts”, northern Chatham Rise. Orange roughly bottom trawl.	Observational, multiple analyses	From surveys in 2001 and 2006, substrate diversity and the amount of intact coral matrix were lower on fished seamounts. Conversely, the proportions of bedrock and coral rubble were higher. No change in the megafaunal assemblage consistent with recovery over 5–10 years on seamounts where trawling had ceased. Some taxa had significantly higher abundance in later surveys. This may be because of their resistance to the direct effects of trawling, their protection in natural refuges, or because these taxa represent the earliest stages of seamount recolonisation.	Clark <i>et al.</i> 2010a&b Williams <i>et al.</i> 2011

An expert based assessment of 65 threats to 62 marine habitats from saltmarsh to the abyss (MacDiarmid *et al.* 2012) concluded that only 7 of the 20 most important threats to New Zealand marine habitats were directly related to human activities within the marine environment. The most important of these was bottom trawling (ranked third equal most important), but invasive species, coastal engineering, and aquaculture were also ranked highly. However, the two top threats, five of the top six threats, and over half of the 26 top threats stemmed largely or completely from human activities external to the marine environment (the most important being ocean acidification, rising sea temperatures, and sedimentation resulting from changes in land-use). The assessment suggested that the number and severity of threats to marine habitats declines with depth, particularly deeper than about 50 m. Shallow coastal habitats face up to 52 non-trivial threats whereas most deep water habitats are threatened by fewer than five. Coastal and estuarine reef, sand, and mud habitats were considered to be the most threatened habitats whereas slope and deep water habitats were among the least threatened.

### **7.3.5. Current research**

Project BEN2007/01 is a 5-year project to assess the effects of fishing on soft sediment habitat, fauna, and processes across the range of habitat types in the TS and EEZ. Sampling and analytical strategies for such broad-scale assessments have been developed and the project has moved into a phase of data collection, collation, and analysis. Two field-based “case studies” in different habitat types will be assessed, and a variety of existing information will be drawn together and analysed to provide a TS & EEZ-wide perspective. The focus of this study is on the relative sensitivities of different habitats in the TS and EEZ to disturbance by mobile bottom fishing methods.


Project DAE2010/04 provides for an annual assessment of the “footprint” of middle depth and deepwater trawl fisheries, including the overlap of the footprint with various depth ranges and habitat classes. Inshore fisheries, including shellfish dredge fisheries, are not covered under this project, so the focus is on offshore fisheries and habitats.

Project DEE2010/05 provides for the development of a suite of ecosystem and environmental indicators for deepwater trawl fisheries. The focus of this study is on developing a cost-effective approach to monitoring ecosystem status (e.g., providing a mechanism to detect ocean climatic changes or regime shifts that could affect fisheries production) or the potential effects of deepwater trawl fisheries (such as changes to benthic invertebrate diversity). The suite could include information that may stem from project DEE2010/06 which provides for a desk-top assessment of the extent to which information can be collected cost-effectively on trends in benthic systems inside and outside of the trawled areas.

Project BEN2012/01 will use existing data and classifications to describe the distribution of benthic habitats, estimate the sensitivity to fishing disturbance of the species within these habitats, and then describe the spatial pattern of fishing using bottom trawls, Danish seine and dredges, to assess the overlap with each habitat class.

Several MBIE-funded projects also have strong linkages with MPI research on benthic impacts. These include “Vulnerable Deep-Sea Communities” (CO1X0906) which is analysing the time series of data from the “Graveyard seamounts” (surveys in 2001, 2006, 2009, all carried out with support from MFish or the cross-departmental Oceans Survey 20/20 programme), as well as evaluating the relative vulnerability of benthic communities in several deep-sea habitats (e.g., seamounts, canyons, continental slope, hydrothermal vents, seeps) and their risk from bottom trawling.

## 7.4. Indicators and trends

Annual number of tows	2010/11 fishing year: 86 024 trawl tows 35 150 shellfish dredge tows				
Trend in number of tows	<p>Trawl effort stable, dredge effort decreasing:</p>  <p>The top chart displays 'Total reported tows' from 1990 to 2012. The y-axis ranges from 0 to 160,000. The legend indicates three categories: TCER (green), CELR (red), and TCEPR (blue). The total tows show a general decline from approximately 110,000 in 1990 to around 80,000 in 2012. The bottom chart displays 'Total reported dredge tows' from 1990 to 2012. The y-axis ranges from 0 to 200,000. The legend indicates two categories: oysters (red) and scallops (blue). The total dredge tows show a significant decline from approximately 190,000 in 1990 to around 35,000 in 2012.</p>				
Annual and cumulative (16 year) overlap of BOMECS habitat classes up to 2004/05	BOMECS class	Area (km <sup>2</sup> )	Min. annual footprint area (km <sup>2</sup> )	Max. annual footprint area (km <sup>2</sup> )	Cumulative (16 yr) proportion overlapped
This analysis will be updated for deepwater trawl fisheries in 2013	A*	27 557	121	4 026	0.42
	B*	12 420	40	484	0.19
	C*	89 710	4 271	11 374	0.58
	D*	27 268	377	1 602	0.30
	E*	60 990	4 046	7 108	0.40
	F	38 608	517	1 391	0.13
	G	6 342	132	833	0.34
	H	138 550	9 583	20 344	0.45
	I	52 224	5 511	18 016	0.70
	J	311 361	10 469	15 975	0.18
	K	1 290	-	2	0.01
	L	198 577	4 238	13 599	0.24
	M	233 825	895	4 390	0.06
	N	493 034	601	1 054	0.02
	O	935 315	2	28	0.00
	TS & EEZ	4 115 806	46 300	90 940	0.08

\* the trawl footprint and proportion overlapped in coastal classes A–E will be grossly underestimated because CELR data are excluded.

## 7.5. References

- Auster PJ; Langton RW (1999). The effects of fishing on fish habitat. *American Fisheries Society Symposium* 22:150–187.
- Barnes PW; Thomas JP (eds) (2005). Benthic habitats and effects of fishing. *American Fisheries Society Symposium* 41. American Fisheries Society, Bethesda, MD.
- Baird SJ; Bagley NW; Wood BA; Dunn A; Beentjes M (2002). The spatial extent and nature of mobile bottom fishing methods within the New Zealand EEZ, 1989–90 to 1998–99. Final Research Report for Objective 1 of project ENV2000/05.
- Baird SJ; Wood BA (2010). Extent of coverage of 15 environmental classes within the New Zealand EEZ by commercial trawling with seafloor contact. Final Research Report for Objective 5 of project BEN2006/01.
- Baird SJ; Wood BA; Bagley NW (2009). The extent of trawling on or near the seafloor in relation to benthic-optimised marine environment classes within the New Zealand EEZ. Final Research Report for Objective 5 of project BEN2006/01.
- Baird SJ; Wood BA; Bagley NW (2011). Nature and extent of commercial fishing effort on or near the seafloor within the New Zealand 200 n. mile Exclusive Economic Zone 1989/90 to 2004/05. *New Zealand Aquatic Environment and Biodiversity Report No. 73*. 144 p.
- Baird SJ; Wood BA; Clark MR; Bagley NW; McKenzie A (2006). Description of the spatial extent and nature of disturbances by bottom trawls in Chatham Rise and Southern Plateau fisheries. Final Research Report for project ENV2003/03. 139 p.
- Blyth RE; Kaiser MJ; Edwards-Jones G; Hart PJB (2004). Implications of a zoned fishery management system for marine benthic communities. *Journal of Applied Ecology* 41: 951–961
- Bowden DA; Compton TJ; Snelder TH; Hewitt JE (2011). Evaluation of the New Zealand Marine Environment Classifications using Ocean Survey 20/20 data from Chatham Rise & Challenger Plateau. *New Zealand Aquatic Environment and Biodiversity Report No. 77*.
- Bradshaw C; Tjensvoll I; Sköld M; Allan IJ; Molvaer J; Magnusson J; Naes K; Nilsson HC (2012). Bottom trawling resuspends sediment and releases bioavailable contaminants in a polluted fjord. *Environmental Pollution* 170: 232–241
- Bradshaw C; Veale LO; Brand AR (2002). The role of scallop-dredge disturbance in long-term changes in Irish Sea benthic communities: a re-analysis of an historical dataset. *Journal of Sea Research* 47: 161–184.
- Brodie S; Clark M (2003). The New Zealand Seamount Management Strategy – steps towards conserving offshore marine habitat. In: Beumer, J.P.; Grant, A.; Smith, D.C. (eds). *Aquatic Protected Areas: what works best and how do we know?* Proceedings of the World Congress on Aquatic Protected Areas, Cairns, Australia, August 2002. Australian Society of Fish Biology, Australia., pp. 664–673. Australian Society of Fish Biology, Cairns, Australia.
- Churchill JH (1989). The effect of commercial trawling on sediment resuspension and transport over the Middle Atlantic Bight Continental Shelf. *Continental Shelf Research* 9: 841–864.
- Clark MR (1999). Fisheries for orange roughy (*Hoplostethus atlanticus*) on seamounts in New Zealand. *Oceanologica Acta* 22: 593–602.
- Clark MR; Bowden DA; Baird SJ; Stewart R (2010). Effects of fishing on the benthic biodiversity of seamounts of the “Graveyard” complex, northern Chatham Rise. *NZ Aquatic Environment and Biodiversity Report No. 46*. 40 p.
- Clark MR; Koslow JA (2007). Impacts of fisheries on seamounts. Chapter 19. p. 413–441 In: Pitcher TJ, Morato T, Hart PJB, Clark MR, Haggan N, Santos RS (eds). *Seamounts: ecology, fisheries, and conservation*. Blackwell Fisheries and Aquatic Resources Series 12. Blackwell Publishing, Oxford. 527 pp.
- Clark M; O'Driscoll R (2003). Deepwater fisheries and aspects of their impact on seamount habitat in New Zealand. *Journal of Northwest Atlantic Fishery Science* 31: 441–458.
- Clark MR; Rowden AA (2009). Effect of deepwater trawling on the macro-invertebrate assemblages of seamounts on the Chatham Rise, New Zealand. *Deep Sea Research* 56: 1540–1554
- Clark MR; Tracey DM; Pallentin A; Schnabel K; Anderson OF; Bowden D (2009). Voyage report of a survey of “seamounts” on the northwest and southeast Chatham Rise (TAN0905). 49 p. (unpublished report available from NIWA, Wellington).
- Clark MR; Watling L; Rowden AA; Guinotte JM; Smith CR (2010). A global seamount classification to aid the scientific design of marine protected area networks. *Journal of Ocean and Coastal Management* 54: 19–36.
- Clark MR; Williams A; Rowden AA; Hobday AJ; Consalvey M (2011). Development of seamount risk assessment: application of the ERAEF approach to Chatham Rise seamount features. *New Zealand Aquatic Environment and Biodiversity Report No. 74*. 18 p.
- Craeymeersch JA; Piet GJ; Rijnsdorp AD; Buijs J (2000). Distribution of macrofauna in relation to the microdistribution of trawling effort. P 187 in Kaiser MJ; de Groot SJ (eds). The effects of fishing on nontarget species and habitats: biological, conservation, and socio-economic issues. Blackwell Scientific, Oxford, UK.
- Cranfield HJ; Michael KP; Doonan IJ (1999). Changes in the distribution of epifaunal reefs and oysters during 130 years of dredging for oysters in Foveaux Strait, southern New Zealand. *Aquatic Conservation of Marine and Freshwater Ecosystems* 9: 461–483
- Cranfield HJ; Carlines G; Michael KP; Dunn A; Stotter DR; Smith DJ (2001). Promising signs of regeneration of blue cod and oyster habitat changed by dredging in Foveaux Strait, southern New Zealand. *New Zealand Journal of Marine and Freshwater Research* 35: 897–908
- Cranfield HJ; Manighetti B; Michael KP; Hill A (2003). Effects of oyster dredging on the distribution of bryozoans biogenic reefs and associated sediments in Foveaux Strait, southern New Zealand. *Continental Shelf Research* 23: 1337–1357
- Cryer M; Coburn R; Hartill BH; O'Shea S; Kendrick T; Doonan I (1999). Scampi stock assessment for 1998 and an analysis of the fish and invertebrate bycatch of scampi trawlers. New Zealand Fisheries Assessment Research Document 99/4. 74 p
- Cryer M; Hartill BH (2002). Relative impacts on benthic invertebrate community structure of trawl fisheries for scampi and finfish in the Bay of Plenty. Final Research Report for Ministry of Fisheries Research Project ENV2000/05 Objective 2. 28 p.
- Cryer M; Hartill BH; O'Shea S (2002). Modification of marine benthos by trawling: toward a generalization for the deep ocean? *Ecological Applications* 12: 1824–1839
- Cryer M; O'Shea S; Gordon DP; Kelly M; Drury JD; Morrison MA; Hill A; Saunders H; Shankar U; Wilkinson M; Foster G (2000). Distribution and structure of benthic invertebrate communities between North Cape and Cape Reinga Final Research Report for Ministry of Fisheries Research Project ENV9805 Objectives 1–4.
- Davies AJ; Guinotte JM (2011). Global habitat suitability for framework-forming cold-water corals. *PLoS One* 6(4).
- Dayton PK; Thrush SF; Agardy MT; Hofman RJ (1995). Environmental effects of fishing. *Aquatic Conservation: Marine and Freshwater Ecosystems* 5:205–232.
- Dayton PK; Thrush SF; Coleman FC (2002). The Ecological Effects of Fishing in Marine Ecosystems of the United States. Arlington, VA: Pew Oceans Comm.
- Dichmont CM; Deng A; Punt AE; Ellis N; Venables WN; Kompas T; Ye Y; Zhou S; Bishop J (2008) Beyond biological performance measures: bringing in economics and the effects of trawling on the benthos. *Fisheries Research* 94: 238–250.

- Ellis N; Pantus F (2001). Management strategy modelling: tools to evaluate trawl management strategies with respect to impacts on benthic biota within the Great Barrier Reef Marine Park area. Cleveland, Qld.: CSIRO Div. Marine Research. Parts 1 & 2, pp132. [http://www.cmar.csiro.au/e-print/open/ellisn\\_2001.pdf](http://www.cmar.csiro.au/e-print/open/ellisn_2001.pdf)
- Ferrier S; Drielsma M; Manion G; Watson G (2002). Extended statistical approaches to modelling spatial pattern in biodiversity in north-east New South Wales: II. Community-level modelling. *Biodiversity and Conservation* 1: 2309–2338.
- Ferrier S; Manion G; Elith J; Richardson K (2007). Using generalised dissimilarity modelling to analyse and predict patterns of beta-diversity in regional biodiversity assessment. *Diversity and Distributions* 13: 252–264.
- Green MO; Vincent CE; McCave IN; Dickson RR; Rees JM; Pearsons ND (1995). Storm sediment transport: observations from the British North Sea shelf. *Continental Shelf Research* 15: 889–912.
- Hall SJ (1994). Physical disturbance and marine benthic communities: life in unconsolidated sediments. *Oceanography and Marine Biology Annual Review* 32: 179–239.
- Hall SJ (1999). The effects of fishing on marine ecosystems and communities. Blackwell Scientific, Oxford, UK.
- Hermesen JM; Collie JS; Valentine PC (2003). Mobile fishing gear reduces benthic megafaunal production on Georges Bank. *Marine Ecology Progress Series* 260: 97–108.
- Hiddink JG; Jennings S; Kaiser MJ; Queiros AM; Duplisea DE; Piet GJ (2006). Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Canadian Journal of Fisheries and Aquatic Sciences* 63:721–36.
- JG Hiddink; AF Johnson; R Kingham; H Hinz (2011). Could our fisheries be more productive? Indirect negative effects of bottom trawl fisheries on fish condition. *Journal of Applied Ecology* 48: 1–9.
- Hinz H; Hiddink JG; Forde J; Kaiser MJ (2008). Large-scale responses of nematode communities to chronic otter-trawl disturbance. *Canadian Journal of Fisheries and Aquatic Sciences* 65: 723–732.
- ICES (1992). Report of the Study Group on Ecosystem Effects of Fishing Activities. ICES CM:1992/G11, 144 pp.
- ICES (2000a). Report of the Advisory Committee on the Marine Environment 2000. ICES Cooperative Research Report #241.
- ICES (2000b). Report of the Working Group on Ecosystem Effects of Fishing Activities. ICES CM 2000/ACME:02.
- Jennings S; Dinmore TA; Duplisea DE; Warr KJ; Lancaster JE (2001). Trawling disturbance can modify benthic production processes. *Journal of Animal Ecology* 70: 459–475.
- Jennings S; Freeman S; Parker R; Duplisea DE; Dinmore TA (2005). *Ecosystem Consequences of Bottom Fishing Disturbance*. American Fisheries Society Symposium 41:73–90.
- Jennings S; Kaiser MJ (1998). The effects of fishing on marine ecosystems. *Advances in Marine Biology and Ecology* 34:201–352.
- Jennings S; Pinnegar JK; Polunin NVC; Warr KJ (2001). Impacts of trawling disturbance on the trophic structure of benthic invertebrate communities. *Marine Ecology Progress Series* 213:127–142.
- Jones JB (1992). Environmental impact of trawling on the seabed: a review. *New Zealand Journal of Marine and Freshwater Research* 26: 59–67.
- Kaiser MJ; Clarke KR; Hinz H; Austen MCV; Somerfield PJ; Karakassis I (2006). Global analysis of the response and recovery of benthic biota to fishing. *Marine Ecology Progress Series* 311: 1–14.
- Kaiser MJ; Collie JS; Hall SJ; Jennings S; Poiner IR (2003). Impacts of fishing gear on marine benthic habitats. Conference on Responsible Fisheries in the Marine Ecosystem. Reykjavik, Iceland, 1–4 October 2001.
- Kaiser MJ; de Groot, SJ (2000). *Effects of fishing on non-target species and habitats: biological, conservation and socio-economic issues*. Blackwell, Oxford, 399 p.
- Leathwick J; Francis M; Julian K (2006). Development of a demersal fish community classification for New Zealand's Exclusive Economic Zone. Prepared for Department of Conservation. NIWA Client Report HAM2006-062. Hamilton.
- Leathwick JR; Rowden AA; Nodder S; Gorman R; Bardsley S; Pinkerton M; Baird SJ; Hadfield M; Currie K; Goh A (2010). Benthic-optimised marine environment classification for New Zealand waters. Final Research Report for Ministry of Fisheries Project BEN2006/01, Objective 5. 52 p. Unpublished report held by Ministry of Fisheries, Wellington.
- Leathwick JR; Snelder T; Chadderton WL; Elith J; Julian K; Ferrier S (2011). Use of generalised dissimilarity modelling to improve the biological discrimination of river and stream classifications. *Freshwater Biology* 56: 21–38.
- Lindeboom H; deGroot SJ (1998). The effects of different types of fisheries on the North Sea and Irish Sea benthic ecosystems. Netherlands Institute of Sea Research, Texel, The Netherlands.
- Lokkeborg S (2005). Impacts of trawling and scallop dredging on benthic habitats and communities. *FAO Tech. Paper* 472.
- Lundquist CJ; Thrush SF; Coco G; Hewitt JE (2010). Interactions between disturbance and dispersal reduce persistence thresholds in a benthic community. *Marine Ecology Progress Series* 413: 217–228.
- MacDiarmid A; McKenzie A; Sturman J; Beaumont J; Mikaloff-Fletcher S; Dunne J (2012). Assessment of anthropogenic threats to New Zealand marine habitats. *New Zealand Aquatic Environment and Biodiversity Report No. 93*. 255 p.
- Michael KP; Kroger K; Richardson K; Hill N (2006). Summary of information in support of the Foveaux Strait Oyster Fishery Plan: the Foveaux Strait ecosystem and effects of oyster dredging. Final Research Report for Ministry of Fisheries project ZBD2005/04. Ministry for the Environment. 2010. Fishing Activity: Fish Stocks Environmental Snapshot. Wellington.
- Ministry of Fisheries 2008. Bottom Fishery Impact Assessment: Bottom Fishing Activities by New Zealand Vessels Fishing in the High Seas in the SPRFMO Area during 2008 and 2009. Ministry of Fisheries, Wellington. 102 pp. Available from <http://www.southpacificrfmo.org/benthic-impact-assessments/>.
- McConnaughey RA; Mier KL; Dew CB (2000). An examination of chronic trawling effects on soft-bottom benthos of the eastern Bering Sea. *ICES Journal of Marine Science* 57: 1377–1388.
- National Marine Fisheries Service (2002). Workshop on the effects of fishing gear on marine habitats off the northeastern United States October 23–25, 2001. *Northeast Fisheries Science Center Reference Document* 02-01.
- National Research Council (2002). Effects of trawling and dredging in seafloor habitat. National Academy Press, Washington DC.
- O'Driscoll RL; Clark MR (2005). Quantifying the relative intensity of fishing on New Zealand seamounts. *New Zealand Journal of Marine and Freshwater Research* 39: 839–850.
- Puig P; Canals M; Company JB; Martin J; Amblas D; Lastras G; Palanques A; Calafat A (2012). Ploughing the deep sea floor. *Nature* 489: 286–289.
- Reiss H; Greenstreet SPR; Siebe K; Ehrich S; Piet GJ; Quirijns F; Robinson L; Wolff WJ; Kronke I (2009). Effects of fishing disturbance on benthic communities and secondary production within an intensively fished area. *Marine Ecology Progress Series* 394: 201–213.
- Rice J (2006). Impacts of mobile bottom gears on seafloor habitats, species, and communities: a review and synthesis of selected international reviews. *Canadian Science Advisory Secretariat Research Document* 2006/057. 35 p. (available from [http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006\\_057\\_e.pdf](http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006_057_e.pdf)).
- Rowden AA; Clark MR; Wright IC (2005). Physical characterisation and a biologically focused classification of seamounts in the New Zealand region. *New Zealand Journal of Marine and Freshwater Research* 39: 1039–1059.

- Rowden AA; Oliver MD; Clark MR; MacKay K (2008). New Zealand's "SEAMOUNT" database: recent updates and its potential use for ecological risk assessment. *Aquatic Environment and Biodiversity Report No. 27*. 49 p.
- Rumohr H (1998). Information on impact of trawling on benthos in Kiel Bay. Annex to Eighth Report of the Benthos Ecology Working Group. ICES, CM: 1989/L19, 80.
- Sainsbury KJ; Punt AE; Smith ADM (2000). Design of operational management strategies for achieving fishery ecosystem objectives. *ICES Journal of Marine Science*, 57: 731–741.
- Schratzberger M; Lampadariou N; Somerfield PJ; Vandepitte L; Vanden Berghe E (2009). The impact of seabed disturbance on nematode communities: linking field and laboratory observations. *Marine Biology* 156:709–724
- Sharp B; Parker S; Smith N. 2009. An impact assessment framework for bottom fishing methods in the CCAMLR Convention area. *CCAMLR Science* 16: 195–210
- Snelder TH; Leathwick JR; Dey KL; Rowden AA; Weatherhead MA; Fenwick GD; Francis MP; Gorman RM; Grieve JM; Hadfield MG; Hewitt JE; Richardson KM; Uddstrom MJ; Zeldis JR (2006). Development of an ecological marine classification in the New Zealand region. *Environmental Management* 39: 12–29.
- Snelder TH; Leathwick JR; Dey KL; Weatherhead MA; Fenwick GD; Francis MP; Gorman RM; Grieve JM; Hadfield MG; Hewitt JE; Hume T; Richardson KM; Rowden AA; Uddstrom MJ; Wild M; Zeldis JR (2005) The New Zealand Marine Environment Classification. Ministry for the Environment. Wellington.
- Snelder TH; Leathwick JR; Image K; Weatherhead MA; Wild M (2004). The New Zealand Marine Environment Classification. NIWA Client Report CHC2004-071. 86 p.
- Thrush SF; Dayton PK (2002). Disturbance to marine benthic habitats by trawling and dredging—implications for marine biodiversity. *Annual Reviews in Ecology and Systematics* 33:449–73
- Thrush SF; Hewitt JE; Cummings VJ; Dayton PK (1995). The impact of habitat disturbance by scallop dredging on marine benthic communities: What can be predicted from the results of experiments? *Marine Ecology Progress Series* 129:141–150.
- Thrush SF; Hewitt JE; Cummings VJ; Dayton PK; Cryer M; Turner SJ; Funnell GA; Budd RG; Milburn CJ; Wilkinson MR (1998). Disturbance of the marine benthic habitat by commercial fishing: impacts at the scale of the fishery. *Ecological Applications* 8: 866–879.
- Tillin HM; Hiddink JG; Jennings S; Kaiser MJ (2006). Chronic bottom trawling alters the functional composition of benthic invertebrate communities on a sea-basin scale. *Marine Ecology Progress Series* 318:31–45
- Tuck ID; Parkinson DM; Dey K; Oldman J; Wadwha S (2006). Information on benthic impacts in support of the Coromandel scallop fisheries plan. Final Research Report for Ministry of Fisheries Research Project ZBD2005/15
- Tuck ID; Drury J; Kelly M; Gerring P (2009) Designing a programme to monitor the recovery of the benthic community between North Cape and Cape Reinga. Final Research Report for Ministry of Fisheries Research Project ENV2005-23.
- Tuck ID; Hewitt JE (2012). Monitoring change in benthic communities in Spirits Bay. Final Research Report for Ministry of Fisheries research project BEN200901. 51 p.
- Tuck ID; Hewitt JE; Handley S; Willis T; Carter M; Hadfield M; Gorman R; Cairney D; Brown S; Palmer A (2011). Assessing the effects of fishing on soft sediment habitat, fauna and processes. Progress Report for Ministry of Fisheries research project BEN200701. (Unpublished report held by MPI, Wellington.).
- Veale LO; Hill AS; Hawkins SJ; Brand AR (2000). Effects of long-term physical disturbance by commercial scallop fishing on subtidal epifaunal assemblages and habitats. *Marine Biology* 137: 325–337
- Warnkena KW; Gilla GA; Dellapennaa TM; Lehmana RD; Harperb DE; Allison MA (2003). The effects of shrimp trawling on sediment oxygen consumption and the fluxes of trace metals and nutrients from estuarine sediments. *Estuarine, Coastal and Shelf Science* 57: 25–42
- Watling L; Norse EA (1998). Disturbance of the seabed by mobile fishing gear: a comparison to forest clearcutting. *Conservation Biology*, 12: 1180–1197.
- Williams A; Dowdney J; Smith ADM; Hobday AJ; Fuller M (2011). Evaluating impacts of fishing on benthic habitats: a risk assessment framework applied to Australian fisheries. *Fisheries Research* 112: 154–167
- Williams A; Schlacher TA; Rowden AA; Althaus F; Clark MR; Bowden DA; Stewart R; Bax NJ; Consalvey M; Kloser RJ (2010). Seamount megabenthic assemblages fail to recover from trawling impacts. *Marine Ecology* 31 (Suppl. 1):183–199.

## **THEME 4: ECOSYSTEM EFFECTS**

## 8. New Zealand Climate and Oceanic Setting


<i>Scope of chapter</i>	Overview of primary productivity, oceanography, benthic-pelagic coupling and oceanic climate trends in the SW Pacific region.
<i>Area covered</i>	New Zealand regional setting
<i>Focal localities</i>	Pan New Zealand waters
<i>Key issues</i>	<ul style="list-style-type: none"> <li>• Climate and oceanographic variability and change are of relevance to fisheries and the broader marine environment</li> <li>• Allows for improved understanding of the links between observed patterns and drivers of biological processes.</li> </ul>
<i>Emerging issues</i>	<ul style="list-style-type: none"> <li>• New Zealand's oceanic climate is changing</li> <li>• Causal mechanisms that link the dynamics of a variable marine environment to variation in biological productivity, particularly of fisheries, are not well understood in New Zealand.</li> <li>• Need for improved understanding of the linkages between the pelagic and benthic environment (i.e., benthic-pelagic coupling).</li> <li>• The cumulative effects of ocean climate change and other anthropogenic stressors on aquatic ecosystems (productivity, structure and function) are likely to be high.</li> <li>• Some long-term trends in the marine environment are available at a national scale but are not reported.</li> </ul> <p>Growing recognition that stressors will act individually &amp; interactively, confounding prediction of net effects of climate change</p>
<i>MPI Research (current)</i>	Projects include ZBD2005-05: Long-term effects of climate variation and human impacts on the structure and functioning of New Zealand shelf ecosystems; ZBD2008-11 Predicting plankton biodiversity & productivity with ocean acidification; ZBD2009-13. Ocean acidification impact on key NZ molluscs; ZBD2010-40. Marine Environmental Monitoring Programme; ZBD2010-41 Deepsea fisheries habitat and ocean acidification.
<i>NZ Research (current)</i>	NIWA Coast & Oceans Centre, Climate Centre; University of Otago-NIWA shelf carbonate geochemistry & bryozoans; Munida time-series transect; Geomarine Services-foraminiferal record of human impact; Regional Council monitoring programmes; Statistics New Zealand Environmental Domain review.
<i>Links to Fisheries 2030 and MPI's Our Strategy</i>	Environmental Outcome Objective 1; environmental principles of Fisheries 2030; MPI's "Our Strategy 2030": two key stated focuses are to maximise export opportunities and improve sector productivity; increase sustainable resource use, and protect from biological risk.
<i>Related issues and/or Chapters</i>	<ul style="list-style-type: none"> <li>• Ocean related climate variability and change are predicted to have major implications for fishstock distributions and abundance, reproductive success, ecosystem goods and services, deepsea coral habitat and Habitats of Particular Significance to Fisheries Management,</li> <li>• A significant warming event occurred in the late 1990s</li> <li>• A regime shift to the negative phase of the IPO occurred in about 2000,</li> </ul>

	<p>which is likely to result in fewer El Niño events for a 20–30 year period, i.e., less zonal westerly winds (already apparent compared to the 1980–2000 period) and increased temperatures; this is the first regime shift to occur since most of our fisheries monitoring time series have started (the previous shift was in the late 1970s), and will likely impact on fish productivity</p> <ul style="list-style-type: none"> <li>• New Zealand trends of increasing air and sea temperatures and ocean acidification are consistent with global trends.</li> </ul>
--	--

## 8.1. Context

Climate and oceanographic conditions play an important role in driving the productivity of our oceans and the abundance and distribution of our fishstocks, and hence fisheries. A full analysis of trends in climate and oceanographic variables in New Zealand is given in Hurst *et al.* (2012) and is now being developed as an Ocean Climate Change Atlas for New Zealand waters (Boyd and Law.2011).

New Zealand is essentially part of a large submerged continent (Figure 8.1).



**Figure 8.1:** New Zealand land mass area 250,000 km<sup>2</sup>; EEZ & territorial sea area (pink) 4,200,000 km<sup>2</sup>; extended continental shelf extension area (light green) 1,700,000 km<sup>2</sup>; Total area of marine jurisdiction 5,900,000 km<sup>2</sup>. The black line shows the boundary of the New Zealand EEZ, the yellow line indicates the extension to New Zealand's legal continental shelf, and the red line the agreed Australia/New Zealand boundary under UNCLOS Article 76. Image courtesy of GNS.

The territorial sea (TS extending from mean low water shore line to 12 nautical miles) and Exclusive Economic Zone (the EEZ, extending from 12 nautical miles to 200 miles offshore) and the extended continental shelf (ECS) combine to produce one of the largest areas of marine jurisdiction in the world, an area of almost 6 million square kilometres, (Figure 8.1). New Zealand waters straddle more than 25 degrees of latitude from 30° S in warm subtropical waters to 56° S in cooler, subantarctic

waters, and 28 degrees of longitude from 161° E in the Tasman Sea to 171° W in the west Pacific Ocean. New Zealand's coastline, with its numerous embayments, is also long, with estimates ranging from 15,000 to 18,000 km, depending on the method used for measurement (Gordon *et al.* 2010).

New Zealand lies across an active subduction zone in the western Pacific plate, tectonic activity and volcanism have resulted in a diverse and varied seascape within the EEZ. The undersea topography comprises a relatively narrow band of continental shelf down to 200 m water depth, extensive continental slope areas from 200 to 1000 m, extensive abyssal plains, submarine canyons and deep sea trenches, ridge systems and numerous seamounts and other underwater topographic features such as hills and knolls. There are three significant submarine plateaus, the Challenger Plateau, the Campbell Plateau in the subantarctic, and the Chatham Rise (Figure 8.2).

Disturbance of current flow across the plateaus and around the New Zealand landmass gives rise to higher ocean productivity than might be expected, given New Zealand's isolated location in the generally oligotrophic western Pacific Ocean (Figure 8.3). Higher ocean colour, reflecting higher levels of productivity, is typically found around the coast and to the east across the Chatham Rise (Figure 8.3; Pinkerton *et al.* 2005). The coastal waters and plateaus support a range of commercial shellfish and finfish fisheries from the shoreline to depths of about 1500 m. Seamounts, seamount chains and ridge structures in suitable depths provide additional localized areas of upwelling and increased productivity sometimes associated with commercial fisheries.






Figure 8.2 Undersea topography of New Zealand (red shallow to blue deep). White dash line shows the EEZ boundary. Image courtesy of NIWA.



**Figure 8.3: SeaWiFS image showing elevated chlorophyll *a* (green) near New Zealand. Image courtesy of NOAA**

Both Figures 8.3 and 8.4 (left panel) show that the strongest chlorophyll *a* and ocean colour are associated with the coastal shelf around New Zealand and the Chatham Rise. Although remote sensing cannot readily distinguish between primary productivity (from phytoplankton) and sediments in freshwater runoff, so interpretation of the relative productivity levels inshore has to be made in conjunction with knowledge of river flow, it is clear that the Chatham Rise has the highest productivity levels in the region. Globally, New Zealand net primary productivity levels in the sea are higher compared with most of Australasia, but lower than most coastal upwelling systems around the world (Willis *et al.*, 2007).




**Figure 8.4: Left panel: Ocean colour in the New Zealand region from satellite imagery. Red shows the highest intensity of ocean colour typically associated with higher primary productivity. Right panel: The relative concentrations of particulate organic carbon (POC) that reach the seafloor. Red shows the highest levels, which are likely to be associated with areas of enhanced benthic productivity (based on the model of Lutz *et al.* (2007)). Images courtesy of NIWA.**

Patterns in surface waters of primary productivity are mirrored to an extent in the amount of “energy” that sinks to the seafloor (Figure 8.4). This POC flux is based on a model which accounts for sinking rates of dead organisms and predation in the water column (Lutz *et al.* 2007). This is a potential

surrogate of benthic production, and indicates where benthic-pelagic coupling may be strong. Highest levels of POC flux match with surface productivity to a large extent, with coastal waters (including around the offshore islands) and the Chatham Rise having high estimated production.

The Tasman Sea (west of New Zealand) is separated from the South Pacific Gyre by the New Zealand landmass (Figure 8.5). The South Pacific Western Boundary Current, the East Australian Current (EAC) flows down the east coast of Australia, before separating from the Australian landmass in a variable eddy field at about 31 or 32°S (Ridgway & Dunn 2003). The bulk of the separated flow crosses the Tasman Sea as the Tasman Front (Stanton 1981; Ridgway & Dunn 2003), before a portion of the flow attaches to New Zealand, flowing down the northeast coast as the East Auckland Current (Stanton *et al.* 1997). In the southern limit of the Tasman Sea is the Subtropical Front, which passes south of Tasmania and approaches New Zealand at the latitude of Fiordland (Stanton 1988), before diverting southward around New Zealand, and then northward up the southeast coast of New Zealand where it is locally called the Southland Front (Heath 1985; Chiswell 1996; Sutton 2003).



**Figure 8.5: Circulation around New Zealand.** TF Tasman Front (large red arrows), WAUC West Auckland Current, EAUC East Auckland Current, NCE North Cape Eddy, ECE East Cape Eddy, ECC East Cape Current, WE Wairarapa Eddy, DC D'Urville Current, WC Westland Current, SC Southland Current, SF Southland Front, STW Subtropical Water, STF Subtropical Front (left diagonal hashed area), SAW Subantarctic Water, SAF Subantarctic Front (right diagonal hashed area), ACC Antarctic Circumpolar Current, CSW Circumpolar Surface Water, DWBC Deep Western Boundary Current (large purple arrows) (after Carter *et al.* 1998).

The water in the eastern central Tasman Sea south of the Tasman Front, east of the influence of the EAC and north of the Subtropical Front is thought to be relatively quiescent. Ridgway & Dunne (2003) show eastward surface flow across the interior of the Tasman Sea sourced from the southernmost limit of the EAC, with the flow bifurcating around Challenger Plateau and, ultimately,

New Zealand. Reid's (1986) analysis indicates that a small anticyclonic gyre exists in the western Tasman Sea at 1000–2500 m depth. This gyre is centred at about 35°S, 155°E on the offshore side of the EAC and west of Challenger Plateau. All indications are that the eastern Tasman region overlying Challenger Plateau is not very energetic.

This is in contrast with the east coast of both the North and South Islands, and Cook Strait, which are highly energetic. Campbell Plateau waters are well mixed though nutrient limited (iron), leading to tight coupling between trophic levels (Bradford-Grieve *et al.* 2003). The Subtropical Front lies along the Chatham Rise and turbulence and upwelling results in relatively high primary productivity in the area.

## 8.2. Indicators and trends

### 8.2.1. Sea temperature

Sea surface temperature (SST), sea surface height (SSH), air temperature and ocean temperature to 800m depth, all exhibit some correlation with each other over seasonal and inter-annual time scales (Hurst *et al.* 2012). Air temperatures have increased by about 1°C since 1900 (Figure 8.6).




Figure 8.6: Annual time series in New Zealand. NOAA annual mean sea surface temperatures (blue line)<sup>25</sup> and NIWA's seven-station annual mean air temperature composite series (red line), expressed as anomalies relative to the 1971-2000 climatological average. Linear trends over the period 1909-2009, in °C/century, are noted under the graph. (Image Source Mullan *et al.* 2010)

Although a linear trend has been fitted to the seven-station temperatures in Figure 8.6, the variations in temperature over time are not completely uniform. For example, a markedly large warming occurred through the periods 1940-1960 and 1990-2010. These higher frequency variations can be related to fluctuations in the prevailing north-south airflow across New Zealand (Mullan *et al.* 2010). Temperatures are higher in years with stronger northerly flow, and are lower in years with stronger southerly flow. One would expect this, since southerly flow transports cool air from the Southern Oceans up over New Zealand.

<sup>25</sup> <http://www.ncdc.noaa.gov/oa/climate/research/sst/ersstv3.php>

The unusually steep warming in the 1940-1960 period is paralleled by an unusually large increase in northerly flow during this same period Mullan *et al.* (2010). On a longer timeframe, there has been a trend towards less northerly flow (more southerly) since about 1960 Mullan *et al.* (2010). However, New Zealand temperatures have continued to increase over this time, albeit at a reduced rate compared with earlier in the 20<sup>th</sup> century. This is consistent with a warming of the whole region of the southwest Pacific within which New Zealand is situated (Mullan *et al.* 2010).

Mullan *et al.* 2010 describe the pattern of warming in New Zealand as consistent with changes in sea surface temperature and prevailing winds. Their review shows enhanced rates of warming (in units of °C/decade) down the Australian coast and to the east of the North Island, and much lower rates of warming south and east of the South Island (Figure 8.7).




Figure 8.7: Trends in sea surface temperature, in °C/decade over the period 1909-2009, calculated from the NOAA\_ERSST\_v3 data-set (provided by NOAA's ESRL Physical Sciences Division, Boulder, Colorado, USA, from their web site at <http://www.esrl.noaa.gov/psd/>). The data values are on a 2° latitude-longitude grid. The box around New Zealand denotes the region used to produce the area-averaged sea temperatures plotted in Figure 2. (Image Source Mullan *et al.* 2010.)

Figure 8.8 gives a broader spatial picture at much higher resolution (but a shorter period, since 1982). It is apparent that sea temperatures are increasing north of about 45°S; they are increasing more slowly, and actually decreasing in recent decades, off the Otago coast and south of New Zealand. This regional pattern of cooling (or only slow warming) to the south, and strong warming in the Tasman and western Pacific can be related to increasing westerly winds and their effect on ocean circulation Mullan *et al.* (2010). Thompson and Solomon (2002) discuss the increase in Southern Hemisphere westerlies and the relationship to global warming; Roemmich *et al.* (2007) describe recent ocean circulation changes; Thompson *et al.* (2009) discuss the consequent effect on sea surface temperatures in the Tasman Sea.




Figure 8.8: Trends in sea surface temperature, in  $^{\circ}\text{C}/\text{decade}$  over the period 1982-2009. The data are again taken from NOAA, but are based on daily interpolated satellite measurements over a much finer  $0.25^{\circ}$  grid. See <http://www.ncdc.noaa.gov/oa/climate/research/sst/oi-daily.php>. This product is the result of an objective analysis, an optimum interpolation rather than a pure satellite retrieval, so as to correct for issues like the effect of the Mt Pinatubo eruption aerosols on satellite detected radiances. It is described in Reynolds *et al.* (2007)

Sea surface temperatures (SST) derived from satellite data have been compared to empirical CTD measurements made from relevant sub-areas of the Chatham Rise and Sub-Antarctic during trawl surveys. This showed good correlations, reassuring us that satellite-derived SST provided a realistic measure of sea surface temperature for these regions in years before CTD data were available (O'Driscoll *et al.* 2011).

Coastal SST data, particularly the longer time series from Leigh and Portobello, have been used in studies attempting to link processes in the marine environment with temperature. The negative relationship between SST and SOI is broadly consistent across the 40 years of data although the pattern is less clear post 1997 (Figure 8.9). The clearest fisheries example of a link between coastal SST and fish recruitment and growth is for northern stocks of snapper (*Pagrus auratus*), where relatively high recruitment and faster growth rates have been correlated with warmer conditions from the Leigh SST series (Francis 1993, 1994a).





Figure 8.9: Sea surface temperature (SST) anomalies from SST measurements at Leigh (Auckland University Marine Laboratory) and Southern Oscillation Index (SOI) anomalies. (Image from Hurst *et al.* 2012.)

Temperature fluctuations also occur at depth in the ocean as demonstrated by changes in temperature down to 800 m in the eastern Tasman Sea between 1992 and 2008 (Figure 8.10).

The ocean temperature between Sydney and Wellington has been sampled about four times per year since 1991. The measurements are made in collaboration with the Scripps Institution of Oceanography. Analyses of the subsurface temperature field using these data include Sutton & Roemmich (2001) and Sutton *et al.* (2005). The index presented for this transect (Figure 8.10) is for the most eastern section closest to New Zealand (161.5°E and 172°E). The eastern Tasman is chosen because it is closer to New Zealand, and because it has less oceanographic variability which can mask subtle interannual changes. The section of the transect shown is along fairly constant latitude is therefore unaffected by latitudinal temperature and seasonal cycle variation. The upper panel shows the temperature averaged along the transect between the surface and 800m and from 1991 to the most recent sampling.



**Figure 8.10: Eastern Tasman ocean temperature: Wellington to Sydney 1991–2008.** Coloured scale to the right is temperature °C. (Image from Hurst *et al.* 2012, after Sutton *et al.* 2005)

The seasonal cycle is clearly visible in the upper 100–150m. There is a more subtle warming signal that occurred through the late 1990s, which is apparent by the isotherms increasing in depth through that time period. This warming was significant in that it extended to the full 800m of the measurements (effectively the full depth of the eastern Tasman Sea). It also began during an El Niño, period when conditions would be expected to be cool. Finally, it was thought to be linked to a large-scale warming event centred on 40°S that had hemispheric and perhaps global implications. This warming has been discussed by Sutton *et al.* (2005) who examined the local signals, Bowen *et al.* (2006) who studied the propagation of the signal into the New Zealand area, and Roemmich *et al.* (2007), who examined the broad-scale signal over the entire South Pacific Ocean. Roemmich *et al.* (2007) hypothesized that the ultimate forcing was due to an increase in high latitude westerly winds effectively speeding up the entire South Pacific gyre.

Other phenomena have led to periods of warming that are not as yet fully understood. In particular a period of widespread warming in the Tasman Sea to depths of at least 800 m, 1996–2002 (Sutton *et*

*al.* 2005). Both stochastic environmental variability and predictable cycles of change influence the productivity and distribution of marine biota in our region.

## 8.2.2. Climate variables

The Interdecadal Pacific Oscillation (IPO) is a Pacific-wide reorganisation of the heat content of the upper ocean and represents large-scale, decadal temperature variability, with changes in phase (or “regime shifts”) over 10–30 year time scales. In the past 100 years, regime shifts occurred in 1925, 1947, 1977 and 2000 (Figure 8.11). The latest shift should result in New Zealand experiencing periods of reduced westerlies, with associated warmer air and sea temperatures and reduced upwelling on western coasts (Hurst *et al.* 2012).




Figure 8.11: Smoothed index of the Interdecadal Pacific Oscillation (IPO) since 1900. (Image source NIWA based on data from the United Kingdom Meteorological Office, UKMO).

The El Niño-Southern Oscillation (ENSO) cycle in the tropical Pacific has a strong influence on New Zealand. ENSO is described here by the Southern Oscillation Index (SOI), a measure of the difference in mean sea-level pressure between Tahiti (east Pacific) and Darwin (west Pacific). When the SOI is strongly positive, a La Niña event is taking place and New Zealand tends to experience more north easterlies, reduced westerly winds, and milder, more settled, warmer anticyclonic weather and warmer sea temperatures (Hurst *et al.* 2012). When the SOI is strongly negative, an El Niño event is taking place and New Zealand tends to experience increased westerly and south-westerly winds and cooler, less settled weather and enhanced along shelf upwelling off the west coast South Island and north east North Island (Shirtcliffe 1990, Zeldis *et al.* 2004, Chang & Mullan 2003). The SOI is available monthly from 1876 (Mullan 1995) (Figure 8.12).




Figure 8.12: Southern Oscillation Index (SOI) 13-month running mean 1876-2010. Red indicates warmer temperatures, blue indicates cooler conditions for New Zealand. (Image courtesy of NIWA.)

### 8.2.3. Ocean acidification

An increase in atmospheric CO<sub>2</sub> since the industrial revolution has been paralleled by an increase in CO<sub>2</sub> concentrations in the upper ocean (Sabine *et al.* 2004), with global ocean uptake on the order of ~2 gigatonnes (Gt) per annum (~30% of global anthropogenic emissions, IPCC, 2001a). The anthropogenic CO<sub>2</sub> signal is apparent to an average depth of ~1000m.

The increasing rate of CO<sub>2</sub> input from the atmosphere has surpassed the ocean's natural buffering capacity and so the surface of the ocean is becoming more acidic. This is because carbon dioxide absorbed by seawater reacts with H<sub>2</sub>O to form carbonic acid, the dissociation of which releases hydrogen ions, so raising the acidity and lowering pH of seawater. Since 1850, average surface ocean pH has decreased by 0.1 units, with a further decrease of 0.4 units to 7.9 predicted by 2100 (Houghton *et al.*, 2001). The pH scale is logarithmic, so a 0.4 pH decrease corresponds to a 150% increase in hydrogen ion concentration. Both the predicted pH in 2100 and the rate of change in pH are outside the range experienced by the oceans for at least half a million years. In the absence of any decrease in CO<sub>2</sub> emissions this trend is proposed by Caldeira & Wickett, (2003) to continue.

In New Zealand, the projected change in surface water pH between 1990 and 2070 is a decrease of 0.15-0.18 pH units (Hobday *et al.* 2006). The only time series of dissolved pCO<sub>2</sub> and pH in NZ waters is the bimonthly sampling of a transect across neritic, subtropical and subantarctic waters off the Otago shelf since 1998 (University of Otago/NIWA Munida Otago Shelf Time Series). Dissolved pCO<sub>2</sub> shows some indication of an increase although this is not linear and does not correlate with rise in atmospheric CO<sub>2</sub> (see Fig 8.13).




Figure 8.13: pCO<sub>2</sub> (partial pressure of CO<sub>2</sub>) in subantarctic surface seawater from the *R.V. Munida* transect, 1998–2012. (Image courtesy of K. Currie, NIWA)

The Munida time-series pH data shows a decline in subantarctic surface waters since 1998 (Fig 8.14). Addition of a sine-wave function to the pH data suggests a) a linear decline in surface water pH and b) that winter time pH values are consistent with that expected from equilibrium with atmospheric CO<sub>2</sub> as recorded at the NIWA Baring Head atmospheric station (K. Hunter (U. Otago) & K. Currie (NIWA), pers comm.). The oscillations are primarily due to seasonal changes in water temperature and biological removal of dissolved carbon in the seawater. The time series sampling period is currently too short to distinguish long-term changes from seasonal and interannual variability, and it is not yet possible to attribute observed changes to uptake of anthropogenic carbon dioxide.





Figure 8.14: pH in Sub- Antarctic surface seawater on the R.V. *Munida* transect, 1998–2006. The blue points and joining lines are the actual measurements, and the red line a best fit to the points using a sine wave function (to represent seasonal change). The black line represents what pH assuming equilibrium with the atmosphere concentration in the Year 1750. The yellow line is the pH assuming equilibrium with actual CO<sub>2</sub> concentrations measured at the NIWA Baring Head Atmospheric Station. pH<sup>25</sup> is the pH measured at 25°C (Image Source: A Southern Hemisphere Time Series for CO<sub>2</sub> Chemistry and pH K. Hunter, K.C. Currie, M.R. Reid, H. Doyle. A presentation made at the International Union of Geodesy and Geophysics (IUGG) General Assembly Meeting, Melbourne June 2011.)

Globally, open ocean seawater pH shows relatively low spatial and temporal variability, compared to coastal waters where pH may vary by up to 1 unit in response to precipitation, biological activity in the seawater and sediment and other coastal processes. Surface pH in the open ocean has been determined on a monthly basis at the BATS (Bermuda Time Series Station) in the North Atlantic since 1983 (Bates 2001, 2007), and at HOT (Hawaii Time Series Station) in the North Pacific since 1988 (Brix *et al.* 2004, Dore *et al.* 2009). Both time series records show long term trends of increasing pCO<sub>2</sub> (partial pressure of CO<sub>2</sub>) and decreasing pH, with the pCO<sub>2</sub> increasing at a rate of 1.25 µatm per year, and pH decreasing by 0.0012 pH units per annum since 1983 at Bermuda (Figure 8.15). Placed in the context of these longer time series of atmospheric CO<sub>2</sub> measurements, the short record of the *Munida* SubAntarctic Water time series shows pCO<sub>2</sub> and pH in surface seawater tracking the atmospheric CO<sub>2</sub> (Figure 8.14). In addition, the regional means of seawater pH differ significantly with temperature, with the South Pacific at the lower end (Feely *et al.* 2009).



**Figure 8.15: Time series of atmospheric carbon dioxide at Moana Loa, seawater carbon dioxide and surface ocean pH at Ocean station ALOHA in the subtropical North Pacific Ocean near Hawaii. pH is shown as *in situ* pH, based on direct measurements and calculated from dissolved inorganic carbon and alkalinity in the surface layer (after Dore *et al.* 2009). (Image directly sourced from Feely *et al.* 2009 with permission.)**

Biological implications of ocean acidification result from increasing dissolved pCO<sub>2</sub>, increasing hydrogen ions (decreasing pH) and decreasing carbonate availability. The concern about ocean acidification is that the resulting reduction in carbonate availability may potentially impact organisms that produce shells or body structures of calcium carbonate, resulting in a redistribution of an organism's metabolic activity and increased physiological stress. Organisms most likely to be affected are those at the base of the food chain (bacteria, protozoa, plankton), coralline algae, rhodoliths, shallow and deepwater corals, echinoderms, molluscs, and possibly cephalopods (e.g., squids) and high-activity pelagic fish (e.g., tunas) (see Feely *et al.* 2004 and references therein; Orr *et al.* 2005, Langer *et al.* 2007). This is particularly of concern for deep-sea habitats such as seamounts, which can support structural reef-like habitat composed of stony corals (Tracey *et al.* 2011) as well as commercial fisheries for species such as orange roughy (Clark 1999). A shoaling carbonate saturation horizon could push such biogenic structures to the tops of seamounts, or cause widespread die-back (e.g., Thresher *et al.* 2012). This has important implications for the structure and function of benthic communities, and perhaps also for the deep-sea ecosystems that support New Zealand's key deepwater fisheries. Conversely some groups, including phytoplankton and sea-grass, may benefit from the increase in dissolved pCO<sub>2</sub> due to increased photosynthesis.

Direct effects of acidification on the physiology and development of fish have also been investigated. This has particularly focussed on the freshwater stages of salmonids (due to the widespread occurrence of pollution-derived acid rain) but increasingly in marine fish, where adverse effects on physiology development have been documented (e.g. Franke, and Clemmesen, 2011). Such studies highlight the potential for increasing acidification to impact larval growth and development, with implications for survival and recruitment of both forage fish and fish harvested commercially.

### 8.3. Ocean climate trends and New Zealand fisheries

This section has been quoted almost directly from the summary in Hurst *et al.* (2012). Some general observations on recent trends in some of the key ocean climate indices that have been found to be correlated with a variety of biological processes among fish (including recruitment fluctuations, growth, distribution, productivity and catch rates) are:

- The Interdecadal Pacific Oscillation (IPO): available from 1900; time scale 10–30 years. The IPO has been found to have been correlated with decadal changes ('regime shifts') in northeast Pacific ecosystems (e.g., Alaska salmon catches). In the New Zealand region, there is evidence of a regime shift into the negative phase of the IPO in about 2000. During the positive phase, from the late 1970's to 2000, New Zealand experienced periods of enhanced westerlies, with associated cooler air and sea temperatures and enhanced upwelling on western coasts. Opposite patterns are expected under a negative phase. For most New Zealand fisheries, monitoring of changes in populations began since the late 1970's, so there is little information on how New Zealand fishstocks might respond to these longer-term climatic fluctuations. Some of the recent changes in fish populations since the mid 1990s, for example, low western stock hoki recruitment indices (Francis 2009) and increases in some elasmobranch abundance indices (Dunn *et al.* 2009) may be shorter-term fluctuations that might be related in some way to regional warming during the period and only longer-term monitoring will establish whether they might be related to longer-term ecosystem changes.
- The Southern Oscillation Index: available from 1876; best represented as annual means. Causal relationships of correlations of SOI with fisheries processes are poorly understood but probably related in some way to one or more of the underlying ocean climate processes such as winds or temperatures. When the index is strongly negative, an El Niño event is taking place and New Zealand tends to experience increased westerly and south-westerly winds, cooler sea surface temperatures and enhanced upwelling in some areas (see, for example, the correlation of monthly SST at Leigh and SOI indices, Figure 8.13). Upwelling has been found to be related to increased nutrient flux and phytoplankton growth in areas such as the west coast South Island, Pelorus Sound and north-east coast of the North Island (Willis *et al.* 2007, Zeldis *et al.* 2008). El Niño events are likely to occur on 3–7 year time scales and are likely to be less frequent during the negative phase of the IPO which began in about 2000. This is likely to impact positively on species that show stronger recruitment under increased temperature regimes (e.g., snapper, Francis 1993, 1994a, b).
- Surface wind and pressure patterns: available from 1940s; variation in patterns can be high over monthly and annual time scales and many of the indices are correlated with each other, and with SOI and IPO indices (e.g., more zonal westerly winds, more frequent or regular cycles in southerlies in the positive IPO, 1977–2000). Correlations with biological process in fish stocks may occur over short time scales (e.g., impact on fish catchability) as well as seasonal and annual scales (e.g., impact on recruitment success). Wind and pressure patterns have been found to be correlated with fish abundance indices for southern gemfish (Renwick *et al.* 1998), hake, red cod and red gurnard (Dunn *et al.* 2008), rock lobster (Booth *et al.* 2000), and southern blue whiting (Willis *et al.* 2007, Hanchet & Renwick, 1999). Causal relationships of these correlations are poorly understood but can be factored into hypothesis testing as wind and pressure patterns affect surface ocean conditions through heat flux, upwelling and nutrient availability on exposed coasts.
- Temperature and sea surface height: available at least monthly over long time scales (air temperatures from 1906) or relatively short time scales (ocean temperatures to 800m, SST and SSH variously from 1987). Ocean temperatures, SST and SSH are all correlated with each other and smoothed air temperatures correlate well with SST in terms of interannual and seasonal variability; there are also some correlations of SST and SSH with surface wind and pressure patterns (see Dunn *et al.* 2009). SST has been found to be correlated with relative fish abundance

indices (derived from fisheries and/or trawl surveys) for elephantfish, southern gemfish, hoki, red cod, red gurnard, school shark, snapper, stargazer and tarakihi (Francis 1994a,b, Renwick et al 1998, Beentjes & Renwick 2001, Gilbert & Taylor 2001, Dunn et al 2009). Air temperatures in New Zealand have increased since 1900; most of the increase occurred since the mid 1940s. Increases from the late 1970s to 2000 may have been moderated by the positive phase of the IPO. Coastal SST records from 1954 (at Portobello) also show a slight increase through the series and, in general, show strong correlations with SOI (i.e., cooler temperatures in El Niño years). Other time series (SSH, ocean temperature to 800m) are comparatively short but show cycles of warmer and cooler periods on 1–6 year time scales. All air and ocean temperature series show the significant warming event during the late 1990s which has been followed by some cooling, but not to the levels of the early 1990s.

- Ocean colour and upwelling: these will be important time series because they potentially have a more direct link to biological processes in the ocean and are more easily incorporated into hypothesis testing. The ocean colour series starts in late 1997, so is not able to track changes that may have occurred since before the late 1990s warming cycle. These indices also need to be analysed with respect to SST, SSH and wind patterns, at similar locations or on similar spatial scales. The preliminary series developed exhibit some important spatial differences and trends that may warrant further investigation in relation to fish abundance indices. Of note are the increased chlorophyll indices off the west and south-west coast of the South Island in spring/summer during the last 5–6 years and the relatively low upwelling indices off the west coast South Island during winter in the late-1990s (Hurst *et al.* 2012).
- Currents: there are no general indices of trends or variability at present. Improvements in monitoring technology (e.g., satellite observations of SSH; CTD; ADCP; ARGO floats) have resulted in more information becoming available to enable numerical models of ocean currents to be developed. On the open ocean scale, there is considerable complexity in the New Zealand zone (e.g., frontal systems, eddy systems of the east coast). In the coastal zone, this is further complicated in coastal areas by the effects of tides, winds and freshwater (river) forcing, and a more limited monitoring capability. Nevertheless, the importance of current systems is starting to become more recognised and incorporated into analysis and modelling of fisheries processes and trends. Recent examples include the retention of rock lobster phyllosoma (mid-stage larvae) in eddy systems (Chiswell & Booth 2005, 2007), the apparent bounding of orange roughy nursery grounds by the presence of a cold-water front (Dunn *et al.* 2009) and the drift of toothfish eggs and larvae (Hanchet *et al.* 2008).
- Acidification: The increase in atmospheric CO<sub>2</sub> has been paralleled by an increase in CO<sub>2</sub> concentrations in the upper ocean, resulting in a decrease in pH. Maintenance of the one existing New Zealand monitoring programme for pH and pCO<sub>2</sub>, and development of new programmes to monitor the impacts of pH on key groups of organisms are critical. Potentially vulnerable groups include organisms that produce shells or body structures of calcium carbonate (corals, molluscs, plankton, coralline algae), and also non-calcifying groups including plankton, squid and high-activity pelagic fishes. Potentially positive impacts of acidification include increased phytoplankton carbon fixation and vertical export and increased productivity of sub-tropical waters due to enhanced nitrogen fixation by cyanobacteria. Secondary effects at the ecosystem level, such as productivity, biomass, community composition and biogeochemical feedbacks, also need to be considered.

Climate change was not specifically addressed as part of the report by Hurst *et al.* (2012), although indices described are an integral part of monitoring the speed and impacts of climate change. As noted under the air temperature section, the slightly increasing trend in temperatures since the mid 1940s is likely to have been moderated by the positive phase of the IPO, from the late 1970s to the late 1990s. With the shift to a negative phase of the IPO in 2000, it is likely that temperatures will increase more steeply. Continued monitoring of the ocean environment and response is critical. This includes not only the impacts on productivity, at all levels, but also on increasing ocean acidification.

For the New Zealand region, key ocean climate drivers in the last decade have been:

- the significant warming event in the late 1990s
- the regime shift to the negative phase of the IPO in about 2000, which is likely to result in fewer El Niño events for a 20–30 year period, i.e., less zonal westerly winds (already apparent compared to the 1980–2000 period) and increased temperatures; this is the first regime shift to occur since most of our fisheries monitoring time series have started (the previous shift was in the late 1970s), and
- global trends of increasing air and sea temperatures and ocean acidification.

## 8.4. References

- Bates, N.R. 2001. Interannual variability of oceanic CO<sub>2</sub> and biogeochemical properties in the Western North Atlantic subtropical gyre. *Deep-Sea Research II* 48:1507–1528.
- Bates, N.R. 2007. Interannual variability of the oceanic CO<sub>2</sub> sink in the subtropical gyre of the North Atlantic Ocean over the last 2 decades. *J. Geophysical Res.* 112, C09013, doi:10.1029/2006JC003759, 2007
- Beentjes, M.P., Renwick, J.A. 2001. The relationship between red cod, *Pseudophycis bachus*, recruitment and environmental variables in New Zealand. *Environmental Biology of Fishes* 61: 315–328.
- Bowen, M. M., Sutton, P.J.H., Roemmich, D. 2006. Wind-driven and steric fluctuations of sea surface height in the southwest Pacific, *Geophys. Res. Lett.*, 33, L14617, doi:10.1029/2006GL026160.
- Brix, H; Gruber, N., Keeling, C.D. 2004. Interannual variability of the upper ocean carbon cycle at station ALOHA near Hawaii. *Global Biogeochemical Cycles* 18(4) GB4019 Nov 24 2004.
- Caldeira, K., Wickett, M.E. 2003. Anthropogenic carbon and ocean pH. *Nature* 425:365.
- Chang, F.H., Mullan, B. 2003. Occurrence of major harmful algal blooms in New Zealand: is there a link with climate variation. *The Climate Update* 53: 4.
- Chiswell SM 1996. Variability in the Southland Current, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 30: 1-17.
- Chiswell, S.M; Booth, J.D. 2005. Distribution of mid- and late-stage *Jasus edwardsii* phyllosomas: implications for larval transport. *New Zealand Journal of Marine and Freshwater Research* 39. 1157–1170.
- Chiswell, S.M; Booth, J.D. 2007. Sources and sinks of larval settlement in *Jasus edwardsii* around New Zealand: Where do larvae come from and where do they go? *Marine Ecology Progress Series* 354: 201-217.
- Clark, M.R. (1999). Fisheries for orange roughy (*Hoplostethus atlanticus*) on seamounts in New Zealand. *Oceanologica Acta* 22(6): 593-602
- Dore, J.E., R. Lukas, D.W. Sadler, M.J. Church, and D.M. Karl. 2009. Physical and biogeochemical modulation of ocean acidification in the central North Pacific. *Proceedings of the National Academy of Sciences of the United States of America* 106(30):12,235–12,240.
- Dunn, M.R.; Hurst, R.; Renwick J.; Francis, R.I.C.C.; Devine, J.; McKenzie, A. 2009. Fish abundance and climate trends in New Zealand. *New Zealand Aquatic Environment and Biodiversity Report No. 31*. 75 p.
- Feely, R. A.; Sabine, Christopher L.; Lee, Kitack; Berelson, Will; Kleypas, Joanie; Fabry, Victoria J.; Millero, Frank J. (2004). "Impact of Anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> System in the Oceans" (abstract). *Science* 305 (5682): 362–366. doi:10.1126/science.1097329. PMID 15256664
- Feely, RA. Doney, SC. Cooley, SR. (2009). Ocean Acidification. Present conditions and future changes in a high CO<sub>2</sub> world. *Oceanography Volume* 22(4): 36-47.
- Francis M.P. 1993. Does water temperature determine year class strength in New Zealand snapper (*Pagrus auratus*, Sparidae)? *Fisheries Oceanography* 2(2): 65–72.
- Francis, M.P. 1994a. Growth of juvenile snapper, *Pagrus auratus* (Sparidae). *New Zealand Journal of Marine and Freshwater Research* 28: 201–218.
- Francis, M.P. 1994b. Duration of larval and spawning periods in *Pagrus auratus* (Sparidae) determined from otolith daily increments. *Environmental biology of Fishes* 39: 137–152.
- Franke, A. and Clemmesen, C. (2011). Effect of ocean acidification on early life stages of Atlantic herring (*Clupea harengus* L.). *Biogeosciences Discuss.*, 8, 7097– 7126.
- Gilbert D.J., Taylor, P.R. 2001. The relationship between snapper (*Pagrus auratus*) year class strength and temperature for SNA2 and SNA7. *New Zealand Fisheries Assessment report* 2001/64. 33p.
- Gordon DP, Beaumont J, MacDiarmid A, Robertson DA, Ahoyong ST (2010) Marine Biodiversity of Aotearoa New Zealand. *PLoS ONE* 5(8): e10905. doi:10.1371/journal.pone.0010905
- Hanchet, S. M., Renwick, J. A. 1999: Prediction of year class strength in southern blue whiting (*Micromesistius australis*) in New Zealand waters. *New Zealand Fisheries Assessment Research Document* 99/51. 24 p.
- Hanchet, S.M., Fenaughty, J.M., Dunn, A., Williams, M.J.H. 2008. A hypothetical life cycle for Antarctic toothfish (*Dissostichus mawsoni*) in the Ross Sea region *CCAMLR Science* 15: 35–53.
- Heath RA 1985. A review of the physical oceanography of the seas around New Zealand.1982. *New Zealand Journal of Marine and Freshwater Research* 19: 79.124.
- Hobday, A.J., Okey, T.A., Poloczanska, E.S., Kunz, T.J., Richardson, A.J. 2006. Impacts of Climate Change on Australian Marine Life. CSIRO Marine and Atmospheric Research report to the Department of the Environment and Heritage
- Houghton J.T., Ding, Y., Griggs, D.J., Noguer, M. van der Linden, P.J., Dai, X., Maskell, K., Johnson, C.A. 2001. Climate change 2001: the scientific basis. Contribution of Working Group I to the third Assessment. Report of the International Panel on Climate Change. Cambridge University Press, Cambridge, UK.
- Hurst, R.J. Renwick, J.A. Sutton, P.J.H Uddstrom, M.J. Kennan, S.C. Law, C.S. Rickard, G.J. Korpela, A. Stewart, C. Evans J. (2012In press). Climate and oceanographic trends of potential relevance to fisheries in the New Zealand fisheriesRegion, 2010. *New Zealand Aquatic Environment and Biodiversity Report No. 90XX*. 202p.
- IPCC Third Assessment Report "Climate Change 2001".
- Langer, G., M. Geisen, K.-H. Baumann, J. Kläs, U. Riebesell, S. Thoms, Young, J. R. 2006. Species-specific responses of calcifying algae to changing seawater carbonate chemistry, *Geochim. Geophys. Geosyst.*, 7 : Q09006, doi:10.1029/2005GC001227

- Lutz MJ, Caldeira K, Dunbar RB, Behrenfeld MJ (2007) Seasonal rhythms of net primary production and particulate organic carbon flux to depth describe the efficiency of biological pump in the global ocean. *J Geophys Res Oceans* 112:C10011 doi: 10.1029/2006JC003706
- Mullan, A. B., 1995. On the linearity and stability of Southern Oscillation-climate relationships for New Zealand. *International Journal of Climatology* 15: 1365–1386.
- Mullan, A.B; Stuart, S.J; Hadfield, M.G; Smith, M.J (2010). Report on the Review of NIWA's 'Seven-Station' Temperature Series NIWA Information Series No. 78. 175 p.
- O'Driscoll, R.L.; Hurst, R.J.; Dunn, M.R.; Gauthier, S.; Ballara, S.L. (2011). Trends in relative mesopelagic biomass using time series of acoustic backscatter data from trawl surveys. *New Zealand Aquatic Environment and Biodiversity Report* 2011/76.
- Orr, J.C., Fabry, V.J., Aumont, O., Bopp, L., Doney, S.C., Feely, R.A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R.M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R.G., Plattner, G-K., Rodgers, K.B., Sabine, C.L., Sarmiento, J.L., Schlitzer, R., Slater, R.D., Totterdell, I.J., Weirig, M-F., Yamanaka, Y., Yool, A. 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437: 681–686. *Global Biogeochemical Cycles* 21: GB2028, doi:10.1029/2006GB002898, 2007
- Philip W. Boyd & Cliff S. Law (2011). An Ocean Climate Change Atlas for New Zealand waters A primer for a major new web-based tool to help predict how oceanic species will be affected by climate change NIWA Information Series No. 79 ISSN 1174-264X
- Pinkerton, M.H., Richardson, K.M., Boyd, P.W., Gall, M.P., Zeldis, J., Oliver, M.D., Murphy, R.J. 2005. Intercomparison of ocean colour band-ratio algorithms for chlorophyll concentration in the Subtropical Front east of New Zealand. *Remote Sensing of Environment* 97: 382–402.
- Reid JL 1986. On the total geostrophic circulation of the South Pacific Ocean: flow patterns, tracers and transports. *Progress in Oceanography* 16: 1-61.
- Renwick, J.A., Hurst, R.J., Kidson, J.W. 1998: Climatic influences on the recruitment of Southern Gemfish (*Rexea solandri*, Gempylidae) in New Zealand waters. *International Journal of Climatology* 18: 1655–1667.
- Reynolds, R.W., Smith, T.M.; Liu, C.; Chelton, D.B.; Casey, K.S.; Schlax, M.G. (2007). Daily High-resolution Blended Analyses for sea surface temperature. *Journal of Climate*, 20, 5473-5496.
- Ridgway KR, Dunn JR 2003. Mesoscale structure of the mean East Australian Current system and its relationship with topography. *Progress in Oceanography* 56: 189-222.
- Roemmich, D., Gilson, J., Davis, R., Sutton, P., Wijffels, S., Riser, S. 2007. Decadal Spin-up of the South Pacific Subtropical Gyre. *Journal of Physical Oceanography* 37: 162–173.
- Sabine, C.L., Feely, R.A., Gruber, N., Key, R.M., Lee, K., Bullister, J.L., Wanninkhof, R., Wong, C.S., Wallace, D.W., Tilbrook, B., Millero, F.J., Peng, T.H., Kozyr, A., Ono, T., Rios, A.F. 2004. The oceanic sink for anthropogenic CO<sub>2</sub>. *Science* 305:367–371.
- Shirtcliffe, G.G.L., Moore, M.I., Cole, A.G., Viner, A.B., Baldwin, R. Chapman, B. 1990. Dynamics of the Cape Farewell upwelling plume, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 24: 555–568.
- Stanton BR 1981. An oceanographic survey of the Tasman Front. *New Zealand Journal of Marine and Freshwater Research* 15: 289-297.
- Stanton BR, Ridgway NM 1988. An oceanographic survey of the subtropical convergence zone in the Tasman Sea. *New Zealand Journal of Marine and Freshwater Research* 22: 583-593.
- Stanton BR, Sutton PJH, Chiswell SM 1997. The East Auckland Current, 1994-95. *New Zealand Journal of Marine and Freshwater Research* 31: 537-549.
- Sutton PJH 2003. The Southland Current: a subantarctic current. *New Zealand Journal of Marine and Freshwater Research* 37: 645–652.
- Sutton, P., Bowen, M., Roemmich, D. 2005. Decadal Temperature changes in the Tasman Sea. *N.Z. Journal of Marine and Freshwater Research* 39(6): 1321–1329.
- Sutton, P.J.H.; Roemmich, D.; 2001. Ocean temperature climate off north-east New Zealand. *N.Z. Journal of Marine and Freshwater Research* 35: 553–565.
- Sutton, R.T.; Dong, B.; Gregory, J.M. (2007). Land/sea warming ratio in response to climate change: IPCC AR4 model results and comparison with observations. *Geophysical Research Letters*, 34, L02701, doi:10.1029/2006GL028164.
- Thompson, D.W.J.; Solomon, S. (2002). Interpretation of recent Southern Hemisphere climate change. *Science*, 296, 895-899.
- Thompson, P.A.; Baird, M.E.; Ingleton, T.; Doblin, M.A. (2009). Long-term changes in temperature Australian waters: implications for phytoplankton. *Marine Ecology Progress Series*, 394, 1-19.
- Thresher, R.E., Guinotte, J., Matear, R., Fallon S. (2012). Adapting to the effects of climate change on Australia's deep marine reserves. Fisheries Research and Development Corporation (FRDC)/National Climate Change Adaptation Research Fund (NCCARF) Report 2010/510, 68 p.
- Tracey, D.M., Rowden, A.A., Mackay, K.A., Compton, T (2011) Habitat-forming cold-water corals show affinity for seamounts in the New Zealand region. *Mar Ecol Prog Ser* 430: 1–22 Francis R.I.C.C. 2009. Assessment of hoki (*Macruronus novaezelandiae*) in 2008. *New Zealand Fisheries Assessment Report* 2009/7. 80p.
- Willis, T.J., Handley, S.J., Chang, F.H., Morrissey, D.J., Mullan, B., Pinkerton, M., Rodgers, K.L., Sutton, P.H.J., Tait, A. 2007. Climate change and the New Zealand marine environment. NIWA Client Report for the Department of Conservation: NEL2007-025 October 2007, NIWA Project: DOC08305
- Zeldis, J.R, Howard-Williams, C, Carter, C.M., Schiel, D.R. 2008. ENSO and riverine control of nutrient loading, phytoplankton biomass and mussel aquaculture yield in Pelorus Sound, New Zealand. <http://www.int-res.com/articles/meps2008/371/m371p131.pdf>
- Zeldis, J.R. 2004. New and remineralised nutrient supply and ecosystem metabolism on the northeastern New Zealand continental shelf. *Continental Shelf Research* 24: 563–581.

## 9. Habitats of particular significance for fisheries management

<i>Scope of chapter</i>	This chapter highlights subject areas that might contribute to the management of HPSFM and hence provides a guide for future research in the absence of an approved policy definition of HPSFM
<i>Area</i>	All of the New Zealand EEZ and territorial sea (inclusive of the freshwater and estuarine areas).
<i>Locality hotspots</i>	None formally defined, but already identified likely candidates include areas of biogenic habitat, e.g. Separation Point and Wairoa Hard, and areas identified with large catches and/or vulnerable populations of juveniles, e.g. Hoki Management Areas, packhorse crayfish legislated closures and toheroa beaches.
<i>Key issues</i>	Defining and identifying likely HPSFM and potential threats to them.
<i>Emerging issues</i>	Connectivity and intra-population behaviour variability, multiple use
<i>MPI Research (current)</i>	Biogenic habitats as areas of particular significance for fisheries management (HAB2007/01), Toheroa abundance (TOH2007/03), Research on Biogenic Habitat-Forming Biota and their functional role in maintaining Biodiversity in the Inshore Region (5-150M Depths) (ZBD2008/01 – this is also part-funded by Oceans Survey 2020, NIWA and MBIE) , Habitats of particular significance for fisheries management: Kaipara Harbour (ENV2009/07), Habitats of particular significance for inshorefinfish fisheries management (ENV2010/03) Spatial Mixing of GMU1 using Otolith Microchemistry (GMU2009/01).
<i>NZ Government Research (current)</i>	Ministry of Business, Innovation and Employment (MBIE) funded programmes (Coastal Conservation Management: protecting the functions of marine coastal habitats that support fish assemblages at local, regional and national scales (C01X0907) Predicting the occurrence of vulnerable marine ecosystems for planning spatial management in the South Pacific region (C01X1229) and Impacts of resource use on vulnerable deep-sea communities (C01X0906).  NIWA Core funding in the 'Managing marine stressors' area under the 'Coasts and Oceans' centre, specifically the programme 'Managing marine resources' and the project 'Measuring mapping and conserving (C01X0505)'
<i>Links to 2030 objectives</i>	Under the Environment Outcome habitats of special significance to fisheries need to be protected.
<i>Related chapters/issues</i>	Land-based impacts on fisheries and supporting biodiversity, bycatch composition, marine environmental monitoring.

### 9.1. Context

The Fisheries Act 1996, in Section 9 (Environmental principles) states that:

*“All persons exercising or performing functions, duties, or powers under this Act, in relation to the utilisation of fisheries resources or ensuring sustainability, shall take into account the following environmental principles:*

- (a) Associated or dependent species should be maintained above a level that ensures their long-term viability;*
- (b) Biological diversity of the aquatic environment should be maintained;*

***(c) Habitat of particular significance for fisheries management should be protected.”***

No policy definition of habitat of particular significance for fisheries management (HPSFM) exists, although work is currently underway to generate one. Some guidance in terms of defining HPSFM is provided by Fisheries 2030 which specifies as an objective under the Environment Outcome that “habitats of special significance to fisheries are protected”. This wording suggests that a specific focus on habitats that are important for fisheries production should be taken rather than a more general focus that might also include other habitats that may be affected by fishing.

Fisheries 2030 re-emphasises that HPSFM should be protected. No specific strategic actions are proposed to implement this protection in Fisheries 2030; although action 6.1 “To implement a revised MPA policy and legal framework” could potentially be relevant to protecting HPSFM. The management of activities other than fishing, such as land-use and vehicle traffic, are outside the control of the Ministry for Primary Industries but Fisheries 2030 specifies actions to “Improve fisheries sector input to processes that manage RMA-controlled effects on the marine and freshwater environment” (Action 8.1) and to “Promote the development and use of RMA national policy statements, environmental standards, and regional coastal and freshwater plans” (Action 8.2). This suggests that the cooperation of other parties outside of the fisheries sector may be necessary in some cases to protect HPSFM.

In the absence of a policy definition of HPSFM this chapter will focus on examples of habitats shown to be important for fisheries and concepts likely to be important to HPSFM. Examples of potential HPSFM include: sources of larvae; larval settlement sites; habitat for juveniles; habitat that supports important prey species; migration corridors; and spawning, pupping or egg-laying grounds. Some of these habitats may be important for only part of the life cycle of an organism, or for part of a year.

The location or relative importance of habitats, compared with other limiting factors, is largely unknown for most stocks. For example, some stocks may be primarily habitat limited, whereas others may be limited by oceanographic variability, food supply, predation rates (especially during juvenile phases), or a mixture of these and other factors. In the case of stocks that are habitat limited, a management goal might be to preserve or improve some aspect of the habitat for the stock.

Hundreds of legislated spatial fisheries restrictions already apply within New Zealand’s territorial sea and exclusive economic zone ([www.nablis.govt.nz](http://www.nablis.govt.nz)), but until further policy work and research is conducted we cannot be sure the contribution they make to protecting HPSFM. Examples of these are listed below:

- Separation Point in Tasman Bay, and the Wairoa Hard in Hawke Bay, were created to protect biogenic habitat which was believed to be important as juvenile habitat for a variety of fish species (Grange *et al.* 2003).
- An area near North Cape is currently closed to packhorse lobster fishing to mitigate sub-legal handling disturbance in this area. This closure was established because of the small size of lobsters caught there and a tagging study that showed movement away from this area into nearby fished areas (Booth 1979).
- The largest legislated closure are the Benthic Protection Areas (BPAs) which protect ~ 1.2 million square km (about 31% of the EEZ) outside the territorial sea from contact of trawl and dredge gear with the bottom (Helson *et al.* 2010).
- Commercial fishers must not use New Zealand fishing vessels or foreign-owned New Zealand fishing vessels over 46 m in overall length for trawling in the territorial sea.

In addition to legislated closures, a number of non-regulatory management measures exist. For example:

- Spatial closures
  - Trawlers greater than 28 m in length are excluded from targeting hoki in four Hoki Management Areas – Cook Strait, Canterbury Banks, Mernoo Bank, and Puysegur Bank (DeepWater Group 2008). These areas were chosen because of the larger number of juveniles caught, relative to adults in these areas.
  - Trawling and pair trawling are both closed around Kapiti Island
- Seasonal closures
  - A closure to trawling exists from November the first until the 30<sup>th</sup> of April each year in Tasman Bay.
  - A closure to commercial potting exists for all of CRA3 for the whole of the month of December each year.

The high-level objectives and actions in Fisheries 2030 have been interpreted in the highly migratory, deepwater and middle-depths (deepwater) inshore national fish plans. The highly migratory fish plan addresses HPSFM in environment outcome 8.1 “Identify and where appropriate protect habitats of particular significance to highly migratory species, especially within New Zealand waters”. In the deepwater fish plan the Ministry proposes in management objective 2.3 “to develop policy guidelines to determine what constitutes HPSFM then apply these policy guidelines to fisheries where necessary”. Inshore fisheries management plans (freshwater, shellfish and finfish) all contain references to identifying and managing HPSFM. These plans recognise that not all impacts stem from fisheries activities, therefore managing them may include trying to influence others to better manage their impacts on HPSFM. Work is underway on a policy definition of HPSFM that will assist implementing these outcomes and objectives.

## **9.2. Global understanding**

This section focuses upon those habitats protected overseas for their value to fisheries and discusses important concepts that may help gauge the importance of any particular habitat to fisheries management. This information may guide future research into HPSFM in New Zealand and any subsequent management action.

### **9.2.1. Habitats protected elsewhere for fisheries management**

Certain habitats have been identified as important for marine species: shallow sea grass meadows, wetlands, seaweed beds, rivers, estuaries, rhodolith beds, rocky reefs, crevices, boulders, bryozoans, submarine canyons, seamounts, coral reefs, shell beds and shallow bays or inlets (Kamenos *et al.* 2004; Caddy 2008, Clark 1999, Morato *et al.* 2010). Discrete habitats (or parts of these) may have extremely important ecological functions, and/or be especially vulnerable to degradation. For example, seabeds with high roughness are important for many fisheries and can be easily damaged by interaction with fishing gear (Caddy 2008). Examples of these include:

1. The *Oculina* coral banks off Florida were protected in 1994 as an experimental reserve in response to their perceived importance for reef fish populations (Rosenberg *et al.* 2000). Later studies confirmed that this area is the only spawning aggregation site for gag (*Mycteroperca microlepis*) and scamp (*M. phenax*) (both grouper species), and other economically important reef fish in that region (Koenig *et al.* 2000). The size of the area within which bottom-tending gears were restricted was subsequently increased based on these findings (Rosenberg *et al.* 2000).

2. *Lophelia* cold-water coral reefs are now protected in at least Norway (Fosså *et al.* 2002), Sweden (Lundälv, & Jonsson 2003) and the United Kingdom (European Commission 2003) due to their importance as habitat for many species of fish (Costello *et al.* 2005).
3. The Western Pacific Regional Fishery Management Council identified all escarpments between 40 m and 280 m as Habitat Areas of Particular Concern (HAPC) for species in the bottom-fish assemblage. The water column to a depth of 1000 m above all shallow seamounts and banks was categorised as HAPC for pelagic species. Certain northwest Hawaiian Island banks shallower than 30 m were categorised as HAPC for crustaceans, and certain Hawaiian Island banks shallower than 30 m were classified as Essential Fish Habitat (EFH) for precious corals. Fishing is closely regulated in the precious-coral EFH, and harvest is only allowed with highly selective gear types which limit impacts, such as manned and unmanned submersibles (West Pacific Fisheries Management Council 1998)

Examples of habitats protected for their freshwater fishery values also exist. For example, the U.S. Atlantic States Interstate fishery management plan (Atlantic States Marine Fisheries Commission 2000) notes the Sargasso Sea is important for spawning, and that seaweed harvesting provides a threat of unknown magnitude to eel spawning. Habitat alteration and destruction are also listed as probably impacting on continental shelves and estuaries/rivers, respectively, but the extent to which these are important is unknown.

It is also possible that HPSFM may be defined by the functional importance of an area to the fishery. For example, large spawning aggregations can happen in mid-water for set periods of time (Schumacher and Kendall 1991, Livingston 1990) these could also potentially qualify as HPSFM.

### 9.2.2. Concepts potentially important for HPSFM

Many nations are now moving towards formalised habitat classifications for their coastal and ocean waters, which may include fish dynamics as part of their structure, and could potentially help to define HPSFM. Such systems help provide formal definitions for management purposes, and to 'rank' habitats in terms of their relative values and vulnerability to threats. Examples include the Essential Fish Habitat (EFH) framework being advanced in North America (Benaka 1999, Diaz *et al.* 2004, Valavanis *et al.* 2008), and in terms of habitat, the developing NOAA Coastal and Marine Ecological Classification Standard for North America (CMECS) (Madden *et al.* 2005, Keefer *et al.* 2008), and the European Marine Life Information Network (MarLIN) framework which has developed habitat classification and sensitivity definitions and rankings (Hiscock and Tyler-Walters 2006).

Habitat connectivity (the movement of species between habitats) operates across a range of spatial scales, and is a rapidly developing area in the understanding of fisheries stocks. These movements link together different habitats into 'habitat chains', which may also include 'habitat bottlenecks', where one or more spatially restricted habitats may act to constrain overall fish production (Werner *et al.* 1984). Human driven degradation or loss of such bottleneck habitats may strongly reduce the overall productivity of populations, and hence ultimately reduce long-term sustainable fisheries yields. The most widely studied of these links is between juvenile nursery habitats and often spatially distant adult population areas. Most studies published have been focussed on species that uses estuaries as juveniles; e.g. blue grouper *Achoerodus viridis* (a large wrasse) (Gillanders and Kingsford 1996) and snapper *Pagrus auratus* (Hamer *et al.* 2005) in Australia; and gag (*Mycteroperca Microlepis*) in the United States (Ross and Moser 1995) which make unidirectional ontogenetic habitat shifts from estuaries and bays out to the open coast as they grow from juveniles to adults. The extent of wetland habitats in the Gulf of Mexico has also been linked to the yield of fishery species dependent on coastal bays and estuaries. Reduced fishery stock production (shrimp and menhaden (a fish)) followed wetland losses and, conversely, stock gains followed increases in the area of wetlands (Turner and Boesch 1987). Juvenile production was limited by the amount of available habitat but, equally, reproduction, larval settlement, juvenile or adult survivorship, or other demographic factors

could also be limited by habitat loss or degradation, and these could have knock-on effects to stock characteristics such as productivity and its variability. Other examples include movements which may be bidirectional and regular in nature e.g., seasonal migrations of adult fish to and from spawning and/or feeding grounds, e.g. grey mullet *Mugil cephalus* off Taiwan (Chang *et al.* 2004).

How habitats are spatially configured to each other is also important to fish usage and associated fisheries production. For example, Nagelkerken *et al.* (2001) showed that the presence of mangroves in tropical systems significantly increases species richness and abundance of fish assemblages in adjacent seagrass beds. Jelbart *et al.* (2007) sampled Australian temperate seagrass beds close to (< 200 m) and distant from (> 500 m) mangroves. They found seagrass beds closer to mangroves had greater fish densities and diversities than more distant beds, especially for juveniles. Conversely, the densities of fish species in seagrass at low tide that were also found in mangroves at high tide were negatively correlated with the distance of the seagrass bed from the mangroves. This shows the important daily habitat connectivity that exists through tidal movements between mangrove and seagrass habitats. Similar dynamics may occur in more sub-tidal coastal systems at larger spatial and temporal scales. For example, Dorenbosch *et al.* (2005) showed that adult densities of coral reef fish, whose juvenile phases were found in mangrove and seagrass nursery habitats, were much reduced or absent on coral reefs located far distant from such nursery habitats, relative to those in closer proximity.

A less studied, but increasingly recognised theme is the existence of intra-population variability in movement and other behavioural traits. Different behavioural phenotypes within a given population have been shown to be very common in land birds, insects, mammals, and other groups. An example of this is a phenomenon known as ‘partial migration’, where part of the overall population migrates each year, often over very large distances, while another component does not move and remains resident. By definition, this partial migration also results in differential use of habitats, often over large spatial scales. Recent work on white perch (*Morone americana*) in the United States shows this population is made up of two behavioural components: a resident natal freshwater contingent; and a dispersive brackish-water contingent (Kerr *et al.* 2010). The divergence appears to be a response to early life history experiences which influence individuals’ growth (Kerr 2008). The proportion of the overall population that becomes dispersive for a given year class ranges from 0% in drought years to 96% in high-flow years. Modelling of how differences in growth rates and recruitment strengths of each component contributed to the overall population found that the resident component contributed to long-term population persistence (stability), whereas the dispersive component contributed to population productivity and resilience (defined as rebuilding capacity) (Kerr *et al.* 2010). Another species winter flounder *Pseudopleuronectes americanus* has also shown intra-population variability in spawning migrations; one group stays coastally resident while a second smaller group migrate into estuaries to spawn (DeCelles & Cadrin 2010). The authors went on to suggest that coastal waters in the Gulf of Maine should merit consideration in the assignment of Essential Fish Habitat for this species.

Kerr and Secor (2009) and Kerr *et al.* (2010) argue that such phenotypic dynamics are probably very common in marine fish populations but have not yet been effectively researched and quantified. The existence of such dynamics would have important implications for fisheries management, including the possibility of spatial depletions of more resident forms and variability in the use of potential HPSFM between years. For instance, recent work on snapper in the Hauraki Gulf has shown that fish on reef habitats are more resident (ie have less propensity to migrate) than those of soft sediment habitats, and can experience higher fishing removals (Parsons *et al.* 2011).

The most effective means of protecting a HPSFM in terms of the benefit to the fishery may differ depending on the life-history characteristics of the fish. A variety of modelling, theoretical, and observational approaches have lead to the conclusion that spatial protection performs best at

enhancing species whose adults are relatively sedentary but whose larvae are broadcast widely (Chiappone and Sealey 2000, Murawski *et al.* 2000, Roberts 2000, Warner *et al.* 2000). The sedentary habit of adults allows the stock to accrue the maximum benefit from the protection, whereas the broadcasting of larvae helps ‘seed’ segments of the population outside the protection. However, the role of spatial protection in directly protecting juveniles after they have settled to seafloor habitats (via habitat protection/recovery, and/or reduced juvenile bycatch), or their interaction with non-fisheries impacts has not yet been explicitly considered.

### **9.3. State of knowledge in New Zealand**

#### **9.3.1. Potential HPSFM in New Zealand**

Important areas for spawning, pupping, and egg-laying are potential HPSFM. These areas (insofar as these are known) have been identified and described using science literature and fisheries databases and summarised within two atlases, one coastal (< 200 m) and one deepwater (> 200 m). Coastally, these HPSFM areas were identified for 35 important fish species by Hurst *et al.* (2000). This report concluded that virtually all coastal areas were important for these functions for one species or other. The report also noted that some coastal species use deeper areas for these functions, either as juveniles, or to spawn (e.g., red cod, giant stargazer) and some coastal areas are important for juveniles of deeper spawning species (e.g., hake and ling). Some species groupings were apparent from this analysis. Elephant fish, rig, and school shark all preferred to pup or lay eggs in shallow water, and very young juveniles of these species were found in shallow coastal areas. Juvenile barracouta, jack mackerel (*Trachurus novaezelandiae*), kahawai, rig, and snapper were all relatively abundant (at least occasionally) in the inner Hauraki Gulf. Important areas for spawning, pupping, and egg-laying were identified for 32 important deepwater fish species (200 to 1500 m depth), 4 pelagic fish species, 45 invertebrate groups, and 5 seaweeds (O'Driscoll *et al.* 2003). This study concluded that all areas to 1500 m deep were important for either spawning or juveniles of one or more species studied. The relative significance of areas was hard to gauge because of the variability in the data, however the Chatham Rise was identified as a “hotspot”.

Areas of high juvenile abundances of certain species may be useful indicators of HPSFM for some species. A third atlas (Hurst *et al.* 2000b) details species distributions (mainly commercial) of adult and immature stages from trawl, midwater trawl and tuna longline where adequate size information was collected. No conclusions are made in this document, and generalisations across species are inherently difficult, therefore like the previous two atlases, this document is probably best examined for potential HPSFM in a species specific way.

Certain locations within New Zealand already seem likely to qualify as HPSFM under any likely definition. The Kaipara Harbour has been identified as particularly important for the SNA 8 stock. Analysis of otolith chemistry showed that, for the 2003 year-class, a very high proportion of new snapper recruits to the SNA 8 stock were sourced as juveniles from the Kaipara Harbour (Morrison *et al.* 2008). This result is likely to be broadly applicable into the future as the Kaipara provides most of the biogenic habitat available for juvenile snapper on this coast. The Kaipara and Raglan harbours also showed large catches of juvenile rig and the Waitemata, Tamaki and Porirua harbours moderate catches (Francis *et al.* 2012). Recent extensive fish-habitat sampling within the harbour in 2010 as part of the MBIE Coastal Conservation Management programme showed juvenile snapper to be strongly associated with sub-tidal seagrass, horse mussels, sponges, and an introduced bryozoan. Negative impacts on such habitats have the potential to have far-field effects in terms of subsequent fisheries yields from coastal locations well distant from the Kaipara Harbour. Beaches that still retain substantive toheroa populations, e.g. Dargaville and Oreti beaches, may also potentially qualify as HPSFM (Beentjes 2010).

Consistent with the international literature, biogenic (living, habitat forming) habitats have been found to be particularly important juvenile habitat for some coastal fish species in New Zealand. For example: bryozoan mounds in Tasman Bay are known nursery grounds for snapper, tarakihi and john dory (Vooren 1975); northern subtidal seagrass meadows fulfil the same role for a range of fish including snapper, trevally, parore, garfish and spotties (Francis *et al.* 2005, Morrison *et al.* 2008, Schwarz *et al.* 2006, Vooren 1975); northern horse mussel beds for snapper and trevally (Morrison *et al.* 2009); and mangrove forests for grey mullet, short-finned eels, and parore (Morrisey *et al.* 2010). Many other types of biogenic habitats exist, and some of their locations are known (e.g. see Davidson *et al.* 2010 for biogenic habitats in the Marlborough Sounds), but their precise role as HPSFM remains to be quantified. Examples include open coast bryozoan fields, rhodoliths, polychaete (worm) species ranging in collective form from low swathes to large high mounds, sea pens and sea whips, sponges, hydroids, gorgonians, and many forms of algae, ranging from low benthic forms such as *Caulerpa* spp. (sea rimu) through to giant kelp (*Macrocystis pyrifera*) forests in cooler southern waters. Similarly, seamounts are well-known to host reef-like formations of deep-sea stony corals (e.g., Tracey *et al.* 2011), as well as being major spawning or feeding areas for commercial deepwater species such as orange roughy and oreos (e.g., Clark 1999, O'Driscoll & Clark 2005). However, the role of these benthic communities on seamounts in supporting fish stocks is uncertain, as spawning aggregations continue to form even if the coral habitat is removed by trawling (Clark & Dunn 2012). Hence the oceanography or physical characteristics of the seamount and water column may be the key drivers of spawning or early life-history stage development, rather than the biogenic habitat.

Freshwater eels are reliant upon rivers as well as coastal and oceanic environments. GIS modelling estimates that for longfin eels, about 30% of longfin habitat in the North Island and 34% in the South Island is either in a reserve or in rarely/non-fished areas, with ~ 49% of the national longfin stock estimate of about 12 000 tonnes being contained in these waterways (Graynoth *et al.* 2008). More regional examination of the situation for eels also exists, e.g., for the Waikato Catchment (Allen 2010). Shortfin eels prefer slower-flowing coastal habitats such as lagoons, estuaries, and lower reaches of rivers (Beentjes *et al.* 2005). In-stream cover (such as logs and debris) has been identified as important habitat, particularly in terms of influencing the survival of large juvenile eels (Graynoth *et al.* 2008). Short-fin eel juveniles and adults have also been found to be relatively common in estuarine mangrove forests, and their abundance positively correlated with structural complexity (seedlings, saplings, and tree densities) (Morrisey *et al.* 2010). In addition oceanic spawning locations are clearly important for eels, the location of these are unknown, although it has been suggested that these may be northeast of Samoa and east of Tonga for shortfins and longfins respectively (Jellyman 1994).

Many of the potential HPSFM are threatened by either fisheries or land-based effects, the reader should look to the land-based effects chapter in this document and the eel section of the Stock assessment plenary report for further details.

### **9.3.2. Habitat classification and prediction of biological characteristics**

Habitat classification schemes focused upon biodiversity protection have been developed in New Zealand at both national and regional scales, these may help identify larger habitats which HPSFM may be selected from, but are unlikely to be useful in isolation for determining HPSFM. The Marine Environment Classification (MEC), the demersal fish MEC and the benthic optimised MEC (BOMECE) are national scale classification schemes have been developed with the goal of aiding biodiversity protection (Leathwick *et al.* 2004, 2006, 2012). A classification scheme also exists for New Zealand's rivers and streams based on their biodiversity values to support the Department of Conservation Waters of National Importance (WONI) project (Leathwick and Julian 2008). Regional

classification schemes also exist such as ones mapping the Marine habitats of Northland, or Canterbury in order to assist in Marine Protected Area planning (Benn 2009; Kerr 2010).

Another tool which may help in terms of identifying HPSFM is the predictions of richness, occurrence and abundance of small fish in New Zealand estuaries (Francis *et al.* 2011). This paper contains richness predictions for 380 estuaries and occurrence predictions for 16 species. This could help minimise the need to undertake expensive field surveys to inform resource management, although environmental sampling may still be needed to drive some models.

### **9.3.3. Current research**

Prior to 2007 research within New Zealand has not been explicitly focused on identifying HPSFM. However, in line with international trends, this situation has changed in recent times, with recognition of some of the wider aspects of fisheries management and the move towards an ecosystem approach foreshadowed in Fisheries 2030.

A number of Ministry and other research projects are underway, or planned, concerning HPSFM in the 2010/11 year. Project ENV200907, “Habitat of particular significance to fisheries management: Kaipara Harbour”, is underway and has the overall objective of identifying and mapping areas and habitats of particular significance in the Kaipara Harbour which support coastal fisheries; and identifying and assessing threats to these habitats. Included in this work is the reconstruction of environmental histories through interviews of long time local residents who have experience of the harbour, and associated collation and integration of historical data sources (e.g., catch records, photographs, diaries, maps, and fishing logs). Another output of this work will be recommendations on the best habitats and methods of monitoring to detect change to HPSFM within Kaipara harbour.

Biogenic habitats on the continental shelf from ~5 to 150 m depths are currently being characterised and mapped through the biodiversity project ZBD2008/01, this will also provide new information on fisheries species utilisation of these habitats. Interviews with 50 retired fishers have provided valuable information on biogenic habitat around New Zealand. A national survey to examine the present occurrences and extents of these biogenic habitats was completed in 2011 in collaboration with Oceans Survey 2020, NIWA and Ministry of Business, Innovation and Employment (MBIE) funding.

A number of other national scale projects are also underway. A desktop review is collating information on the importance of biogenic habitats to fisheries across the entire Territorial Sea and Exclusive Economic Zone (project HAB2007/01). A project has been approved to review the literature and recommend the relative urgency of research on habitats of particular significance for inshore finfish species (project ENV2010/03).

The Ministry of Business, Innovation and Employment (MBIE) funded project Coastal Conservation Management started in 2009 and runs for six years. This programme aims to integrate and add to existing fish-habitat association work to develop a national scale marine fish-habitat classification and predictive model framework. This project will also attempt to develop threat assessments at local, regional and national scales. MPI is maximising the synergies between its planned research and this project. As part of that synergy, work on the connectivity and stock structure of grey mullet (*Mugil cephalus*) is underway in collaboration with MFish project GMU2009/01. Otolith chemistry is being assessed for its utility in partitioning the GMU 1 stock into more biologically meaningful management units, and in quantifying the suspected existence of source and sink dynamics between the various estuaries that hold juvenile grey mullet nursery habitats.

MBIE also funded in 2012 the three year project delivered by NIWA entitled Predicting the occurrence of vulnerable marine ecosystems for planning spatial management in the South Pacific region. The development of predictive models of species occurrence under this project may also aid in

identifying HPSFM. Identification of biogenic habitat has been part of the MBIE project “Vulnerable deep-sea communities” since 2009 (and its predecessor seamount programme) which includes surveys of a range of habitats that may be important for various life-history stages of commercial fish species: seamounts, canyons, continental slope, hydrothermal vents, seeps.

## 9.4. Indicators and trends

As no HPSFM are defined this section cannot be completed.

## 9.5. References

- Allen M., Rosell R., Evans D. 2006. Predicting catches for the Lough Neagh (Northern Ireland) eel fishery based on stock inputs, effort and environmental variables. *Fisheries Management and Ecology* (13): 251-260.
- Allen D. 2010. Eels in the Waikato Catchment. Client report prepared for Mighty River Power Ltd. 105p.
- Atlantic States Marine Fisheries Commission 2000. Interstate Fishery Management Plan for American Eel. Fishery Management Report No. 36 of the Atlantic States Marine Fisheries Commission. 93p.
- Benaka L. (Ed.) 1999. Fish habitat: essential fish habitat and rehabilitation, American Fisheries Society, Bethesda, MD. 45pp.
- Beentjes M. 2010. Toheroa survey of Oreti Beach, 2009, and review of historical surveys. New Zealand Fisheries Assessment Report 2010/6, ed. 40p.
- Beentjes M., Boubée J., Jellyman J.D., Graynoth E. 2005. Non-fishing mortality of freshwater eels (*Anguilla* spp.). New Zealand Fisheries Assessment Report 2005/34. 38p.
- Benn, L. 2009. Marine Protected Areas (MPA): Habitat Maps for Canterbury. Internal Report for the Canterbury Conservancy.
- Booth J. 1979. North Cape - a 'nursery area' for the packhorse rock lobster, *Jasus verreauxi* (Decapoda: Palinuridae). New Zealand journal of Marine & Freshwater Research (13): 521-528.
- Caddy J.F. 2008. The importance of "Cover" in the life histories of demersal and benthic marine resources: A neglected issue in fisheries assessment and management. *Bulletin of Marine Science* (83): 7-52.
- Chang C., Iizyuka Y., Tzeng W. 2004. Migratory environmental history of the grey mullet *Mugil cephalus* as revealed by otolith Sr:Ca ratios. *Marine Ecology Progress Series*. 269: 277-288.
- Chiappone M., Sealey K.M.S. 2000. Marine reserve design criteria and measures of success: Lessons learned from the Exuma Cays Land and Sea Park, Bahamas. *Bulletin of Marine Science* (66): 691-705.
- Clark MR (1999) Fisheries for orange roughy (*Hoplostethus atlanticus*) on seamounts in New Zealand. *Oceanologica Acta* 22: 593-602.
- Clark MR, Rowden AA (2009) Effect of deepwater trawling on the macro-invertebrate assemblages of seamounts on the Chatham Rise, New Zealand. *Deep Sea Research* 56: 1540-1554.
- Clark, M.R.; Dunn, M.R. (2012). Spatial management of deep-sea seamount fisheries: balancing exploitation and habitat conservation. *Environmental Conservation* 39(2): . doi:10.1017/S0376892912000021
- Costello, M., McCrea, M., Freiwald, A., Lundälv, T., Jonsson, L., Bett, B., van Weering, T., de Haas, H., Roberts, J., Allen, D., 2005. Role of cold-water *Lophelia pertusa* coral reefs as fish habitat in the NE Atlantic, in: Freiwald, A., JM, R. (Eds.), *Cold-water Corals and Ecosystems*. Springer-Verlag Berlin pp. pp 771-805.
- Davidson, R., Richards, L., Duffy, C., Kerr, V., Freeman, D., D'Archino, R., Read, G., Abel, W., 2010. Location and biological attributes of biogenic habitats located on soft substrata in the Marlborough Sounds. , Prepared by Davidson Environmental Ltd. for Department of Conservation and Marlborough District Council. Survey and monitoring report no. 575.
- Diaz R., Solan M., Valente R. 2004. A review of approaches for classifying benthic habitats and evaluating habitat quality. *Journal of Environmental Management* 73: 165-181.
- Deep Water Group 2008. Operational Procedures: Hoki Fishery. 27p.
- Dorenbosch M., Grol M., Christianen M., Nagelkerken I. van der Velde G. 2005. Indo-Pacific seagrass beds and mangroves contribute to fish density coral and diversity on adjacent reefs. *Marine Ecology Progress Series*. 302: 63-76.
- Ellingsen, K., Hewitt, J., Thrush, S. 2007. Rare species, habitat diversity and functional redundancy in marine benthos. *Journal of Sea Research* 58: 291-301.
- European Commission, 2003. Commission Regulation (EC) No 1475/2003 of 20 August 2003 on the protection of deep-water coral reefs from the effects of trawling in an area north west of Scotland. *Official Journal L* 211, 14-15.
- Fosså, J., Mortensen, P., Furevik, D., 2002. The deep-water coral *Lophelia pertusa* in Norwegian waters: distribution and fisheries impacts. *Hydrobiologia* 471, 1-12.
- Francis R., Hadfield M., Bradford-Grieve J., Renwick J., Sutton P. 2005. Environmental predictors of hoki year-class strengths: an update.
- Francis, M., Morrison, M., Leathwick, J., Walsh, C., 2011. Predicting patterns of richness, occurrence and abundance of small fish in New Zealand estuaries. *Marine and Freshwater Research* 62, 1327-1341.
- Francis, M., Lyon, W., Jones, E., Notman, P., Parkinson, D., Getzlaff, C., 2012. Rig nursery grounds in New Zealand: a review and survey, New Zealand Aquatic Environment and Biodiversity Report No. 95.
- Gillanders B. Kingsford M. 1986. Elements in otoliths may elucidate the contribution of estuarine recruitment to sustaining coastal reef populations of a temperate reef fish. *Marine Ecology Progress Series* 141: 13 - 20.
- Grange K., Tovey A., Hill A.F. 2003. The spatial extent and nature of the bryozoan communities at Separation Point, Tasman Bay. 22 p.
- Graynoth E., Francis R., Jellyman D. 2008. Factors influencing juvenile eel (*Anguilla* spp.) survival in lowland New Zealand streams New Zealand Journal of Marine and Freshwater Research [N. Z. J. Mar. Freshw. Res.]. Vol. 42, no. 2, pp. 153-172. Jun 2008. (42): 153-172.
- Hall-Spencer J., Kelly J. Maggs C. 2008. Assessment of maerl beds in the OSPAR area and the development of a monitoring program. Department of Environment, Heritage and Local Government: Ireland.
- Hamer P., Jenkins G., Gillanders B. 2005. Chemical tags in otoliths indicate the importance of local and distant settlement areas to populations of a temperate sparid, *Pagrus auratus*. *Canadian Journal of Fisheries and Aquatic Sciences*. 62: 623-630.
- Haro A., Richkus W., Whalen K., Hoar A., Busch W.D., Lary S., Brush T., Dixon D. 2000. Population decline of the American eel: Implications for research and management. *Fisheries* (25): 7-16.

- Helson J., Leslie S., Clement G., Wells R., Wood R. 2010. Private rights, public benefits: Industry-driven seabed protection. *Marine Policy* (34): 557-566.
- Hewitt J.E., Thrush S.F., Halliday J., Duffy C. 2005. The importance of small-scale habitat structure for maintaining beta diversity. *Ecology* (86): 1619-1626.
- Hiscock K., Tyler-Walters H. 2006. Assessing the seabed species and biotopes- the Marine Life Information Network (MarLIN). *Hydrobiologica*. 555: 309-320.
- Hurst R., Bagley N., Anderson O., Francis M., Griggs L., Clark M., Paul L., Taylor P. 2000a. Atlas of juvenile and adult fish and squid distributions from bottom and midwater trawls and tuna longlines in New Zealand waters.
- Hurst R., Stevenson M., Bagley N., Griggs L., Morrison M., Francis M. 2000b. Areas of importance for spawning, pupping or egg-laying and juveniles of new Zealand coastal fish. . 56.
- Inland Fisheries Service Tasmania 2009. Tasmanian Freshwater Eel Fishery: Application to the Department of the Environment, Water, Heritage and Arts for the re-assessment of the Tasmanian Freshwater Eel Fishery. 22.
- Jelbart J., Ross P., Connolly R. 2007. Fish assemblages in seagrass beds are influenced by the proximity of mangrove forests. *Marine Biology* 150, 993-1002.
- Jellyman D. 1994. The fishery for freshwater eels (*Anguilla* spp.) in New Zealand New Zealand Fisheries Assessment Research Document 94/14. 25p.
- Kamenos, N., Moore, P., Hall-Spencer, J. 2004. Small-scale distribution of juvenile gadoids in shallow inshore waters; what role does maerl play? *ICES Journal of Marine Science* 61,422-429.
- Keefer M., Peery C., Wright N., Daigle W., Caudill C., Clabough T., Griffith D., Zacharias M. 2008. Evaluating the NOAA Coastal and Marine Ecological Classification Standard in estuarine systems: A Columbia River Estuary case study. *Estuarine, Coastal and Shelf Science*. 78: 89-106.
- Kerr L. 2008. Cause, consequence, and prevalence of spatial structure of white perch (*Morone americana*) populations in the Chesapeake Bay. Dissertation. University of Maryland, College Park, Maryland, USA.
- Kerr L., Secor D. 2009. Bioenergetic trajectories underlying partial migration in Patuxent River (Chesapeake Bay) white perch *Morone americana*. *Canadian Journal of Fisheries and Aquatic Sciences* 66:602-612.
- Kerr L.A., Cadrin S.X., Secor D.H. 2010. The role of spatial dynamics in the stability, resilience, and productivity of an estuarine fish population. *Ecological Applications* (20): 497-507.
- Kerr V. 2010. Marine habitat map of Northland: mangawhai to Ahipara (vers 1).
- Koenig C., Coleman F., Grimes C., FitzHugh G., Scanlon K., Gledhill C., Grace M. 2000. Protection of fish spawning habitat for the conservation of warm-temperate reef-fish fisheries of shelf-edge reefs of Florida. *Bulletin of Marine Science* (66): 593-616.
- Leathwick J., Image K., Snelder T., Weatherhead M., Wild M. 2004. Definition and tests of the Marine Environment Classifications of New Zealand's Exclusive Economic Zone and the Hauraki Gulf. NIWA Client Report CHC2004-085. 64pp.
- Leathwick J.R., Dey K., Julian K. 2006. Development of a demersal fish community classification for New Zealand's Exclusive Economic Zone. . 38.
- Leathwick J.R., Julian K. 2008. Updated conservation rankings for New Zealand's rivers and streams NIWA Report for Department of Conservation. 39.
- Leathwick J., Rowden A., Nodder S., Gorman, R., Bardsley S., Pinkerton, M., Baird, S., Hadfield, M., Currie, K., Goh, A. 2012. Benthic optimised marine environment classification for New Zealand waters. *Aquatic Environment and Biodiversity Report No. 89*. 52pp.
- Livingston M. 1990. Spawning hoki (*Macrurus novaezelandiae* Hector) concentrations in Cook Strait and off the east coast of the South Island, New Zealand, August-September 1987. *New Zealand Journal of Marine and Freshwater Research* (24):
- Lundälv, T., Jonsson, L., 2003. Mapping of deep-water corals and fishery impacts in the north-east Skagerrak, using acoustical and ROV survey techniques. *Proc 6th Underwater Sci Symp, Aberdeen, April 2003*.
- Madden C., Grossman D., Goodin K. 2005. Coastal and marine ecosystems of North America; framework for an ecological classification standard: Version II. NatureServe, Virginia, Arlington. 48 pp.
- Morato, T., Hoyle, S.D., Allain, V. and Nicol, S.J. (2010a). Seamounts are hotspots of pelagic biodiversity in the open ocean. *Proceedings of the National Academy of Science U S A*, 107, 9707-9711. doi:10.1073/pnas.0910290107
- Morato, T., Hoyle, S.D., Allain, V. and Nicol, S.J. (2010b). Tuna longline fishing around West and Central Pacific seamounts. *PloS ONE* 5(12): e14453. doi:10.1371/journal.pone.0014453
- Morrisey, D., Swales A., Dittmann S., Morrison M., Lovelock, C., Beard, C. 2010. The ecology and management of temperate mangroves. *Oceanography and Marine Biology* 48: 43-160.
- Morrison M., Jones E., Consalvey M., Berkenbusch K. 2008. Biogenic habitats and their value to New Zealand fisheries. *Water & Atmosphere* (16): 20-21.
- Morrison M., Lowe M., Parsons D., Usmar N., McLeod I. 2009. A review of land-based effects on coastal fisheries and supporting biodiversity in New Zealand. 100p.
- Murawski S.A., Brown R., Lai H.L., Rago P.J., Hendrickson L. 2000. Large-scale closed areas as a fishery-management tool in temperate marine systems: The Georges Bank experience. *Bulletin of Marine Science* (66): 775-798.
- Nagelkerken I., Kleijnen S., Klop T., van den Brand, R., de la Moriniere, E., van der Velde G. 2001. Dependence of Caribbean reef fishes on mangroves and seagrass beds as nursery habitats: a comparison of fish faunas between bays with and without mangroves/seagrass beds. *Marine Ecology Progress Series* 214: 225-235.
- North Pacific Fishery Management Council. 1998. Omnibus essential fish habitat amendments for groundfish in the Bering Sea, Aleutian Islands, and Gulf of Alaska, king and tanner crab in the Bering Sea and Aleutian Islands, Alaska scallop, and Alaska salmon fishery management plans. North Pacific Fishery Management Council, Alaska.
- O'Driscoll R., Booth J., Bagley N., Anderson O., Griggs L., Stevenson M., Francis M. 2003. Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand deepwater fish, pelagic fish, and invertebrates. NIWA Technical Report 119. 377.
- O'Driscoll, R.L.; Clark, M.R. (2005). Quantifying the relative intensity of fishing on New Zealand seamounts. *New Zealand Journal of Marine and Freshwater Research* 39: 839-850.
- Parsons D., Morrison M., McKenzie J., Hartill, B, Bian R., Francis RICC (2011) A fisheries perspective of behavioural variability: differences in movement behavior and extraction rate of an exploited sparid, snapper (*Pagrus auratus*). *Canadian Journal of Fisheries and Aquatic Sciences*. 68: 632-642.
- Roberts C.M. 2000. Selecting marine reserve locations: Optimality versus opportunism. *Bulletin of Marine Science* (66): 581-592.
- Rosenberg A., Bigford T., Leathery S., Hill R., Bickers K. 2000. Ecosystem approach to fishery management through essential fish habitat. *Bulletin of Marine Science* (66): 535-542.
- Ross S., Moser M. 1995. Life-history of juvenile gag, *Mycteroperca microlepis*, in North-Carolina estuaries. *Bulletin of Marine Science*. 56; 222-237.

- Schumacher J.D., Kendall A.W. 1991. Some interactions between young walleye Pollock and their environment in the western gulf of Alaska 40p.
- Schwarz A.-M., Morrison M., Hawes I., Halliday J. 2006. Physical and biological characteristics of a rare marine habitat: subtidal seagrass beds of offshore islands. 39.
- Thrush S.F., Gray J.S., Hewitt J.E., Uglund K.I. 2006. Predicting the effects of habitat homogenization on marine biodiversity. *Ecological Applications* (16): 1636-1642.
- Tracey, D.M., Rowden, A.A., Mackay, K.A., Compton, T (2011) Habitat-forming cold-water corals show affinity for seamounts in the New Zealand region. *Marine Ecology Progress Series* 430: 1–22.
- Turner G.E., Boesch D. 1987. Aquatic animal production and wetland relationships: insights gleaned following wetland loss or gain. In: Hooks D. (Ed.) *Ecology and management of wetlands*. Croon Helms, Ltd., Beckenham, Kent, U.K. 25-39.
- Turner S., Schwarz A.-M. 2006. Management and conservation of seagrass in New Zealand: an introduction. (264): 90.
- Vooren C. 1975. Nursery grounds of tarakihi (Teleostei: Cheilodactylidae) around New Zealand. *New Zealand Journal of Marine and Freshwater Research* (9): 121-158.
- Warner R.R., Swearer S.E., Caselle J.E. 2000. Larval accumulation and retention: Implications for the design of marine reserves and essential fish habitat. *Bulletin of Marine Science* (66): 821-830.
- Western Pacific Regional Fishery Management Council 1998. Magnuson-Stevens Act definitions and required provisions. Western Pacific Regional Fishery Management Council, Hawaii.

## 10. Land-based effects on fisheries, aquaculture and supporting biodiversity

<i>Scope of chapter</i>	This chapter outlines the main known threats from land-based activities to fisheries, aquaculture and supporting biodiversity. It also describes the present status and trends in land-based impacts.
<i>Area</i>	All of the New Zealand freshwater, EEZ and territorial sea.
<i>Focal localities</i>	Freshwater habitats and areas closest to the coast are likely to be most impacted; this will be exacerbated in areas with low water movement. Anthropogenically increased sediment run-off is particularly high from the Waipua and Waipaoa river catchments on the east coast of the North Island. Areas of intense urbanisation or agricultural use of catchments are also likely to be impacted by bacteria, viruses, heavy metals and nutrients.
<i>Key issues</i>	Habitat modification, sedimentation, aquaculture, shellfish, terrestrial land-use change (particularly for urbanisation, forestry or agriculture) water quality and quantity, contamination, consequences to seafood production of increased pollutants, freshwater management and demand.
<i>Emerging issues</i>	Impacts on habitats of particular significance to fisheries management (HPSFM), linkages through rainfall patterns to climate change, shellfish bed closures, habitat remediation, domestic animal diseases in protected marine species, proposed aquaculture expansion, water abstraction impacts.
<i>MPI Research (current)</i>	Habitats of particular significance for fisheries management: Kaipara Harbour (ENV2009/07), Toheroa abundance (TOH2007/03), Biogenic habitats as areas of particular significance for fisheries management (HAB2007/01), Research on Biogenic Habitat-Forming Biota and their functional role in maintaining Biodiversity in the Inshore Region, 5-150m depths (ZBD2008/01 – this is also part-funded by Oceans Survey 2020, NIWA and MBIE).
<i>NZ Government Research (current)</i>	Ministry of Business, Innovation and Employment (MBIE) funded programs: (After the outfall: recovery from eutrophication in degraded New Zealand estuaries (UOCX0902).  NIWA Core funding in two areas. Firstly, The 'Managing marine stressors' area under the 'Coasts and Oceans' centre, specifically the programme 'Managing marine resources' and the project 'Measuring mapping and conserving (C01X0505)'. Secondly, in the 'Fisheries' Centre programme 3 which deals with ecosystem-based management approaches in conjunction with the 'Coasts and Oceans' centre.
<i>Links to 2030 objectives</i>	Objective 8: Improve RMA fisheries interface. Objective 4: Support aquaculture development
<i>Related chapters/issues</i>	Habitats of particular significance for fisheries management (HPSFM), marine environmental monitoring.

### 10.1. Context

It has been acknowledged for some time now that land-based activities can have important effects on seafood production. The main threats to the quality and use of the world's oceans are (GESAMP 2001):

- alteration and destruction of habitats and ecosystems;
- effects of sewage on human health;

- widespread and increased eutrophication;
- decline of fish stocks and other renewable resources; and
- changes in sediment flows due to hydrological changes

. Coastal development is projected to impact 91% of all inhabited coasts by 2050 and will contribute to more than 80% of all marine pollution (Nellemann *et al.* 2008).

Aquaculture and land-based activities that may have impacts on seafood production are primarily regulated under the Resource Management Act 1991 (and subsequent amendments). Fisheries are controlled under the Fisheries Act 1996. Fisheries 2030 is a long-term policy strategy and direction paper of the Ministry for Primary Industries. It was released in 2009 and states that improving the Fisheries/Resource Management Act interface is a priority (objective 8). Strategic actions to achieve this priority are listed as:

- 8.1 Improve fisheries sector input to processes that manage RMA-controlled effects on the marine and freshwater environment.
- 8.2 Promote the development and use of RMA national policy statements, environmental standards, and regional coastal and freshwater plans

The Government's 'Fresh Start for Freshwater Programme'<sup>26</sup> (lead by MfE and MPI) is addressing a range of issues through a water reform strategy that includes governance, setting objectives and limits, managing within limits (quality and quantity) and that better reflects Maori/Iwi rights and interests in water management. The Coastal Policy Statement (2010) also has relevance to matters of fisheries interest, e.g. Policy 20(1) (*paraphrased*) controls the use of vehicles on beaches where (b) harm to shellfish beds may result. MPI also works with other agencies, principally DOC, MfE and regional councils and through the Natural Resource Cluster to influence these processes to ensure consideration of land-based impacts upon seafood production.

Land-based effects on seafood production and supporting biodiversity in this context are defined as resulting either from the inputs of contaminants from terrestrial sources or through engineering structures (e.g., breakwaters, causeways, bridges) that change the nature and characteristics of coastal habitats and modify hydrodynamics. The major route for entry of land-based contaminants into the marine environment is associated with freshwater flows (rivers, streams, direct runoff and ground water), although contaminants may enter the marine environment via direct inputs (e.g., landslides) or atmospheric transport processes.

The most important land-based effect in New Zealand is arguably increased sediment deposition around our coasts (Morrison *et al.* 2009). This deposition has been accelerated due to increased erosion from land-use, which causes gully and channel erosion and landslides (Glade 2003). Inputs of sediments to our coastal zone, although naturally high in places due to our high rainfall and rates of tectonic uplift (Carter 1975), have been accelerated by human activities (Goff 1997). Sediment inputs are now high by world standards and make up ~1% of the estimated global detrital input to the oceans (Carter *et al.* 1996). By contrast New Zealand represents only ~0.3% of the land area that drains into the oceans (Griffiths and Glasby 1985, Milliman and Syvitski 1992).

Different land use effects act over different scales; for example localised effects act on small streams and adjacent estuarine habitats, large scale effects extend to coastal embayments and shelf ecosystems. Associated risks will vary according to location and depend on the relevant ecosystem services (e.g. high value commercial fishery stocks) and their perceived sensitivities. The risk from stormwater pollutants will be more important near urban areas and the effects of nutrient enrichment will be more important near intensively farmed rural areas.

---

<sup>26</sup> <http://www.mfe.govt.nz/issues/water/freshwater/fresh-start-for-fresh-water/>

The risk from land-based impacts for seafood production is that they will limit the productivity of a stock or stocks. For example, the bryozoan beds around Separation Point in Golden Bay, were protected from fishing, amongst other reasons, due to their perceived role as nursery grounds for a variety of coastal fish species in 1980 (Grange *et al.* 2003). Recent work has suggested the main threat to these bryozoans is now sedimentation from the Motueka River, which may inhibit recovery of any damaged bryozoans (Grange *et al.* 2003, Morrison *et al.* 2009). Any declines in this bryozoan bed and associated ecological communities could also affect the productivity of adjacent fishery stocks.

The New Zealand aquaculture industry has an objective of developing into a billion dollar industry by 2025 (Aquaculture New Zealand 2012). Government supports well-planned and sustainable aquaculture through its *Aquaculture Strategy* and *Five-year Plan*. One of the desired outcomes of actions by the New Zealand Government is to enable more space to be made available for aquaculture. This outcome is likely to heighten the potential for conflict between aquaculture proponents and those creating negative land-based effects.

MPI mainly manage in the marine environment, therefore this topic area will be dealt with first. MPI also manages the freshwater eel fishery; this will be dealt with latterly within relevant sections.

## **10.2. Global understanding**

### **10.2.1. Land-based influences**

The importance of different land-based influences differ regionally but the South Pacific Regional Environmental Programme (SPREP, which includes New Zealand) defines waste management and pollution control as one of its four strategic priorities for 2011-2015 (SPREP 2010). “

Influences, including land-based influences, seldom work in isolation; for example the development of farming and fishing over the last hundred years has meant that increased sediment and nutrient runoff has to some degree occurred simultaneously with increased fishing pressure. However, the impact of these influences has often been studied in isolation. In a review on coastal eutrophication, Cloern (2001) stated that “*Our view of the problem [eutrophication] is narrow because it continues to focus on one signal of change in the coastal zone, as though nutrient enrichment operates as an independent stressor; it does not reflect a broad ecosystem-scale view that considers nutrient enrichment in the context of all the other stressors that cause change in coastal ecosystems*”. These influences (in isolation or combination) can also cause indirect effects, such as decreasing species diversity which then lessens resistance to invasion by non-indigenous species or species with different life-history strategies (Balata *et al.* 2007, Kneitel and Perrault 2006, Piola and Johnston 2008). Studies that research a realistic mix of influences are rare.

Sediment deposition can be an important influence, particularly in areas of high rainfall, tectonic uplift, and forest clearances, or areas where these activities coincide. Sediments are known to erode from the land at an increased rate in response to human use, for example, estimates from a largely deforested tropical highland suggest erosion rates 10-100 times faster than pre-clearance rates (Hewawasam *et al.* 2003). Increased sediment either deposited on the seafloor or suspended in the water column can negatively impact upon invertebrates in a number of ways including: burial, scour, inhibiting settlement, decreasing filter-feeding efficiency and decreasing light penetration, generally leading to less diverse communities, with a decrease in suspension feeders (Thrush *et al.* 2004). These impacts can affect the structure, composition and dynamics of benthic communities (Airolidi 2003, Thrush *et al.* 2004). Effects of this increased sediment movement and deposition on finfish are mostly known from freshwater fish and can range from behavioural (such as decreased feeding rates) to sublethal (e.g., gill tissue disruption) and lethal as well as having effects on habitat important to fishes (Morrison *et al.* 2009). These effects differ by species and life-stages and are dependant upon factors that include the

duration, frequency and magnitude of exposure, temperature, and other environmental variables (Servizi and Martens 1992).

Increased nutrient addition to the aquatic environment can initially increase production, but with increasing nutrients there is an increasing likelihood of harmful algal blooms and cascades of effects damaging to most communities above the level of the plankton (Kennish 2002; Heisler *et al.* 2008). This excess of nutrients is termed eutrophication. Eutrophication can stimulate phytoplankton growth which can decrease the light availability and subsequently lead to losses in benthic production from seagrass, microalgae or macroalgae and their associated animal communities. Algal blooms then die and their decay depletes oxygen and blankets the seafloor. The lack of oxygen in the bed and water column can lead to losses of finfish and benthic communities. These effects are likely to be location specific and are influenced by a number of factors including: water transparency, distribution of vascular plants and biomass of macroalgae, sediment biogeochemistry and nutrient cycling, nutrient ratios and their regulation of phytoplankton community composition, frequency of toxic/harmful algal blooms, habitat quality for metazoans, reproduction/growth/survival of pelagic and benthic invertebrates, and subtle changes such as shifts in the seasonality of ecosystems (Cloern 2001). These effects of eutrophication abound in the literature, for example, the formation of dead (or anoxic) zones is exacerbated by eutrophication, although oceanographic conditions also play a key role (Diaz and Rosenberg 2008). Dead zones have now been reported from more than 400 systems, affecting a total area of more than 245,000 square kilometres (Diaz and Rosenberg 2008). This includes anoxic events from New Zealand in coastal north-eastern New Zealand and Stewart Island (Taylor *et al.* 1985, Morrissey 2000).

Other pollutants such as heavy metals and organic chemicals can have severe effects, but are more localised in extent than sediment or nutrient pollution (Castro and Huber 2003, Kennish 2002). Fortunately the concentration of these pollutants in most New Zealand aquatic environments is relatively low, with a few known exceptions. Examples of this include naturally elevated levels of arsenic in Northland<sup>27</sup>, Cadmium levels in Foveaux Strait oysters (Frew *et al.* 1996) and levels of Nickel and chromium within the Motueka river plume in Tasman Bay (Forrest *et al.* 2007). The Cadmium levels have caused market access issues for Foveaux Strait Oysters. Some anthropogenically generated pollutants such as copper, lead, zinc and PCBs are high in localised hotspots within urban watersheds. In the Auckland region these hotspots tend to be in muddy estuarine sites and tidal creeks that receive runoff from older urban catchments<sup>28</sup>. There is a lack of knowledge on the impacts of these pollutants upon fisheries.

Climate change is likely to interact with the effect of land-based impacts as the main delivery of land-based influences is through rainfall and subsequent freshwater flows. Global climate change projections include changes in the amount and regional distribution of rainfall over New Zealand (IPCC 2007). More regional predictions include increasing frequency of heavy rainfall events over New Zealand (Whetton *et al.* 1996). This is likely to exacerbate the impact of some land-based influences as delivery peaks at times of high rainfall, e.g. sediment delivery (Morrison *et al.* 2009).

Physical alterations of the coast are generally, but not exclusively (i.e. wetland reclamation for agriculture), concentrated around urban areas and can have a number of consequences on the marine environment (Bulleri and Chapman 2010). Changes in diversity, habitat fragmentation or loss and increased invasion susceptibility have all been identified as consequences of physical alteration. The effects of physical alterations upon fisheries remain largely unquantified; however the habitat loss or alteration portion of physical alterations will be dealt with under the habitats of particular significance for fisheries management (HPSFM) section.

---

<sup>27</sup> Accessible on the [www.os2020.org.nz](http://www.os2020.org.nz) website.

<sup>28</sup> Available from the State of the Auckland Region report 2010, Chapter 4.4 Marine, at <http://www.arc.govt.nz/albany/index.cfm?FD6A3403-145E-173C-986A-A0E3C199B8C5>

An area of emerging interest internationally is infectious diseases from land-based animals affecting marine populations. Perhaps the most well-known example of this is the canine distemper outbreak in Caspian seals that cause a mass mortality in the Caspian sea in 2000 (Kennedy *et al.* 2000)

### **10.2.2. Habitat restoration**

Habitat restoration or rehabilitation has been the subject of much recent research. Habitat restoration or rehabilitation rarely, if ever, replaces what was lost and is most applicable in estuarine or enclosed coastal areas as opposed to exposed coastal or open ocean habitats (Elliott *et al.* 2007). Connectivity of populations is a key consideration when evaluating the effectiveness of any marine restoration or rehabilitation (Lipcius *et al.* 2008). In the marine area, seagrass replanting methodologies are being developed to ensure the best survival success (Bell *et al.* 2008) and artificial reefs can improve fisheries catches, although whether artificial reefs boost population numbers or merely attract fish is unclear (Seaman 2007). In addition, The incorporation of habitat elements in engineering structures, e.g., artificial rockpools in seawalls, shows promise in terms of ameliorating impacts of physical alterations (Bulleri 2006). Spatial approaches to managing land-use impacts, such as marine reserves, will be covered under the section about HPSFM.

Freshwater rehabilitation has been reviewed by Roni *et al.* (2008). Habitat reconnection, floodplain rehabilitation and instream habitat improvement are all suggested to result in improved habitat and local fish abundances. Riparian rehabilitation, sediment reduction, dam removal, and restoration of natural flood regimes have shown promise for restoring natural processes that create and maintain habitats, but there is a lack of long-term studies to gauge their success. Wild eel fisheries in America and Europe have declined over time (Allen *et al.* 2006, Atlantic States Marine Fisheries Commission 2000, Haro *et al.* 2000). Declines in wild eel fisheries have been linked to a number of factors including: barriers to migration; hydro turbine mortality; and habitat loss or alteration. Information to quantitatively assess these linkages is however often lacking (Haro *et al.* 2000).

## **10.3. State of knowledge in New Zealand**

Land-based effects will be most pronounced closest to the land, therefore it is freshwater, estuarine, coastal, middle depths and deepwater fisheries, in decreasing order, that will be most affected. The scale of land-use effects will, however, differ depending upon the particular influence. The most localised of these are likely to be direct physical impacts; for example, the replacement of natural shorelines with seawalls; although even direct physical impacts can have larger scale impacts, such as affecting sediment transport and subsequently beach erosion, or contributing to cumulative effects upon ecosystem responses. Point-source discharges are likely to have a variable scale of influence, and this influence is likely to increase where a number of point-sources discharge, particularly when this occurs into an embayed, low-current environment. An example of this is the multiple stormwater discharges into the Waitemata harbour in Auckland (Hayward *et al.* 2006). The largest influence can be from diffuse-source discharges such as nutrients or sediment (Kennish 2002). For example, the influence of diffuse-source materials from the Motueka river catchment in Golden Bay on subtidal sediments and assemblages and shellfish quality can extend up to tens of kilometres offshore (Tuckey *et al.* 2006; Forrest *et al.* 2007), with even a moderate storm event extending a plume greater than 6km offshore (Cornelisen *et al.* 2011). Terrestrial influences on New Zealand's marine environment can, at times be detected by satellites from differences in ocean colour and turbidity extending many kilometres offshore from river mouths (Gibbs *et al.* 2006).

All coastal areas are unlikely to suffer from land-based impacts in the same way. The quantities of pollutants or structures differ spatially. Stormwater pollutants, seawalls and jetties are more likely to be concentrated around urban areas. Nutrient inputs are likely to be concentrated either around sewage

outlets or associated with areas of intensive agriculture or horticulture. Sediment production has been mapped around the country and is greatest around the west coast of the South Island and the East coast of the North Island (Griffiths and Glasby 1985, Hicks and Shankar 2003, Hicks *et al.* 2011). Notably the catchments where improved land management may result in the biggest changes to sediment delivery to coastal environments are likely to be the Waiapu and Waipaoa river catchments on the East coast of the North Island. In addition to this, the sensitivity of receiving environments is also likely to differ; this will be covered in subsequent sections.

A MPI funded survey of scientific experts (MacDiarmid *et al.* 2012) addressed the vulnerability to a number of threats of marine habitat types within the New Zealand's Territorial Sea and Exclusive Economic Zone (EEZ). Each vulnerability score was based on an assessment of five factors including the spatial scale, frequency and functional impact of the threat in the given habitat as well as the susceptibility of the habitat to the threat and the recovery time of the habitat following disturbance from that threat. The study found that the number of threats and their severity were generally considered to decrease with depth, particularly below 50m. Reef, sand, and mud habitats in harbours and estuaries and along sheltered and exposed coasts were considered to be the most highly threatened habitats. The study also reported that over half of the twenty-six top threats fully, or in part, stemmed from human activities external to the marine environment itself. The top six threats in order were:

1. ocean acidification,
2. rising sea temperatures resulting from global climate change,
- 3<sup>rd</sup> equal bottom trawling fishing,
- 3<sup>rd</sup> equal increased sediment loadings from river inputs
- 5<sup>th</sup> equal change in currents from climate change
- 5<sup>th</sup> equal increased storminess from climate change

The reader is guided to MacDiarmid *et al.* (2012) for more detail including tables of threats-by-habitat and habitats-by-threat. Climate change and ocean acidification, although they can be considered land-based effects, are covered under the Chapters in this document called “New Zealand Regional climate and oceanic setting” and “Biodiversity”.

The protozoan *Toxoplasma gondii* has been identified as the cause of death for 7 of 28 Hector's and Maui's dolphins examined since 2007 (W. Roe, Massey University, unpubl. data, 31 July 2012). Land-based runoff containing cat faeces is believed to be the means by which *Toxoplasma gondii* enters the marine environment (Hill & Dubey 2002). A Hector's dolphin has also tested positive for *Brucella abortus* (or a similar organism) a pathogen of terrestrial mammals that can cause late pregnancy abortion, and has been seen in a range of cetacean species elsewhere<sup>29</sup>.

### 10.3.1. Completed research

A MPI funded project (IPA2007/07) reviewed the impacts of land based influences on coastal biodiversity and fisheries (Morrison *et al.* 2009). This review used a number of lines of evidence to conclude that in this context, sedimentation is probably New Zealand's most important pollutant. The negative impacts of sediment include decreasing efficiency of filter-feeding shellfish (such as cockles, pipi, and scallops), reduced settlement success and survival of larval and juvenile phases (e.g., paua, kina), and reductions in the foraging abilities of finfish (e.g., juvenile snapper). Indirect effects include the modification or loss of important nursery habitats, particularly biogenic habitats (green-lipped and horse mussel beds, seagrass meadows, bryozoan and tubeworm mounds, sponge gardens,

---

<sup>29</sup><http://www.doc.govt.nz/Documents/conservation/native-animals/marine-mammals/maui-tmp/mauis-tmp-discussion-document-full.pdf>

kelps/seaweeds, and a range of other structurally complex species). Inshore filter-feeding bivalves and biogenic habitats were identified as the most likely to be adversely affected by sedimentation. Eutrophication was also identified as a potential threat from experience overseas.

Marine restoration studies published in New Zealand have focused on the New Zealand cockle *Austrovenus stutchburyi*. The first of these studies identified a tagging methodology to aid relocation of transplanted individuals (Stewart and Creese 1998). Subsequent studies stressed the use of adults in restoration and the importance of site selection, either from theoretical or modelling viewpoints (Lundquist *et al.* 2009, Marsden and Adkins 2009). Detailed restoration methodology has been investigated in Whangarei Harbour and recommends replanting adults at densities between 222 and 832 m<sup>-2</sup> (Cummings *et al.* 2007).

Multiple influences in areas relevant to seafood production in New Zealand have been addressed by three studies. A field experiment near Auckland showed greater effects of three heavy metals (Copper, lead and Zinc) in combination compared to isolation on infaunal colonisation of intertidal estuarine sediments (Fukunaga *et al.* 2010). A survey approach looking at the interaction of sediment grain size, organic content and heavy metal contamination upon densities of 46 macrofaunal taxa across the Auckland region also showed a predominance of multiplicative effects (Thrush *et al.* 2008). Although influences can work in unexpected directions; as in a study on large suspension feeding bivalves off estuary mouths where the anticipated negative impacts from sediment were not observed and these species benefited from food resources generated from those estuaries (Thrush *et al.* In Press).

Toheroa populations are currently closed to all but customary harvesting but have failed to recover to former population levels even though periodic (and sometimes substantial) pulses in young recruits have been detected in both Northland and Southland (Beentjes 2010, Morrison and Parkinson 2008). Current thinking suggests a mix of influences are probably responsible for these declines including over-harvesting, land-use changes leading to changes in freshwater seeps on the beaches and vehicle traffic (Morrison *et al.* 2009). A number of discrete pieces of research have been completed in this area. A review of the wider impact of vehicles on beaches and sandy dunes has been completed, and suggested more research was needed on the impacts of vehicle traffic on the intertidal (Stephenson 1999). A four day study over a fishing contest on 90 mile beach showed the potential of traffic to produce immediate mortalities of juvenile toheroa, but the temporal importance of this could not be gauged (Hooker and Redfearn 1998). Mortalities of toheroa from the Burt Munro Classic motorcycle race on Oreti beach have been quantified and recommendations made for how to minimise these, but again the importance of vehicle traffic for toheroa survival over longer time periods was unclear (Moller *et al.* 2009).

The effects of large-scale habitat loss and modification on eels in New Zealand are clearly significant, but difficult to quantify (Beentjes *et al.* 2005). Significant non-fisheries mortality of New Zealand freshwater longfin and shortfin eels are caused by mechanical clearance of drainage channels, and damage by hydro-electric turbines and flood control pumping. Eels prefer habitat that offers cover and in modified drains aquatic weed provides both daytime cover and nighttime foraging areas. Loss of weed and natural debris can thus result in significant displacement of eels to other areas. In addition, wetlands drainage has resulted in greatly reduced available habitat for eels, particularly shortfins which prefer slower-flowing coastal habitats such as lagoons, estuaries, and lower reaches of rivers. Water abstraction is one of a number of information requirements identified in this paper to better define the effects on eel populations.

Rhodolith beds have been surveyed in the Bay of Islands and high diversity reported even in areas of abundant fine sediments (Nelson *et al.* 2012). It is unclear if the increasing sedimentation occurring in the Te Rawhiti Reach is negatively impacting rhodoliths and whether this atypical rhodolith bed (i.e., with abundant fine sediments) is at risk if current sedimentation and mobilisation rates continue.

A number of Integrated Catchment Management (ICM) projects are underway in New Zealand. These take a holistic view to land management incorporating aquatic effects; this approach could help restore water quality of both fresh and coastal waters. An overview of these projects is given in a Ministry for the Environment Report on integrated catchment management (Environmental Communications Limited 2010). Many of these projects employ restoration techniques such as riparian planting, but few assessments of the effectiveness of riparian planting exist. One assessment of the effect of nine riparian zone planting schemes in the North Island on water quality, physical and ecological indicators concluded that riparian planting could improve stream quality; in particular rapid improvements were seen in terms of visual clarity and channel stability (Parkyn *et al.* 2003). Nutrient and faecal contamination results were more variable. Improvement in macroinvertebrate communities did not occur in most streams and the three factors needed for these were canopy closure (which decreased stream temperature), long lengths of riparian planting and protection of headwater tributaries. A modelling study also demonstrated the long time lag needed to grow large trees which then provide wood debris to structure channels which achieves the best stream rehabilitation results (Davies-Colley *et al.* 2009). Although some of these studies extend into the marine realm (at least in terms of monitoring) it is difficult to gauge the impact of these activities upon fisheries or aquaculture, particularly on wider scales because ICM studies have been localised at small scales.

The review of land based effects (Morrison *et al.* 2009) identified knowledge gaps and made suggestions for more relevant research on these influences:

- identification of fisheries species/habitat associations for different life stages, including consideration of how changing habitat landscapes may change fisheries production;
- better knowledge of connectivity between habitats and ecosystems at large spatial scales;
- the role of river plumes;
- the effects of land-based influences both directly on fished species, and indirectly through impacts on nursery habitats;
- a better spatially-based understanding, mapping and synthesis of the integrated impacts of land-based and marine-based influences on coastal marine ecosystems.

The locations where addressing land-based impacts is likely to result in a lowering in risk to seafood production or increased seafood production, excluding those already mentioned, are undefined.

### 10.3.2. Current research

A number of ongoing research projects exist that will improve the knowledge of land-based impacts upon seafood production. Project ENV2009/07 investigates habitats of particular significance for fisheries management within the Kaipara Harbour and one objective is to assess fishing and land-based threats to these habitats. Current research is investigating the impact of a range of influences upon toheroa at Ninety-Mile Beach (project TOH2007/03). Environmental factors, including land-based impacts (particularly vehicle use and changing land-use patterns) are implicated in poor recovery of this population since the closure of this commercial and recreational fishery in the 1960s. A MPI biodiversity project also has components that address land-based effects; the threats to biogenic habitats are addressed in project ZBD2008/01.

Research is also ongoing on land-use effects at a national scale. A national scale threat analysis is also being carried out for biogenic habitats, given their likely importance for fisheries management (project HAB2007/01). A Ministry of Business, Innovation and Employment (MBIE) funded project<sup>30</sup> of particular relevance is (project number and lead agencies in brackets): Nitrogen reduction and benthic recovery (UOCX0902, University of Canterbury). This research aims to determine the trajectories and thresholds of coastal ecosystem recovery following removal of excessive nutrient

---

<sup>30</sup> <http://www.msi.govt.nz/update-me/who-got-funded/>

loading (called "eutrophication") and earthquake impacts. This will be achieved by monitoring the effects of diverting all of Christchurch's treated wastewater discharge from the eutrophied Avon-Heathcote (Ihutai) Estuary and the subsequent earthquake induced disturbances to this diversion.

Although not current research, the Department of Conservations suggested research priorities in the "*Review of the Maui's dolphin Threat management plan: Consultation paper*" include objectives to determine the presence, pathways and possible mitigation of the threat from *Toxoplasmosis gondii*<sup>31</sup>.

## 10.4. Indicators and trends

A national view of the impacts of land-based influences upon seafood production does not exist; this could be facilitated by better coordination and planning of the many disparate marine monitoring programmes running around the country. Monitoring of marine water quality and associated communities is carried out through a variety of organisations, including, universities, regional councils and aquaculture or shellfisheries operations. Regional council monitoring of water quality and associated biological communities is often reported through web sites such as the Auckland Regional Council environmental monitoring data which is available on the internet<sup>32</sup> or summary reports such as the Hauraki Gulf state of the Environment 2011 report<sup>33</sup>. Water quality and associated communities may also be monitored for a regional council as part of a consent application or as a stipulation for a particular marine development. The data from aquaculture and shellfisheries water quality monitoring is not generally available. Improved coordination and planning of marine monitoring has been achieved in some places, e.g., the United Kingdom<sup>34</sup>. The Marine Environmental Monitoring Programme (ZBD2010-42), is a step towards this goal, more information is available on this project in the *Biodiversity* chapter of this document. Possible national scale proxies for coastal faecal contamination may exist after collating information from sanitation area monitoring for shellfish harvesting or shellfish harvesting closure information.

Marine water quality indicators are available nationally from 407 coastal bathing beaches which have been monitored for human health issues, rather than environmental purposes, over the last six years<sup>35</sup>. No temporal trends were detectable in this relatively short time period, however changes in sites monitored over this time may have confounded this analysis. Over the 2007-8 and 2008-9 summers, 79% of the swimming sites met the guidelines for contact recreation almost all the time. At least 95% of the samples at these sites had safe *Enterococci* levels (which is an indicator of human and animal sewage). Two percent of the sites (located within the Manukau harbour and on the West coast of Auckland), breached the guidelines more than 25% of the time. In general, the most polluted sites were embayed locations with poor natural flushing.

The Ministry for the Environment (MfE) also reports on freshwater quality. River water quality indicators that have been assessed have direct relevance to the eel, and other freshwater fisheries, and this water will flow through estuaries and enter the marine environment. The National River Water Quality Network (NRWQN) has national coverage, and has been running for over 20 years and has recently reported upon the following 8 variables: temperature, dissolved oxygen, visual clarity, dissolved reactive and total phosphorous, and ammoniacal, oxidised and total nitrogen (Ballantine and

<sup>31</sup> <http://www.doc.govt.nz/Documents/conservation/native-animals/marine-mammals/maui-tmp/mauis-tmp-discussion-document-full.pdf>

<sup>32</sup> <http://maps.auckland.govt.nz/aucklandregionviewer/?widgets=HYDROTEL>

<sup>33</sup> <http://www.arc.govt.nz/albany/fms/main/Documents/Environment/Coastal%20and%20marine/hgstateoftheenvreport2011.pdf>

<sup>34</sup> [http://www.cefas.co.uk/data/marine-monitoring/national-marine-monitoring-programme-\(nmmp\).aspx](http://www.cefas.co.uk/data/marine-monitoring/national-marine-monitoring-programme-(nmmp).aspx)

<sup>35</sup> <http://www.mfe.govt.nz/environmental-reporting/freshwater/recreational/snapshot/coastal.html#results>

Davies-Colley 2009). Dissolved oxygen showed few meaningful trends and the ammoniacal nitrogen data suffered from a processing artefact. An upward, although not significant trend in temperature and an improvement of water clarity were seen at the national scale. However, a negative correlation was seen between water clarity and percent of catchment in pasture, which suggests any expansion of pasture lands may have impacts on clarity. Strong increasing trends over time were seen in oxidised nitrogen, total nitrogen, total phosphorous and dissolved reactive phosphorous. These latter trends all signify deteriorating water quality and are mainly attributable to increased diffuse-source pollution from the expansion and intensification of pastoral agriculture.

Total Nitrogen and Phosphorous loads to the coast in New Zealand have been modelled and were estimated at 167,300 and 63,100 t yr<sup>-1</sup>, respectively (Elliot *et al.* 2005)<sup>36</sup>. The main sources of Nitrogen and Phosphorous were from pastoralism (70%) and erosion (53%), respectively. The dairy herd in New Zealand has approximately doubled (increased 211%) since 1981 (whilst other grazer numbers have been relatively stable or declining)<sup>9</sup>. The amount of Urea and Superphosphate (New Zealand's most common nitrogen and phosphorous fertiliser) have increased 27.7 and 1.6 fold, respectively over the same period<sup>37</sup>. The use of Urea is currently around 100 kg.ha<sup>-1</sup> for dairying and ~10 kg.ha<sup>-1</sup> for sheep and beef farms (MPI 2012). The area in dairy farming is ~ 2 million hectares compared to 3.6 million hectares for sheep and 2 million hectares in beef farming (MPI 2012). Therefore Urea use in New Zealand is dominated by the dairy industry. These statistics provide strong circumstantial evidence that the expansion in dairying is primarily responsible for these declines in water quality from agricultural sources.

High faecal coliform counts (primarily from mammal or bird faeces) can impact upon the value gained from shellfish fisheries and aquaculture. Area closures to commercial harvesting usually depend on an areas rainfall/runoff relationship and areas closer to significant farming areas or urban concentrations are likely to be closed more frequently, due to high faecal coliform counts, than areas where the catchment is unfarmed or not heavily populated, e.g. Inner Pelorus sound is likely to be closed more frequently than outer Pelorus Sound (Marlborough Sounds)<sup>38</sup>. For coastal areas of the Marlborough Sounds, the Coromandel Peninsula and Northland closures can range from a few days to over 50 percent of the time in a given year<sup>39</sup>. Certain fisheries may in practice be limited by the amount of time where water quality is sufficient to allow harvesting, e.g. the cockle fishery in COC1A (Snake bank in Whangarei harbour) was closed for 101, 96, 167, 96 and 117 days for the 2006-7, 2007-8, 2008-9, 2009-10 and 2010-11 fishing years, respectively due to high faecal coliform counts from sewage spills or runoff<sup>40</sup>. Models also now exist that allow real-time prediction of *E. coli* pulses associated with storm events, e.g. Wilkinson *et al.* 2011, which may help harvesters to better cope with water quality issues.

## 10.5. References

- Airolidi L. 2003. The effects of sedimentation on rocky coast assemblages. In: *Oceanography and Marine Biology*. 161-236.
- Allen M., Rosell R., Evans D. 2006. Predicting catches for the Lough Neagh (Northern Ireland) eel fishery based on stock inputs, effort and environmental variables. *Fisheries Management and Ecology* (13): 251-260.
- Atlantic States Marine Fisheries Commission 2000. Interstate Fishery Management Plan for American Eel. Fishery Management Report No. 36 of the Atlantic States Marine Fisheries Commission. 93p.
- Balata D., Piazzzi L., Cinelli F. 2007. Increase of sedimentation in a subtidal system: Effects on the structure and diversity of macroalgal assemblages. *Journal of Experimental Marine Biology and Ecology* (351): 73-82.
- Ballantine D., Davies-Colley R. 2009. Water quality trends at National River Water Quality Network sites for 1989–2007. Prepared for Ministry for the Environment, NIWA Client Report: HAM2009-026. 43p.
- Beentjes M., Boubée J., Jellyman J.D., Graynoth E. 2005. Non-fishing mortality of freshwater eels (*Anguilla* spp.). New Zealand Fisheries Assessment Report 2005/34. 38p.

<sup>36</sup> This is an underestimate because streams with catchments less than 10km<sup>2</sup> were excluded from this calculation.

<sup>8</sup> <http://www.stats.govt.nz/infoshare>

<sup>10</sup> Pers. Comms. Brian Roughan, New Zealand Food Safety Authority.

<sup>11</sup> Pers. Comms. Brian Roughan, New Zealand Food Safety Authority.

<sup>12</sup> Statistics supplied by New Zealand Food Safety Authority in Whangarei.

- Beentjes M. 2010. Toheroa survey of Oreti Beach, 2009, and review of historical surveys. 40p.
- Bell S.S., Tewfik A., Hall M.O., Fonseca M.S. 2008. Evaluation of seagrass planting and monitoring techniques: Implications for assessing restoration success and habitat equivalency. *Restoration Ecology* (16): 407-416.
- Bulleri F. 2006. Is it time for urban ecology to include the marine realm? *Trends in Ecology & Evolution* (21): 658-659.
- Bulleri F., Chapman M. 2010. The introduction of coastal infrastructure as a driver of change in marine environments. *Journal of Applied Ecology* (47): 26-35.
- Carter L. 1975. Sedimentation on the continental terrace around New Zealand: A Review. *Marine Geology* (19): 209-237.
- Carter L., Carter R., McCave I., Gamble J. 1996. Regional sediment recycling in the abyssal Southwest Pacific Ocean. *Geology* (24): 735-738.
- Castro P., Huber M. 2003. *Marine Biology*. (Vol) 4. New York, McGraw Hill Higher Education.
- Cloern J. 2001. Our evolving conceptual model of the coastal eutrophication problem *Marine Ecology Progress Series* (210): 223-253.
- Cornelisen, C., Gillespie, P., Kirs, M., Young, R., Forrest, R., Barter, P., Knight, B., Harwood, V., 2011. Motueka River plume facilitates transport of ruminant faecal contaminants into shellfish growing waters, Tasman Bay, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 45, 477-495.
- Cummings V., Hewitt J., Halliday J., Mackay G. 2007. Optimizing the success of *Austrovenus stutchburyi* restoration: preliminary investigations in a new Zealand estuary. *Journal of Shellfish Research* (26): 89-100.
- Davies-Colley R.J., Meleason M.A., Hall G.M.J., Rutherford J.C. 2009. Modelling the time course of shade, temperature, and wood recovery in streams with riparian forest restoration. *New Zealand Journal of Marine and Freshwater Research* (43): 673-688.
- Diaz R.J., Rosenberg R. 2008. Spreading dead zones and consequences for marine ecosystems. *Science* (321): 926-929.
- Elliott A., Alexander R., Schwarz G., Shankar U., Sukias J., McBride G. 2005. Estimation of nutrient sources and transport for New Zealand using the hybrid mechanistic-statistical model SPARROW. *Journal of Hydrology: New Zealand* (44): 1-27.
- Elliott M., Burdon D., Hemingway K.L., Apitz S.E. 2007. Estuarine, coastal and marine ecosystem restoration: Confusing management and science - A revision of concepts. *Estuarine Coastal and Shelf Science* (74): 349-366.
- Environmental Communications Limited 2010. Integrated catchment management - a review of literature and practice. A report for the Ministry for the Environment. . 158p.
- Forrest B., Gillespie P., Cornelisen C., Rogers K. 2007. Multiple indicators reveal river plume influence on sediments and benthos in a New Zealand coastal embayment. *New Zealand Journal of Marine and Freshwater Research* (41): 13-24.
- Frew R., Hunter K., Beyer R. 1996. Cadmium in Oysters and Sediments From Foveaux Strait, New Zealand. *Proceedings of the Trace Element Group of New Zealand*, 1996 Waikato University.,
- Fukunaga A., Anderson M.J., Webster-Brown J.G., Ford R.B. 2010. Individual and combined effects of heavy metals on estuarine infaunal communities. *Marine Ecology-Progress Series* (402): 123-136.
- GESAMP 2001. Protecting the oceans from land-based activities. Land-based sources and activities affecting the quality and uses of the marine, coastal and associated freshwater environment. Nairobi, United Nations Environment Program. 168.
- Glade T. 2003. Landslide occurrence as a response to land use change: A review of evidence from New Zealand. *Catena* (51): 297-314.
- Goff J.R. 1997. A chronology of natural and anthropogenic influences on coastal sedimentation, New Zealand. *Marine Geology* (138): 105-117.
- Grange K., Tovey A., Hill A.F. 2003. The spatial extent and nature of the bryozoan communities at Separation Point, Tasman Bay. 22 p.
- Griffiths G., Glasby G. 1985. Input of River-derived sediments to the New Zealand Continental Shelf: I. Mass. *Estuarine, Coastal and Shelf Science* (21): 773-784.
- Halpern B., Selkoe K., Micheli F., Kappel C. 2007. Evaluating and ranking the vulnerability of global marine ecosystems to anthropogenic threats. *Conservation Biology* (21): 1301-1315.
- Haro A., Richkus W., Whalen K., Hoar A., Busch W.D., Lary S., Brush T., Dixon D. 2000. Population decline of the American eel: Implications for research and management. *Fisheries* (25): 7-16.
- Hayward B.W., Grenfell H.R., Sabaa A.T., Morley M.S., Horrocks M. 2006. Effect and timing of increased freshwater runoff into sheltered harbor environments around Auckland City, New Zealand. *Estuaries and Coasts* (29): 165-182.
- Hewawasam T., von Blanckenburg F., Schaller M., Kubik P. 2003. Increase of human over natural erosion rates in tropical highlands constrained by cosmogenic nuclides. *Geology* (31): 597-600.
- Hicks, D., Shankar, U., McKerchar, A.I., Basher, L., Jessen, Lynn, I., Page, M., 2011. Suspended sediment yields from New Zealand rivers. *Journal of Hydrology (NZ)* 50, 81-142.
- Hicks, D., Shankar, U., 2003. Sediment from New Zealand rivers, NIWA Chart, Miscellaneous Series No.79.
- Hill, D.; Dubey, J.P. 2002: *Toxoplasma gondii*: transmission, diagnosis and prevention. *Clinical Microbiology and Infection* 8(10): 634–640.
- Hooker S., Redfearn P. 1998. Preliminary survey of toheroa (*Paphies ventricosa*) populations on Ninety Mile Beach and possible impacts of vehicle traffic. NIWA Client Report: AK98042. 34p.
- IPCC 2007. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Pachauri R. and Reisinger A.E., eds. Geneva, Switzerland., pp 104.
- Kennish M.J. 2002. Environmental threats and environmental future of estuaries. *Environmental Conservation* (29): 78-107.
- Kneitel J.M., Perrault D. 2006. Disturbance-induced changes in community composition increase species invasion success. *Community Ecology* (7): 245-252.
- Lipcius R.N., Eggleston D.B., Schreiber S.J., Seitz R.D., Shen J., Sisson M., Stockhausen W.T., Wang H.V. 2008. Importance of metapopulation connectivity to restocking and restoration of marine species. *Reviews in Fisheries Science* (16): 101-110.
- Lundquist C.J., Oldman J.W., Lewis M.J. 2009. Predicting suitability of cockle *Austrovenus stutchburyi* restoration sites using hydrodynamic models of larval dispersal. *New Zealand Journal of Marine and Freshwater Research* (43): 735-748.
- Marsden I., Adkins S. 2009. Current status of cockle bed restoration in New Zealand *Aquaculture international* ISSN: 0967-6120 (Print) 1573-0143X (Online).
- Milliman J., Syvitski J. 1992. Geomorphic/Tectonic Control of Sediment Discharge to the Ocean: The Importance of Small Mountainous Rivers. *The Journal of Geology* (100): 525-544.
- Moller J., Moller S., Futter J., Moller J., Harvey J., White H., Stirling F., Moller H. 2009. Potential impacts of vehicle traffic on recruitment of Toheroa (*Paphies ventricosa*) on Oreti Beach, Southland, New Zealand. Dunedin, University of Otago. 61p.
- Morrisey D.J. 2000. Predicting impacts and recovery of marine farm sites in Stewart Island New Zealand, from the Findlay-Watling model. *Aquaculture* (185): 257-271.
- Morrison M., Lowe M., Parsons D., Usmar N., McLeod I. 2009. A review of land-based effects on coastal fisheries and supporting biodiversity in New Zealand. 100p.
- Morrison M., Parkinson D. 2008. Distribution and abundance of toheroa (*Paphies ventricosa*) on Ninety Mile Beach, March 2006. *New Zealand Fisheries Assessment Report 2008/26*, ed. 27p.

- Nellemann C., Hain S., Alder J.E. 2008. In Dead Water - Merging of climate change with pollution, over-harvest, and infestations in the world's fishing grounds. United Nations Environment Programme, GRID-Arendal., 64p.
- Parkyn S., Davies-Colley R., Halliday N., KJ C., GF C. 2003. Planted Riparian Buffer Zones in New Zealand: Do They Live Up to Expectations? *Restoration Ecology* (11): 436-447.
- Piola R.F., Johnston E.L. 2008. Pollution reduces native diversity and increases invader dominance in marine hard-substrate communities. *Diversity and Distributions* (14): 329-342.
- Roni P., Hanson K., Beechie T. 2008. Global review of the physical and biological effectiveness of stream habitat rehabilitation techniques. *North American Journal of Fisheries Management* (28): 856-890.
- Seaman W. 2007. Artificial habitats and the restoration of degraded marine ecosystems and fisheries. *Hydrobiologia* (580): 143-155.
- SPREP 2010. South Pacific Regional Environment Programme, Strategic Plan 2011-2015: Draft for comment. available at [www.sprep.org](http://www.sprep.org), ed. 29p.
- Stephenson G. 1999. Vehicle impacts on the biota of sandy beaches and coastal dunes: A review from a New Zealand perspective. *Science for Conservation* (121): 48p.
- Stewart M.J., Creese R.G. 1998. Restoration of the New Zealand little necked clam, *Austrovenus stutchburyi*. *Journal of Shellfish Research* (17): [np].
- Taylor F.J., Taylor N.J., Walsby J.R. 1985. A bloom of planktonic diatom *Ceratulina pelagica* off the coastal northeastern New Zealand in 1983, and its contribution to an associated mortality of fish and benthic fauna. *Internationale Revue der gesamten Hydrobiologie und Hydrographie*. 70: 773-795.
- Thrush S., Hewitt J., Cummings V., Ellis J., Hatton C., Lohrer A., Norkko A. 2004. Muddy waters: elevating sediment input to coastal and estuarine habitats. *Frontiers in ecology and the environment* (2): 299-306.
- Thrush S.F., Hewitt J.E., Hickey C.W., Kelly S. 2008. Multiple stressor effects identified from species abundance distributions: Interactions between urban contaminants and species habitat relationships. *Journal of Experimental Marine Biology and Ecology* (366): 160-168.
- Whetton P., Mullan A.B., Pittock A. 1996. Climate change scenarios for Australia and New Zealand. In: Bouma W., Pearman G. and Manning M. (Eds.) *Greenhouse: coping with climate change*. CSIRO Publishing, Collingwood, Australia. 145-168.
- Wilkinson, R., McKergow, L., Davies-Colley, R., Ballantine, D., Young, R., 2011. Modelling storm-event *E. coli* pulses from the Motueka and Sherry Rivers in the South Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 45, 369-393.

## **THEME 5: MARINE BIODIVERSITY**

## 11. Biodiversity

<i>Scope of chapter</i>	Provide an overview of the MPI Biodiversity Programme and address: National and global context of NZ marine biodiversity research; Research findings and progress of the MPI Biodiversity Research Programme from 2000–2012; including one-off whole-of-government research initiatives administered under this programme (e.g. Ocean Survey 20/20 Biodiversity and Fisheries projects; International Polar Year Census of Antarctic Marine life project 2007)
<i>Geographic area</i>	New Zealand Territorial Seas, EEZ and Continental shelf extension (BioInfo); South-west Pacific Region associated with South Pacific Regional Fisheries Management Organisation (SPRFMO); Antarctic Ross Sea region (BioRoss)
<i>Focal issues</i>	New Zealand waters have globally significant levels of marine biodiversity, and productivity particularly coastal habitats, offshore island habitats and underwater topographical features such as seamounts, and canyons. With the exception of shallow sea ice impacted coastal habitats, these features apply also to the Ross Sea region. Adjacent international waters in the SPRFMO area contain areas likely to constitute Vulnerable Marine Ecosystems (VMEs),
<i>Key progress 2011-12</i>	<ul style="list-style-type: none"> <li>• Predictive habitat modelling has identified potential areas of VMEs in SPRFMO areas</li> <li>• Significant progress has been made on mapping deepsea fisheries habitat at risk from ocean acidification; research on shellfish has identified thermal stress and ocean acidification as two areas of concern for New Zealand in an increasing CO<sub>2</sub> world.</li> <li>• Progress has been made towards developing a national Marine Environmental Monitoring Programme</li> <li>• A major project on changes in marine shelf systems over the past 1000 years has almost reached completion.</li> <li>• IPY and Chatham Challenger completed with many outputs and leveraging opportunities</li> </ul>
<i>Emerging issues</i>	<ul style="list-style-type: none"> <li>• The combined effects of multiple stressors arising from climate change and a range of other anthropogenic activities on biodiversity and marine ecosystems (structure and function) are likely to be large and complex.</li> <li>• Keen interest in the development of ecosystem approaches to marine resource management is developing.</li> <li>• The nature and functional role of marine microbial biodiversity in large scale biogeochemical and ecosystem processes are important but not well understood.</li> <li>• Genetic and life-history stage connectivity between and within large scale habitats may be important to the size and placement of protection zones.</li> <li>• Apart from fisheries data, long-term (eg decadal to millenia) observations of variability and change in the marine environment (including biodiversity) are not yet generally available at geographic scales appropriate for national reporting .</li> <li>• Metrics for assessing the effectiveness of current protection measures in safeguarding marine biodiversity and aquatic ecosystem health in NZ and Ross Sea region are inadequate.</li> <li>• Economic value of ecosystem goods and services provided by marine biodiversity to current and future generations are not addressed in extractive business models.</li> <li>• Marine biodiversity and its monitoring, loss reduction and enhancement</li> </ul>

	<p>are emerging requirements for signatories (including New Zealand) to the CBD Aichi-Nagoya Agreement 2010</p> <ul style="list-style-type: none"> <li>• Geo-engineering methods including ocean fertilisation continues to be advocated in some areas of international climate change mitigation</li> <li>• Meeting New Zealand responsibilities participate in international data collection programmes, e.g., IMOS, SOCPR ARGO, BIO-ARGO,</li> </ul>
<i>MPI Research (current)</i>	<p>55 biodiversity projects commissioned over the period 2000-12; Currently in 4th year of a 5 year programme to address seven science objectives in the Biodiversity Programme: 1 characterisation and description; 2 ecosystem scale biodiversity; 3 functional role of biodiversity; 4 genetics; 5 ocean climate effects; 6 indicators; 7 threats to biodiversity. MPI biodiversity research has strong synergies with marine research funded by MPI Aquatic and Environment Working Group (AEWG), Ministry of Business Innovation and Employment (MBIE), Department of Conservation (DOC), Land Information New Zealand (LINZ), other sections within the Ministry for Primary Industries (MPI), Ministry for the Environment (MfE), Statistics New Zealand (Stats NZ), Te Papa and Crown Research Institutes</p>
<i>NZ Research and associated initiatives (current)</i>	<p>Research programmes and database initiatives on Marine Biodiversity are run at University of Auckland (World Register of Marine Species (WoRMS), marine reserves, rocky reef ecology, Ross Sea meroplankton, genetics); Auckland University of Technology, University of Waikato (soft sediment functional ecology and biodiversity), Victoria University of Wellington (monitoring marine reserves, population genetics), University of Canterbury (intertidal and subtidal ecology, kelp forests and biodiversity), University of Otago (land-use effects, bryozoans, inshore ecology, ocean acidification), National Institute of water and Atmospheric Research (NIWA) and Cawthron Institute. Former MBIE programmes i.e., Coasts &amp; Oceans OBI C01X0501, Marine Biodiversity &amp; Biosecurity OBI C01X0502, are now part of Core Funding managed by NIWA through the Coast and Oceans Centre; Protecting Ross Sea Ecosystems C01X1001, Climate Change Effects in the Ross Sea C01X1226, Coastal Conservation Management C01X0907, Impacts of resource use on vulnerable deep-sea communities C01X0906; DOC, MPI, NIWA and Landcare Research - NZ Organisms Register.</p>
<i>Links to Fisheries 2030 and MPI's Our Strategy 2030</i>	<p>Fisheries 2030 Environmental Outcome Objective 1; environmental principles of Fisheries 2030 include: Ecosystem-based approach, Conserve biodiversity: Environmental bottom lines, Precautionary approach, Responsible international citizen, Inter-generational equity, Best available information, Respect rights and interests (MPI 2009). MPI's Strategy "Our Strategy 2030": two key stated focuses are to maximise export opportunities and improve sector productivity; increase sustainable resource use, and protect from biological risk</p>
<i>Related chapters/issues</i>	<p>Multiple use, land-based effects, variability and change, marine monitoring, cumulative effects of use and extraction in the marine environment, protected areas; benthic impacts, ecosystem approaches to fisheries and marine resource management.</p>

## 11.1. Introduction

This chapter summarises the development and progress of the MPI Marine Biodiversity Research Programme 2000-2012 and reviews the work commissioned in the context of national and global concerns about biodiversity and the maintenance of the marine ecosystem in a healthy functioning state, as identified by the New Zealand Biodiversity Strategy (NZBS, Anon 2000).

### 11.1.1. Halting the decline in biodiversity

In June 2000, the '*New Zealand Biodiversity Strategy– Our Chance to Turn the Tide*' (NZBS) with the over-arching objectives “to halt the decline of biodiversity in New Zealand and protect and enhance the environment” was launched as part of New Zealand’s commitment to the international Convention on Biological Diversity 1993 (Anon 2000). To meet long-term goals of the NZBS, a comprehensive plan, with stated objectives and actions<sup>1</sup>, was developed to address biodiversity issues in terrestrial, freshwater and marine systems. The Desired Outcomes by 2020 for the marine environment (Coasts and Oceans, Theme 3) in the NZBS were stated as:

- “New Zealand's natural marine habitats and ecosystems are maintained in a healthy functioning state and degraded marine habitats are recovering.
- A full range of marine habitats and ecosystems representative of New Zealand's indigenous marine biodiversity is protected.
- No human-induced extinctions of marine species within New Zealand's marine environment have occurred.
- Rare or threatened marine species are adequately protected from harvesting and other human threats, enabling them to recover.
- Marine biodiversity is appreciated, and any harvesting or marine development is done in an informed, controlled and ecologically sustainable manner.”

In the marine environment, biodiversity decline is characterised not only by extinctions or reduction in species richness and abundance, but also by environmental degradation such as species invasion and hybridisations, habitats that have been diminished or removed, and the disruption of ecosystem structure and function, as well as ecological processes (e.g. biological cycling of water, nutrients and energy). Measuring the decline of marine biodiversity is complicated by the ‘shifting baseline syndrome’, a common obstacle to useful biodiversity assessment and monitoring<sup>2</sup>. Furthermore the size range of organisms sampled is often limited to macroscopic. Changes (declines) in biodiversity metrics at a macroscopic level may not detect potentially large changes in biodiversity in smaller sized organisms below our sampling threshold that may also be critical to marine ecosystem health and well-being.

Responsibility for addressing Theme 3 of the Biodiversity Strategy was allocated across government departments with active roles in the management of the marine environment, including the Department of Conservation (DOC), the Ministry for Environment (MfE), and the Ministry of Fisheries (now MPI)<sup>3</sup>.

---

<sup>1</sup> The New Zealand Biodiversity Strategy with its stated goals, objectives and actions can be viewed at <http://www.biodiversity.govt.nz>

<sup>2</sup> A National Approach to Addressing Marine Biodiversity Decline (Australian Government-available on line at [www.environment.gov.au/coasts/publications/marine-diversity-decline/index.html](http://www.environment.gov.au/coasts/publications/marine-diversity-decline/index.html)

<sup>3</sup> <https://www.biodiversity.govt.nz/picture/doing/programmes/index.html>

### 11.1.2. Defining biodiversity

New Zealand's Biodiversity Strategy defines biodiversity as:

*“The variability among living organisms from all sources including inter alia, terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are a part [as defined by the CBD]; this includes diversity within species, between species and of ecosystems [as further disaggregated for New Zealand purposes]. Components include:*

- *Genetic diversity: the variability in the genetic make-up among individuals within a single species. In more technical terms, it is the genetic differences among populations of a single species and those among individuals within a population.*
- *Species diversity – the variety of species—whether wild or domesticated— within a particular geographic area.*
- *Ecological diversity – the variety of ecosystem types (such as forests, deserts, grasslands, streams, lakes wetlands and oceans) and their biological communities that interact with one another and their non-living environments.”*

MPI's Biodiversity programme is concerned primarily with research to underpin NZBS Theme 3: Biodiversity in Coastal and Marine Ecosystems:

*“Coastal and marine ecosystems include estuaries, inshore coastal areas and offshore areas, and all the resident and migratory marine species that live in them.*

New Zealand's ocean territory (including territorial sea and the recent continental shelf extension<sup>4</sup>) is very large relative to the area of land<sup>5</sup> and includes some 15-18,000 kilometres of coastline extending from the sub-tropical north to the cool Subantarctic waters to the south. New Zealand also has a rich marine biodiversity that has been recognised as being globally significant with up to 44% estimated as endemic and comprising up to 10% of global marine biodiversity Gordon *et al.* 2010.

An estimated 34,400 marine species and associated ecosystems around New Zealand deliver a wide range of environmental goods and services that sustain considerable fishing, aquaculture and tourism industries as well as drive major biogeochemical and ecological processes. Several factors would suggest that this estimate of marine species number is conservative. Such factors include the region's size, the depth range, geomorphological and hydrological complexity as well as limited water column sampling and limited benthic sampling, especially below 1500 metres. If recent indications of massive oceanic microbial diversity are taken into account (e.g. Sogin *et al.* 2006) then the number above is certainly conservative.

New Zealand's marine biodiversity is affected by many uses of the marine environment, particularly fishing, aquaculture, shipping, petroleum and mineral extraction, renewable energy, tourism and recreation<sup>6</sup>. Impacts from changing land use, including agricultural, urban run-off and coastal development can also affect marine biodiversity (Morrison *et al.* 2009). The potential loss of marine biodiversity and possible functionality caused by climate change and ocean acidification are of increasing concern worldwide (e.g., Guinotte *et al.*, 2006; Ramirez-Llodra *et al.* 2011; as well as in New Zealand—see NZ Royal Society Workshop papers<sup>7</sup>). The growing arrival of non-indigenous

<sup>4</sup> <http://www.mfat.govt.nz/Treaties-and-International-Law/04-Law-of-the-Sea-and-Fisheries/NZ-Continental-Shelf-and-Maritime-Boundaries.php>

<sup>5</sup> NZ sea area is ~5.8 million km<sup>2</sup> including TS, EEZ and continental shelf extension; 4<sup>th</sup> largest in the world; [www.lin.govt.nz](http://www.lin.govt.nz)

<sup>6</sup> <http://www.royalsociety.org.nz/media/Future-Marine-Resource-Use-web.pdf>

[http://www.stats.govt.nz/browse\\_for\\_stats/environment/natural\\_resources/fish.aspx](http://www.stats.govt.nz/browse_for_stats/environment/natural_resources/fish.aspx)

<sup>7</sup> : <http://www.royalsociety.org.nz/publications/policy/yr2009/ocean-acidification-workshop/>

(sometimes invasive) marine species is also a threat to local biodiversity (e.g., Coutts and Dodgshun 2003, Cranfield *et al.* 2003, Gould *et al.* 2008 Russel *et al.* 2008, Williams *et al.* 2008).

Understanding about New Zealand's coastal marine environment and its land-sea interactions has progressed although knowledge about the state of the marine environment and marine biodiversity on a national scale remains limited. Current knowledge about New Zealand's and the Ross Sea's marine biodiversity suggests that it may generally be in better shape than that of many other countries (Costello *et al.* 2010, Gordon *et al.* 2010). However, New Zealand is less well placed when it comes to understanding the threats to marine biodiversity (Costello *et al.* 2010, MacDiarmid *et al.* 2012) and the nature of their impacts. There are significant concerns with the decline of some key species (MfE 2007), localised impacts on habitats and conditions (Thrush and Dayton 2002, Cryer *et al.* 2002, Clark *et al.* 2010a., Gordon *et al.* 2010, Clark & Dunn 2012) and emerging threats to the marine environment (MacDiarmid *et al.* 2012) despite the combined efforts of New Zealand's government and stakeholders. Global scale threats associated with the potential effects of ocean acidification on microbial diversity and their roles in biogeochemical processes have yet to be quantified but could have EEZ wide implications (Bostock *et al.* 2012).

New Zealanders increasingly value environmental, economic and social aspects of marine biodiversity and the ecosystem services that a healthy marine environment provides. They also value the need to sustainably manage the use of coastal and marine environments and maintain biological diversity as reflected by recent policy statements by the New Zealand Government.<sup>8,9</sup> A broad range of legislation, regulations and policies are in place to manage and regulate uses of the marine environment, to protect marine biodiversity, to improve management of the coastal and marine environment and to meet world-wide consumer demands for improved sustainability. The most recent introduction is the Exclusive Economic Zone and Continental Shelf (Environmental Effects) Act 2012 which will come into effect once the first set of regulations is promulgated. However, progress on an integrated oceans policy and strategic direction for implementation of New Zealand's Biodiversity Strategy has been slow compared with other countries such as Canada, the UK, the USA and Australia (Peart *et al.* 2011).

### 11.1.3. Implementation of New Zealand's Biodiversity Strategy

A number of initiatives have been supported by MPI to meet the goals of the NZBS. Commitments include the creation of NABIS (the National Aquatic Biodiversity Information System)<sup>10</sup>, the administration of the MPI Biodiversity Research Programme, convening and chairing the Biodiversity Research Advisory Group<sup>11</sup>, and developing a Marine Protected Area policy with DOC. DOC also surveys and monitors aspects of marine biodiversity, particularly in marine reserves<sup>12</sup>. MfE has encouraged Regional Councils to develop coastal monitoring programmes and with MPI and DOC, initiated an approach to Marine Environmental Classification<sup>13</sup>. Biodiversity related research has also been carried out through MPI's Biosecurity Science Strategy. One result includes mapping and valuation of marine biodiversity around New Zealand's coastline<sup>14</sup>.

---

<sup>8</sup> MfE Proposed National Policy Statement on Indigenous Biological Diversity (biodiversity) under the Resource Management Act 1991 [www.mfe.govt.nz/publications/biodiversity/indigenous-biodiversity/proposed-national-policy-statement/statement.pdf](http://www.mfe.govt.nz/publications/biodiversity/indigenous-biodiversity/proposed-national-policy-statement/statement.pdf)

<sup>9</sup> New Zealand Coastal Policy Statement 2010 [www.doc.govt.nz/conservation/marine-and-coastal/coastal-management/nz-coastal-policy-statement/](http://www.doc.govt.nz/conservation/marine-and-coastal/coastal-management/nz-coastal-policy-statement/)

<sup>10</sup> NABIS is an interactive database accessible at [www.nabis.govt.nz](http://www.nabis.govt.nz)

<sup>11</sup> [www.fish.govt.nz/en-nz/Research+Services/Background+Information/Biodiversity+background.htm](http://www.fish.govt.nz/en-nz/Research+Services/Background+Information/Biodiversity+background.htm)

<sup>12</sup> [www.doc.govt.nz](http://www.doc.govt.nz)

<sup>13</sup> [www.mfe.govt.nz/issues/biodiversity/initiatives/marine.html#regional](http://www.mfe.govt.nz/issues/biodiversity/initiatives/marine.html#regional)

<sup>14</sup> [www.biosecurity.govt.nz/biosec/research](http://www.biosecurity.govt.nz/biosec/research)

Marine biodiversity research is also supported through public good funding and is conducted mainly by Universities and CRIs. Both have contributed to New Zealand's high profile on the international scientific network for marine biodiversity through participation in global initiatives such as the Census of Marine Life as well as to local programmes that have improved understanding of the role of biodiversity in the marine ecosystem. The Museums of Auckland, Canterbury, Otago and Wellington (Te Papa) also conduct biodiversity sampling expeditions and national collections of specimens have been set up within Museums and also at NIWA. Regional Councils give effect to NZBS; Coastal Biodiversity Policy Statement 2011, protected areas and spatial planning.

#### **11.1.4. New challenges and agendas**

Since the launch of the Biodiversity Strategy, there have been substantial changes in Government goals for New Zealand. In July 2009, the Minister of Science set an overarching goal for research science and technology<sup>15</sup>:

“to improve New Zealand’s economic performance while continuing to strengthen our society and protect our environment”.

This goal is reflected in first progress report on “Building Natural Resources” as part of the Business Growth Agenda<sup>16</sup> released December 2012. The Business Growth Agenda sets an ambitious goal of increasing the ratio of exports to GDP to 40% by 2025. Meeting the target will require the value of our exports to double in real terms by 2025. The report states that one of the goals is to “Make the most of the considerable opportunities for New Zealand to gain much greater value from its extensive marine and aquaculture resources”.

The biological economy of the sea (currently largely fisheries and aquaculture, oil and gas, minerals) is a significant part of the overall economy and may have potential for growth (e.g. unlocking the potential of the fisheries sector–Fisheries 2030 (MPI 2009<sup>17</sup>). It is essential that the aquatic environment and biodiversity on which industry depends are not adversely affected by these or other impacting activities.

Bodies such as the Marine Stewardship Council (MSC<sup>18</sup>) require fisheries to satisfy stringent environmental requirements to achieve certification. Many fisheries management systems throughout the globe have begun to develop policies that are ecosystem based. Implementation has met with varied success, and measurement of success is a challenge.

The large scale threats to the marine environment posed by increasing global impacts of anthropogenic stressors such as climate change and ocean acidification, increasing exploitation of resources (living or non-living) and the cumulative effect of multiple uses of the marine environment (e.g., renewable energy, commercial fisheries, recreational fisheries, aquaculture, hydrocarbon and mineral extraction) remain.

Scientific research has provided information about the predicted distribution and abundance of marine biodiversity in some areas of New Zealand’s coasts and oceans, but progress on validation in areas that remain unsampled has been slow. The structure and function of biodiversity of macrofauna within some New Zealand and Ross Sea marine ecosystems is well understood and available information has

---

<sup>15</sup> MoRST feedback document on New Zealand’s research science and technology:  
[www.morst.govt.nz/Documents/publications/policy](http://www.morst.govt.nz/Documents/publications/policy)

<sup>16</sup> <https://www.mbie.govt.nz/what-we-do/business-growth-agenda/pdf-folder/BGA-Natural-Resources-report-December-2012.pdf>

<sup>17</sup> MFish (2009). Fisheries 2030 report. New Zealanders maximising benefits from the use of fisheries within environmental limits available from <http://www.fish.govt.nz/en-nz/Fisheries+2030/default.htm>

<sup>18</sup> Marine Stewardship Council [www.msc.org](http://www.msc.org)

been used to assess the habitat types at greatest risk from disturbance, particularly fishing. However, the proportions of marine habitat types should be or can be protected to maintain a healthy aquatic environment is unknown.

There is growing awareness of the likely importance of the huge diversity, biomass and species mix of micro-organisms, nano- and pico-plankton, and is a fast developing field of research. The rate of change and the resilience of biodiversity to the cumulative effect of multiple stressors across large spatial scales (e.g. ocean acidification, temperature increase and oxygen depletion), particularly as utilisation of marine resources increases, remain semi-quantified (Ramirez-Llodra *et al.* 2011). Understanding the dynamics of climate change and predicting the impacts on food webs and fisheries are only just being investigated (e.g., Fulton 2004, Brown *et al.* 2010, Garcia and Rosenberg 2010).

## **11.2. Global understanding and developments**

In April 2002, the Parties to the Convention on Biological Diversity (CBD) committed to achieve by 2010 a significant reduction of the current rate of biodiversity loss at the global, regional and national level as a contribution to poverty alleviation and to the benefit of all life on Earth. This target was subsequently endorsed by the World Summit on Sustainable Development and the United Nations General Assembly and was incorporated as a target under the Millennium Development Goals<sup>19</sup>.

The third edition of the Global Biodiversity Outlook confirmed that the 2010 biodiversity target had not been met, and the CBD 2010 Strategic Plan notes that “actions [to achieve the 2010 target] have not been on a scale sufficient to address the pressures on biodiversity<sup>20</sup>. Moreover there has been insufficient integration of biodiversity issues into broader policies, strategies, programmes and actions, and therefore the underlying drivers of biodiversity loss have not been significantly reduced”. The Strategic Plan includes a new series of targets for 2020 under the heading “*Taking action now to decrease the direct pressures on biodiversity*”. The Strategic Plan for 2011–2020 was updated, revised and adopted by over 200 countries, including New Zealand<sup>21</sup>.

The eleventh meeting of the Conference of the Parties to the Convention on Biological Diversity (held 8-19 Oct 2012)<sup>22</sup> generated some agreed outcomes of relevance for New Zealand, in particular:

- There was confirmation that the application of the scientific criteria for EBSAs and the selection of conservation and management measures is a matter for states and relevant inter-governmental bodies but that it is an open and evolving process that should continue to allow ongoing improvement and updating as new information comes to hand
- It was recognised that there was a need to promote additional research and monitoring in accordance with national and international laws, to improve the ecological or biological information in each region with a view to facilitating the further description of the areas described
- There is a tentative schedule of further regional workshops to facilitate the description of areas meeting the criteria for EBSAs.

---

<sup>19</sup> UNEP's work to promote environmental sustainability, the object of Millennium Development Goal 7, underpins global efforts to achieve all of the Goals agreed by world leaders at the Millennium Summit <http://www.unep.org/MDGs/>

<sup>20</sup> [www.cbd.int/2010-target](http://www.cbd.int/2010-target)

<sup>21</sup> draft updated and revised Strategic Plan for the Convention on Biological Diversity for the post-2010 period (UNEP/CBD/WG-RI/3/3) <http://www.cbd.int/nagoya/outcomes/>

<sup>22</sup> <http://www.cbd.int/doc/?meeting=cop-11> UNEP/CBD/COP/11/23 Marine and Coastal Biodiversity: Revised Voluntary Guidelines for the Consideration of Biodiversity in Environmental Impact Assessments and Strategic Environmental Assessments in Marine and Coastal Areas

New Zealand government agencies will need to consider how to update the NZBS to better align with the Aichi Biodiversity targets.

### **11.2.1. The decade of biodiversity 2011-2020**

The United Nations General Assembly at its 65th session declared the period 2011-2020 to be “the United Nations Decade on Biodiversity, with a view to contributing to the implementation of the Strategic Plan for Biodiversity for the period 2011-2020” (Resolution 65/161). It will serve to support and promote implementation of the objectives of the Strategic Plan for Biodiversity and the Aichi-Nagoya Biodiversity Targets. The principal instruments for implementation are to be National Biodiversity Strategies and Action Plans or equivalent instruments (NBSAPs). CBD signatory nations are expected to revise their NBSAPs and to “ensure that this strategy is mainstreamed into the planning and activities of all those sectors whose activities can have an impact (positive and negative) on biodiversity” (<http://www.cbd.int/nbsap/>). Throughout the United Nations Decade on Biodiversity, governments are encouraged to develop, implement and communicate the results of progress on their NBSAPs as they implement the CBD Strategic Plan for Biodiversity.

There are five strategic goals and 20 ambitious yet achievable targets. Collectively known as the Aichi Targets, they are part the Strategic Plan for Biodiversity. The five Strategic Goals are:

- Goal A - Address the underlying causes of biodiversity loss by mainstreaming biodiversity (NBSAPs) across government and society
- Goal B - Reduce the direct pressures on biodiversity and promote sustainable use
- Goal C - Improve the status of biodiversity by safeguarding ecosystems, species and genetic diversity
- Goal D - Enhance the benefits to all from biodiversity and ecosystem services
- Goal E - Enhance implementation through participatory planning, knowledge management and capacity building

Targets 6-11 specifically refer to fisheries and marine ecosystems and are provided in Appendix 1 to this Chapter.

The CBD also calls for renewed efforts specifically on coastal and marine biodiversity: “The road ahead for coastal areas lies in better and more effective implementation of integrated marine and coastal area management in the context of the Convention’s ecosystem approach. This includes putting in place marine and coastal protected areas to promote the recovery of biodiversity and fisheries resources and controlling land-based sources of pollution. For open ocean and deep sea areas, sustainability can only be achieved through increased international cooperation to protect vulnerable habitats and species.”<sup>23</sup> The CBD held regional workshops during 2011 to identify information sources that might inform the location of Ecologically or Biologically Sensitive Areas (EBSAs). New Zealand participated in the SW Pacific workshop, and EBSAs were identified<sup>24</sup> The criteria for identifying EBSAs and Vulnerable Marine Ecosystems as recommended through UNGA and managed by Regional Fisheries Management Organisations<sup>25</sup>. The 2012 SPRFMO Science Working Group noted that the differing approaches to identifying VMEs and EBSAs could lead to conflicts in how areas possible in need of protection are defined.

---

<sup>23</sup> [www.cbd.int/marine/done.shtml](http://www.cbd.int/marine/done.shtml)

<sup>24</sup> [www.cbd.int/doc/meetings/cop/cop-11/official/cop-11-03-en.doc](http://www.cbd.int/doc/meetings/cop/cop-11/official/cop-11-03-en.doc)

<sup>25</sup> [http://www.un.org/Depts/los/consultative\\_process/documents/no4\\_spc2.pdf](http://www.un.org/Depts/los/consultative_process/documents/no4_spc2.pdf)

### 11.2.2. Global marine assessment

The biological diversity of the 72% of the planet covered by seawater is a crucial component of global resource security, ecosystem function and to climate dynamics. The Marine Biodiversity Outlook Reports and Summaries prepared by UNEP's Regional Seas Programme for the 10th Conference of Parties of the Convention on Biological Diversity (CBD) held in 2010 provide the first systematic overview at a sub-global scale of the state of knowledge of marine biodiversity, the pressures it faces currently and the management frameworks in place for addressing those pressures<sup>26</sup>.

The regional reports reflect a poor outlook for the continuing well being of marine biodiversity, which faces increasing pressures in all regions from land sourced pollution, ship sourced pollution and impacts of fishing. These pressures are serious and generally increasing despite measures in place to address them. They are amplified by predicted impacts of ocean warming, acidification and habitat change arising from climate and atmospheric change. Without significant management intervention marine biological diversity is likely to deteriorate substantially in the next 20 years with growing consequences for resource and physical security of coastal nations.

With respect to fisheries, the main findings of the reports are that in most regions fisheries peaked at some point between the mid-1980s and mid-2000s that catch expansion is not possible in many cases and that increased exploitation levels would lead to lower catch levels.

All regions report increases in shipping at levels which generally reflect annual economic growth. All regions report progress in the establishment of Marine Protected Areas but current levels of 1.2% of global ocean surface or 4.3% of continental shelf areas fall far short of the 10% target set by CBD COP7 in 2004. It is likely to be many years before this target is reached. The figures do not include some managed fishery areas that have objectives consistent with multiple sustainable use and overall objectives for conservation but even if these are taken into account the proportion managed with objectives that explicitly address sustainability of biodiversity or ecosystem processes is inadequate. The need to plan and implement ecosystem scale and ecosystem-based management of the seas is urgent.

After many years of international negotiations on the need to strengthen the science-policy interface on biodiversity and ecosystem services at all levels, more than 90 governments (including New Zealand) agreed in April 2012 to officially establish the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES)<sup>27</sup>. It will be a leading global body providing scientifically sound and relevant information to support more informed decisions on how biodiversity and ecosystem services are conserved and used around the world.

The United Nations Conference on Sustainable Development (UNCSD), also known as the Rio+20 Conference (June 2011)<sup>28</sup> had a strong sustainability focus and generated an outcome document entitled "the future we want" which had a section on oceans (para 158 - 177) including:

- Support for the Regular Process of Global Reporting and Assessment of the State of the Marine Environment established under the General Assembly and looked forward to the completion of the first global integrated assessment of the state of the marine environment by 2014<sup>29</sup>.

---

<sup>26</sup> UNEP (2003) Global Marine Assessments: a survey of global and regional marine environmental assessments and related scientific activities. UNEP-WCMC/UNEP/UNESCO-IOC. 132p available online at [www.unep-wcmc.org/resources/publications/ss1/GMA\\_Review.pdf](http://www.unep-wcmc.org/resources/publications/ss1/GMA_Review.pdf)

<sup>27</sup> <http://www.iucn.org/what/>

<sup>28</sup> <http://sustainabledevelopment.un.org/futurewewant.html> Rio +20 outcome document

<sup>29</sup> Integrated assessment of the state of the marine environment by 2014.

[http://www.un.org/depts/los/global\\_reporting/Santiago\\_Regular\\_Proceess\\_Workshop\\_Presentations/GRAME\\_Outline\\_of\\_the\\_First\\_Integrated\\_Assessment\\_Report.pdf](http://www.un.org/depts/los/global_reporting/Santiago_Regular_Proceess_Workshop_Presentations/GRAME_Outline_of_the_First_Integrated_Assessment_Report.pdf)

- The ongoing work of the Ad Hoc Open-ended Informal Working Group on Study Issues Relating to the Conservation and Sustainable Use of Marine Biodiversity Beyond Areas of National Jurisdiction and the wish to, by the end of the 69th session (2014) make a decision about the development of an international instrument under UNCLOS.
- A concern about the health of oceans and marine biodiversity and the work of the IMO and relevant conventions including initiatives like the London Protocol on ocean fertilisation and the global programme of action for the protection of the marine environment from land based activities.
- The Rio+20 outcome also endorsed a process to develop sustainable development goals (to apply to all countries) which will include oceans issues. (This is still in its nascent stage and a clear work programme will be finalised by Sept 2013).

### 11.2.3. Ocean climate change and ocean acidification

Ocean climate change at the global scale overshadows the existing challenges of managing local impacts causing declines in marine biodiversity in the face of current levels of human use and impact. The projected increases in temperature, acidity, severe storm incidence and sea level present major challenges for biodiversity management. This is reflected in changes at the Great Barrier Reef in Australia, which is a globally iconic marine ecosystem that has been subject to adaptive scientifically-based ecosystem-based management for more than 30 years. An Outlook Report by the Great Barrier Reef Marine Park Authority (2009) concluded that “without significant additional management intervention, some components of the ecosystem will deteriorate in the next 20 years and only a few areas are likely to be healthy and resilient in 50 years.” Without strong ecosystem based management the global threats to marine biodiversity may be similar and their implications for food and physical security could be substantial.

The Outlook Report provides a reasonable understanding of the nature and extent of the problems facing marine biodiversity and marine resources. There are examples of effective actions to address some of these problems but management performance is generally insufficient and inadequately coordinated to address the growing problems of marine biodiversity decline and ecosystem change.

Climate change can adversely impact on the spatial patterns of marine biodiversity and ecosystem function through changes in species distributions, species mix and habitat availability, particularly at critical stages of species life histories. A study of the global patterns of climate change impacts on ocean biodiversity projected the distributional ranges of a sample of 1066 exploited marine fish and invertebrates for 2050 using a newly developed dynamic bioclimate envelope model which showed that climate change may lead to numerous local extinctions in the sub-polar regions, the tropics and semi-enclosed seas (Cheung *et al.* 2009). Simultaneously, species invasion is projected to be most intense in the Arctic and the Southern Ocean. With these elements taken together, the model predicted dramatic species turnovers of over 60% of the present biodiversity, implying ecological disturbances that potentially disrupt ecosystem services (Cheung *et al.* 2009).

The World Bank, together with IUCN and Environmental Services Association released a brief for decision-makers entitled, "Capturing and Conserving Natural Coastal Carbon – Building Mitigation, Advancing Adaptation"<sup>30</sup>. This brief highlights the crucial importance of carbon sequestered in coastal wetlands and in submerged vegetated habitats such as seagrass beds, for climate change mitigation.

---

<sup>30</sup> UNFCCC COP-16 event. Cancun Messe, Jaguar. 'Blue Carbon: Valuing CO<sub>2</sub> Mitigation by Coastal Marine Systems. Sequestration of Carbon Along Our Coasts: Are We Missing Major Sinks and Sources?'

The Intergovernmental Panel on Climate Change (IPCC) is preparing material for the 5<sup>th</sup> IPCC Report in 2014 and for the first time includes chapters to explicitly address ocean climate change issues<sup>31</sup>. The Working Group I and Working Group II Contributions to the Fifth Assessment Report include chapters on the ocean (WG I) and Climate Change 2014: Impacts, Adaptation, and Vulnerability including Chapters on Coastal and Oceans ecosystems, and sections on biodiversity(WGII). Working Group I will consider "Ocean biogeochemical changes, including ocean acidification" in their Chapter 3 (Observations - Ocean), and "Processes and understanding of changes, including ocean acidification" in their Chapter 6 on "Carbon and other biogeochemical cycles". Working Group II will consider "Water property changes, including temperature and ocean acidification" in their Chapter 6 on "Ocean Systems". In addition, "Carbon Cycle including Ocean Acidification" has been identified as a "Cross-Cutting Theme" across (predominantly) WG1 and WG2.

Hobday *et al.* (2006) reported on the relative risks and likely impacts of ocean climate change and ocean acidification to marine life in Australian waters (Figure 11.1). This approach was extremely useful for summarising risks and threats of climate change on marine systems to policy makers and the subsequent development of the Commonwealth Environment Research Facilities (CERF) Marine Biodiversity Hub in Australia<sup>32</sup>.

The Hub analysed patterns and dynamics of marine biodiversity through four research programmes to determine the appropriate units and models for effectively predicting Australia's marine biodiversity. These programmes were designed to develop and deliver tools needed to manage Australia's marine biodiversity in a changing ocean climate. The final report from three years intense research is available at the website<sup>33</sup>. Australia also has The Marine Adaptation Network that comprises a framework of five connecting marine themes (integration; biodiversity and resources; communities; markets and policy) that cut across climate change risk, marine biodiversity and resources, socio-economics, policy and governance, and includes ecosystems and species from the tropics to Australian Antarctic waters<sup>34</sup>.

In late June 2011, two science-based reports heightened concerns about the critical state of the world's oceans in response to ocean climate change. One focuses on the potential impacts of ocean acidification on fisheries and higher trophic level ecology and takes a modelling approach to scaling from physiology to ecology (Le Quesne and Pinnegar 2011) and the other assesses the critical state of the world's oceans in relation to climate change and other stressors (Rogers and Laffoley (2011).

### 11.2.1. Census of Marine Life 2000–2010

In 2010, the international initiative to conduct a Census of Marine Life<sup>35</sup> was concluded after ten years of accessing and databasing existing records, sampling and exploration around the globe. The Census is an unprecedented collaboration among researchers from more than 80 nations to assess and explain the diversity, distribution, and abundance of life in the oceans. During the last decade, the 2,700 scientists involved in the Census have mounted 540 expeditions, identified more than 6,000 potentially new species, catalogued upward of 31 million distribution records, and generated 2,600 scientific publications. NIWA scientists were part of the team that led CenSeam<sup>36</sup>, the seamount component of the Census of Marine Life, and scientists from NIWA and the University of Auckland played significant roles in a number of other programmes. The New Zealand IPY-CAML voyage to the Ross Sea in 2008 was a major contribution to CAML.

<sup>31</sup> <http://www.global-greenhouse-warming.com/IPCC-5th-Report.html>

<sup>32</sup> [www.marinehub.org/](http://www.marinehub.org/)

<sup>33</sup> [www.marinehub.org/](http://www.marinehub.org/)

<sup>34</sup> [arnmbr.org/content/index.php/site/aboutus/](http://arnmbr.org/content/index.php/site/aboutus/)

<sup>35</sup> [www.coml.org/results-publications](http://www.coml.org/results-publications)

<sup>36</sup> [www.coml.org/global-census-marine-life-seamounts-censeam](http://www.coml.org/global-census-marine-life-seamounts-censeam)

Groups	Distribution/ Abundance	Phenology	Physiology/ Morphology/ Behaviour	Impacts on biological communities	Examples of impacts
Phytoplankton	High	High	Medium	High	Temperate phytoplankton province will shrink considerably
Zooplankton	High	High	Medium	High	Acidification will dissolve planktonic molluscs
Seagrasses	Medium	Low	High	Medium	Increased dissolved carbon dioxide may increase productivity
Mangroves	Medium	Low	Medium	High	Sea level rise will destroy mangrove habitat
Kelp	High	Medium	High	High	Ranges will shift southwards as SST warms
Rocky reefs	High	Medium	High	Low	Ranges will shift southwards as temperature warms
Coral reefs	High	Medium	High	High	Acidification and warming will cause calcification problems and coral bleaching
Cold water corals	High	Low	Low	High	Ocean acidification will dissolve reefs
Soft bottom dwelling fauna	Medium	Medium	Medium	Medium	Modified plankton communities or productivity will reduce benthic secondary production
Seafloor dwelling and demersal fishes	High	Medium	Medium	High	Southward movement of species along the east and west coast of Australia
Pelagic fishes	Medium	Low	Medium	Low	Pelagic tunas will move south with warming
Turtles	High	Medium	High	Low	Warming will skew turtle sex ratios
Seabirds	Medium	Medium	Low	Low	Shift in timing of peak breeding season as temperatures warm
Total number of high impact habitats or species groups	8	2	5	7	High impacts are expected for distribution, physiology and community processes

**Figure 11.1: Potential biological impacts of climate change on Australian marine life.** The ratings in this table are based on the expected responses to predicted changes in Sea Surface Temperature (SST), salinity, wind, pH, mixed layer depth and sea level, and from literature reviews for each species group. The implicit assumption underlying this table is that Australian marine species will respond in similar ways to their counterparts throughout the world (Hobday *et al.* 2006.) Note: phenology means life cycle.

The Census increased the total number of known marine species by about 20,000, from 230,000 in 2000 to about 250,000 in 2010. Among the millions of specimens collected in both familiar and seldom-explored waters, the Census found more than 6,000 potentially new species and completed formal descriptions of more than 1,200 of them. It also found that some species considered to be rare are more common than previously thought (Ausubel *et al.* 2010). The digital archive (the Ocean Biogeographic Information System OBIS (<http://www.iobis.org/>)) has now grown to 31 million observations, and the Census compiled the first regional and global comparisons of marine species diversity. It helped to create the first comprehensive list of the known marine species, and also helped to compose web pages for more than 80,000 species in the Encyclopaedia of Life<sup>37</sup>.

Applying genetic analysis on an unprecedented scale to a dataset of 35,000 species from widely differing major groupings of marine life, the Census graphed the proximity and distance of relations

<sup>37</sup> [www.eol.org/](http://www.eol.org/)

among distinct species, providing new insight into the genetic structure of marine diversity. With the genetic analysis often called barcoding, the Census sometimes decreased diversity but generally its analyses expanded the number of species, especially the number of different microbes, including bacteria and archaea.

The Census has overwhelmingly demonstrated that the total number of species in the ocean remain largely unknown. The Census also demonstrated that evidence of human impacts on the oceans extends to all depths and habitats and that we still have much to learn to integrate use of resources with stewardship of a healthy marine ecosystem. The Census results could logically extrapolate to at least a million kinds of eukaryotic marine life that earn the rank of species and to tens or even hundreds of millions of kinds of microbes.

A summary of the overall state of knowledge about marine biodiversity after the Census by Costello *et al.* (2010) places New Zealand 6<sup>th</sup> out of 18 national regions based on the collective knowledge assembled by the Census National and Regional Implementation Committees (NRIC) and comparing the Spearman rank correlation coefficients between known diversity (total species richness, alien species, and endemics) and available resources, such as numbers of taxonomic guides and experts. (Figure 11.2).




Figure 11.2: The regions are ranked by their state-of-knowledge index (mean  $\pm$  standard error) across taxa. Dashed line represents the overall mean. (Image Source Costello *et al.* 2010).

All NRICs reported what they considered the main threats to marine biodiversity in their region, citing published data and expert opinions. Although the reports were not standardised, the threats identified were grouped into several overarching issues. We integrated these data on biodiversity threats so as to rank each threat from 1 (very low) to 5 (very high threat) in each region. New Zealand was placed 12<sup>th</sup> out of 18 regions in terms of overall threat levels to biodiversity, overfishing and alien species invasion. Habitat loss and ocean acidification were identified as the biggest threats to marine biodiversity in New Zealand (Costello *et al.* 2010).

### 11.2.2. Global monitoring and indicators for marine biodiversity

There are numerous schemes within and between nations to monitor the marine environment, including physical, chemical and biological components. Marine biodiversity indicators have been developed for the UK and the EU<sup>38</sup>. Marine environmental monitoring networks have been developed in the USA, Canada, Australia and South Africa. Global networks include the Global Ocean Observing System (GOOS) which is a permanent global system for observations, modelling and analysis of marine and ocean variables; Global Climate Observing System (GCOS<sup>39</sup>) which stimulates, encourages, coordinates and otherwise facilitates observations by national or international organizations. A Southern Ocean Observing System (SOOS) is under development.<sup>40</sup>

Others include:

- ARGO an international deepwater monitoring system of free floating buoys that are part of the integrated global observation strategy<sup>41</sup>.
- The Ocean Observation Systems (OOS) in Canada have demonstrated many positive benefits.
- The Continuous Plankton Recorder (CPR) Surveys have been collecting data from the North Atlantic and the North Sea on the ecology and biogeography of plankton since 1931<sup>42</sup>. Sister CPR surveys around the globe include the SCAR SO-CPR Survey established in 1991 by the Australian Antarctic Division to map the spatial-temporal patterns of zooplankton and then to use the sensitivity of plankton to environmental change as early warning indicators of the health of the Southern Ocean. It also serves as reference for other monitoring programs such as CCAMLR's Ecosystem Monitoring Program C- EMP and the developing Southern Ocean Observing System<sup>43</sup>.
- The Marine Environmental Change Network (MECN) is a collaboration between organisations in England, Scotland, Wales, Isle of Man and Northern Ireland collecting long-term time series information for marine waters<sup>44</sup>.
- The MECN has developed links with other networks coordinating long-term data collection and time series. These networks include the Marine Biodiversity and Ecosystem Functioning European Union Network of Excellence (MarBEF<sup>45</sup>) which coordinates long-term marine biodiversity monitoring at a European level.
- New Zealand has now formed a partnership with Australia's Integrated Marine Observing System (IMOS<sup>46</sup>) was established in 2007. IMOS is designed to be a fully integrated national array of observing equipment to monitor the open oceans and coastal marine environment around Australasia, covering physical, chemical and biological variables. All IMOS data is freely and openly available through the IMOS Ocean Portal for the benefit of Australian and New Zealand marine and climate science as a whole. Oceans 2025<sup>47</sup> is an initiative of the Natural Environment Research Council (NERC) funded Marine Research Centres. This addresses environmental issues that require sustained long-term observations.

A challenge for MPI and New Zealand researchers is how to assimilate any or all of the above monitoring approaches as a means of measuring biodiversity baseline levels and the nature and extent

<sup>38</sup> <http://jncc.defra.gov.uk/page-4233>

<sup>39</sup> [www.ioc-goos.org/index.php?option=com\\_content&view=article&id=12&Itemid=26&lang=en](http://www.ioc-goos.org/index.php?option=com_content&view=article&id=12&Itemid=26&lang=en)

<sup>40</sup> <http://www.scar.org/soos/>

<sup>41</sup> <http://www.qc.dfo-mpo.gc.ca/publications/science/evaluation-assessment-eng.asp>.

<sup>42</sup> [www.sahfos.ac.uk/](http://www.sahfos.ac.uk/)

<sup>43</sup> [www.sahfos.ac.uk/sister-survey/sister-surveys/-southern-ocean-continuous-plankton-recorder-survey-\(scar\).aspx](http://www.sahfos.ac.uk/sister-survey/sister-surveys/-southern-ocean-continuous-plankton-recorder-survey-(scar).aspx)

<sup>44</sup> <http://www.mba.ac.uk/MECN/>

<sup>45</sup> <http://www.marbef.org/>

<sup>46</sup> <http://imos.org.au/>

<sup>47</sup> <http://www.oceans2025.org/>

of biodiversity changes, especially as a means of assessing the effectiveness of management measures to protect or enhance biodiversity or halt its decline.

### **11.2.3. Economic valuation of biodiversity**

The national and global responsibility for New Zealand to maintain a strong environmental record in fisheries and other marine-based industries is increasing. There is growing awareness of international treaties and agreements that New Zealand is party to. Global markets are becoming increasingly sensitive to our national environmental record. Fishing companies who meet rigorous standards receive Marine Stewardship Council Certification for certain fisheries (currently, hoki trawl, southern blue whiting pelagic trawl and albacore tuna troll fisheries). Proposals to exploit other living marine resources or extract non-living marine resources are increasingly under scrutiny to ensure that such activities do not adversely degrade the marine environment or impact on marine living resource industries.

The invisibility of biodiversity values has often encouraged inefficient use or even destruction of the natural capital that is the foundation of our economies. A recent international initiative “The Economics of Ecosystems and Biodiversity” (TEEB)<sup>48</sup> demonstrates the application of economic thinking to the use of biodiversity and ecosystem services. This can help clarify why prosperity and poverty reduction depend on maintaining the flow of benefits from ecosystems; and why successful environmental protection needs to be grounded in sound economics, including explicit recognition, efficient allocation, and fair distribution of the costs and benefits of conservation and sustainable use of natural resources. Valuation is seen as a tool to help recalibrate the faulty economic compass that has led to decisions about the environment (and biodiversity) that are prejudicial to both current well-being and that of future generations.

## **11.3. State of knowledge in New Zealand**

The past 750 years of human activity have impacted on marine environments. For example, depletion of fur seals and sea lions occurred from the earliest days of human settlement, not just with European arrival (Smith 2005, 2011). There was also a pulse of sedimentation coinciding with the initial clearance of 40% of NZ forests within 200 years of Polynesian settlement (McWethy *et al.* 2010). Impacts have occurred near population centres, as well as more remote areas and to depths in excess of 1000 metres (MacDiarmid *et al.* 2012, Ministry for Primary Industries 2012). In some cases by looking back over historical records it becomes apparent how much biodiversity loss has occurred. Over long time spans incremental impacts can lead to major shifts in biodiversity composition. An analysis of marine biodiversity decline over a couple of decades could miss the major changes that can occur incrementally over long periods.

While New Zealand has reasonable archaeological, historical and contemporary data on the decline in abundance of individual marine species, in some cases over a period of 750 years (e.g., MacDiarmid *et al.* in press), current trends in the status of New Zealand’s marine biodiversity are difficult to determine for several reasons. These include a lack of both pre-disturbance baseline and recent information, and a lack of a nationally coordinated approach to assessing and monitoring marine biodiversity

A re-evaluation of the threat status of New Zealand's marine invertebrates was undertaken by the Department of Conservation in 2009, and identified no taxa that had improved in threat status as a result of past or ongoing conservation management action, nor any taxa that had worsened in threat

---

<sup>48</sup> TEEB (2010) The Economics of Ecosystems and Biodiversity: Mainstreaming the Economics of Nature: A synthesis of the approach, conclusions and recommendations of TEEB. [www.teebweb.org/](http://www.teebweb.org/)

status because of known changes in their distribution, abundance or rate of population decline (Freeman *et al.* 2010). The authors cautioned however that only a small fraction of New Zealand's marine invertebrate fauna had been evaluated for their threat status and that many taxa remain 'data deficient' or unlisted.

A re-evaluation of marine mammal threat status found that relative to the previous listing, the threat status of two species worsened: the NZ sea lion (*Phocarctos hookeri*) was uplisted to Nationally Critical and the bottlenose dolphin (*Tursiops truncatus*) was uplisted to Nationally Endangered. No species was considered to have an improved status (See Chapter on marine mammals and also Baker *et al.* 2010).

The most recent State of the Environment Report in New Zealand (MfE 2007) covers terrestrial and freshwater organisms in its Biodiversity section<sup>49</sup>. Comment on marine biodiversity is provided in the Oceans section which states:

“Of the almost 16,000 known marine species in New Zealand, 444 are listed as threatened. Well-known species of particular concern include both subspecies of Hector's dolphin, New Zealand sea lion, southern right whale, Fiordland crested penguin, and New Zealand fairy tern. Land-based pressures on the inshore marine environment, as well as pressures on fisheries stocks, can be expected to persist and, therefore, continue to pose a challenge to the health of the marine environment. The increasing number of introduced species brought to New Zealand through marine-based trade and travel, and climate change may exacerbate existing pressures. Further information about our marine environment is needed if we are to help set priorities for future use and protection of our oceans”.

Two major knowledge gaps identified by MfE 2007 that hinder resource management are sparse biodiversity baseline information; and the lack of a systematic national-scale approach to monitoring biodiversity trends (i.e. by comparing subsequent studies to the baseline information) in New Zealand.

The most recent summary of knowledge about marine biodiversity in New Zealand is provided by Gordon (2009, 2010, 2012) and Gordon *et al.* (2010). Figure 11.3 gives a tally of 17,058 living species in the EEZ, including 4,320 known undescribed species in collections.

The Hub analysed patterns and dynamics of marine biodiversity through four research programmes to determine the appropriate units and models for effectively predicting Australia's marine biodiversity. These programmes were designed to develop and deliver tools needed to manage Australia's marine biodiversity in a changing ocean climate. The final report from three years intense research is available at the website<sup>50</sup>. Australia also has The Marine Adaptation Network that comprises a framework of five connecting marine themes (integration; biodiversity and resources; communities; markets and policy) that cut across climate change risk, marine biodiversity and resources, socio-economics, policy and governance, and includes ecosystems and species from the tropics to Australian Antarctic waters<sup>51</sup>.

Species diversity for the most intensively studied animal phyla (Cnidaria, Mollusca, Brachiopoda, Bryozoa, Kinorhyncha, Echinodermata, Chordata) is more or less equivalent to that in the ERMS (European Register of Marine Species) region, an area 5.5 times larger than the New Zealand EEZ (Gordon *et al.* 2010), suggesting that the NZ region biodiversity is proportionately richer than the ERMS region (Figure 11.3).

---

<sup>49</sup> State of the Environment MfE 2007.

<sup>50</sup> [www.marinehub.org/](http://www.marinehub.org/)

<sup>51</sup> [arnmbr.org/content/index.php/site/aboutus/](http://arnmbr.org/content/index.php/site/aboutus/)

Taxonomic group	No. Species <sup>1</sup>	State of knowledge	No. Introduced species	No. Experts	No. ID guides <sup>2</sup>
<b>Domain Prokaryota<sup>3</sup></b>	<b>79</b>	<b>3</b>	<b>&gt;1</b>	<b>3</b>	<b>1</b>
Cyanobacteria	40	2	1	1	1
<b>Domain Eukaryota<sup>3</sup></b>	<b>16,979</b>	<b>3-4</b>	<b>159</b>	<b>58</b>	<b>75</b>
<b>Kingdom Chromista<sup>3</sup></b>	<b>2,643</b>	<b>3-4</b>	<b>11</b>	<b>7</b>	<b>2</b>
Ochrophyta	858	4-5	11	1	1
Myzozoa incl. Dinoflagellata	249	3-4	0	2	0
Foraminifera	1141	4-5	3	2	2
<b>Kingdom Plantae<sup>3</sup></b>	<b>702</b>	<b>4-5</b>	<b>12</b>	<b>12</b>	<b>3</b>
Chlorophyta	156	3-4	0	12	1
Rhodophyta	541	3-4	12	0	1
Tracheophyta	5	5	0	3	2
<b>Kingdom Protozoa<sup>3</sup></b>	<b>43</b>	<b>2-3</b>	<b>4</b>	<b>5</b>	<b>4</b>
<b>Kingdom Fungi</b>	<b>89</b>	<b>3</b>	<b>0</b>	<b>1</b>	<b>0</b>
<b>Kingdom Animalia<sup>3</sup></b>	<b>13,502</b>	<b>3-4</b>	<b>150</b>	<b>40</b>	<b>66</b>
Porifera	742	3	7	1	4
Cnidaria	1,116	4	23	0	6
Platyhelminthes	323	2	2	1	1
Mollusca	3,813	4	14	1	3
Annelida	792	4	32	1	2
Arthropoda (esp. Crustacea)	2,926	3-4	27	13	17
Bryozoa	953	4	24	1	4
Echinodermata	623	5	0	3	6
Tunicata	192	5	0	3	6
Other invertebrates	456	2-5	3	7	12
Vertebrata (Pisces)	1,387	4-5	6	6	7
Other vertebrates	179	5	0	4	4
<b>TOTAL REGIONAL DIVERSITY<sup>3</sup></b>	<b>17,058</b>	<b>3-4</b>	<b>177</b>	<b>62</b>	<b>76</b>
<sup>1</sup> Sources of the tallies: scientific literature, books, field guides, technical reports, museum collections.					
<sup>2</sup> Identification guides cited in Gordon <i>et al.</i> 2010.					
<sup>3</sup> Totals from Gordon 2010, 2012 and Gordon <i>et al.</i> 2010 and unpublished NIWA data.					

**Figure 11.3: Diversity of marine species found in the New Zealand region (after Gordon 2010, 2012; Gordon *et al.* 2010 and current unpublished NIWA data).**

In late June 2011, two science-based reports heightened concerns about the critical state of the world's oceans in response to ocean climate change. One focuses on the potential impacts of ocean acidification on fisheries and higher trophic level ecology and takes a modelling approach to scaling from physiology to ecology (Le Quesne and Pinnegar 2011) and the other assesses the critical state of the world's oceans in relation to climate change and other stressors (Rogers and Laffoley (2011).

In New Zealand, new marine research projects initiated in 2012 include 'Marine Futures' that aims to develop an agreed decision-making framework, enabling participation of all stakeholders (public, iwi, industry, government), that facilitates economic growth, improves marine stewardship and ensures that cumulative stresses placed on the environment do not degrade the ecosystem beyond its ecological adaptive capacity.( C01X1227). The 'Ross Sea Climate & Ecosystem'will model likely future changes in the physical environment of the region and potential consequences of these changes on the ecosystem in terms of functional links between the environment and the marine food web. (C01X1226). 'Management of offshore mining'will develop a clear framework that will guide appropriate and robust environmental impact assessments and the development of integrated environmental management plans for the marine-mining sector, other resource users and resource management agencies. (C01X1228)

Core purpose funding within the Coasts and Oceans Centre at NIWA include "Managing Marine Stressors: Quantifying and predicting the effects of natural variability, climate change and anthropogenic stressors to enable ecosystem-based approaches to the management of New Zealand's marine resources" and within the Fisheries Centre, "Ecosystem Approaches to Fisheries Management: Determine the impact of fisheries on the aquatic environment to inform an ecosystem-based approach

to fisheries management and contribute to broader ecosystem-based management approaches in conjunction with the Coasts & Oceans Centre.

### 11.3.1. The MPI Biodiversity Research Programme

The recognition of increasing societal expectation to use fisheries management measures that will achieve biodiversity conservation has signalled by MPI through Fisheries 2030<sup>52</sup> in its long-term commitment to– “ecosystem based fisheries management” and to ensuring that “biodiversity and the function of ecological systems, including trophic linkages, are conserved”. While New Zealand’s environmental record with regard to fishing is perceived to be relatively high on an international scale, the Ministry is not complacent about the ongoing requirement to monitor and provide evidence that measures to achieve biodiversity conservation needs are being met. This is particularly true of the need to better understand and mitigate the effects of fishing on the areas impacted by fishing. The effects of fishing on the aquatic environment and risks to biodiversity and marine ecosystems are recognised in Fisheries Plans. Research continues to be supported through the Deepwater Research Plan, as well as the Aquatic Environment and Biodiversity Research Programmes.

There are also a range of societal values beyond commercial, customary and recreational take from the sea that are recognised as part of “strengthening our society” (see footnote 12). These include aesthetic and cultural values as well as other economic values such as tourism and marine recreation other than fishing<sup>53</sup>. To link socio-economic values of biodiversity to science supporting fisheries management will require a multi-disciplinary approach only just beginning in New Zealand.

MPI responded to the NZBS in 2000 with the establishment of the MPI Biodiversity Programme which has run successfully for more than 10 years with 55 research projects and a large number of published outputs, presentations and contributions to NZ and CCAMLR management measures.

The Ministry is one of several New Zealand government agencies with a strong interest and a statutory management mandate in the Ross Sea region of Antarctica through the Antarctic Marine Living Resources Act 1981. MPI Antarctic science contributes strongly to New Zealand’s whole-of-government involvement in contributions to the Commission for the Convention on Antarctic Marine Living Resources (CCAMLR) and the Antarctic Treaty. Research conducted under the MPI Antarctic Biodiversity Programme seeks to help New Zealand deliver on its international obligations to support an ecosystem-based approach to management in Antarctic waters. There are strong links with the MPI Antarctic Working Group research and with other Ross Sea ecosystems research carried out under NIWA core purpose Fisheries, and Coast and Oceans Centres (e.g., Sharp *et al.* 2010).

The biodiversity research programme set up under the NZBS was established with a multi-stakeholder biodiversity research advisory group (BRAG), chaired by the former Ministry of Fisheries (now MPI). The research commissioned for the period 2001–2005 reflected goals set by the NZBS and the BRAG, while remaining compatible with the Ministry of Fisheries Statements of Intent (SOIs). During the first three years of this period, MPI also commissioned marine biosecurity research under NZBS, but this was transferred to Biosecurity New Zealand (MAFBNZ) in 2004. From 2006 to 2010, the programme evolved further with the development of a new 5-year work programme to address shortcomings identified in the review of the NZBS by Clark and Green (2006). An overview of the Biodiversity Programme at a glance is given in Figure 11.4.

---

<sup>52</sup> Fisheries 2030 The full document can be downloaded from [www.fish.govt.nz/en-nz/Fisheries+2030](http://www.fish.govt.nz/en-nz/Fisheries+2030)

<sup>53</sup> MARBEF: The Valencia Declaration 2008 [www.marbef.org/worldconference](http://www.marbef.org/worldconference)











	BIODIVERSITY THEMES	KEY QUESTIONS
         	<p><b>BIODIVERSITY PATTERNS &amp; DISTRIBUTION</b></p> <ul style="list-style-type: none"> <li>Fauna and flora (taxonomy, biosystematics)</li> <li>Distribution &amp; abundance of major groups</li> <li>Reviews of existing knowledge</li> <li>Biogeography</li> <li>Drivers of observed patterns</li> </ul> <p><b>HABITAT DIVERSITY</b></p> <ul style="list-style-type: none"> <li>Biogenic reefs</li> <li>Rocky reefs</li> <li>Rhodolith beds</li> <li>Seamounts</li> <li>Soft sediments</li> <li>Habitat mapping EEZ</li> <li>Deepsea habitats</li> <li>Physical and biological characterisation</li> </ul> <p><b>FUNCTIONAL DIVERSITY</b></p> <ul style="list-style-type: none"> <li>The role of different animal/plant groups in the ecosystem</li> <li>Trophic processes</li> <li>Benthic-pelagic processes</li> </ul> <p><b>GENETIC DIVERSITY</b></p> <ul style="list-style-type: none"> <li>Barcode of Life</li> <li>Connectivity (populations, areas)</li> </ul> <p><b>THREATS TO BIODIVERSITY</b></p> <ul style="list-style-type: none"> <li>Climate change and variability</li> <li>Invasive organisms; fishing</li> <li>Land-use effects</li> <li>Cumulative effects</li> </ul> <p><b>METHODS</b></p> <ul style="list-style-type: none"> <li>Measuring biodiversity</li> <li>Classification</li> <li>Predictive modelling</li> <li>Biodiversity indicators</li> <li>Monitoring biodiversity</li> <li>Ecosystem approaches</li> </ul> <p><b>BIOROSS/ &amp; IPY RESEARCH</b></p> <ul style="list-style-type: none"> <li>Bioross coastal biodiversity</li> <li>Subtidal ice-sea interface</li> <li>Census of Antarctic marine Life survey for IPY, Ross Sea</li> <li>Trophic modelling Ross Sea</li> <li>Balleny Islands survey for MPA</li> <li>Functional habitats</li> </ul>	<ul style="list-style-type: none"> <li>What is the abundance and distribution of marine biodiversity in NZ?</li> <li>What are the key drivers of observed patterns in biodiversity?</li> <li>How much marine endemism is there in NZ waters?</li> <li>What is the organism size distribution?</li> <li>How do patterns in biodiversity change over time?</li> </ul> <ul style="list-style-type: none"> <li>What are the relative goods and services offered by each habitat to aquatic environment health?</li> <li>Can the assemblages and biodiversity of marine habitats in the EEZ be predicted by modelling?</li> <li>Which habitats are at greatest risk from extraction practices?</li> <li>What proportion of a given habitat needs to remain intact for healthy ecosystem functioning?</li> </ul> <ul style="list-style-type: none"> <li>How does biodiversity contribute to the resilience of ecosystems to perturbation?</li> <li>Can we use ecosystem function to classify biodiversity?</li> <li>Which key processes need to be retained?</li> </ul> <ul style="list-style-type: none"> <li>What barriers drive connectivity within species?</li> <li>What is the role of endemism in characterising the evolutionary history and taxonomy?</li> </ul> <ul style="list-style-type: none"> <li>What are the key threats?</li> <li>Does biodiversity increase resilience to climate change?</li> <li>Which components of the ecosystem will be most at risk from climate change?</li> </ul> <ul style="list-style-type: none"> <li>How can we best measure and portray biodiversity?</li> <li>How scalable are results from a local scale to an ecosystem scale?</li> <li>What do we need to monitor to measure risks and change to ecosystem health?</li> <li>How can we measure the economic value of biodiversity and ecosystem services?</li> </ul> <ul style="list-style-type: none"> <li>What is the connectivity between biodiversity in the Ross Sea and NZ?</li> <li>How are biota adapted to polar conditions and what is their sensitivity to perturbation?</li> <li>Are MPAs a useful protection tool for the Ross Sea?</li> <li>Are climate change effects on the ocean already impacting on the Ross Sea biota?</li> </ul>

Figure 11.4: Summary of MPI Biodiversity Research Programme 2000–2012.

ACHIEVEMENTS & KNOWLEDGE TO DATE	CURRENT WORK
<ul style="list-style-type: none"> <li>• Taxonomy of coralline algae and bryozoans ( 2 ID Guides)</li> <li>• New species from surveys added to benthic ID Guides</li> <li>• Review of macroalgae distribution on soft sediments</li> <li>• Contribution to several books on marine biodiversity in NZ</li> <li>• EEZ surveys on Fjordland, Spirit's Bay, Kermadec seamounts, Farewell Spit, Norfolk Ridge, Chatham Rise and Challenger Plateau.</li> <li>• Links to MAFBNZ biodiversity mapping; MEC, MFish BOMECE</li> <li>• Extensive new data sets and specimen collections obtained</li> </ul>	<ul style="list-style-type: none"> <li>• Ongoing taxonomic work in relation to deep sea corals (VMEs)</li> <li>• Ongoing taxonomic work on specimens collected from the Chatham-Challenger project and from the IPY –CAML project.</li> </ul>
<ul style="list-style-type: none"> <li>• Ecological input to improve MEC (fish, benthic invertebrates)</li> <li>• Deep-sea habitats , biogenic habitat and soft-sediment reviewed</li> <li>• Ocean Survey 20/20 habitats mapped Chatham-Challenger</li> <li>• Biodiversity of Kermadec and Chatham Rise seamounts mapped</li> <li>• Foveaux Strait habitats mapped</li> <li>• Classification of seamounts and VMEs developed</li> <li>• Testing of MEC with Chatham Challenger data</li> <li>• Rhodolith beds as havens of biodiversity in NZ</li> </ul>	<ul style="list-style-type: none"> <li>• Mapping biogenic structures</li> <li>• Mapping deepsea fisheries habitats in relation to ocean acidification threats from changing saturation horizons</li> <li>• Modelling benthic impacts</li> </ul>
<ul style="list-style-type: none"> <li>• Rocky reef ecosystem function studied</li> <li>• Chatham Rise fish feeding study completed</li> <li>• Productivity in horse mussel and echinoderm benthic communities determined</li> <li>• Bioindicators in estuarine systems in Otago determined</li> <li>• Chatham-Challenger functional component analysis completed</li> <li>• Shellhash habitat function in the coastal zone</li> </ul>	<ul style="list-style-type: none"> <li>• Ocean acidification on shellfish</li> <li>• Response and recovery of seabed to disturbance-modelling project</li> </ul>
<ul style="list-style-type: none"> <li>• Molecular ID of certain fish and plankton determined</li> <li>• EEZ and Ross Seaspecies added to Barcode of Life Database,</li> <li>• Genetic assessment of ocean microbe diversity</li> <li>• Seamount connectivity reviewed</li> </ul>	<ul style="list-style-type: none"> <li>• Connectivity among coastal fish populations</li> </ul>
<ul style="list-style-type: none"> <li>• Threats and impacts to biodiversity and ecosystem functioning beyond natural environmental variation identified</li> <li>• Monitoring of plankton on transect NZ to Ross Sea annually</li> <li>• Changes in coccolithophore diversity and abundance in NZ waters and predicted change as temp and acidification increase assessed</li> <li>• Long-term effects of climate change on shelf ecosystems determined</li> </ul>	<ul style="list-style-type: none"> <li>• Experimental response of shellfish pH and temp.</li> <li>• CPR monitoring</li> <li>• Initial appraisal for MEMP</li> <li>• Acidification in deepwater fish habitat</li> </ul>
<ul style="list-style-type: none"> <li>• Diversity metrics and other indicators to monitor change developed</li> <li>• Large-scale sampling protocols for habitat mapping determined</li> <li>• Acoustic habitat mapping tools developed</li> <li>• Workshop held on qualitative modelling and marine environment monitoring</li> <li>• Development of "OFOP" and DTIS-visual analytical methods</li> <li>• Predictive modelling techniques progressed for biodiversity on different scales</li> <li>• Development of data to end-user portal interfaced with NABIS</li> </ul>	<ul style="list-style-type: none"> <li>• Development of functional biota model for habitat classification</li> <li>• Qualitative and quantitative modelling of rocky reef ecosystem</li> <li>• Predictive modelling VMEs</li> <li>• Measuring risk and resilience (Chat-Chall objective)</li> </ul>
<ul style="list-style-type: none"> <li>• Latitudinal gradient project and ICECUBE completed in Ross sea</li> <li>• Fish taxonomy and ID guide developed for the Ross Sea</li> <li>• Foodweb and role of silverfish vs krill studied</li> <li>• IPY-CAML 2008; Ross Sea 2006, BioRoss 2004 surveys done</li> <li>• Subtidal and offshore biodiversity sampled, Balleny Islands 2006</li> <li>• Seaweed diversity determined at Balleny Islands</li> <li>• Bioregionalisation of the Ross Sea region completed</li> </ul>	<ul style="list-style-type: none"> <li>• finalisation IPY analyses</li> <li>• Uptake of biodiversity results to CCAMLR trophic modelling and biomass estimation, VMEs</li> <li>• New spp logged for CAML</li> <li>• Review of squids, octopus</li> </ul>




Figure 11.4: Continued Summary of MPI Biodiversity Research Programme 2000–2012.

### **11.3.2. Overall progress in MPI marine biodiversity research**

The MPI Biodiversity Research programme has three overarching science goals:

- To describe and characterise the distribution and abundance of fauna and flora, as expressed through measures of biodiversity, and improving understanding about the drivers of the spatial and temporal patterns observed.
- To determine the functional role of different organisms or groups of organisms in marine ecosystems, and assess the role of marine biodiversity in mitigating the impacts of anthropogenic disturbance on healthy ecosystem functioning.
- To identify which components of biodiversity are required to ensure the sustainability of healthy marine ecosystems as well as to meet societal values on biodiversity.

More specific Science Objectives developed below have been modified by BRAG over time and are used to focus the research commissioned:

1. To classify and characterise the biodiversity, including the description and documentation of biota, associated with nearshore and offshore marine habitats in New Zealand.
2. To develop ecosystem-scale understanding of biodiversity in the New Zealand marine environment.
3. To investigate the role of biodiversity in the functional ecology of nearshore and offshore marine communities.
4. To assess developments in all aspects of diversity, including genetic marine biodiversity and identify key topics for research.
5. To determine the effects of climate change and increased ocean acidification on marine biodiversity, as well as effects of incursions of non-indigenous species, and other threats and impacts.
6. To develop appropriate diversity metrics and other indicators of biodiversity that can be used to monitor change.
7. To identify threats and impacts to biodiversity and ecosystem functioning beyond natural environmental variation.

To date, 55 research projects have been commissioned. Early studies focused primarily on Objectives 1 and 2 and resulted in reviews, Identification Guides, habitat and community characterisations, and revised taxonomy for certain groups of organisms. These objectives have also resulted in large collaborative ship-based surveys that have contributed to improved seabed classification in New Zealand waters and the exploration of new habitats in the region and in Antarctic waters. Over time, the complexity and scale of studies has increased with projects on the functional ecology of marine ecosystems from localised experimental manipulation to broad-scale observations across 100s km<sup>2</sup> under Objective 3. Such studies have also pursued the development of improved measures of biodiversity and indicators under Objectives 6 and 7. A study on changes in shelf ecosystems over the past 1000 years is yielding insights into the effects of long-term climate change, land-use effects and fishing on marine ecosystems while more recently, some studies have begun to address the effects of ocean acidification on marine biodiversity under Objective 5. A study underway has reviewed genetic variation in the New Zealand marine environment and is conducting field observations on several species to examine genetic variation across latitudinal gradients. Aspects of the seven Objectives have also been addressed through a range of biodiversity projects in the Ross Sea region including the International Polar Year Census of Antarctic Marine Life project (IPY-CAML). A key to study findings is consideration of biodiversity within the context of the carrying capacity of the system and the natural assemblages of biota supported by that system in the absence of human disturbance. Progress in the MPI Biodiversity Programme is summarised in Figure 11.5.

<i>Progression of research understanding</i>	<i>Science objective<sup>†</sup></i>	<i>Estuarine/ Coastal 0-30 m</i>	<i>Shelf 30-200 m</i>	<i>Slope 200-1500 m</i>	<i>Deep/Abyss &gt;1500 m</i>	<i>Antarctica All depths</i>
1. Review extent of knowledge of biodiversity (desktop)	1-7					
2. Identify & characterise species and habitat diversity (field work, qualitative analysis, taxonomy & systematics)	1					
3. Quantify biodiversity distribution, abundance (replication, purpose designed surveys)	1					
4. Model and predict biodiversity distribution and abundance	1					
5. Assess or measure functional processes in healthy marine ecosystems (experiments, process studies)	2, 3					
6. Assess the role of genetic diversity	4					
7. Assess interactions and connectivity on ecosystem scale, (genetics, modelling)	2, 5					
8. Develop indicators and measures to monitor bio-diversity, ecosystem health	6					
9. Define key risks and threats to biodiversity	5, 7					
10. Define standards for maintaining biodiversity and healthy ecosystem functioning	6					
11. Examine strategies to mitigate remedy or avoid threats to biodiversity	6					
12. Monitor risks and compliance with standards	6					

**Figure 11.5. Progress on biodiversity research commissioned by MPI 2000–2010. Dark grey: Significant progress (several projects completed and results emerging from research underway). Light grey: Limited progress (some results emerging, more research needed). White: no substantive research. Diagonal-hatch: progress linked to large whole-of-government projects (e.g. Ocean Survey 2020) and/or other funding outside MPI (e.g. MBIE (MSI) funded Outcome Based Investment projects, DOC Marine Coastal Services, MAFBNZ marine biosecurity research).**

<sup>†</sup> Science objectives are- 1 characterisation and description; 2 ecosystem scale biodiversity; 3 functional role of biodiversity; 4 genetics; 5 ocean climate effects; 6 indicators; 7 threats to biodiversity. The objectives are detailed in MPI Biodiversity Programme: Part 2. Medium Term Research Plan 2011-2014.

The chart depicts a logical flow down the page of increasing conceptual complexity from cataloguing of biodiversity to increasingly complex understanding of environmental drivers and functionality of biodiversity; and ultimately methods to develop standards and protection of biodiversity. Across the

chart, the marine environment is graded from the coastline to offshore regions, and Antarctica. A full list of projects can be obtained from the MPI Biodiversity Medium Term research programme 2010-2014.

Greatest progress has been made in the shallower inshore parts of the marine environment, not least because of cost and ease of access. However, by leveraging from existing offshore projects, significant progress has also been made to depths of 1500 m.

MPI Biodiversity research based in Antarctica lags behind EEZ-based research, simply because of the difficulty in securing additional funding to access and work in such a remote and hostile marine environment. While the top left side of the figure shows the area of greatest progress, it would be a mistake to conclude that biodiversity work is completed here.

### **11.3.1. Progress on Science Objective 1. Characterisation and Classification of Biodiversity**

The characterisation and classification of biodiversity requires an assessment of the abundance and distribution of marine life. Building on earlier research to map fish and squid species (Anderson *et al.* 1998, Bagley *et al.* 2000) and the biodiversity of the New Zealand ecoregion (Arnold 2004), literature reviews, taxonomic studies and habitat mapping surveys have been undertaken.

#### **Reviews and books**

The following lists scientific reviews and books on biodiversity that were commissioned by the programme:

- ZBD2000-01 A review of current knowledge describing the biodiversity of the Ross Sea region (Bradford-Grieve and Fenwick 2001, 2002; Fenwick and Bradford-Grieve 2002a, 2002b, Varian 2005)
- ZBD2000-06 “*The Living Reef: The Ecology of New Zealand's Rocky Reefs*” (eds. Andrew and Francis 2003)
- ZBD2000-08 A review of current knowledge describing New Zealand’s Deepwater Benthic Biodiversity (Key 2002),
- ZBD2000-09 Antarctic fish taxonomy (Roberts and Stewart 2001)
- ZBD2001-02 Documentation of New Zealand Seaweed (Nelson *et al.* 2002)
- ZBD2001-04 “Deep Sea New Zealand” (Batson 2003)
- ZBD2001-05 Crustose coralline algae of New Zealand (Harvey *et al.* 2005, Farr *et al.* 2009, Broom *et al.* 2008)
- ZBD2001-06 Biodiversity of New Zealand’s soft-sediment communities (Rowden *et al.* 2011)
- ZBD2003-09 Macquarie Ridge Complex Research Review (Grayling 2004)
- ZBD2008-27 Scoping investigation into New Zealand abyss and trench biodiversity (Lörz *et al.* 2012).

In addition a major work which includes marine species – “The New Zealand Inventory of Biodiversity” (Gordon 2009, Gordon 2010, Gordon 2012), has been completed. Field identification guides have also been published by MPI on deepsea invertebrates (–projects ENV2005-20 and ZBD2010-39, Tracey *et al.* 2005, 2007, 2011), bryozoans (project IPA2009/14 Smith and Gordon 2011) and on fish species (IDG2006-01 MacMillan *et al.* (2011 a, b, c) which further contribute to the accurate monitoring and identification of biodiversity in New Zealand waters.

#### **Projects**

Several hundred new species of marine organisms have been discovered, and the known range of species extended, through exploratory surveys such as the NORFANZ project ZBD2002-16 (Clark

and Roberts 2008); MSI's Seamount Programme, mainly commissioned through public-good science, supplemented by MPI projects ZBD2000-04, e.g., Rowden *et al.* 2002 and 2003, ZBD2001-10 (Rowden *et al.* 2004), ZBD2004-01 (Rowden *et al.* 2010) and MPI projects ENV2005-15, ENV2005-16 (Clark *et al.* 2010, Rowden *et al.* 2008) and the Ocean Survey 20/20 programme (Clark *et al.* 2009); inshore surveys of bryozoans at Tasman Bay ZBD2000-03 (Grange *et al.* 2003); Farewell Spit, ZBD2002-18 (Battley *et al.* 2005), Fiordland, ZBD2003-04 (Wing 2005); coralline algae ZBD2001-05, ZBD2004-07 (Harvey *et al.* 2005, Farr *et al.* 2009); soft sediment environments ZBD2003-08 (Neill *et al.* 2011); rhodolith community study ZBD2009-03 (Nelson *et al.* 2012); offshore surveys of the Chatham Rise and Challenger Plateau funded through Ocean Survey 20/20 programme, ZBD2006-04 (Nodder 2008) and ZBD2007-01 (Nodder *et al.* 2011; Hewitt *et al.* 2011; Bowden 2011, Bowden and Hewitt 2012; Bowden *et al.* 2011b; Bowden *et al.* in press).

Research in the Ross Sea Region (BioRoss projects) have also generated records of new species including MPI projects ZBD2000-02 (Page *et al.* 2001), ZBD2001-03 (Norkko *et al.* 2002), ZBD2002-02 (Sewell *et al.* 2006, Sewell 2005, 2006), ZBD2003-02 (Cummings *et al.* 2003, 2006), ZBD2003-03 (Rowden *et al.* 2012a, Rowden *et al.* in press), ZBD2005-03 (MacDiarmid and Stewart 2012), ZBD2006-03 (Cummings *et al.* 2003, 2006; Norkko *et al.* 2002), ZBD2008-23 (Nelson *et al.* 2010) and IPY2007-01 (Bowden *et al.* 2011a, Clark *et al.* 2010, Eakin *et al.* 2009, Hanchet, *et al.* 2008a Hanchet 2008b, Hanchet 2008c, Hanchet *et al.* 2008d. Hanchet 2009, Hanchet 2010, Koubbi *et al.* in press, Lörz and Coleman 2009, Lörz in press, Lörz *et al.* in press, Mitchell 2008, O'Driscoll *et al.* 2009, O'Driscoll 2009, O'Driscoll, *et al.* 2010, O'Loughlin *et al.* 2010)

### Habitat diversity, classification and characterisation

The development of the Marine Environment Classification or "MEC" (Snelder *et al.* 2006) was an important step in the delineation of areas with similar environmental attributes in the offshore environment. However, significant environmental drivers of variability in marine biodiversity, such as substrate type for seafloor organisms, were absent from the classification. In 2005, DOC and MPI jointly commissioned a project to optimise the MEC using fish distribution data. This project (ZBD2005-02) demonstrated a substantial improvement in the MEC classification for offshore habitats (Leathwick *et al.* 2006a, b, c). In 2006, three projects to map coastal biodiversity were completed in the Coromandel scallop, Foveaux Strait oyster and southern blue whiting fisheries as part of fishery plan development for these fisheries (ZBD2005-04, ZBD2005-15, ZBD2005-16). These projects found that the biological distribution of organisms and their habitats were not well predicted by the MEC. MPI project (BEN2006-01) aimed to further optimise the MEC by producing a methodology for a Benthic Optimised MEC (Leathwick *et al.* 2009). MPI Ecological studies to improve habitat classification and vulnerability indices have also been completed through MPI AEWG projects on seamounts (ENV2005-15, ENV2005-16) (e.g., Clark *et al.* 2010), and to supplement other studies funded by MPI, and MSI (e.g. ZBD2004-01, ZBD2001-10, ZBD2000-04, and CO1X0508).

Distribution maps providing indicative abundance and characterisation of biodiversity are now emerging and have been produced through projects using predictive modelling tools e.g., Compton *et al.* 2012; the fish optimised MEC in project ZBD2005-02 (Leathwick *et al.* 2006a, 2006b, 2006c), the benthic optimised MEC (Leathwick *et al.* 2009) and Chatham-Challenger project ZBD2007-01 (Hewitt *et al.* 2011, Bowden *et al.* 2012, Compton *et al.* in press).

Progress has advanced considerably in recent years with the introduction of the whole-of-government Ocean Survey 20/20 Programme and Biosecurity New Zealand mapping projects (Beaumont *et al.* 2008, 2010) In addition, MPI implemented spatial management tools (Benthic Protection Areas<sup>54</sup>) implemented on the basis of the Marine Environment Classification<sup>55 56</sup> to address broader statutory responsibilities on the environmental effects of fishing on biodiversity.

<sup>54</sup> [www.fish.govt.nz/en-nz/Environmental/Seabed+Protection+and+Research/Benthic+Protection+Areas.htm](http://www.fish.govt.nz/en-nz/Environmental/Seabed+Protection+and+Research/Benthic+Protection+Areas.htm)

<sup>55</sup> Marine Environmental Classification. (2005). Can be viewed online at <http://www.mfe.govt.nz/publications/ser/marine-environment-classification-jun05/index.html>

**ZBD2007-01 Chatham-Challenger seabed habitats-post voyage analyses.**

This large project has been completed. Progress for each objective is as follows:

1. To count, measure, and identify to species level (where possible, otherwise to genus) all macro invertebrates (>2 mm) and fish collected during Oceans Survey 20/20 voyages. Completed (Figure 6, Bowden 2011).
2. To count, measure and identify to species-level (where possible, otherwise to genus or family) all meiofauna (>45µm to <500 µm) from multicore samples collected during the Oceans Survey 20/20 voyages. [Collaborative venture MBIE-Otago University]. Completed (Leduc *et al.* 2012)
3. To count, measure and identify to species- level (where possible, otherwise to genus or family) all fauna collected by hyper-benthic sled during the Oceans Survey 20/20 voyages. Completed. (Lorz 2011, Bowden 2011).
4. To count, measure, and identify to species-level (where possible, otherwise to genus or family) all macrofauna observed on DTIS images collected during the Oceans Survey 20/20 voyages. The number of biogenic features (burrows/mounds) and habitat (spatial) complexity should also be estimated. Completed. (Bowden 2011, Compton *et al.* 2012).
5. To count, measure, and identify to species-level (where possible, otherwise to genus or family) all macrofauna observed on DTIS video footage collected during the Oceans Survey 20/20 voyages. Completed. (Bowden 2011, Compton *et al.* 2012).
6. To calculate and compare the performance of a suite of diversity measures (species and taxonomic based) at varying levels of resolution. Completed. (Hewitt *et al.* 2011a).
7. To estimate particle size composition and organic content of sediment samples. Sediment samples should be aggregated over the top 5 cm of sediment. Completed. (Nodder *et al.* 2011).
8. To measure the bacterial biomass (top 2 cm) of the sediment and in the sediment surface water samples, collected during the Oceans Survey 20/20 voyages. Completed. (Nodder *et al.* 2011).
9. To elucidate the relationships, patterns and contrasts in species composition, assemblages, habitats, biodiversity and biomass (abundance) both within and between stations, strata and areas. Completed. (Floerl *et al.* 2012).
10. To define habitats (biotic) encountered during the survey and assess their relative sensitivity to modification by physical disturbance, their recoverability and their importance to ecosystem function / production. Completed (Hewitt *et al.* 2011b).
11. *To quantify the productivity, energy flow (trophic networks) and the energetic coupling (benthic pelagic or otherwise) of the area surveyed areas at various levels of resolution. Objective withdrawn*
12. To assess the extent to which patterns of species distributions and communities can be predicted using environmental data (including fishing) collected during the Ocean 20/20 voyages or held in other databases. Modelling approaches as well as standard statistical procedures are anticipated. (Compton *et al.* 2012).
13. To provide an interactive, high resolution mapping facility for displaying and plotting all data collected and derived indices. This would include environmental data, the abundance of individual species, indices of biomass or diversity, and statistically derived groupings. Completed in conjunction with Bay of Islands Ocean Survey 20/20 Portal<sup>57</sup>.
14. To assess the extent to which acoustic, environmental, or other remote-sensed data can provide cost-effective, reliable means of assessing biodiversity at the scale of the Oceans

<sup>56</sup> <http://seafoodindustry.co.nz/bpa> and use of MEC (2005)

<sup>57</sup> <http://www.os2020.org.nz/>

Survey 20/20 samples. Completed. (Bowden *et al.* in press, Bowden *et al.* 2011b, Compton *et al.* 2012).

15. To assess the extent to which the 2005 MEC and subsequent variants can provide cost-effective, reliable means of assessing biodiversity at the scale of the Oceans 20/20 surveys. Completed. (Bowden *et al.* 2011b).
16. Collating all information and analysis from all objectives, devise a series of statistically supported recommendations for surveying marine biodiversity in the future. This should include, but may not be limited to, statistical analyses and modelling. Bowden and Hewitt 2011).

**ZBD2008-05 Macroalgal diversity associated with soft sediment habitats.**

Although macroalgae normally require hard substrata for attachment and occur less frequently in soft sediment environments they contribute to biodiversity in a range of soft sediment environments providing structural complexity, modifying flow and sediment regimes, and contributing to productivity. Soft sediment habitats where macroalgae are found are physically highly diverse, ranging from harbours and estuaries (with varying sediment types and sizes, freshwater influence, tidal flushing, current flows), to coarse stabilised sediments (shell fragments, cobbles, coarse gravels), and biogenic habitats such as worm tubes, horse mussel beds, brachiopod beds, mangrove forests, rhodolith (maerl) beds and seagrass meadows.

The state of knowledge of macroalgal diversity, distribution and abundance is poor, and there are few examples of targeted collecting programmes for macroalgal assemblages, particularly in soft sediment habitats. This research conducted (a) a targeted collection programme across diverse soft sediment environments to develop a permanent reference collection of representative macroalgae, and (b) examined algal distribution in soft sediment habitats in relation to selected environmental variables.

Macroalgal sampling trips to Kaipara (1), Whangarei (3) and Otago (4) Harbours were completed. Further sampling trips were planned for 2010, however, no further collections will be made in Kaipara Harbour. Approximately 2400 collections of algae were made from soft sediments in these harbours. In Whangarei and Otago Harbours, collections were made from a range of soft sediment habitats including mud, sand, shell gravel, sea grass, scallop, pipi and horse mussel beds. At each site algae were collected opportunistically, quantitatively (i.e. by quadrats), or by both methods. Standard ecological methods (e.g. species area curves, count frequencies) were used to assess the appropriateness of the methods.

A database was developed for information about specimens and collection sites. Information was gathered on environmental variables within the target harbours. Identified algal distributions were analysed relative to these environmental variables.

Collections were made from three harbours with the primary focus on Whangarei and Otago Harbours where seasonal sampling programmes were conducted in spring and in autumn. In the Kaipara Harbour sampling was conducted only in spring. Two hundred and forty four taxa sampled from intertidal and subtidal sites and a range of habitats: 146 (112 spring, 102 autumn) from Whangarei, 43 Kaipara, 150 (107 spring, 115 autumn) from Otago. Diversity indices indicate that the collecting was not saturated and predict that there is higher diversity of macroalgae in these harbours than found in the samples obtained.

The flora composition in the harbours was found to differ markedly e.g., only 67 taxa (45%) of the Whangarei flora were found to be in common with Otago Harbour collections; 17 taxa (39% ) of the Kaipara flora was in common with the Otago flora, in common (39% of K found in O); 27 taxa (63%) of the Kaipara flora was also found in Whangarei. 19 non-indigenous species were found in the harbours, including two new records for the New

Zealand algal flora (confirmed by sequence data), *Hypnea cornuta* and *Polysiphonia morrowii*. In Whangarei Harbour 8 non-indigenous species were found (4 new records for harbour including *Hypnea*), in Kaipara Harbour 4 species were found including 2 new records for the harbour, and in Otago Harbour 11 non-indigenous species were found including 1 new record as well as *P. morrowii*. More taxa were collected in the subtidal (107) in Whangarei Harbour than in the intertidal (84), compared with Otago where numbers of intertidal taxa (120) exceeded the subtidal taxa collected (83).

Two methods were employed to enable high resolution sampling and these provided differing outcomes in the two main harbours sampled, clearly indicating that there was value in collecting by both methods in order to adequately sample the diversity: Whangarei Harbour 90 taxa were collected in quadrat sampling compared with 118 taxa via opportunistic collections, and in the Otago Harbour 107 taxa were collected in quadrat sampling and 118 taxa via opportunistic collections.

***ZBD2008-27 Review of deep-sea benthic biodiversity associated with trench, canyon and abyssal habitats below 1500 m depth in New Zealand waters***

The state of knowledge of benthic biodiversity and ecosystem functioning in deep-sea abyssal, canyon and trench habitats in the New Zealand Exclusive Economic Zone and the Ross Sea region, was summarised and recommendations for future deep-sea research in depths exceeding 1500 m were made. All biological information in scientific papers and reports from New Zealand below 1500 m was reviewed and an exhaustive search of multiple data sources was conducted.

The area of the deep seafloor below 1500 m covers more than 65% of New Zealand's Exclusive Economic Zone. A total of 1489 benthic gear deployments have been conducted by New Zealand-based sampling initiatives since 1955, most of which were focused on obtaining geological samples. Less than 0.002 % of New Zealand's deep-sea environment (i.e. in terms of seabed area) below 1500 m has been sampled. All taxonomy-based studies of all taxa reported in New Zealand waters below 1500 m have been reviewed. To date, 8 species of Bacteria, 293 species of Protozoa, 785 species of invertebrates, and 56 fish species have been recorded from water depths greater than 1500 m.

More than 8000 images are known to have been taken of the seafloor below 1500 m in the New Zealand region, covering an area of approximately 0.016 km<sup>2</sup>. Over 4000 of the images held at NIWA exist either as paper prints or negatives and ideally should be digitised for future storage and access for analyses. Analysis of these photographic images should yield considerable information about deep-sea biodiversity and ecosystem function in the New Zealand region and could be used to answer a number of research questions (especially around deep-sea benthic biodiversity).

Recommendations on how to potentially further analyse existing data from images, databases and actual specimens were provided. The technical challenges, including gear requirements to sample deep-sea New Zealand benthos and potential future investments, were summarised. (see Coleman and Lörz 2010; Lörz 2011a, 2011b; Lörz *et al.* 2012a, 2012b).

***ZBD2008-50 Chatham Rise biodiversity hotspots.***

This survey covered the “Graveyard Seamount Complex” and “Andes Seamount Complex” on the Chatham Rise. Objectives were to monitor changes over time on Graveyard hills subject to differing management regimes (some open to fishing, some closed), as well as to compare seamount biodiversity between different regions of the Rise. It was linked to the CoML CenSeam programme, and the former FRST Seamounts research, now under the MBIE Vulnerable Deep-sea Communities project. The data from that survey are being

worked up under the latter project (see Clark *et al.* 2009).). Analyses comparing the 3 surveys of the Graveyard complex between 2001 and 2009 indicate there are changes in some taxa following cessation of fishing operations on one of the features in 2001, but little sign of any recovery of stony coral species and associated benthic communities. Preliminary results were presented at the 2012 Deep Sea Biology Symposium (Clark *et al.* 2012).

**ZBD2009-03 The vulnerability of rhodoliths to environmental stressors and characterisation of associated biodiversity.**

Rhodoliths are free-living calcified red algae. They occur worldwide, forming structurally and functionally complex benthic marine habitats. Rhodolith beds form a unique ecosystem with a high benthic biodiversity supporting many species, including some that are rare and unusual. Recent international studies show that these fragile algae are at risk from the impacts of a range of human activities e.g., physical disruption, reduction in water quality, alterations to water movement, and aquaculture installations. Impacts of fragmentation may be critical in terms of biodiversity and abundance associated with rhodolith beds.

The focus of this programme was to improve knowledge about the location, extent or ecosystem functioning of rhodolith beds in New Zealand. The ecology of subtidal rhodolith beds was been investigated for the first time in New Zealand, characterising two rhodolith species, *Lithothamnion crispatum* and *Sporolithon durum*, examining the structure and physical characteristics of beds at two locations and documenting their associated biodiversity. In addition the responses of these rhodolith species to environmental stressors were investigated for the first time.

This study documented high biodiversity in two subtidal rhodolith beds sited in relatively close proximity in the coastal zone, with significant differences in biotic composition. The rhodolith beds studied (located in the Bay of Islands) differed significantly in terms of water motion, sediment characteristics and light levels. Biodiversity of the rhodolith beds was investigated sampling (1) invertebrates at three levels of association (epifauna, infauna, cryptofauna), (2) macroalgae, (3) fishes, as well as recording the biogenic and non-biogenic substrates:

- a number of undescribed taxa were discovered as well as new records for the New Zealand region, and range extensions of species known elsewhere,
- more than double the number of invertebrate taxa were present in the rhodolith beds than found outside the beds,
- both rhodolith beds harboured high diversity of associated macroalgae and invertebrates but with markedly different species composition,
- the floral and faunal composition differed significantly between sites.

Both species of rhodolith were found to be vulnerable to the impacts of increasing temperature and decreasing pH. There was a significant difference between the effects of treatments on the two species and further statistical analysis showed significant interaction between temperature and pH level on growth. Overall the greatest effect on growth rate came with the combination of high temperature (25° C) and low pH (7.65) on *Lithothamnion crispatum* which showed negative growth, indicating probable dissolution. In experiments investigating other environmental stressors, temperature was found to be more important for the survival and growth of the rhodolith species examined than the effects of burial, light and fragmentation.

The extent of rhodolith beds in other parts of the New Zealand region remain to be documented, including those in coastal areas (including intertidal beds) and subtidal beds on the shelf.

**ZBD2010-40 Predictive modelling of the distribution of vulnerable marine ecosystems in the South Pacific Ocean region.**

In January 2010 New Zealand and the United States held their second Joint Commission meeting (JCM) on Scientific and Technological Cooperation. The meeting was to share knowledge about common interests and capabilities and identify areas for future collaboration. The JCM consisted of six workshops held simultaneously around the North Island and an officials meeting held in Wellington. One of the six workshops, ocean and marine sciences, identified an area of interest in a joint project in the South Pacific Regional Fisheries Management Organisation (SPRFMO) area to map and groundtruth vulnerable marine ecosystem (VME) distribution.

The 3rd New Zealand and United States Joint Commission on Science and Technology Cooperation (JCM) met on 19 and 20 September 2012 in Washington. Building on several recommendations from the previous JCM (held in January 2010), and on the Marine Conservation Think Tank VME Workshop report 3: Science requirements for effective High Seas governance, held on 2-5 December 2011 (Lundquist et al 2012) 1, Topic 1 for the Oceans and Marine Workshop and the 3rd JCM meeting was again Vulnerable Marine Ecosystems (VMEs). Several actions were developed at the workshop and these included: 'Increase in situ deep-sea exploration and VME studies in regions of common interest by exploring options for NOAA/WHOI participation (including use of ROV/AUV technologies) in New Zealand funded initiative and voyage to explore and ground-truth VMEs in the South Pacific' and 'Facilitate U.S. researcher involvement in NIWA Louisville Ridge Exploration.'

There are relatively few data available on the distribution of VME species or taxa in the South Pacific Ocean (Parker *et al.* 2009) although studies have been conducted in Antarctica (Tracey *et al.* 2010, Parker *et al.* 2009) to use for the objective planning of spatial protection measures to protect those taxa, particularly in the SPRFMO Area. It is therefore becoming increasingly important to develop robust predictions of where VMEs are likely to occur, using habitat prediction and species distribution models. Such models have recently been developed and/or are in the process of being refined for certain VME taxa on a global scale (e.g. Actinaria, Guinotte *et al.* 2006; Scleractinia, Tittensor *et al.* 2009). However, the spatial resolution of existing models is coarse (larger than the scale of the topographic features typically targeted during demersal high seas fishing), and the level of uncertainty around the predictions is variable or still unknown.

Phase 1 a project to use modelling to predict the location of VMEs in the SPRFMO area was initiated between the US and New Zealand (ZBD2010-40) and has now been completed. The objectives of the project were to:

1. To develop and test spatial habitat modelling approaches for predicting distribution patterns of vulnerable marine ecosystems in the Convention Area of the South Pacific Regional Fisheries Management Organisation with agreed international partners.
2. To collate data sets and evaluate modelling approaches which are likely to be useful to predict the distribution of vulnerable marine ecosystems in the South Pacific Ocean region.

Data for ten Vulnerable Marine Ecosystems (VME) taxa were compiled from different data sources to produce a single groomed dataset for VME indicator taxa in the New Zealand region. Regional-tuned environmental data layers and global environmental data layers were obtained from available data sources. Using these data, three types of predictive models were made for each VME indicator taxon. Two models were made using regional-tuned environmental data layers, using maximum entropy analysis (MaxEnt) and boosted regression tree (BRT) techniques to provide a comparison of the different model approaches. The third type of model made used the MaxEnt approach, but using globally available environmental

data layers. Having a third model meant that model performance could be compared based on the use of different environmental data layers. Three model types for all VME taxa have been completed and the performance of the different modelling approaches and usefulness of the environmental data sets described.

The next phases of the project will be undertaken as part of a MBIE-funded project that will revise models that predict the sites of Vulnerable Marine Ecosystems from existing data by conducting a ground truthing survey of benthic biodiversity on the Lewisville Ridge in 2013/14 (Ministry of Business Innovation and Employment project code C01X1229). This will be used to inform New Zealand and South Pacific Regional Fisheries Management Organisation initiatives on spatial management in the South Pacific region, and potentially the New Zealand EEZ.

Other research relevant or specifically linked to the projects above, is listed in Table 11.1.

**Table 11.1: Other research linked to Objective 1 habitat classification and characterisation.**

<b>MPI</b>	HAB2007-01 Biogenic habitats as areas of particular significance for fisheries management ZBD2006-02 NABIS ongoing development Useful data related to defining potential VMEs are collected by MPI scientific fisheries observers working on NZ authorised fishing vessels that operate on the high seas in the South Pacific.
<b>CRI core purpose funding</b>	C01X501 Coasts & oceans Centre (NIWA) ecosystem based management, habitat model development with Auckland Regional Council C01X0907 Coastal Conservation Management (fish habitat classification) (NIWA) C01X502 Biodiversity & Biosecurity (NIWA) C01X0508 Seamount fisheries (linking acoustic backscatter to habitat type and biota) (NIWA) C01X0906 Vulnerable deep-sea communities (mapping and sampling a range of deep-sea habitats (seamounts, slope, canyons, seeps, vents) (NIWA) C01X0702 Kermadec Arc minerals (mapping and sampling the biodiversity of several Kermadec Arc seamounts) (NIWA)
<b>DOC</b>	MEC development and application to MPAs, Regional surveys
<b>OTHER</b>	University studies, Regional Council studies
<b>ZBD2010-40</b>	Mapping VMEs in the SPRFMO area Part 1. Predictive modelling desktop study
<b>EMERGING ISSUES</b>	
What portion of a given habitat type should remain intact to support sustainable ecosystems? What are the most effective predictive tools for predicting biodiversity in areas as yet unsampled? Can ecological mapping used in OS20/20 projects to date be extended to other areas of New Zealand?	

### 11.3.2. Progress on Science Objective 2. Ecosystem-scale research

Marine ecosystems influence, and are influenced by, a wide array of oceanic, climatic, and ecological processes across a broad range of spatial and temporal scales. Marine communities are generally dynamic, can occur over large areas and have strong links to other communities through processes such as migration and long-distance physical transport (e.g. of larvae, nutrients, and biomass). Patterns observed on a small scale can interact with larger and longer-scale processes that in turn result in large scale patterns. Marine food webs are usually complex and dynamic over time (Link 1999). To distinguish useful descriptors of long-term ecosystem change from short-term fluctuations requires innovative approaches to integrate broad-scale correlative studies from smaller scale manipulative experiments (Hewitt *et al.* 1998, 2007).

Recent theoretical and technical advances show great promise toward the goal of understanding the role of biodiversity in ecosystems. Technologies for remote sensing and deepwater surveying, combined with powerful integrative and interpretive tools such as GIS, climate modelling, qualitative ecosystem modelling, and trophic ecosystem modelling, will contribute to the development of an

ecosystem-based approach to management (Thrush *et al.* 1997, 2000), with potential benefits for marine conservation and management. Ecosystem modelling of species distribution (and habitats) with respect to known and projected environmental parameters will improve predictability for both broad and fine-scale biodiversity distribution. This has already resulted in improved definition of environmental classifications addressing biodiversity assessment. It is also important to make progress in establishing the links between biodiversity and the long-term viability of fish stocks under various harvesting strategies. It is also important that modellers consider processes from all ecosystem function perspectives i.e., top-down effects such as predation (e.g. trophic modelling), bottom-up effects such as the environment (e.g., habitat classification based on environmental variable), and wasp-waisted systems where there are major effects in both directions.

## Projects

**ZBD2002-06A: Impacts of terrestrial run-off on the biodiversity of rocky reefs** Completed.  
(Schwarz *et al.* 2006).

**ZBD2004-02: Ecosystem scale trophic relationships of fish on the Chatham Rise**, Completed.  
(Connell *et al.* 2010, Dunn 2009, Dunn *et al.* in press, Dunn *et al.* 2010a, b, c, Eakin *et al.* 2009, Forman and Dunn 2010, Horn *et al.* 2010, Stevens and Dunn 2010. Follow-up research on isotope signatures to improve the trophic data from ZBD2004-02 has been incorporated into the NIWA's Coast and Ocean programme and trophic modelling is underway in this programme.

**ZBD2004-08 Sea-grass meadows as biodiversity and connectivity hotspots.**

This contract links closely with the MBIE project Coastal Conservation Management (CO1X0907). National scale sampling across North and South Island seagrass meadows in a range of estuarine and coastal settings has shown that seagrass meadows overall consistently supported higher species richness, biomass, and productivity of invertebrates (infaunal and epifaunal). Associated sampling of small fish assemblages found that while seagrass meadows provided a nursery function to a number of species, this function was most pronounced in northern New Zealand systems, where relatively high numbers of juvenile snapper, trevally, spotties, parore, and garfish/piper were caught. However, there was strongly spatial variation across different estuary and coast settings (MBIE91B).

**ZBD2004-19 Ecological function and critical trophic linkages in New Zealand softsediment habitats**, Project completed. (see Lohrer *et al.* 2010.)

**ZBD2005-05 Effects of climate variation and human impacts on the structure and functioning of New Zealand shelf ecosystems.**

The project is a multidisciplinary study to utilise archeological, paleoecological, and historical data to retrospectively model ecosystem states during different historical and prehistoric time periods. The project is collaborating with the international History of Marine Animal Populations (HMAP) project, itself a part of the Census of Marine Life (CoML) programme. The data have been used as inputs to a mass balance model of the shelf ecosystem starting with the present day Hauraki Gulf. A short video about the NZ Taking Stock project was made by HMAP staff and is currently available on the HMAP website <http://hmapcoml.org/projects/nz/>. Several presentations have been made at NZ and international conferences as results have emerged.

**ZBD2008-01 Inshore biogenic habitats.**

Existing knowledge on biogenic habitat-formers in the <5 – 200 m depth zone of New Zealand's continental shelf, from sources including structured fisher interviews ("Local Ecological Knowledge" LEK), primary and grey literature, and other sources have been integrated to generate maps of key biogenic habitats in New Zealand coastal waters.

Over 600 targets of interest were identified and marked on marine charts, with more than 200 of these targets being biogenic in nature. Fieldwork has been completed to verify and quantify biodiversity in biogenic habitats using Ocean Survey 20/20 vessel days on Tangaroa and a new MSI project to extend the survey potential of the project. New biogenic habitats have been identified, including extensive worm tube ‘meadows’ off the east coast of the South Island (“the Hay Paddock” and “Wire-weed”), with associated relatively high epi-faunal invertebrate diversity compared to adjacent bare sediments. Over 60 new species were also collected (dominated by sponges), along with range extensions of many other species. Analyses are underway for key selected areas included in the Tangaroa voyages, including offshore North Taranaki Bight, Ranfurly Bank, the polychaete meadows mentioned above, and the Otago Peninsula bryozoan fields.

**IPA2009-11. Trophic Review.**

This project publishes a report prepared on the feeding habits of New Zealand fishes 1960 to 2000 (Stevens *et al.* 2011)

Other research relevant or specifically linked to the projects above, is listed in Table 11.2.

**Table 11.2: Other research linked to ecosystem scale understanding of biodiversity in the marine environment.**

<b>MPI</b>	ENV2006-04 Ecosystem indicators for New Zealand fisheries ENV2007-04 Climate and oceanographic trends relevant to New Zealand fisheries ENV2007-06 Trophic relationships of commercial middle depth species on the Chatham Rise
<b>CRI Core purposes</b>	C01X501 coasts & oceans productivity plankton-mesopelagic fish trophic relations Chatham Rise IO 2. Second Fisheries Oceanography voyage to Chatham Rise: mesopelagics and hyperbenthics
<b>OTHER</b>	AUT deepsea and subtidal food web dynamics; offshore & coastal biodiversity post graduate studies

**11.3.3. Progress on Science Objective 3. The role of biodiversity in the functional ecology of nearshore and offshore communities.**

An identified outcome of the Biodiversity Strategy is that by 2020 “*New Zealand’s natural marine habitats and ecosystems are maintained in a healthy functioning state. Degraded marine habitats are recovering.*” Sustaining ecosystem integrity in marine habitats requires a thorough understanding of the ecological and anthropogenic drivers affecting biodiversity and ecosystem function, and the ability to manage human impacts in marine environments.

Near-shore environments range from wetlands to estuaries, coasts and continental shelf ecosystems, they contain a variety of habitats and often contain species that are particularly important, either for cultural, recreational, and commercial reasons, or because the species exerts disproportionate influence on community structure and ecosystem function. Near-shore ecosystems are the multi-use ecosystems most subjected to multiple stressors. Due to ocean-coast and land-coast interactions these ecosystems will be subjected to the greatest range of stresses associated with global warming. Near-shore environments may also contain habitats that are particularly important for biodiversity in other environments, for instance by providing larval/juvenile nursery areas or by exporting nutrients. The MPI Biodiversity Programme has directed funds into research examining the implications of environmental and human impacts on the functional ecology of these key species and habitats.

Near-shore ecosystems are complex and changes in diversity and community composition may be driven by multiple variables. Interactions between variables are likely to be non-linear, with disturbance thresholds and the potential for multiple stable states. As a consequence, it is often difficult to distinguish ‘natural’ from ‘anthropogenic’ impacts affecting ecosystem dynamics. MPI

BioInfo research seeks to help disentangle this complexity, recognising that there will be contributions to this from both biodiversity research and Fisheries Services research.

Regional Councils and universities support some research projects and survey programmes in coastal and estuarine waters by investigating the effects of sedimentation, pollution, ocean outfalls, sand dredge spoils, sand mining and nutrient enrichment on the marine ecosystem<sup>58</sup>. Although this workstream applies to offshore areas as well as near-shore, research to date has focussed on the near-shore.

## Projects

### ***ZBD2005-09 Rocky reef ecosystems - how do they function?***

The draft report for this project has been submitted and reviewed (Beaumont *et al.* In press). The Hauraki Gulf in north-eastern New Zealand offers one of the best opportunities to investigate how rocky reef ecosystems function and what impact fishing and other human activities may have on them. This study took advantage of these circumstances to first review the extensive literature to set the parameters of a model of how north-eastern New Zealand reef ecosystems function. The study used the model to identify key species and interactions, and explore the impacts of fishing. Field work was then undertaken across the range of reefs within the Hauraki Gulf to test the model predictions, describe spatial variation in patterns of abundance of key species, determine trophic relationships and investigate the linkages of reefs to other habitats.

A qualitative model of northeast New Zealand rocky reef ecosystems was developed to explore the complexity of interactions amongst New Zealand rocky reef species and the impacts of exploitation. This model was developed on the basis of a review and summary of interactions among reef components. A key modelling outcome was the highly predictive but opposite responses by small lobsters and large predatory invertebrates to changes in the abundance of a range of other groups. This suggests that these two groups are ideal candidates as variables for monitoring reef ecosystem responses to perturbations. The modelling agreed with a well-documented example of responses to a perturbation in fishing pressure in the Leigh Marine Reserve. However, the predictability was low for all responses. This implies, for example, that the reduction of kina in the Leigh Marine Reserve and the subsequent increase in macro-algae consequent to an increase in lobster abundance may not necessarily occur in another area.

Field sampling at ten rocky reef sites across the Hauraki Gulf revealed differences among sites in community structure of macroalgae and invertebrates within all habitat strata. Of the environmental factors available, depth followed by a measure of water clarity (mean secchi) explained the most variation in the dependent variables (invertebrate taxa) from the quadrat data. Fish abundance data showed a similar, though weaker, trend across sites with depth, distance across the Gulf, and water clarity being the most important factors. The strong association between depth and water clarity and abundances of key taxa was expected and is similar to that found in earlier studies. With the exception of crayfish, there was no apparent overall relationship between invertebrate and fish abundances and marine reserve status of study sites, though the baited underwater video data showed snapper to be significantly larger within marine reserve sites than at fished sites.

Stable isotope analysis of tissue samples collected from key species from all study sites allowed insight into the functional relationships among species as well as dietary sources of carbon. Many of the study taxa, from the primary producers through to the predators, had the most depleted  $\delta^{13}\text{C}$  values at the furthest inshore and offshore sites (e.g. Poor Knights and Long Bay)

---

<sup>58</sup> See MFish Biodiversity Research Programme 2010: Part 4. Reference Materials and Other research

and the highest  $\delta^{13}\text{C}$  values at the coastal sites (e.g. Leigh, Tawharanui and Kawau). Without direct modelling of end point source signatures we cannot definitively determine the percentage contribution of each carbon source. However, we suggest that the depleted  $\delta^{13}\text{C}$  of taxa from offshore sites is the result of a pelagic source of C and the enriched  $\delta^{13}\text{C}$  at coastal sites is the result of a more benthic input of C than at offshore sites, with sources including kelp detritus. Taxa at the inner gulf sites are also likely to be subjected to a proportion of benthically-derived enriched  $\delta^{13}\text{C}$ . There were no obvious effects of marine reserve status on the isotopic signatures of study taxa with the exception of slightly enriched  $\delta^{13}\text{C}$  of kina and snapper at Leigh, and of kina at Tawharanui.

Otolith microchemistry results for parore and snapper indicate strong connectivity between reef and non-reef systems within the wider Hauraki Gulf ecosystem. The majority of fishes sampled (both species) were likely to have originated as juveniles from lower salinity water environments such as estuaries fringing the Gulf. For snapper, our data suggest that only a small percentage of juveniles derive from reefs themselves. However, greater sampling replication is now required across a range of reef sites to better define the ratio of reef- versus estuary-derived juveniles, given the low percentage of reef-derived snapper.

A qualitative model of northeast New Zealand rocky reef ecosystems was developed to explore the complexity of interactions amongst New Zealand rocky reef species and the impacts of exploitation. This model was developed on the basis of a review and summary of interactions among reef components. A key modelling outcome was the highly predictive but opposite responses by small lobsters and large predatory invertebrates to changes in the abundance of a range of other groups. This suggests that these two groups are ideal candidates as variables for monitoring reef ecosystem responses to perturbations. The modelling agreed with a well documented example of responses to a perturbation in fishing pressure in the Leigh Marine Reserve. However, the predictability was low for all responses. This implies, for example, that the reduction of kina in the Leigh Marine Reserve and the subsequent increase in macro-algae consequent to an increase in lobster abundance may not necessarily occur in another area.

Field sampling at ten rocky reef sites across the Hauraki Gulf revealed differences among sites in community structure of macroalgae and invertebrates within all habitat strata. Of the environmental factors available, depth followed by a measure of water clarity (mean secchi) explained the most variation in the dependent variables (invertebrate taxa) from the quadrat data. Fish abundance data showed a similar, though weaker, trend across sites with depth, distance across the Gulf, and water clarity being the most important factors. The strong association between depth and water clarity and abundances of key taxa was expected and is similar to that found in earlier studies. With the exception of crayfish, there was no apparent overall relationship between invertebrate and fish abundances and marine reserve status of study sites, though the baited underwater video data showed snapper to be significantly larger within marine reserve sites than at fished sites.

Stable isotope analysis of tissue samples collected from key species from all study sites allowed insight into the functional relationships among species as well as dietary sources of carbon. Many of the study taxa, from the primary producers through to the predators, had the most depleted  $\delta^{13}\text{C}$  values at the furthest inshore and offshore sites (e.g. Poor Knights and Long Bay) and the highest  $\delta^{13}\text{C}$  values at the coastal sites (e.g. Leigh, Tawharanui and Kawau). Without direct modelling of end point source signatures we cannot definitively determine the percentage contribution of each carbon source. However, we suggest that the depleted  $\delta^{13}\text{C}$  of taxa from offshore sites is the result of a pelagic source of C and the enriched  $\delta^{13}\text{C}$  at coastal sites is the result of a more benthic input of C than at offshore sites, with sources including kelp detritus. Taxa at the inner gulf sites are also likely to be subjected to a proportion of benthically-derived enriched  $\delta^{13}\text{C}$ . There were no obvious effects of marine reserve status on the isotopic signatures of study taxa with the exception of slightly enriched  $\delta^{13}\text{C}$  of kina and snapper at Leigh, and of kina at Tawharanui.

Otolith microchemistry results for parore and snapper indicate strong connectivity between reef and non-reef systems within the wider Hauraki Gulf ecosystem. The majority of fishes sampled (both species) were likely to have originated as juveniles from lower salinity water environments such as estuaries fringing the Gulf. For snapper, our data suggest that only a small percentage of juveniles derive from reefs themselves. However, greater sampling replication is now required across a range of reef sites to better define the ratio of reef- versus estuary-derived juveniles, given the low percentage of reef-derived snapper.

***ZBD2008-07 Carbonate Sediments: The positive and negative effects of land-coast interactions on functional diversity (complete):***

Land-coast interactions can profoundly influence coastal biodiversity and ecosystem function. Estuarine primary productivity derived from phytoplankton, resuspended phytobenthos, aquatic vegetation and fringing habitat plant material is exported to the adjacent coast on outgoing tides and contributes to secondary production in the vicinity of the estuary mouth. However, land-derived sediments and contaminants that are discharged from estuaries can also stress open coastal populations. The balance of these competing processes was evaluated using a combination of laboratory and field investigations. A survey of two coastal locations (outside Whangapoua and Tairua harbours on the Coromandel Peninsula, New Zealand) quantified shifts in community structure in mollusc-dominated habitats and demonstrated that both distance from the mouth of the estuary and the size and density of large shellfish living in the sediments affect the composition and functionality seafloor communities. Tracing the importance of different estuary-derived food resources (seagrass, mangrove, estuarine phytoplankton and phytobenthos) using stable isotopes emphasized the importance of estuarine productivity to coastal bivalve. The work in the field has been supplemented with laboratory feeding trials, with the goal of verifying isotopic uptake rates in bivalve body tissues in a carefully controlled experimental setting. Trophic connections have important effects on coastal biodiversity. Understanding ecosystem processes and dynamics and their implications for functional biodiversity emphasises the importance of shifting the management focus from exploitation to resilience. Enhancing or maintaining this biodiversity will require more integrative ecosystem-based management focused on maintaining the resilience of coastal ecosystems.

Other research relevant or specifically linked to the projects above, are listed in Table 11.3.

**Table 11.3: Other research linked to investigation of the role of biodiversity in the functional ecology of nearshore and offshore marine communities.**

<b>MPI</b>	ZBD2005-04 Information on benthic impacts in support of the Foveaux Strait Oyster Fishery Plan ZBD2005-15 Information on benthic impacts in support of the Coromandel Scallops Fishery Plan ENV2005-23 Monitoring recovery of the benthic community between North Cape and Cape Reinga BEN2007-01 Assessing the effects of fishing on soft sediment habitat, fauna, and processes HAB2007-01 Biogenic habitats as areas of particular significance for fisheries management
<b>CRI Core purpose</b>	C01X1005— Management Of Cumulative Effects Of Stressors On Aquatic Ecosystems ; CO1X0907 Coastal Conservation Management, Freshwater and Estuaries and Coasts and Oceans
<b>DOC</b>	Conservancy surveys
<b>BNZ</b>	Biosecurity surveys
<b>OTHER</b>	Universities
<b>EMERGING ISSUES</b>	
Cumulative footprint of human activities; understanding cumulative impacts and risks; marine spatial planning	
Land-base effects on marine biodiversity and inshore/offshore habitats; pollution in offshore	
Ecosystem-based management and integrative governance	
Defining marine ecosystem services, linking them to ecosystem function and societal values	

#### **11.3.4. Progress on Science Objective 4. Marine genetic biodiversity**

Genetic biodiversity can be measured directly by measurement made at the genes and chromosomes scale or indirectly by measuring physical features at the organism scale (assuming they have a genetic basis).

Genetic diversity is fundamental to the long-term survival, stability and success of a species. Central to this is the “metapopulation” concept where populations are sufficiently genetically distinct from each other to be identifiable as individual units. A low level of recruitment between populations counters the effects of random genetic drift and inbreeding depression of genetic diversity.

Human activities can profoundly affect genetic diversity both within populations and between populations. For example, shipping activity (movement across the globe) and aquaculture practices (transfer of organisms to different areas) can increase population connectivity such that genetic biodiversity may decrease between populations. In extreme cases, populations can become the same genetically (homogeneous) although considerable within population diversity may remain. In the event of increased genetic connectivity, a species may become more susceptible to extinction through biological or catastrophic stochasticity. That is, in the absence of between population diversity there is insufficient genetic variance to adapt to the effects of climate change, disease epidemics and so on.

In contrast, under the much more common scenario of habitat fragmentation caused by human activities (fishing, pollution), decreased connectivity between populations will result in greater between-population diversity, but a reduction of within-population diversity. This also results in a decrease in a species survival (fitness) because fragmented or isolated populations may become extinct through environmental and genetic stochasticity or localised depletion. Periodic fluctuations in annual temperature for example can lead to small scale population extinction, which in the absence of recruitment between populations will result, over time, in the demise of all populations.

To reduce the risk of species loss information about the genetic diversity both within populations (population isolation) and between populations (population connectivity) is needed. Without such information, the effects of perturbation on a species persistence and survival cannot be predicted. Furthermore, the links between genetic diversity, the dispersal capacity (mode of reproduction and life history development) of a species and the minimum viable population (MVP) size required in the marine environment to ensure population persistence, are little understood. For example, the MVP size for a species with a large dispersal capacity is likely to be quite different from that of a species with a relatively restricted dispersal capacity. Examining the connectivity between populations in the marine environment is fundamental to resolving some of the central challenges in ecology and has almost been ignored in the management of New Zealand fisheries or protection of biodiversity.

#### **Projects**

##### ***ZBD2002-12 Molecular identification of cryptogenic/invasive marine species – gobies.***

Project complete. (Lavery *et al.* 2006.)

##### ***ZBD2009/10 Multi-species analysis of coastal marine connectivity.***

An extensive literature review of published and unpublished information about connectivity of New Zealand coastal biota has been completed. Reviews were made of 58 studies of 42 taxa to identify the taxon or taxa studied, the habitat where each study took place, and geographic location of sampling sites used by each study. From these data, gaps in knowledge about taxa, habitats and spatial coverage of sampling were identified. Recommendations about four species to be studied, habitats that they should be collected from, and location of sampling sites were made. Recommendations included a standardised collecting protocol and

for the development and application of microsatellite markers to quantify the population genetic structure and the coastal connectivity of these taxa (Gardner *et al.* 2010).

Two PhD students are carrying out field work, genetic analyses, and writing up (in the form of two theses) of this research. Both studies are underway. Fieldwork has begun on two flatfish species and two species of shellfish. The project been extended to incorporate a subtidal species of shellfish. A new component of the coastal connectivity project has been added to include work on the New Zealand scallop, *Pecten novaezelandiae*. This work focuses on population genetic structure and genetic connectivity at two different spatial scales and uses microsatellite markers (consistent with the use of microsatellite markers for the 4 species already under investigation in the original ZBD2009-10 project). First, the extension work focusses on scallops in the Hauraki Gulf and Coromandel Peninsula region. Scallops have been collected from several populations in this region and further samples will be added in the next two years. Second, the extension work focuses on scallops across New Zealand (the full range of this species' distribution). Samples have been sourced from several regions including the fiords, the far north, and central New Zealand. In both cases, genetic connectivity will be assessed to determine linkages among populations at the two different spatial scales. The smaller spatial scale information will be of particular relevance to the scallop fishery in the Hauraki Gulf and Coromandel Peninsula region, whereas the larger scale work will complement ongoing studies of coastal connectivity at the national scale already under examination as part of the project. A PhD student has been recruited for this work and a suite of microsatellite markers has been developed for the New Zealand scallop and testing of population genetic variation is underway

Other research relevant or specifically linked to the projects above, are listed in Table 11.4.

**Table 11.4: Other research linked to marine genetic biodiversity.**

<b>MPI</b>	ENH2007-01 Stock enhancement of blackfoot paua GEN2007-01 Genetic population profile of blackfoot paua ENH2007-02 Outbreeding depression in invertebrate populations IPY2007-01 Objective 11. Barcode of life
<b>MBIE</b>	C01X0502 Biodiversity& Biosecurity
<b>MPI</b>	Base line surveys for non-indigenous species
<b>OTHER</b>	Universities [?]
<b>BRAG PROJECTS FOR 2011-12</b>	
Extension to ZBD2009-10 to include subtidal shellfish	
<b>EMERGING ISSUES</b>	
Can genetics combined with hydrographic models usefully contribute to the identification of biodiversity hot-spots and/or to source-sink relationships within ecosystems?	

### **11.3.5. Progress on Science Objective 5. Effects of climate change and variability on marine biodiversity**

Cyclical changes or trends in climate and oceanography and associated effects such as increased ocean acidification and how they affect the marine ecosystem as a whole have long-term implications for trophic interactions and biodiversity, as well as functional aspects of the system e.g. biogeochemical processes. With significant improvement in remote sensing tools and global monitoring of climate change, new patterns are emerging indicating that there are long-term cycles. Examples include the Interdecadal Pacific Oscillation as well as shorter periods of change in relation to the El Niño Southern Oscillation that affect ocean ecosystems. Further, physical phenomena such as the deep subtropical gyre 'spin-up' in the South Pacific which resulted in a warmer ocean around New Zealand from 1996–2002, can have flow-on effects on ecosystem functioning.

A new report was launched in 2010 by the United Nations on ocean acidification<sup>59</sup> Among other findings, the study shows that increasing ocean acidification will mean that by 2100 some 70% of cold water corals, (a key refuge and feeding ground for some commercial fish species), will be exposed to corrosive waters (see also Tracey *et al.* 2011). In addition, given the current greenhouse gas emission rates, it is predicted that the surface water of the highly productive Arctic Ocean will become under-saturated with respect to essential carbonate minerals by the year 2032, and the Southern Ocean by 2050 with disruptions to large components of the marine food source, in particular those calcifying species, such as foraminifera, pteropods, coccolithophores, which rely on calcium carbonate.

Emerging research suggests that many of the effects of ocean acidification on marine organisms and ecosystems will be variable and complex and will affect different species in different ways. Evidence from naturally acidified locations confirms, however, that although some species may benefit, biological communities in acidified seawater conditions are less diverse and calcifying (calcium-reliant) species are absent whereas algae tend to dominate.

Many questions remain regarding the biological and biogeochemical consequences of ocean acidification for marine biodiversity and ecosystems, and the impacts of these changes on ecosystems and the services they provide, for example, in fisheries, coastal protection, tourism, carbon sequestration and climate regulation.

Studies to predict changes in biodiversity in relation to climate change in more than a rudimentary way are beyond the state of current knowledge in New Zealand. Nevertheless, surveys of biodiversity that have occurred or are planned will provide a snapshot against which future research results or trends can be compared.

Meeting the challenges of climate change and identifying crucial issues for marine biodiversity is an area of high political interest internationally<sup>60</sup> and has been identified as a gap in biodiversity research in New Zealand<sup>61</sup>

## Projects

### ***ZBD2005-05 Long-term effects of climate variation and human impacts on the structure and functioning of New Zealand shelf ecosystems.***

This is a large scale project to investigate changes in shelf ecosystems over a 1000 year time-scale to provide context and perspective on issues of natural variation versus human impacts on marine biodiversity.

The project is a multidisciplinary study to collate and synthesise paleoecological, archaeological, historical, and contemporary data relating to changes in the structure and functioning of New Zealand shelf ecosystems since human arrival about 750 years ago. The data have been used to model present and four past states of the Hauraki Gulf ecosystem over the last 1000 years.

The project is collaborating with the international History of Marine Animal Populations (HMAP) project, itself a part of the Census of Marine Life (CoML) programme. A short video about the NZ Taking Stock project was made by HMAP staff and is currently available on the HMAP website <http://hmapcoml.org/projects/nz/>.

---

<sup>59</sup> <http://www.un.org/apps/news/story.asp?NewsID=36941&Cr=emissions&Cr1> Downloadable Report The Environmental Consequences of Ocean Acidification

<sup>60</sup> <http://biodiversity-l.iisd.org/news/ungas-second-committee-considers-biodiversity-and-sustainable-development/>

<sup>61</sup> Green, W.; Clarkson, B. (2006). Review of the New Zealand Biodiversity Strategy Themes

Fifteen reports stemming from this project have been submitted to the Ministry and are at various stages of review, acceptance and publication. Four reports are still to be delivered. The report most relevant to this section is Pinkerton (In press). Other reports published to date are Carroll *et al.* (In press); Jackson *et al.* (In Press); Lalas *et al.* (In Press) a; b; Lalas & MacDiarmid (In Press); Lorrey *et al.* (In Press); MacDiarmid *et al.* (In Press a; b); Maxwell & MacDiarmid (In Press); Neil *et al.* (In Press); Paul (2012); Parsons *et al.* (In Press); Smith (2011).

**ZBD2008-11 Predicting plankton biodiversity & productivity with ocean acidification.**

This multi-year project is inter-linked with the Coasts and Oceans OBI and has the following objectives:

1. To document the spatial and inter-annual variability of coccolithophore abundance and biomass, and assess in terms of the phytoplankton abundance, biomass and community composition in sub-tropical and sub-Antarctic water.
2. To document the seasonal and inter-annual variability of foraminifera and pteropod abundance and biomass at fixed locations in sub-tropical and sub-Antarctic water by analysis of sediment trap material from time-series data collection.
3. To document the spatial and seasonal distribution of the key coccolithophore species, *Emiliana huxleyi*, using both archived and ongoing ingestion of satellite images of Ocean Colour, and ground-truth the reflectance algorithm for *E. huxleyi* for future application in New Zealand waters
4. To determine the sensitivity of, and response of *E. huxleyi* and other EEZ coccolithophores to pH under a range of realistic atmospheric CO<sub>2</sub> concentrations in perturbation experiments, using monocultures and mixed populations from in situ sampling.
5. To document the spatial variability of diazotrophs (nitrogen-fixing organisms) and associated nitrogen fixation rate, and assess in terms of phytoplankton abundance, biomass and community composition in sub-tropical waters north of New Zealand.
6. To determine the sensitivity of diazotrophs to ocean acidification composition in sub-tropical waters north of New Zealand.

The project is proceeding according to plan and is still primarily in the sample collection phase with some data analysis but limited interpretation to date. The biodiversity record of coccolithophore species in New Zealand waters has been extended, with a transect across the Tasman Sea and a number of transects across the Chatham Rise. A bloom of the coccolithophore *Emiliana huxleyi* on the Chatham Rise was extensively characterised in terms of surface water biogeochemistry, and subsequently successfully cultured in the lab. Seasonal and interannual variability of *E. huxleyi* blooms were further characterised by extending the true colour satellite image analysis of presence/absence of coccolithophore blooms in the New Zealand EEZ. This was augmented by sample collection for ground-truthing of published calcite algorithms (for satellite detection of coccolithophore blooms) and application of a published calcite algorithm to New Zealand waters for 2002-3. Coccolithophore acidification sensitivity experiments were run in the Tasman Sea and the Chatham Rise region, with preliminary analysis indicating a decline in coccolithophore abundance under high CO<sub>2</sub>, but not when accompanied by elevated temperature as predicted under future climate change scenarios. Analysis of sediment trap samples for pteropod and foraminifera identification and abundance was completed for 2000-2010, with significant interannual variability noted in both, but also some indication of a recent decline in pteropod abundance in Sub-Antarctic water. Sample analysis from the 2010 Tasman Sea voyage identified the presence of nitrogen-fixing unicellular cyanobacteria and significant nitrogen fixation south of the Tasman Front, in contrast to previous observations. In acidification sensitivity experiments on this voyage nitrogen fixation did not change or decreased under high CO<sub>2</sub> concentrations, in contrast to published data. Outputs to date include Boyd *et al.* (in press).

**ZBD2009-13 Ocean acidification impact on key NZ molluscs.**

Ocean acidification associated with increased atmospheric CO<sub>2</sub> levels is a pressing threat to coastal and oceanic ecosystems. The chemical reaction which occurs when this CO<sub>2</sub> is dissolved in seawater results in a well documented decrease in seawater pH (and an increase in seawater

acidity), which may physically dissolve  $\text{CaCO}_3$  shells and/or skeletons and affect the shell/skeleton generation, as well as influencing many other physiological processes. Flow on effects to the viability of populations and the economic benefit that can be derived from commercially important species are likely. There is very little information on how key NZ calcifying species will respond to this change.

This project is using laboratory experiments to quantify responses of key New Zealand mollusc species (paua, *Haliotis iris*, cockles, *Austrovenus stutchburyi*, and oysters *Tiostrea chileensis*) to levels of ocean  $\text{CO}_2$  saturation predicted to occur in NZ waters over the following decades. Results will be combined with information on the role of these key species in influencing ecosystem structure and function, to assess local and ecosystem-scale implications of acidification of NZ coastal waters expected in the following decades.

**ZBD2010-41. Potential effects of ocean acidification on habitat forming deep-sea corals in the New Zealand region.**

Specific Objectives of this research were to 1. Determine the carbonate mineralogy of selected deep-sea corals found in the New Zealand region, 2. Assess the distribution of deep-sea coral species in the region relative to improved knowledge of current and predicted aragonite (ASH) and calcite saturation horizons (CSH), and 3. Assess potential locations vulnerable to deepwater upwelling and areas of key deep-water fishery habitat. Through a literature search and analysis, the project aimed to determine the most appropriate tools to age corals and measure the effects of ocean acidification on deep-sea habitat-forming corals, and recommend the best approach for future assessments of the direct effects of declining ocean pH on these key fauna.

Under Objective 1, new results of investigations into the carbonate mineralogy of selected deep-sea corals found in the New Zealand region were presented, and previous work on coral mineralogy summarised. The mineralogy and trace element concentration (Sr and Mg) of the five branching stony coral species (Order: Scleractinia) *Goniocorella dumosa*, *Solenosmilia variabilis*, *Enallopsammia rostrata*, *Madrepora oculata*, and the endemic *Oculina virgosa*, and for the key habitat forming gorgonian coral species (Order: Alcyonacea) *Keratoisis* spp., *Lepidisis* spp., *Paragorgia* spp. and *Primnoa* sp., was ascertained. Stony branching corals are all aragonitic with high Sr and low Mg while most of the gorgonian corals are made of high Mg and low Sr, with high Mg calcite (>8 mol% Mg). The gorgonian sea fan, *Primnoa* sp., is aragonitic.

Under Specific Objective 2, up to date position and depth data were used to produce distribution maps for the study species. Data compare well with previous publications from biodiversity research, research trawl, and observer sampling effort on wide regional distribution, but individual species display variations within the region. The peak depth distributions are unimodal at about 800-1000 m for most of the above species, but *G. dumosa*, *E. rostrata*, and *Lepidisis* spp. show bi-modal distributions and *O. virgosa* occurs primarily in shallow depths. In the second year of the project these distribution data will be compared with existing and predicted aragonite and calcite saturation horizons, particularly in areas of key deepwater fishery habitat.

Also under Specific Objective 2, on-going opportunistic water sampling analyses are being carried out to determine alkalinity and dissolved inorganic carbon (DIC), and modelling to determine aragonite (ASH) and calcite saturation horizon (CSH) data is in progress. The aim is to compare water carbonate chemistry with regional biogeochemistry models and future scenarios to identify areas potentially at risk from ocean acidification.

Under Specific Objective 3, at-sea sampling of live corals for aquarium studies has been carried out to investigate the feasibility of keeping the corals alive for growth and ocean acidification experiments. The corals collected in April, 2012 are still alive in the laboratory and include one small colony of *S. variabilis*. A literature search and analysis to determine the most appropriate tools to age and measure the effects of ocean acidification on deep-sea habitat-forming corals is

in progress. From these trials and reviews, recommendations on the best approach for future assessments of the direct effects of changes in ocean pH on these key fauna will be made.

Other research relevant or specifically linked to the projects above, are listed in Table 11.5.

**Table 11.5: Other research linked to effects of climate change and variability on marine biodiversity.**

<b>MPI</b>	SAM2005-02 Effects of climate on commercial fish abundance ENV2007-04 Climate and oceanographic trends relevant to New Zealand fisheries
<b>MBIE</b>	C01X502 Coasts & Oceans Centre
<b>DOC</b>	Baseline surveys; protected deepsea corals (Tracey <i>et al.</i> 2011; Baird <i>et al.</i> 2012)
<b>OTHER</b>	University of Otago-NIWA shelf carbonate geochemistry and bryozoans Geomarine Services-foraminiferal record of human impact Regional Council monitoring programmes
<b>EMERGING ISSUES (this objective)</b>	
What papers can be generated on the effects of climate change on marine biodiversity in NZ in time for 5 <sup>th</sup> IPCC report?	
How does climate change influence marine microbial diversity, species mix and biogeochemical roles?	
How will harmful toxic algal blooms be affected by warming seas? (e.g. Chang 2003, Chang <i>et al.</i> 2003)	

### 11.3.6. Progress on Science Objective 6. Biodiversity metrics and other indicators for monitoring change

In the mid 1990s, monitoring of marine biodiversity and the marine environment was a topic of considerable discussion, yielding several reports on developing MfE indicators<sup>62</sup> However, since the publication of MfE's indicators in 2001, a much reduced set of core indicators that relate to the marine environment have been reported on<sup>63</sup>. A new international initiative launched in 2010 "Biodiversity Indicators Partnership"<sup>64</sup> provides guidelines and examples of biodiversity indicators developed around the globe, however, Oceania does not appear to have any partnership identified. The link between this initiative and OECD environmental indicators is unclear.

A serious gap identified by Green and Clarkson (2006)<sup>65</sup> in their review of progress on implementation of the NZBS was the lack of development of an integrated national monitoring system (see Biodiversity Research Programme 2010: Part 4). Efforts to respond to this gap within the Biodiversity Programme resulted in the immediate initiation of a 5-year Continuous Plankton Recorder project, and a series of workshops to determine how best to approach monitoring on a national scale (ZBD2008-14). [One objective of monitoring would be to test the effectiveness of management measures.]

#### Projects

**ZBD2004-10 Development of bioindicators in coastal ecosystems.**

<sup>62</sup> Downloadable MfE reports [Confirmed indicators for the marine environment](#) 2001, ME398; [An analysis of potential indicators for marine biodiversity](#) 1998 TR44; [Environmental Performance Indicators: an analysis of potential indicators for fishing impacts](#) 1998 TR43; [Environmental Performance Indicators: Summary of Proposed Indicators for the Marine Environment](#) 1998, ME296; [Environmental Performance Indicators: Marine environment potential indicators for physical and chemical processes, and human uses and values](#) 1998 TR45; [Potential coastal and estuarine indicators - a review of current research and data](#) 1997 TR40; [Monitoring and indicators of the coastal and estuarine environment - a literature review](#) 1997 TR39

<sup>63</sup> <http://www.mfe.govt.nz/environmental-reporting/about/tools-guidelines/indicators/core-indicators.html>

<sup>64</sup> [www.bipnational.net/IndicatorInitiatives](http://www.bipnational.net/IndicatorInitiatives)

<sup>65</sup> Green, W.; Clarkson, B. (2006). Review of the New Zealand Biodiversity Strategy Themes.

Project complete (Savage 2009). Agricultural and urban development can increase run-off and lead to excessive nutrient loadings in fragile coastal environments that are nursery grounds for a diverse array of coastal and estuarine species, as well as other resident organisms. This project investigated the development of bioindicators to strengthen the ability of managers to detect and quantify changes in anthropogenic nitrogen inputs to coastal and estuarine ecosystems by comparing six study sites with different levels of development ranging from pristine through to fully urban. The results show a strong positive relationship between the percent agricultural land in surrounding catchments and total nitrogen (TN) loading to nearshore environments.

These results also hint at differences in dissolved and particulate nitrogen source pools, and highlight the importance of using complementary components of food webs and high spatial replication to show linkages between watershed land use and chemical markers in biota. The effects of nutrient enrichment were transmitted up the food web, with growth of secondary consumers, *Notolabrus celidotus* (spotties) and *Grahamina nigripenne* (estuarine triplefins) generally enhanced in nutrient enriched coastal areas. Benthic prey dominated the diets of these fish species, with amphipods and brachyurans being the most important prey items for triplefins and spotties, respectively. However, there were site-specific differences in prey importance and diet diversity. Both triplefins and spotties consumed considerably more diverse prey items at pristine than nutrient-enriched coastal areas. Food web models based on stomach content analyses and dual isotope ratios suggest that there are shifts in the relative importance of the different organic matter sources supporting food structure among the different coastal ecosystems due to nutrient enhancement from land-based activities. [how might these results be used in a biodiversity management context?]

**ZBD2008-14 What and where should we monitor to detect long-term marine biodiversity and environmental changes?**

Two workshops and a follow up meeting were held with stakeholders in 2008/09 to discuss a marine environmental monitoring programme (MEMP) for New Zealand, to detect long-term changes in the marine environment, building on existing time series and data collection (Livingston 2009). The MEMP was formulated into a developmental project staged over 3 years and submitted to the former Ministry of Research Science and Technology's Cross Departmental Research Pool (CDRP) for funding starting July 2010. Since that time, CDRP funding has been withdrawn. Instead a call for proposals taking a more modest approach to developing MEMP beginning with collation of all potential data series into a metadata database, a scientific evaluation of the existing time series as to their 'fit to purpose' for MEMP was made and tender evaluations are underway.

Monitoring change in the marine environment is the only way we can measure long-term trends, mitigate risk and provide evidence of changes which may require policy or management practice response. DOC has since been developing an integrated approach to monitoring biodiversity particularly on the land but also in marine reserves<sup>66</sup>.

**ZBD2008-15 Continuous Plankton Recorder Project: implementation and identification.**

This project adopts the methods used in a long-term programme that has proved highly relevant to measuring biological changes in the ocean, i.e., the Continuous Plankton Recorder Programme in the North Atlantic (SAHFOS) and more recently the Southern Ocean<sup>67</sup>. This 5-year MPI project aims to map changes in the quantitative distribution of epipelagic plankton, including phytoplankton, zooplankton and euphausiid (krill) life stages annually when vessels depart and return from New Zealand on their journey to the Ross Sea toothfish fishery each year in November/December, and February/March traversing key water masses and ocean fronts in New

<sup>66</sup> The Department of Conservation Biodiversity Monitoring and Reporting System Fact Sheet July 2010

<sup>67</sup> Southern Ocean CPR programme <http://data.aad.gov.au/aadc/cpr/>

Zealand's EEZ as well as south to the Ross Sea. Four years the transects have been collected, staff have been trained in plankton ID work and over three years of samples analysed.

#### **ZBD2010-42 Marine Environmental Monitoring Programme.**

This project continues from ZBD2008-14. A starting point to the assessment and reporting of broad-scale changes in New Zealand's marine environment is to define basic criteria and locate all existing and past time series of marine environmental data to improve awareness and access to these data. After this, these data can be evaluated as to their fitness-for-purpose for contributing towards a national Marine Environmental Monitoring Programme (MEMP). To date an online catalogue has been designed and a portal to this is available at <http://geodata.govt.nz>. Questionnaires were developed to determine what marine environmental time series data were available within New Zealand. Information to date gives us 131 databases, 50% of these are listed as having ongoing funding (although not necessarily for all locations), and another 19% are listed as likely to continue. Over 70% are publically available. Most cover more than one location, although this is dependent on how the databases are constructed, e.g., DOC at present has a separate database for each marine reserve, while regional councils tend to have separate databases for different subjects (e.g., contaminant monitoring, ecological monitoring). Around 95 estuaries and harbours are being sampled, which is not surprising given that the majority of the information comes from Regional Councils (Figure 1). There are 78 coastal locations and 33 marine reserves.

The second phase, determining fitness-for-purpose, was begun at a workshop held at NIWA on 11th June (see objective 3). Priority variables for inclusion in a national monitoring programme have been identified from responses to a questionnaire sent to scientific experts and central and regional government departments involved in monitoring and/or reporting. Core reference sites and major gaps in the spatial network are presently being determined and the requirements for spatial and temporal sampling determined. The project is due to be completed by June 2013.

Other research relevant or specifically linked to the projects above, are listed in Table 11.6.

**Table 11.6: Other research linked to biodiversity metrics and other indicators for monitoring change.**

<b>MPI</b>	ENV2006-15: Database and fishing indicator on seamount habitats (Rowden <i>et al</i> 2008) BEN2009-02 (Tuck <i>et al.</i> 2010) ENV2006-04: Fisheries indicators from trawl surveys (Tuck 2009) DEE2010-05
<b>MBIE</b>	Core funding for Coasts and Oceans Centre
<b>DOC</b>	Conservancy projects-Hawke's Bay;
<b>OTHER</b>	Regional Councils, Universities
<b>EMERGING ISSUES</b>	
Monitoring coastal waters and New Zealand's oceans to report on a national scale remains a major gap	
There is little longterm commitment to direct monitoring the marine environment	

### **11.3.7. Scientific Objective 7. Identifying threats and impacts to biodiversity and ecosystem functioning**

Many marine ecosystems in New Zealand have been modified in some way through the harvesting of marine biota, the selective reduction of certain species and size/age classes, modification of food webs, including the detrital components and habitat destruction. Benthic communities including seamount communities, volcanic vent communities, bryozoans, corals, hydroids and sponges are vulnerable to human disturbance. The mechanical disturbance of marine habitats that occurs with some activities such as trawling, dredging, dumping, and oil, gas and mineral exploration and extraction; can substantially change the structure and composition of benthic communities. The

invasion of alien species into New Zealand waters is also a real threat, with evidence of nuisance species already well established<sup>68</sup>

A number of inshore marine ecosystems (especially estuaries and other sheltered waters) have been modified by sediment, contaminants and nutrients derived from human land use activities (Morrison *et al.* 2009). Coastal margin development has had a major impact on some inshore marine communities.

A recent project commissioned by the MPI Aquatic Environment Programme, identifies key threats to the marine environment (BEN2007-05) is complete and has listed and ranked the top threats to New Zealand's marine environment, as perceived by expert opinion. Relevant findings are that the highest ranking threats are ocean acidification, increasing sea water temperatures and bottom trawling (across all habitats) and that the most threatened habitats are intertidal reef systems in harbours and estuaries (MacDiarmid *et al.* 2012). Ecological risk assessment (ERA) methods have also been reviewed (under ENV200515, Rowden *et al.* 2008), and a trial Level 2+ assessment completed on Chatham Rise seamounts to estimate the relative risk to seamount benthic habitat from bottom trawling (under ENV200516, Clark *et al.* 2011). An MPI project (DEE2010-04) has resulted in a new ecological risk assessment being developed that is tailored for New Zealand deepwater fisheries (Clark *et al.* in press).

## Projects

**ZBD2009-25 Predicting impacts of increasing rates of disturbance on functional diversity in marine benthic ecosystems.** The objectives of this project are to:

1. Further develop landscape/seascapes ecological model of disturbance/recovery dynamics in marine benthic communities, incorporating habitat connectivity, based on existing model by Lundquist *et al.* (2010).
2. Predict impacts of increasing rates of disturbance on rare species abundance, functional diversity, relative importance of biogenic habitat structure, and ecosystem productivity.
3. Use literature and expert knowledge to quantify rare species abundance, biomass, functional diversity, habitat structure, and productivity of various successional community types in the model.
4. Field test predictions of the model in appropriate marine benthic communities where historical rates of disturbance are known, and benthic communities have been sampled.

The baseline model, incorporating connectivity, has been created in Matlab. Objective 2 (predictions for functional biodiversity based on model) is underway. Some progress has been made on objective 3 (quantify functional biodiversity from existing data) through familiarisation of the programmers with the datasets of the Ocean Survey 2020 Chatham/Challenger project (ZBD2007-01) and biodiversity analyses to date for objective 8 of that project. Objective 4 is in process, with the majority of the field test funded by BEN2007-01. Researchers from both projects have met to discuss and modify the draft sampling design in order to best allocate sampling to test the predictions of the functional diversity model. The field testing took place in March-April 2010 in Tasman/Golden Bay.

Other research relevant or specifically linked to the projects above, are listed in Table 11.7.

---

<sup>68</sup> <http://www.biosecurity.govt.nz/biosec/camp-acts/marine>  
<http://www.biosecurity.govt.nz/pests/salt-freshwater/saltwater>  
<http://www.biosecurity.govt.nz/about-us/our-publications/technical-papers>

**Table 11.7: Other research linked to threats to and impacts on biodiversity.**

<b>MPI</b>	BEN2007-05 Assessment of anthropogenic threats to New Zealand marine habitats. MacDiarmid <i>et al</i> 2012 DEE2010-04
<b>MBIE</b>	CO1X0906 Vulnerable deep-sea communities (mapping and sampling a range of deep-sea habitats (seamounts, slope, canyons, seeps, vents), and determining relative risk to their benthic communities from human activities
<b>EMERGING ISSUES</b>	
The socio-economic valuation of biodiversity in NZ has not been adequately addressed.	
The cumulative footprint of anthropogenic activities on the NZ marine environment has not been assessed. Potential development of seabed mining makes this a priority in deepwater environments as well as coastal.	

### 11.3.8. Biodiversity in Antarctica: BioRoss Project Summaries and Progress

The objectives of BioRoss are to improve understanding of the biodiversity and functional ecology of selected marine communities in the Ross Sea. These objectives are being achieved by commissioning directed research on the diversity and function of selected marine communities in the Ross Sea region. BioRoss is committed to linking with ongoing Ross Sea ecosystems research through the Antarctic Working Group, and supporting climate change related research, especially at high latitudes.

Data acquisition from the Antarctic marine environment is logistically difficult and expensive. Nevertheless, the seven biodiversity Science Objectives listed above also drive BioRoss research projects. The BioRoss survey in 2004 and the Latitudinal Gradient Project ICECUBE have provided significant new information on biodiversity, species abundance and distribution that are now facilitating research into functional ecology and longer term monitoring programmes. This research has the potential to lead into other research on genetic diversity, climate variability and the development of indicators. The research results are also being used in the MPI Antarctic Research Programme projects on ecosystem modelling of the Ross Sea.

The MPI Antarctic Research and BioRoss Programmes are also directly involved in supporting the development of protection measures around the Balleny Islands. In 2005 MPI scientists and Ministry of Foreign Affairs and Trade (MFAT) personnel prepared a paper for submission to CCAMLR justifying MPA designation around the islands to protect ecosystem processes occurring there that may be important for the stability and function of the wider Ross Sea regional ecosystem.

To collect data in support of the MPA proposal, MPI BioRoss funded a targeted research voyage to the Balleny Islands in February 2006 (ZBD2005-01), and also provided supplementary funding to carry out opportunistic biological sampling at the Balleny Islands on a voyage to the Ross Sea that was primarily funded by LINZ to do bathymetric mapping.

The field sampling of these projects were successful, both providing important data and specimens from the Balleny Islands area and supplementary information for the Antarctic Working Group Research Programme. The results will inform research planning for subsequent projects. Support for Ross Sea region biodiversity will remain a high priority for future research in the BioRoss Programme.

In addition, BioRoss funded a further ICECUBE project to sample the Antarctic coastline during the summer season of 2006/07 (ZBD2006-03). ICECUBE is a key part of the international Latitudinal Gradient Project to explore hypotheses about environmental drivers of structure and function in sub-tidal ecosystems along the western Ross Sea coastline (Cummings *et al.* 2008 ). This project acquired funding for three seasons (2007/08, 08/09, 09/10) as part of the MBIE IPY contestable round (see also Cummings *et al.* 2011 and Thrush and Cummings 2011). Published reports and papers from the MPI

Ross Sea coastal projects include Cummings *et al.* 2003, 2006, 2008, 2010, 2011. De Domenico *et al.* 2006, Grotti *et al.* 2008, Guidetti *et al.* 2006, Norkko *et al.* 2002, 2004, 2005, 2007; Pinkerton *et al.* 2006, Schwarz *et al.* 2003, 2005, Sharp *et al.* 2010, Sutherland 2008, Thrush *et al.* 2006, 2010 and in press.

The New Zealand Government provided one-off funding for a Census of Antarctic Marine Life (CAML) survey to the Ross Sea from *R.V. Tangaroa* as part of New Zealand's involvement in the 2007-08 International Polar Year activities. The CAML Voyage was a large cooperative research effort under the banner of Ocean Survey 20/20 with considerable international collaboration, simultaneously utilising a number of different vessels with different strengths and capabilities. Progress on the two projects IPY2007-01 and IPY2007-02, is detailed below.

## Projects

**ZBD2002-02 *Whose larvae is that? Molecular identification of planktonic larvae of the Ross Sea.*** Completed. (See Sewell *et al.* 2006, Sewell 2005, Sewell 2006.)

**ZBD2003-03 *Biodiversity of deepwater invertebrates and fish communities of the north western Ross Sea.*** Completed. Two AEBR reports were produced by Rowden *et al.* (2012a, in press) and a Voyage Report, Mitchell and Clark 2004. A number of papers have also been published in the scientific literature using specimens or data from the 2004 biodiversity survey (e.g. De Domenico *et al.* 2006, Schiaparelli *et al.* 2006, Rehm *et al.* 2007, Kröger & Rowden 2008, Clark *et al.* 2010)

**ZBD2005-01 *Balleny Islands Ecology Research, Tiama Voyage (2006).***

This voyage collected a large amount of new data from the Balleny Islands and surrounding waters using a range of methods, including bird and mammal observations, whale biopsy sampling, shore-based penguin colony surveys, SCUBA dive quadrats and transects, tissue collections for stable isotope analyses, and continuous acoustic/bathymetric data collection (Smith 2006). Some of the specimens and data have been used for other studies.

**ZBD2005-03 *Opportunistic biological data during 2006 Ross Sea voyage utilising Tangaroa.***

This project is complete (MacDiarmid and Stewart 2012). In brief it proved feasible to assess demersal fish abundance using the camera and lights. Because sampling was restricted to areas outside the main fishery, no toothfish were observed. The camera system, (a predecessor to the deep towed imaging system (DTIS)) proved capable of characterizing the demersal fish habitat associations. Sampling using a variety of methods yielded specimens and tissue samples of a wide variety of benthic and pelagic organisms. The acoustic information collected on water column organisms was less useful than desired because of interference from the bottom profiling aspects of the voyage. Marine mammals and seabirds were routinely recorded and automated sampling of the surface waters using a continuous plankton recorder and instruments to record sea surface temperature, salinity and chlorophyll-a concentration was successful.

**ZBD2008-23 *Macroalgae diversity and benthic community structure at the Balleny Islands.***

Project complete. As a result of this study, the known macroalgal flora of the Balleny Islands has increased from 13 to 27 species, and there are 2 new records for the Ross Sea in addition to the 3 new records reported by Page *et al.* (2001). The biodiversity however remains poorly known, and detailed comparisons with other parts of the Antarctic region would be premature. A high proportion of the taxa reported here are known from only one collection, with a further group of taxa known from either two or three collections. Many of the taxa cannot be fully documented as there is insufficient mature material available.

The samples collected as part of a benthic survey at Borradaile Island, one of the Balleny Islands group, during the 2006 *Tiama* expedition have been analysed to provide an assessment of benthic community structure. The Borradaile Island sites were located in a high energy environment, sediments had relatively high organic and chlorophyll *a* content, and considerably lower

concentrations of degraded plant material (phaeophytin) than noted in previously surveyed southern Ross Sea locations. Borradaile Island macrofaunal diversity was within the range noted for the more southern sites; macrofaunal abundance however, was more variable. Epifaunal diversity was very low, with the seastar *Odontaster validus* the only large epifaunal taxon found. In contrast, the Borradaile Island dive sites had high macroalgal diversity. Although not observed at these dive sites, the *Tiama* voyage researchers noted shallow water areas with high diversities of encrusting organisms. This study has provided the first analysis of shallow water benthic communities of the Balleny Islands. While it has shown some interesting similarities and contrasts in benthic diversity with other coastal Ross Sea locations, this information from Borradaile Island may not be representative of the entire Balleny area, and further surveys from other sites within the Balleny group are recommended (Nelson *et al.* 2010).

**ZBD2008-20 Ross Sea Ecosystem function: predicting consequences of shifts in food supply.**

Project complete. Detailed information on the uptake and incorporation of different primary food sources to key epibenthic species help predict consequences of potential environmental change. Over a two year period, *in situ* investigations into responses to, and utilisation of, primary food sources by a common ophiuroid, were conducted at two contrasting coastal Ross Sea locations, Granite Harbour and New Harbour. At both locations, benthic net primary production was measured and the contributions of large macrobenthic organisms to ecosystem functions such as organic matter processing and nutrient recycling were quantified. Granite Harbour benthic soft-sediments supplied overlying waters with regenerated ammonium and phosphate, and the ophiuroid significantly increased the rates of nutrient release. Ultimately, the nutrients will be used by microalgae in the water column and under the ice. Detrital algae (phaeophytin) were present in sediments at greater concentrations than fresh microalgal material (chlorophyll a), and appears to be functionally important; it was a significant predictor of dissolved oxygen, phosphate, ammonium and nitrate-plus-nitrite flux. Benthic organisms in predominantly ice covered Ross Sea locations such as Granite Harbour probably feed on degraded detrital algae for much of year, given the limited amount of fresh microalgae available due to the dimly lit environment, and the consequently low rates of *in situ* benthic primary production. Results of the New Harbour investigations contrast those of Granite Harbour, reflecting the very different ice conditions at these two locations (Cummings *et al.* 2010; Lohrer *et al.* 2012).

**IPY2007-01 NZ International Polar Year Census of Antarctic Marine Life**

Overall science objectives for the Project were developed by MPI, NIWA and other interested and participatory parties in discussions held through the Ocean Survey 20/20 Science Working Group.

1. To measure and describe the relationships between patterns of marine organisms, their biodiversity and environmental variables between longitudes ~170°E and ~175°W, and depths down to ~3500-4000m in the Ross Sea region.
2. To assess the trophic interrelationships of the major functional groups in the Ross Sea and regional ecosystem, with particular reference to improving inputs to ecosystem modelling.
3. To obtain baseline measures of the marine environment and identify a suite of ecosystem or environmental indicators that could potentially be used to monitor change in response to environmental or anthropogenic forcing in the Ross Sea region

All specific objectives apart from objective 2 have now been completed.

*Specific Objective 1:* To measure seabed depth and rugosity using the multibeam system (whenever possible) to identify topographic features such as bottom type, iceberg scouring, seamounts etc and to determine areas for targeted benthic fauna sampling. (not funded in this project). Objective Completed. (Mitchell 2008, Hanchet *et al.* 2008)

*Specific Objective 2:* To continue the analysis of opportunistic seabird and marine mammal distribution observations from this and previous BioRoss voyages and published records, and in relation to environmental variables. (Draft report completed.)

The distributions of the seabird and marine mammal taxa reported from two RV Tangaroa voyages (TAN200602 and TAN200802) have been mapped. These represent the count data of seabirds recorded during the 2006 Ross Sea voyage and the locations of images of seabird taxa (recorded opportunistically) from the 2008 IPY-CAML voyage and records from observers from the toothfish fishery. The distributions include the presence data of taxa over waters south of about 60° S to the Ross Sea. Additional work to explain the distribution of the most common seabirds in relation to environmental variables has been proposed but has not yet started.

*Specific Objective 3:* To identify and determine near-surface spatial distribution, diversity and abundance of phytoplankton, and zooplankton, based on Continuous Plankton Recorder samples collected during transit to and from the Ross Sea.

The Continuous Plankton Recorder (CPR) was deployed during the IPY voyage, both during the transit to and from Wellington, and within the Ross Sea itself. CPR silks collected during transit were preserved in formalin and sent to Australian Antarctic Division where they were analyzed for zooplankton species composition and abundance. CPR silks collected within the Ross Sea were preserved in ethanol for the analysis of epipelagic meroplankton. In addition to the zooplankton, sampling, water samples were collected for phytoplankton analysis using the underway water sampling system from a depth of 7 m, corresponding to the approximate depth of CPR sampling. In addition to the work described above, ICOMM (International census of marine microbes) samples collected during the IPY-CAML survey (10 m depth x 4 stations) have been analysed by collaborators in the USA (Ghiglione *et al.* 2012).

*Specific Objective 4:* To analyse underway and station data collected on salinity, temperature and chlorophyll *a* data, spot optical measurements with the SeaWiFS Profiling Multichannel Radiometer (SPMR), surface samples for chlorophyll *a*, nutrients and particle analysis as well as underway nutrient observations to allow ground-truthing of data collection from satellites and identify water masses (e.g. surface seawater temperature, and chlorophyll concentration).

This objective addressed background physical and surface biological conditions at the time of the IPY-CAML survey. The objective was split into two parts 1. characterisation of the biological environment and bio-optical regime using continuous underway sampling, and 2. identification of thermohaline fronts using discrete and underway sampling of temperature, salinity and nutrient profiles. The combined dataset was used to validate satellite data of temperature and surface chlorophyll distributions, providing a synoptic overview of physical and biological conditions during the survey.

*Specific Objective 5:* To identify and determine the spatial distribution, abundance (biomass), diversity, and size structure of epipelagic, mesopelagic (and possibly bathypelagic) species using acoustics data, target strength estimation techniques and net sampling.

This objective addressed samples collected using the mesopelagic trawl and acoustic data collected from midwater marks using the ship's echosounders. Results were presented at five conferences: 1) CAML-IPY Symposium in Genoa, Italy, May 2009; 2) CCAMLR SG-ASAM meeting in Genoa, Italy, May 2009; 3) Antarctic New Zealand conference in Auckland, July 2009; New Zealand Marine Sciences' Society conference in Stewart Island, July 2011; and International Polar Year Symposium, Montreal, Canada, April 2012. Results were also presented to the Ross Sea Bioregionalisation workshop in Wellington in June 2009 (see below) and were incorporated in the bioregionalisation reports prepared for CCAMLR (SC-CAMLR-XXIV-BG-25) and the Antarctic Treaty Consultative Meeting (ATCM). Reports include those by Koubbi *et al.* (2011), O'Driscoll (2009), O'Driscoll *et al.* (2009, 2011), Pinkerton *et al.* (in press), and Hanchet *et al.* (in press).

*Specific Objective 6:* To identify and measure diversity, distribution and densities of mesozooplankton, macrozooplankton and meroplankton.

This objective addressed the samples taken by Multiple Opening/Closing Net and Environmental Sampling System (MOCNESS) from the sea surface to the sea floor. The samples were quantitatively divided at sea to allow several complementary analyses to be performed. In terms of the mesozooplankton community in the Ross Sea, copepods were the dominant zooplankton collected in most samples, and this was primarily calanoids and cyclopoids (i.e., *Oithona* spp.). However, in certain cases pteropods (*Limacina helicina antarctica*) and salps (*Salpa thompsoni*) made important contributions to mesozooplankton abundance. Total water column mesozooplankton biomass ranged between 0.6-9.1 g C m<sup>-2</sup> and was usually highest close to the surface. Mesozooplankton biomass in the Ross Sea was generally higher than expected, and can rival that of productive subantarctic regions (e.g., South Georgia). Salps were the main macrozooplankton species recorded in the MOCNESS samples and a paper describing the population ecology and distribution of *Salpa thompsoni* on the continental slope and around the seamounts to the north of the Ross Sea has been published by Pakhamov *et al.* (2011).

Samples were also preserved in ethanol for the analysis of meroplankton species composition and DNA sequencing. Larvae from at least eight phyla were found, with a remarkable dominance of annelids in both abundance and diversity. Overall, larval abundances observed were lower than other Antarctic studies, likely attributable to the late summer sampling, months after Ross Sea's phytoplankton bloom and the main trigger of spawning in many benthic invertebrates. Analysis of variation in meroplankton community composition showed significant differences among geographic regions (Shelf, Slope and waters of the Antarctic Circumpolar Current - ACC), among water masses (Shelf Water, Antarctic Surface Water, and Circumpolar Deep Water), and among depth strata (upper, midwater and bottom). Overall, near surface waters showed greater larval abundances, and these values decreased from the continental shelf to the slope, declining further in the deeper waters of the ACC. Differences between these locations were due not only to the presence or absence of certain taxa, but also a result of changes in OTU abundance.

*Specific Objective 7:* To determine diversity, distribution and densities of viral, bacterial, phytoplankton and microzooplankton species in the water column.

The full data sets have been completed and loaded into an MPI database and to the South western Pacific OBIS node (Gordon 2000). Phytoplankton and nanoplankton cell counts have revealed that there is a significant difference between shelf and abyssal site water column assemblages, both in terms of cell numbers, diversity and density. These data now have to be integrated with the water column data to help understand what may be driving the changes in these compositions.

*Specific Objective 8:* To determine the spatial distribution, abundance (biomass), diversity, and size structure of shelf and slope demersal fish species and associated invertebrate species using a demersal survey.

This objective had three key tasks; (i) to identify specimens, update the Ross Sea species list and determine biodiversity, (ii) to identify fish assemblages and relate them to environmental data, and (iii) to compare estimates of fish density and abundance between trawls, visual (video & still images) and acoustic sampling techniques. A fourth key task, to determine density and abundance of demersal fish using a bottom trawl survey, was funded under MPI project ANT2007-02. Results have been published as three scientific journal papers with an additional paper in review, and have been submitted to several CCAMLR working group meetings.

A paper on the distribution and diversity of demersal and pelagic fish species in the Ross Sea region including results from both the BioRoss and IPY surveys and collections from the toothfish fishery will soon be published (Hanchet *et al.* in press). A diverse collection of over 2,500 fish

specimens was obtained from the BioRoss and IPY-CAML surveys representing 110 species in 21 families. When combined with previous documented material this gave a total species list of 175, of which 137 were from the Ross Sea shelf and slope (to the 2,000 m isobath). Demersal species richness, diversity and evenness indices all decreased going from the shelf to the slope and the seamounts. In contrast, indices for pelagic species were similar for the slope and seamounts/abyss but were much lower for the shelf.

A paper on the variation of demersal fish assemblages in the western Ross Sea including results from both the BioRoss and IPY surveys has been published (Clark *et al.* 2010). The distribution and abundance of 96 species able to be identified to species level collected in these surveys were examined to determine if demersal fish communities varied throughout the area, and what environmental factors might influence this. Three broad assemblages were identified, in the southern Ross Sea (south of 74°S), central–northern Ross Sea (between latitudes 71°–74°S), and the seamounts further north (65°–68°S) where some species more typical of sub-Antarctic latitudes were observed. Multivariate analyses indicated that environmental factors of seafloor rugosity (roughness), temperature, depth, and current speed were the main variables determining patterns in demersal fish communities.

Acoustic data collected during the demersal survey suggest that there may be potential to use fisheries acoustic methods to obtain estimates of grenadier abundance (O’Driscoll *et al.* in press). The acoustic target strength distribution of single targets close to the bottom was very similar to that predicted based on the measured size range of grenadiers. There are also positive correlations between acoustic backscatter and trawl catches of grenadiers.

Photographic data collected using NIWA’s Deep Towed Imaging System (DTIS) suggest that there may be potential to use photographic methods to obtain estimates of community structure and grenadier abundance (Bowden *et al.* in prep.).

Twenty-three sites spanning the continental shelf, northern continental slope, abyssal plain, and two seamounts were sampled using the towed camera and either demersal trawl or beam trawl, allowing direct comparisons between sampling methods. Patterns of species turnover between sites were similar across all methods. Estimates of fish population densities from the towed camera and beam trawl data were also comparable but those from the demersal trawl were consistently lower than for the other methods. *Macrourus* spp. grenadiers were ca. eight times less abundant in the demersal trawl than the video data but more large individuals were sampled by the trawl than the video and biomass estimates were similar.

*Specific Objective 9:* To determine the diversity, abundance/density, spatial distribution, and physical habitat associations of benthic assemblages across a body size spectrum from megafauna to bacteria, for shelf, slope, seamounts, and abyssal sites in the Ross Sea.

Using cameras, corers, epibenthic sleds, and trawls, benthic bacteria, macro-infauna, macro-hyperbenthic fauna, and mega-epifauna were sampled at sites on the continental shelf and previously unsampled areas on the northern continental slope of the Ross Sea, the abyssal plain, and seamounts to the north. Photographic data from seamounts in the northern Ross Sea region revealed a diverse and abundant fauna. Particularly striking were benthic communities comprised of stalked crinoids and brachiopods on Admiralty Seamount and the flanks of Scott Island which are reminiscent of an archaic fauna that may have survived through the isolation of these seamounts and reduced predator species (Bowden *et al.* 2010).

Taxonomists in New Zealand and around the world identified more than 150,000 individual specimens representing more than 700 species, many undescribed, across sixteen phyla for the mega-epifauna groups alone (e.g. Lörz 2009, 2010, Eléaume *et al.* 2011). At least three genera and sixty-two species are new to science. All eukaryotic components of the benthic fauna showed similar broad-scale distributional trends across the study region. Total abundances and numbers of

taxa were orders of magnitude higher on the continental shelf than on the slope or abyss plain, and shelf, slope, and abyssal samples were distinct from each other in multivariate analyses. Diversity, however, was comparable between shelf and abyssal sites and lowest on the slope. Bacterial diversity was highest in abyssal and slope samples, but abundance, biomass, production, and activity of all enzymes except proteinase, which was highest in the abyss, were significantly higher in shelf samples. Benthic mega-epifaunal community composition was more strongly correlated with depth and seabed current speed than either water column productivity or seasonal ice cover, indicating that local hydrodynamics and their influence on advection of primary production are more important in determining distributions across the shelf than are local variations in production. Fauna on the seamounts were distinct from all other samples and were comprised of both Antarctic and Southern Ocean species, including remarkable populations of a new hyocrinid species on Admiralty seamount (Bowden *et al.* 2011, Eléaume *et al.* 2011).

Published research to date has provided new insights into the distributions of several taxonomic groups (Lörz *et al.* 2009), raised questions about the history of the northern seamount fauna over evolutionary time (Bowden *et al.* 2011), and contributed to a meta-analysis of the relationship between productivity and diversity in the deep sea (Leduc *et al.* 2012). In combination with molecular phylogenies and existing data from around Antarctica, results from this project represent a major contribution to knowledge of the Antarctic marine ecosystem.

*Specific Objective 10:* To describe trophic/ecosystem relationships in the Ross Sea ecosystem (pelagic and benthic, fish and invertebrates).

Progress has been made on obtaining data from which to elucidate trophic relationships between organisms in the Ross Sector of Antarctica collected on the IPY-CAML survey in February–March 2008. Two methods have been used. First, 1081 stomachs from 22 species of Antarctic fish were examined and the contents of the full or partially-full stomachs (comprising 776 fish) were identified to 68 prey codes. Index of Relative Importance (IRI) has been calculated from these data and diet overlap between fish species is presented. Second, stable isotope and elemental composition analysis of samples were carried out for carbon and nitrogen. In total, nearly 2000 samples were analysed. Samples include:

- Fish (N=662 muscle, N=377 liver samples, 22 species);
- Cephalopods (N=193);
- Pelagic invertebrates (N=407);
- Benthic sediments (N=36);
- Phytoplankton (N=92);
- Benthic invertebrates (N=200 completed, 95 pending analysis);

Results have already been used to assist in parameterising and validating the quantitative model of the food web of the Ross Sea (paper accepted by CCAMLR Science). Research on the shrinkage of Antarctic silverfish carried out as part of this objective has contributed to a paper presented to the Ministry of Fisheries Antarctic Fisheries Working Group and accepted for submission to the CCAMLR working group on fisheries assessment in September 2010 (Pinkerton *et al.* 2007, 2009a, 2009b).

*Specific Objective 11:* Assess molecular taxonomy and population genetics of selected Antarctic fauna and flora to estimate evolutionary divergence within and among ocean basins in circumpolar species. Provide DNA barcoding for all fish and multi-cellular invertebrate species by sequencing reference specimens in conjunction with Canadian Barcoding Centre, for specimen identification in gut content, plankton, and in taxonomic and population genetic projects.

DNA data sets generated for selected Ross Sea taxa were combined with parallel data sets generated by other Institutes in order to estimate divergence within and among regions in the Southern Ocean. High levels of divergence, indicative of cryptic speciation, were found in all

major groups tested to date. Fishes: DNA sequencing of the COI gene revealed four well supported clades among the three recognized species of *Macrourus* in the Southern Ocean, indicating the presence of an undescribed species (Smith *et al.* 2011). A conclusion subsequently supported by meristic and morphometric examination of specimens with the description of a new species by McMillan *et al.* (2012). DNA barcodes also showed high sequence divergence among specimens of the slender codling *Halargyreus johnsonii* from New Zealand and the Southern Ocean, indicative of a cryptic species in this cosmopolitan species (Smith *et al.* 2011). A study of snailfishes collected during the IPY survey and from the toothfish fishery showed high species diversity with more than 34 Ross Sea liparid species in three genera; 18 of them new to science divergence (Stein 2012).

Invertebrates: A combined NZ-BAS data set on the octopod genus *Pareledone* provided one of the largest barcoding studies on a Southern Ocean genus. Ross Sea specimens provisionally identified as *Pareledone aequipapillae* appeared in a discrete clade to specimens from the Antarctic Peninsula, with a barrier to gene flow to the west of the Antarctic Peninsula (Allcock *et al.* 2010). Large numbers of echinoderms have been tissue sampled and sequenced for COI and include the Asteroidea, Ophiuroidea, Echinoidea, Holothuroidea, and the crinoids (Dettai *et al.* in press). In the Ophiuroidea two dominant patterns emerged: a. widely distributed species showing shallow divergence by location and b. species with deeper divergence associated with location or depth, that represent cryptic species. A similar pattern emerged in the smaller set of Asteroid sequences, with deep divergences within some Ross Sea taxa. Preliminary results for the amphipod genus *Rhacotropis* showed 5 well supported clades, indicative of cryptic taxa; while for the genus *Epimeria* (27 specimens from the Ross Sea) there were two well supported clades for specimens identified as *Epimeria robusta*, and likewise for specimens identified as *E. schiaparelli*, indicative of cryptic taxa (Lörz 2009, 2010, Lörz *et al.* in press). These taxa show shallow morphological differences.

#### **IPY2007-02 NZ IPY-CAML Cephalopoda.**

This project will report on the diversity of Antarctic Cephalopoda (Octopus and Squid), including a complete inventory of taxa, and reports on ontogenetic and sexual variation in species, their systematics, diversity, distribution, life histories, and trophic importance. A MAppSc thesis has been completed as part of this project (Garcia 2010).

Other research relevant or specifically linked to the projects above, are listed in Table 11.8.

**Table 11.8: Other research linked to MPI Ross Sea Antarctica biodiversity programme.**

<b>MPI</b>	ANT2011-01 Stock modelling, fishery effects and ecosystems of the Ross Sea
<b>MBIE</b>	C01X1001 Protecting Ross Sea Ecosystems. Comparative distribution and ecology of <i>Macrourus caml</i> and <i>M. whitsoni</i> in the Ross Sea region; feeding relationships of fish species in the Ross Sea region; Spatial processes, including spatial marine protection; Ecosystem modelling of the Ross Sea region).(Pinkerton <i>et al.</i> 2012a,b; Murphy <i>et al.</i> 2012)
<b>DOC</b>	Leigh Torres NIWA/Alison
<b>OTHER</b>	Universities NIWA;Lincoln, Canterbury, Otago, Auckland, Waikato
<b>EMERGING ISSUES</b>	
Coastal research and functional ecology-ongoing need	
Taxonomic issues for fish and invertebrates (from IPY)ANT 2005-02	
Water samples from throughout water column to assess microbial content (from IPY) check with Els	

## 11.4. Progress and re-alignment

Given that the MPI Biodiversity programme has been running for 11+ years, and that a number of new strategic documents and directions are emerging across government, it is time to look both back and forward and review the programme to ensure its alignment with more recent strategic documents.

In 2000, five strategic outcomes were built into the MPI Biodiversity Research Programme:

*That by 2010:*

- i) *the MPI Biodiversity programme will have become an integral part of the research effort devoted to understanding New Zealand's marine environment.*
- ii) *research planning will benefit from close cooperative relationships within the Ministry of Fisheries, with other government agencies, and with external stakeholders.*
- iii) *mutually beneficial collaborative research projects will be carried out alongside other New Zealand and international research providers, especially for vessel-based research.*
- iv) *MPI Biodiversity projects will have contributed substantially to an improved understanding of New Zealand's marine biodiversity and its role in marine ecosystem function, yielding scientifically rigorous outputs for a national and international professional audience.*
- v) *results generated by MPI Biodiversity projects will be incorporated into management policy, with clear benefits for the New Zealand marine environment.*

The Biodiversity Programme has been highly effective in delivering on the first 4 and part of the 5<sup>th</sup> of the five outcomes. A missing element is some measure of “*clear benefits for the New Zealand marine environment*”. In recent years, significant all-of-government projects have been administered through the programme, and one-off funding applications made jointly with other stakeholders have been successful. The Programme has made a significant contribution to increasing understanding about biodiversity in the marine environment. Achievements in each outcome are addressed below.

*i) Has the Biodiversity Research Programme become integrated with New Zealand's research effort to understand the marine environment?*

Seven science objectives were developed by multiple stakeholders through the Biodiversity Research Advisory Group. The agreed objectives include ecosystem-scale studies in the New Zealand marine environment, the classification and characterisation of the biodiversity of nearshore and offshore marine habitats, the role of biodiversity in the functional ecology of marine communities, connectivity and genetic marine biodiversity, the assessment of the effects of climate change and increased ocean acidification, identification of indicators of biodiversity that can be used to monitor change, identification of key threats to biodiversity, identification of threats and impacts to biodiversity and ecosystem functioning beyond natural environmental variation.

Projects ranged from localised experiments on seabed communities of shellfish and echinoderms, to integrated studies of rocky reef systems and offshore fishery-scale trophic studies. The effects of ocean climate change (temperature, acidification) are being explored on shellfish, rhodolith communities, plankton productivity and the microbial productivity engines of polar waters. A major project to investigate shelf communities in relation to climate over the past 1000 years has resulted in the development of new methods and insights to past changes and human impact on New Zealand's marine environment.

A total of 55 projects were commissioned and managed within this 10 year period, yielding over 100 final research reports, most of which have been published through MPI Publications (Marine Biosecurity and Biodiversity Reports and Aquatic Environment and Biodiversity Reports), books, Identification Guides and mainstream scientific literature. A number of other publications are still in preparation. In addition, several workshops have been run through the Programme, including qualitative modelling techniques, how to set up a marine monitoring programme and predictive

modelling. A large number of science providers, including NIWA, Cawthron Institute, University of Auckland, Auckland University of Technology, University of Waikato, Victoria University of Wellington, University of Otago, University of Canterbury and Massey University have been directly commissioned or sub-contracted to take part in or conduct research projects through the Programme during the 10-year period. For some, the projects have provided critical synergies with MBIE funded OBIs or projects, while others have provided one-off opportunities for marine biodiversity investigation or opportunistic leveraging for research voyages.

Research into the biodiversity of habitats such as seamounts has been completed and new methods to assess the vulnerability of seabed habitats have been developed. The land-sea interface is being investigated and projects have shown how land use in a given catchment can affect nutrient transfer and the living conditions and impact diversity and functioning of estuarine and coastal organisms. Publication and presentation of the results from these projects has resulted in widespread contribution to the development of Marine Science in New Zealand. Partnership with overseas researchers and presentations to international meetings and conferences has added to the growing global initiatives on marine biodiversity research questions.

Feedback from stakeholders has indicated that the move to a 5 year research planning horizon was welcomed by research providers, but some stakeholders felt that Requests for Proposals should be at a higher level than individual projects to safeguard intellectual property on new ideas and methods.

ii) *Does research planning now benefit from close cooperative relationships within the Ministry of Fisheries, with other government agencies, and with external stakeholders?*

The Biodiversity Programme is very co-operative. Of 38 projects underway in the last 5 years, 14 have formal collaborative components across government departments, with other stakeholders or multiple research providers and 10 have formal linkages to international research programmes. Within MPI and with other stakeholders (NGOs, industry, other government departments), the Biodiversity Projects have contributed to discussions about Marine Stewardship Council (MSC) certification, to decision papers on aspects of Antarctic management under CAMLR, fulfilling MPI commitments to the NZ Biodiversity Strategy, and to MPI progress towards recognising the role of the ecosystem in underpinning sustainable and healthy fisheries production. There are many other examples, e.g. the programme has towards DOC and MPI decisions on marine protected areas. The interaction at the research and policy advice stages of resource management feeds back into the BRAG planning for future research.

There are close links with the MPI Aquatic Environment research programme, the National Aquatic Biodiversity Information System (NABIS), an MPI web-based interactive data access and mapping tool) and the MPI Antarctic Research programme. These and other links have enabled contributions resulting from progress on land-sea interface research, habitats of significance to fisheries management, trophic studies (MSC Certification), climate change (effects on shellfish) and habitat classification (fish optimised MEC, testing of MEC and BOMECE). The successful involvement of the Biodiversity Programme in major all-of-government projects such as Ocean Survey 20/20 and IPY-CAML, has also raised the profile of MPI and the research it has commissioned both across New Zealand and internationally.

Datasets, voucher specimens and samples from all biodiversity research projects have resulted in a substantial amount of material that has been physically preserved and housed in the Te Papa Fish Collection and NIWA National Invertebrate Collection. All data are held in databases either at MPI, NIWA or Te Papa, and accessibility is being improved. The recent Bay of Islands Ocean Survey 20/20 Portal was very well received and nominated for NZ Govt Open Source awards. It will also incorporate data access from Chatham Challenger and IPY projects. Data from a number of MPI biodiversity projects have also been entered into international biodiversity databases such as OBIS and from there into the Global Biodiversity Information Facility (GBIF).

Biodiversity Research planning receives regular input from DOC, SeaFIC, MfE, Cawthron Institute, NIWA, GNS, LINZ, MAFBNZ, Te Papa, University of Auckland, AUT, University of Otago, MoRST, MFAT, Regional Councils and others. Research planning for 2011-12 and beyond will include a re-alignment of the current research programme to take account of new developments such as Fisheries 2030, MfE's National Monitoring programme, DOC's integrated coastal monitoring programme, Statistic New Zealand's Environmental Domain Plan<sup>69</sup>, and international commitments such as the recent CBD COP10 Aichi-Nagoya Agreement.

Feedback and support for projects by external stakeholders has shown that the Programme has been effective in promoting inter-agency collaboration. The Programme has also had close links with Research Data Management and the Observer Programme for certain projects (e.g trophic studies on the Chatham Rise, ZBD2004-02). With the former restructure of MPI and now the merger with MAF, and the move to Fisheries 2030 and Fisheries Plans, it important that the Programme develops strong relationships with the Fisheries Management and Strategy (International) groups within MPI and at MAF.

iii) *Have mutually beneficial collaborative research projects been carried out alongside other New Zealand and international research providers, especially for vessel-based research?*

As discussed above, collaborative research projects across government and among research providers have resulted in many mutually beneficial data and specimen collection, surveys of New Zealand marine biodiversity in NZ territorial seas, the EEZ and the Ross Sea, groundbreaking research into seamount biodiversity and the identification of VMEs, and research for international collaboration, particularly vessel based studies. Large scale vessel dependent oceanic research projects have made significant gains in baseline knowledge about the distribution and abundance of biodiversity in the EEZ/Ross Sea region. Vessel-based projects include: NORFANZ (Norfolk Island-Australia-New Zealand survey of biodiversity on Norfolk Ridge and Lord Howe Rise); BioRoss (MPI-LINZ, first NZ survey of biodiversity in the Ross Sea); Chatham-Challenger (LINZ-MPI-NIWA-DOC first Ocean Survey 20/20 project), NZ IPY-CAML (MPI-LINZ-NIWA (with international and NZ wide collaboration) survey of the Ross Sea as part of International Polar Year; Biodiversity of seamounts (MPI-NIWA-LINZ-MBIE voyages to the Kermadec Arc and on the Chatham Rise). These projects have generated huge geo-referenced datasets and thousands of specimens for Te Papa and National Invertebrate Collections. They have also resulted in the identification of new species, new genera and new families, as well as new records extending the known distribution of species. These surveys have contributed to habitat classification, identified areas of high biodiversity and challenged paradigms on the environmental drivers that determine biodiversity. More recently they have provided new information on the effects of ocean acidification on the productivity of polar seas, and in New Zealand waters.

Vessel dependent coastal projects have also generated significant new understanding about the distribution of inshore biota, and the role they play in maintaining a healthy ecosystem. Experimental field work on the productivity of the seabed has been carried out in NZ waters (Fiordland, Otago, Bay of Islands, Hauraki Gulf, Kaipara and Manukau Harbours), and along the west coast of the Ross Sea. The impact of land practices on the land-sea interface has also highlighted real downstream effects on the productivity of the coastal environment. These projects have provided new insights into the connectivity between different species groups, and data are being used in a number of ways to assist with spatial planning by RMAs.

Feedback from stakeholders has indicated that the collaborative voyages administered through the Programme have successfully created synergy and opportunity for New Zealand scientists as well as facilitating new international collaborations.

---

<sup>69</sup>[http://www.stats.govt.nz/browse\\_for\\_stats/environment/natural\\_resources/environment-domain-plan-stocktake-paper.aspx](http://www.stats.govt.nz/browse_for_stats/environment/natural_resources/environment-domain-plan-stocktake-paper.aspx)

iv) *Have MPI [MFish] Biodiversity projects contributed substantially to an improved understanding of New Zealand's marine biodiversity and its role in marine ecosystem function, yielding scientifically rigorous outputs for a national and international professional audience?*

In the early years, the Programme focussed primarily on taxonomy and the description of marine biodiversity. As the Programme matured, projects to address biodiversity roles in ecosystem function were introduced. Some were experimental and on a local scale while others were on a regional scale. Recent projects have addressed patterns of marine biodiversity in relation to environmental drivers with ecosystem function. This enabled modelling to predict the distribution of biodiversity in unsurveyed areas of ocean, and evaluation of the vulnerability of biodiversity to perturbations such as climate change, as well as the modelling of trophic interactions among key fish species. Presentations of research results have been made to numerous overseas and New Zealand science audiences, and publications in the mainstream literature have been encouraged. IPY, Chat chall, Alison s etc CBD-FAO Int seabed authority

v) *Have results generated by MPI [MFish] Biodiversity projects been incorporated into management policy, with clear benefits for the New Zealand marine environment?*

Examples of incorporation into management policy with clear benefits for the marine environment include the increased awareness of research topics initiated in the biodiversity programme by policy analysts to core Aquatic Environment research projects and Fishery Plans, (land-use effects, climate change in the ocean, habitat classification); links to the Antarctic research programme and uptake into CCAMLR (ecotrophic studies, ecosystem baselines, VME risk assessment, bioregionalisation), spatial management (seamount closures, BPAs, MPAs, RMAs), the need by MfE to report on the marine environment at a national scale (plankton recording programme, Marine Environmental Monitoring Programme). MPI biodiversity advice is frequently requested to contribute to cross-government initiatives including Ocean Survey 20/20, DOC Sub-Antarctic Islands Forum National Monitoring, Stats New Zealand Tier 1 statistic review and Environmental Domain Stocktake, International Year of Biodiversity, OECD and CBD reports, International Oceans Issues, SPRFMO, NRS marine issues paper, the Antarctic Science Framework, Ocean Fertilisation and IPCC Finally, the programme has contributed to New Zealand's efforts in the international Census of Marine Life and an ongoing assessment of New Zealand's progress in Marine Biodiversity has been proposed as a new Tier 1 Environmental Statistic. However, the benefits to the marine environment are more inferred than demonstrated. There is substantially increased awareness within MPI and across government, that the health of fisheries and other valued uses of the sea depend on intact ecosystem services provided by the diversity of organisms, the diversity of habitats and the genetic diversity found in the marine environment. Statements of intent and long-term strategic documents such as Fisheries 2030 and Fish Plans have biodiversity protection and an ecosystem approach to fisheries management objectives explicitly stated. Future research questions will also need to address follow-up of management decisions to assess whether and to what extent the objectives have been achieved.

In 2000, the concept of research on marine biodiversity was hotly debated among stakeholders and the benefit of the research (other than to scientists) was not widely accepted. In 2010, it is clear that much of the research in this biodiversity programme has been about defining and mapping the biological diversity of the sea, its roles in marine ecosystem function, threats to these roles and how best biodiversity and its successful protection can be measured. Huge advances have been made in providing new identification tools for major groups (e.g. Coralline algae ...). Much progress has been made, and the programme has successfully raised the profile of biodiversity in coastal and ocean environmental management, in particular fisheries management, and biodiversity research uptake into policy and management decisions within MPI and across government.

#### **11.4.1. Concluding remarks**

New Zealand is moving into an era of unprecedented and increasing interest in the utilisation of marine resources. Mineral, petroleum and gas resources are estimated to be worth billions of dollars to the economy (Glasby and Wright 1990), and new environmental legislation has been drafted (the Exclusive Economic Zone and Continental Shelf (Environmental Effects) Act 2012). Changes inshore are also taking effect with the Environmental Protection Authority Act passed by Parliament on 11 May 2011. This Act establishes a new Environmental Protection Authority (EPA) as a standalone crown agent from 1 July 2011. The newly released Coastal Policy statement and proposed Policy Statement on Indigenous Biodiversity demonstrates an awareness by Government that much of New Zealand's primary production based economy is dependent on clean "green" policies supporting effective environmental management both on land, freshwater and in the sea.

New Zealand is also a signatory to the CBD Aichi-Nagoya Agreement with a new International Decade for Biodiversity that runs 2011-2020 and New Zealand's contribution to the identification of EBSAs in the SW Pacific, and to GOBI. Progress in our knowledge of the marine biodiversity and ecosystem services provided by the marine environment has clearly been made over the last decade. However, we need a more co-ordinated approach across government to link science to policy needs. For example, there is a compelling need for large-scale projects such as mapping seafloor habitats and establishing long-term nation-wide monitoring and reporting schemes to measure the effects of ocean climate change, regular assessment of the cumulative effects of anthropogenic activities and multiple stressors in the ocean and the effectiveness of their management. Without these, we face the risks that New Zealand's "green" branding will be increasingly challenged, and that tipping points in the health of the aquatic environment may be reached too soon for evasive action to be taken.

## 11.5. References

- Allcock A, Norman M, Smith P, Steinke D, Stevens D, Strugnell J (2010) Cryptic speciation and the circumpolarity debate: A case study on endemic Southern Ocean octopuses using the COI barcode of life. *Deep-Sea Research II* (2010), doi:10.1016/j.dsr2.2010.05.016
- Anderson OF, Bagley NW, Hurst RJ, Francis MP, Clark MR, *et al.* (1998). Atlas of New Zealand fish and squid distributions from research bottom trawls. *NIWA Tech Rep*.42:1–303.
- Andrew, N. Francis, M. (eds.) (2003). *The Living Reef: The Ecology of New Zealand's Rocky Reefs*. Craig Potton Publishing, Nelson, New Zealand.
- Anon (2000). The New Zealand Biodiversity Strategy. Our chance to turn the tide. Department of Conservation and Ministry for the Environment. 146 p.
- Arnold A, (editor). (2004) Wellington: WWF-New Zealand; Shining a spotlight on the biodiversity of New Zealand's Marine Ecoregion.
- Asnaghi, V., Bertolotto, R., Giussani, V., Mangialajo, L., Hewitt, J., Thrush, S., Moretto, P., Castellano, M., Rossi, A., Povero, P., Cattaneo-Vietti, R. & Chiantore, M. (2012). Interannual variability in *Ostreopsis ovata* bloom dynamic along Genoa coast (North-western Mediterranean): a preliminary modeling approach. *Cryptogamie Algologie*,33:181-189.
- Ausubel, JH, Crist DT, Waggoner PE. (Eds) (2010). First census of marine life 2010 highlights of a decade of discovery Contributors: 2,700 marine scientists from around the globe. A publication of the Census of Marine Life Census of Marine Life International Secretariat Consortium for Ocean Leadership Washington, DC USA www.coml.org
- Baco AR, Rowden AA, Levin LA, Smith CR, Bowden D. (2009). Initial characterization of cold seep faunal communities on the New Zealand margin. *Mar Geol.* doi: 10.1016/j.margeo.2009.06.015.
- Bagley NW, Anderson OF, Hurst RJ, Francis MP, Taylor PJ. (2000). Atlas of New Zealand fish and squid distributions from midwater trawls, tuna longline sets, and aerial sightings. *NIWA Tech Rep.*;72:1–171.
- Baird, S.J., Tracey, D., Mormede, S., Clark, M. (2012). The distribution of protected corals in New Zealand waters. NIWA Client Report No: WLG2012-43 prepared for Department of Conservation, Wellington. 93 p.
- Baker CS, Chilvers BL, Constantine R, DuFresne S, Mattlin RH, van Helden A, Hitchmough R (2010). Conservation status of New Zealand marine mammals (suborders Cetacea and Pinnipedia), 2009. *New Zealand Journal of Marine and Freshwater Research* 44: 101–115.
- Batson, P. (2003). Deep New Zealand: Blue Water, Black Abyss. Christchurch:Canterbury University Press.
- Battley, P.F., Melville, D.S., Schuckard, R., Ballance, P.F. (2005). Quantitative survey of the intertidal benthos of Farewell Spit, Golden Bay. *Marine Biodiversity Biosecurity Report No. 7*. 19 p.
- Beaumont J, MacDiarmid A, D'Archino R. (2010 ). Mapping the environmental values of New Zealand's marine ecosystem: A meta-analysis. *Biosecurity NZ Tech Pap.*;2010:1–76.
- Beaumont J, Oliver M, MacDiarmid A. (2008). Mapping the values of New Zealand's coastal waters. 1. Environmental values. *Biosecurity NZ Tech Pap.*;2008/16:1–89.
- Beaumont, J., MacDiarmid, A.B., Morrison, M. (in press). Rocky reef ecosystems – how do they function? Final Research Report for project ZBD200509 to MPI 296 p.
- Bostock, H., Tracey, D., Cummings, V., Mikaloff-Fletcher, S., Williams, M., Guy, C., Neil, H., Currie, K. (2012). Ocean acidification in fisheries habitat (ZBD201041). Research Progress Report prepared for the Ministry of Primary Industries Fisheries. 39 p.
- Bowden, D.A. (2011). Benthic invertebrate samples and data from the Ocean Survey 20/20 voyages to the Chatham Rise and Challenger Plateau, 2007. *New Zealand Aquatic Environment and Biodiversity Report No. 65*. 40 p.
- Bowden D.A., Schiaparelli S., Clark M.R., Rickard G.J. (2011) A lost World? Archaic crinoid-dominated assemblages on an Antarctic seamount. *Deep Sea Research Part II. Topical Studies in Oceanography* 58:119-127
- Bowden DA, Hewitt J, Verdier A-L, Pallentin A (in press) (Project ZBD200701 Objective 14, The potential of multibeam echosounder data for predicting benthic invertebrate assemblages across Chatham Rise and Challenger Plateau. Draft Aquatic Environment and Biodiversity Report, Ministry of Fisheries, New Zealand, Wellington
- Bowden DA, Hewitt J. (2012). Recommendations for surveys of marine benthic biodiversity: outcomes from the Chatham-Challenger Ocean Survey 20/20 Post-Voyage Analyses Project New Zealand Aquatic Environment and Biodiversity Report No. xx91. 34 p.
- Bowden, D.A.; Compton, T.J.; Snelder, T.H.; Hewitt, J.E. (2011b). Evaluation of the New Zealand Marine Environment Classifications using Ocean Survey 20/20 data from Chatham Rise and Challenger Plateau. *New Zealand Aquatic Environment and Biodiversity Report No. 77*. 27 p.
- Bowden, DA, Hanchet, SM, Marriott, PM (in prep.). Population estimates of Ross Sea demersal fish: a comparison between video and trawl methods. (Submitted as short note to *Fisheries Research*)
- Boyd, P.W., Law, C.S. and S. C. Doney (in press). A climate change atlas for the Ocean. *Oceanography*.
- Bradford-Grieve J. Absence of government leadership is damaging the health of systematics and taxonomy in the UK. *NZ Sci Rev.* 2008;65:84–88.
- Bradford-Grieve, J., Fenwick, G. (2001). A review of the current knowledge describing the biodiversity of the Ross Sea region.FRR Ministry of Fisheries Final Research Report for Project
- Bradford-Grieve, J., Fenwick, G. (2002). A review of the current knowledge describing the biodiversity of the Balleny Islands. *FRR ZBD2000/01*
- Bradford-Grieve J, Probert K, Lewis K, Sutton P, Zeldis J. (2006). New Zealand shelf region In: Robinson AR, Brink KH, editors. *The global coastal ocean: regional studies and syntheses. The Sea 14*. New York: John Wiley & Sons, Inc. pp. 1451–1492.
- Broom, J.E.S.; Hart, D.R.; Farr, T.J.; Nelson, W.A.; Neill, K.F.; Harvey, A.H.; Woelkerling, W.J. (2008). Utility of psbA and nSSU for phylogenetic reconstruction in the Corallinales based on New Zealand taxa. *Molecular Phylogenetics & Evolution* 46: 958–973.
- Brown, CJ, Fulton, EA, Hobday, AJ, Matear, RJ, Possingham, HP, Bulman, C, Christensen, V, Forrest, RE, Gehrke, PC, Gribble, NA, Griffiths, SP, Lozano-Montes, H, Martin, JM, Metcalf, S, Okey, TA, Watson, R and Richardson, AJ (2010) Effects of climate-driven primary production change on marine food webs: implications for fisheries and conservation. *Global Change Biology*, 16 4: 1194-1212
- Carroll, E.; J.A Jackson, JA; Paton, D.; Smith, T (in press). Estimating 19<sup>th</sup> and 20<sup>th</sup> century right whale catches and removals around east Australia and New Zealand. Final Research Report ZBD200505, MS12 Part C.
- Chang FH, Mullan B. 2003. Occurrence of major harmful algal blooms in New Zealand: is there a link with climate variations? *Climate Update.*;53(11):4.

- Chang FH, Zeldis J, Gall M, Hall J. (2003). Seasonal and spatial variation of phytoplankton assemblages, biomass and cell size from spring to summer across the north-eastern New Zealand continental shelf. *J Plankt Res.*;25:737–758.
- Chang, F.H.; Williams, M.J.M.; Schwarz, J.; Hall, J.; Stewart, R.; Maas, E.W. (2012). Spatial variation of phytoplankton assemblages and biomass in the New Zealand sector of the Southern Ocean during the late austral summer 2008 Polar Biology.
- Cheung L, Lam VWY, Sarmiento JL, Kearney K, Watson R, Pauly D, (2009). Projecting global marine biodiversity impacts under climate change scenarios. *Fish and Fisheries* 10(3):235-251
- Clark M, O'Shea S. 2001 Hydrothermal vent and seamount fauna from the southern Kermadec Ridge, New Zealand. *InterRidge News.*;10(2):14–17.
- Clark MR, O'Driscoll R. (2003). Deepwater fisheries and aspects of their impact on seamount habitat in New Zealand. *J Northw Atlantic Fish Sci.*;31:441–458.
- Clark MR, Rowden AA. (2009). Effect of deepwater trawling on the macro-invertebrate assemblages of seamounts on the Chatham Rise, New Zealand. *Deep-Sea Res I*;56:1540–54.
- Clark, M.R., Dunn, M.R., McMillan, P.J., Pinkerton, M.H., Stewart, A., Hanchet, S.M. (2010b). Latitudinal variation of demersal fish assemblages in the western Ross Sea. CCAMLR document WG-FSA-10/P3. 13 p. (Antarctic Science, 2010. <http://dx.doi.org/10.1017/S0954102010000441>)
- Clark, M.R., Bowden, D.A., Baird, S.J., Stewart, R. (2010a). Effects of fishing on the benthic biodiversity of seamounts of the “Graveyard” complex, northern Chatham Rise. New Zealand Aquatic Environment and Biodiversity Report No. 46. 40 p.
- Clark, M.R., Roberts, C.D. (2008). Fish and invertebrate biodiversity on the Norfolk Ridge and Lord Howe Rise, Tasman Sea (NORFANZ voyage, 2003). New Zealand Aquatic Environment and Biodiversity Report No. 28. 131 p.
- Clark, M.R., Stokes, K., Baird, S.J. (in press). Development of a methodology for ecological risk assessments for New Zealand deepwater fisheries. *New Zealand Aquatic Environment and Biodiversity Report*.
- Clark, M.R.; Tittensor, D.P. (2010). An index to assess the risk to stony corals from bottom trawling on seamounts. *Marine Ecology* 31(suppl 1): 200–211.
- Clark, M.R.; Tracey, D.M.; Pallentin, A.; Schnabel, K.; Anderson, O.F.; Bowden, D. (2009). Voyage report of a survey of “seamounts” on the northwest and southeast Chatham Rise (TAN0905). 49 p. (unpublished report available from NIWA, Wellington).
- Clark, M.R.; Watling, L.; Rowden, A.A.; Guinotte, J.M.; Smith, C.R (2010). A global seamount classification to aid the scientific design of marine protected area networks. *Journal of Ocean and Coastal Management* 54: 19–36.
- Clark, M.R.; Williams, A.; Rowden, A.A.; Hobday, A.J.; Consalvey, M. (2011). Development of seamount risk assessment: application of the ERAEF approach to Chatham Rise seamount features. New Zealand Aquatic Environment and Biodiversity Report No. 74. 18p.
- Coleman, C.O.; Lörz, A.-N. (2010). A new species of Camacho (Crustacea, Amphipoda, Aoridae) from the Chatham Rise, New Zealand. *Zoosystematics and Evolution* 86(1): 33–40.
- Compton TJ, Julian K, Leathwick JR, Bowden DA (Project ZBD200701 Objective 12, in press) Modelling distributions of benthic invertebrate taxa across Chatham Rise and Challenger Plateau in relation to environmental variables. Ministry of Fisheries Wellington, New Zealand
- Compton, T. J., D. A. Bowden, C. Roland Pitcher, J. E. Hewitt, and N. Ellis. (2012). Biophysical patterns in benthic assemblage composition across contrasting continental margins off New Zealand. *Journal of Biogeography* DOI 10.1111/j.1365-2699.2012.02761.x
- Connell, AM, Dunn, MR and Forman, J. (2010). Diet and dietary variation of New Zealand hoki *Macruronus novaezelandiae*. *New Zealand Journal of Marine and Freshwater Research* 44(4):289 - 308
- Costello MJ, Coll M, Danovaro R, Halpin P, Ojaveer H, *et al.* (2010) A Census of Marine Biodiversity Knowledge, Resources, and Future Challenges. *PLoS ONE* 5(8): e12110. doi:10.1371/journal.pone.0012110
- Coutts ADM, Dodgshun TJ. (1998). The nature and extent of organisms in vessel sea-chests: a protected mechanism for marine bioinvasions. *Mar Poll Bull.* 2007;54:875–886.
- Cranfield HJ, Gordon DP, Willan RC, Marshall BA, Battershill CN. Adventive marine species in New Zealand. NIWA Technical Report;34:1–48.
- Cranfield HJ, Manighetti B, Michael KP, Hill A (2003). Effects of oyster dredging on the distribution of bryozoan biogenic reefs and associated sediments in Foveaux Strait, southern New Zealand. *Continental Shelf Res.*;23:1337–1357. Cryer M, Hartill B, O'Shea S. 2002 Modification of marine benthos by trawling: toward a generalization for the deep ocean? *Ecol Applications.*;12:1824–1839.
- Cranfield, H. J.; Gordon, D. P.; Willan, R. C.; Marshall, B. A.; Battershill, C. N.; Francis, M. P.; Nelson, W. A.; Glasby, C. J.; Read, G. B. (1998). Adventive marine species in New Zealand. NIWA Technical Report 34: 1-48.
- Cryer M, Hartill B, O'Shea S. 2002 Modification of marine benthos by trawling: toward a generalization for the deep ocean? *Ecol Applications.*;12:1824–1839.
- Cummings, V., Thrush, S., Andrew, N., Norkko, A., Funnell, G., Budd, R., Hewitt, J., Gibbs, M., Mercer, S., Marriott, P., Anderson, O. (2003). Ecology and biodiversity of coastal benthic communities in McMurdo Sound, Ross Sea: emerging results Final Research Report ZBD2002-01
- Cummings, V., Hewitt, J., Van Rooyen, A., Currie, K., Beard, S., Thrush, S., Norkko, J., Barr, N., Heath, P., Halliday, J., Sedcole, R., Gomez, A., McGraw, C., Metcalf, V. (2011). Ocean acidification at high latitudes: potential effects on functioning of the Antarctic bivalve *Laternula elliptica*. *PLoS ONE* 6(1):e16069.
- Cummings, V., Thrush, S., Andrew, N., Norkko, A., Funnell, G., Budd, R., Gibbs, M., Hewitt, J., S. Mercer, S., Marriott, P., Anderson, O. (2003). Ecology and biodiversity of coastal benthic communities in McMurdo Sound, Ross Sea: emerging results. Final Research Report for Ministry of Fisheries Research Project ZBD2002/01 Objectives 1 & 2. National Institute of Water and Atmospheric Research. 105 p.
- Cummings, V., Thrush, S., Schwarz, A.-M., Funnell, G., Budd, R. (2006). Ecology of coastal benthic communities of the northwestern Ross Sea. *New Zealand Aquatic Environment and Biodiversity Report* No 6. 67 p.
- Cummings, V.J., Thrush, S., Chiantore, M., Hewitt, J. & Cattaneo-Vietti, R. (2010). Macrobenthic communities of the north-western Ross Sea shelf: links to depth, sediment characteristics and latitude. *Antarctic Science*, 22:793-804.
- Cummings, V.J., Thrush, S.F., Marriott, P.M., Funnell, G.A., Norkko, A., Budd, R.G. (2008). Antarctic coastal marine ecosystems (ICECUBE). Final Research Report for Ministry of Fisheries Research Project ZBD2006-03 73 p.
- Cummings, V.J., Thrush, S.F., Norkko, A., Andrew, N.L., Hewitt, J.E., Funnell, G.A., Schwarz, A.-M. (2006). Accounting for local scale variability in benthos: implications for future assessments of latitudinal trends in the coastal Ross Sea. *Antarctic Science* 18(4): 633-644.
- D'Archino, R.; Neill, K.F.; Nelson, W.A. (in press) Recognition and distribution of *Polysiphonia morrowii* (Rhodomelaceae, Rhodophyta) in New Zealand. *Botanica Marina*.

- D'Archino, R.; Sutherland, J.E. (in press) First record of the genus *Dudresnaya* (Dumontiaceae, Rhodophyta) in New Zealand waters. *Phycological Research*.
- De Domenico, F., Chiantore, M., Buongiovanni, S., Paola Ferranti, M., Ghione, S., Thrush, S., Cummings, V., Hewitt, J., Kroeger, K., Cattaneo-Vietti, R. (2006). Latitude versus local effects on echinoderm assemblages along the Victoria Land coast, Ross Sea, Antarctica. *Antarctic Science* 18(4): 655-662.
- Dettai *et al.* (+45 authors). (2011). DNA barcoding and molecular systematics of the benthic and demersal organisms of the CEAMARC survey. *Polar Science* 5: 298-312.
- Dodgshun T, Coutts A. (2003). Opening the lid on sea chests. *Seafood NZ*;11:35.
- Dunn MR (2009) Feeding habits of the ommastrephid squid *Nototodarus sloanii* on the Chatham Rise, New Zealand. *NZ J Mar Freshw Res* 43:1103-1113
- Dunn MR (2009) Feeding habits of the ommastrephid squid *Nototodarus sloanii* on the Chatham Rise, New Zealand. *NZ J Mar Freshw Res* 43:1103-1113
- Dunn MR, Connell AM, Forman J, Stevens DW, Horn PL (2010a) Diet of Two Large Sympatric Teleosts, the Ling (*Genypterus blacodes*) and Hake (*Merluccius australis*). *PLoS ONE* 5(10):
- Dunn MR, Griggs, L, Forman J, Horn PL (2010b). Feeding habits and niche separation among the deep-sea chimaeroid fishes *Harriotta raleighana*, *Hydrolagus bemisi* and *Hydrolagus novaezealandiae* Marine Ecology Progress Series 407: 209-225
- Dunn, MR, Horn PL, Connell, AM, Stevens, DW, Forman, J, Pinkerton, M, Griggs, L, Notman, P, Wood B. (in press). Ecosystem-scale trophic relationships: diet composition and guild structure of middle-depth fish on the Chatham Rise. *AEBA series*
- Dunn, MR, Szabo, A, McVeagh, MS, Smith, PJ. (2010c). The diet of deepwater sharks and the benefits of using DNA identification of prey. *Deep-Sea Research Part 1*. 57(7): 923-930
- Eakin, R.A.; Eastman, J.T.; Near, T.J. (2009). A new species and a molecular phylogenetic analysis of the Antarctic Fish Genus *Pogonophryne* (Nototheniidae: Artedidraconidae). *Copeia* 2009, No. 4, 705-713.
- Eleaume M, Hemery LG, Bowden DA, Roux M (in press) A large new species of the genus *Ptilocrinus* (Echinodermata, stalked Crinoidea, Hyocrinidae) from Antarctic seamounts. *Polar Biology* DOI: 10.1007/s00300-011-0993-2
- Farr, T.; Broom, J.; Hart, D.; Neill, K.; Nelson, W. (2009). Common coralline algae of northern New Zealand: an identification guide. *NIWA Information Series* No. 70.
- Fenwick, G., Bradford-Grieve, J. (2002a). Recommendations for future directed research to describe the biodiversity of the Ross Sea region. *FRR ZBD2000/01*
- Fenwick, G., Bradford-Grieve, J. (2002b). Human pressures on Ross Sea region marine communities: recommendations for future research. *FRR ZBD2000/01*
- Floerl O, & Hewitt J. (2012). Chatham-Challenger OS 20/20 Post Voyage Analyses: Objective 9- Patterns in species composition. New Zealand Aquatic Environment and Biodiversity Report No. 97. 40 p.
- Forman; JS, Dunn MR. (2010). The influence of ontogeny and environment on the diet of lookdown dory, *Cyttus traversi* New Zealand Journal of Marine and Freshwater Research Volume 44(4): 329 - 342
- Freeman DJ, Marshall BA, Ah Yong ST, Wing SR, Hitchmough RA (2010). The conservation status of New Zealand marine invertebrates, 2009. *New Zealand Journal of Marine and Freshwater Research* 44: 129-148.
- Frost, M. T, Jefferson, R. & Hawkins, S. J. (Editors). (2006). The evaluation of time series: their scientific value and contribution to policy needs. Report prepared by the Marine Environmental Change Network (MECN) for the Department for Environment, Food and Rural Affairs (DEFRA). Marine Biological Association, Plymouth. Contract CDEP 84/5/311. Marine Biological Association Occasional Publications No. 22. 94pp.
- Fulton, E.A.; Smith, A.D.M.; Webb, H.; Slater, J. (2004). Ecological indicators for the impacts of fishing on non-target species, communities and ecosystems: Review of potential indicators. *AFMA Final Research report R99/1546*: 119pp.
- Garcia, A. (2010). Comparative study of the morphology and anatomy of octopuses of the family Octopodidae. (AUT MAppSc). 247 p.
- Garcia, S.M., Rosenberg, A.A. (2010). Food security and marine capture fisheries: characteristics, trends, drivers and future perspectives. *Phil. Trans. R. Soc. B* 365, 2869-2880 doi:10.1098/rstb.2010.0171
- Gardner, J.P.A., Bell, J.J., Constable, H.B., Hannan, D., Ritchie, P.A., Zuccarello, G.C. (2010). Multi-species coastal marine connectivity: a literature review with recommendations for further research. *New Zealand Aquatic Environment and Biodiversity Report* No. 58. 47 p.
- Ghiglione, J.-F.; Galand, P.E.; Pommier, T.; Pedrós-Alió, C.; Maas, E.W.; Bakker, K.; Bertilson, S.; Kirchman, D.L.; Lovejoy, C.; Yager, P.L.; Murray, A.E. (2012). Pole-to-pole biogeography of surface and deep marine bacterial communities. *Proceedings of the National Academy of Sciences*. <http://dx.doi.org/10.1073/pnas.1208160109>
- Ghiglione, J.-F.; Galand, P.E.; Pommier, T.; Pedros-Alio, C.; Maas, E.W.; Bakker, K.; Bertilson, S.; Kirchman, D.L.; Lovejoy, C.; Yager, P.L.; Murray, A.E. (2012). Pole to pole biogeography of surface and deep marine bacterial communities. [www.pnas.org/cgi/doi/10.1073/pnas.1208160109](http://www.pnas.org/cgi/doi/10.1073/pnas.1208160109).
- Glasby GP, Wright IC. (1990). Marine mineral potential in New Zealand's Exclusive Economic Zone. *Mar Mining*;9:403-427.
- Gordon DP (2000). The Pacific Ocean and global OBIS: a New Zealand perspective. *Oceanography*;13:41-47.
- Gordon DP (Ed.) (2009). *New Zealand Inventory of Biodiversity Volume One. Kingdom Animalia: Radiata, Lophotrochozoa, Deuterostomia*. Canterbury University Press, Christchurch. 568+16 p.
- Gordon DP (Ed.) (2010) *New Zealand Inventory of Biodiversity Volume Two. Kingdom Animalia: Chaetognatha, Ecdysozoa and Ichnofossils*. Canterbury University Press, Christchurch. 528+16 p.
- Gordon DP (Ed.) (2012) *New Zealand Inventory of Biodiversity Volume Three. Kingdoms Bacteria, Protozoa, Chromista, Plantae, Fungi*. Canterbury University Press, Christchurch. 616+16 p.
- Gordon DP, editor. (2009). Volume 1. *Kingdom Animalia: Radiata, Lophotrochozoa, Deuterostomia*. Christchurch: Canterbury University Press; New Zealand inventory of biodiversity. p. 568+16.
- Gordon DP, Beaumont J, MacDiarmid A, Robertson DA, Ah Yong ST (2010) Marine Biodiversity of Aotearoa New Zealand. *PLoS ONE* 5(8): e10905. doi:10.1371/journal.pone.0010905
- Gordon DP, Bisby FA. (2009) Introduction. In: Gordon DP, editor. *New Zealand inventory of biodiversity. Volume 1. Kingdom Animalia: Radiata, Lophotrochozoa, Deuterostomia*. Christchurch: Canterbury University Press; pp. 9-12.
- Gordon DP, Hosie AM, Carter MC. (2008) Post-2000 detection of warm-water alien bryozoan species in New Zealand — the significance of recreational vessels. *Virginia Mus Nat Hist Spec Pub*;15:37-48.
- Gould B, Ah Yong ST. (2008) Marine Invasive Taxonomic Service. *Biosecurity*;85:18-19.
- Grange, K.R., Tovey, A., Hill, A.E. (2003). The spatial extent and nature of the bryozoan communities at Separation Point, Tasman Bay. *Marine Biodiversity Biosecurity Report* No. 4. 22 p.
- Grayling, S. (2004). A review of scientific studies conducted on the Macquarie Ridge. Final Research Report (ZBD2003-09) held at MPI. 33p.
- Grotti, M.; Soggia, F.; Lagomarsino, C.; Dalla Riva, S.; Goessler, W., Francesconi, K.A. (2008). Natural variability and distribution of trace elements in marine organisms from Antarctic coastal environments. *Antarctic Science* 20(1): 39-51. (peer)

- Guidetti, M. Marcato, S. Chiantore, M. Patarnello, T. Albertelli, G. Cattaneo-Vietti, R. (2006). Exchange between populations of *Adamussium colbecki* (Mollusca: Bivalvia) in the Ross Sea. *Antarctic Science* 18(4): 645-653.
- Guinotte, J.M., J. D. Bartley, A. Iqbal, D. G. Fautin, R. W. Buddemeier (2006). Modeling habitat distribution from organism occurrences and environmental data: case study using anemonefishes and their sea anemone hosts. *Marine Ecology Progress Series*, 316: 269-283.
- Hanchet, S.M. (Compiler) (2009). New Zealand IPY-CAML Progress Report. August 2009. 76 p.
- Hanchet, S.M. (Compiler) (2010). New Zealand IPY-CAML Progress Report. July 2010. p.
- Hanchet, S.M., Fu, D., Dunn, A. (2008a). Indicative estimates of biomass and yield of Whitson's grenadier (*M. whitsoni*) on the continental slope of the Ross Sea in Subareas 88.1 and 88.2. WG-FSA-08/32.
- Hanchet, S.M., Stewart, A.L., McMillan, P.J., Clark, M., O'Driscoll, R.L., Stevenson, M.L. (in press). Diversity, relative abundance, new locality records, and updated fish fauna of the Ross Sea, Antarctica. *Antarctic Science*.
- Hanchet, S.M.; Mitchell, J.; Bowden, D.; Clark, M.; Hall, J.; O'Driscoll, R.; Pinkerton, M.; Robertson, D. (2008b). Preliminary report of the New Zealand RV Tangaroa IPY-CAML survey of the Ross Sea region, Antarctica in February-March 2008. Unpublished NIWA report to CCAMLR working group on ecosystem monitoring and management. WG-EMM-08/18. 15 p.
- Hanchet, S.M.; Mitchell, J.; Bowden, D.; Clark, M.; Hall, J.; O'Driscoll, R. (2008c). Ocean survey 20/20: New Zealand IPY-CAML Final Voyage Report. NIWA Client Report: WLG2008-74, October 2008. 193 p.
- Hanchet, S.M.; Stevenson, M.L., Jones, C., Marriott, P.M., McMillan, P.J. O'Driscoll, R.L. Stevens, D., Stewart, A.L., Wood, B.A (2008d). Biomass estimates and size distributions of demersal finfish on the Ross Sea shelf and slope from the New Zealand IPY-CAML survey, February-March 2008. WG-FSA-08/31.
- Harvey, A.S., Woelkerling, W.J., Farr, T.J., Neill, K.F., Nelson, W.A., (2005). Coralline algae of central New Zealand: an identification guide to common 'crustose' species. NIWA Information Series No. 57.
- Heimeier, D. Lavery, S. Sewell MA. (2010). Using DNA barcoding and phylogenetics to identify Antarctic invertebrate larvae: Lessons from a large scale study *Marine Genomics* 3:165-177
- Hewitt J, Lundquist CJ, Bowden DA (Project ZBD2007-01 Objective 6, in review) Chatham-Challenger Ocean Survey 20/20 Post Voyage Analyses: Diversity Metrics. Ministry of Fisheries, New Zealand, Wellington
- Hewitt J, Thrush S, Lohrer A, Townsend M (2010). A latent threat to biodiversity: consequences of small-scale heterogeneity loss. *Biodiversity and Conservation* 19:1315-1323
- Hewitt J., Julian, K; Bone, E K; (2010). Chatham-Challenger Ocean Survey 20/20 Post-Voyage Analyses: Objective 10 – Biotic habitats and their sensitivity to physical disturbance New Zealand Aquatic Environment and Biodiversity Report No.
- Hewitt JE, Anderson MJ, Hickey C, Kelly S, Thrush SF (2009). Enhancing the ecological significance of contamination guidelines through integration with community analysis. *Environmental Science and Technology* 43:2118-2123
- Hewitt JE, Thrush SF, Cummings VJ, Turner SJ (1998) The effect of changing sampling scales on our ability to detect effects of large-scale processes on communities. *Journal of Experimental Marine Biology and Ecology* 227: 251-264
- Hewitt JE, Thrush SF, Dayton PD (2008). Habitat variation, species diversity and ecological functioning in a marine system. *Journal of Experimental Marine Biology and Ecology* 366:116-122
- Hewitt, J., Thrush, S., Lohrer, A. & Townsend, M. (2010). A latent threat to biodiversity: consequences of small-scale heterogeneity loss. *Biodiversity and Conservation*, 19:1315-1323.
- Hewitt, J.E. & Thrush, S.F. (2010). Empirical evidence of an approaching alternate state produced by intrinsic community dynamics, climatic variability and management actions. *Marine Ecology Progress Series*, 413:267-276.
- Hewitt, J.E., Thrush, S.F., Dayton, P.K., Bonsdorff, E. (2007). The effect of spatial and temporal heterogeneity on the design and analysis of empirical studies of scale-dependent systems. *The American Naturalist* 168 (3): 398-408.
- Hobday, AJ, Okey, TA, Poloczanska, ES, Kunz, TJ & Richardson, AJ (eds) (2006) Impacts of Climate Change on Australian Marine Life, CSIRO Marine and Atmospheric research report to the Australian Greenhouse Office, Canberra
- Horn, P. L. Forman J. Dunn M. R. (2010). Feeding habits of alfonsoino *Beryx splendens*. *Journal of Fish Biology* 76:2382-2400
- Horn, P.L., Dunn, M.R. (2010). Inter-annual variability in the diets of hoki, hake, and ling on the Chatham Rise from 1990 to 2009. New Zealand Aquatic Environment and Biodiversity Report No. 54. 57 p.
- Hurst RJ, Bagley NW, Anderson OF, Francis MJ, Griggs LH, (2000). Atlas of juvenile and adult fish and squid distributions from bottom and midwater trawls and tuna longlines in New Zealand waters. NIWA Tech Rep;84:1-612.
- Jackson, J; Carroll, E; Smith, TD; Patenaude, N; Baker, CS (in press). Taking Stock: the historical demography of the New Zealand right whale (the *Tohoro*)1830-2008. Final Research Report ZBD200505, MS12 Part D.
- Key, J.M. (2002). A review of current knowledge describing New Zealand's deepwater benthic biodiversity. *Marine Biodiversity Biosecurity Report No. 1*. 25 p.
- Koubbi, P.; Masato, M.; Duhamel, G.; Goarant, A.; Hulley, P-A; O'Driscoll, R.; Takashi, I.; Pruvost, P.; Tavenier, E.; Hosie, G. (2011). Ecological importance of micronektonic fish for the ecoregionalisation of the Indo-Pacific sector of the Southern Ocean: role of myctophids. *Deep Sea Research II* 58: 170-180. Koubbi, P.; Masato, M.; Duhamel, G.; Goarant, A.; Hulley, P-A; O'Driscoll, R.; Takashi, I.; Pruvost, P.; Tavenier, E.; Hosie, G. (in press). Ecological importance of micronektonic fish for the ecoregionalisation of the Indo-Pacific sector of the Southern Ocean: role of myctophids. *Deep Sea Research II*.
- Lalas, L; MacDiarmid AB (in press). Rapid re-colonisation of south-eastern South Island by New Zealand fur seals *Arctocephalus forsteri*. Final Research Report ZBD200505 MS12 Part E.
- Lalas, L; MacDiarmid AB; Abraham, E. (in press). : Estimates of annual food consumption by a population of New Zealand fur seals. Final Research Report ZBD200505 MS12 Part F.
- Lalas, L; MacDiarmid AB; Abraham, E. (in press). : Estimates of annual food consumption by a population of New Zealand sea lions. Final Research Report ZBD200505 MS12 Part G.
- Lavery, S., Clements, K., Hickey, A. (2006). Molecular identification of cryptogenic/invasive gobies in New Zealand. New Zealand Aquatic Environment and Biodiversity Report No 5. 48 p.
- Le Quesne, W. J. F. and Pinnegar, J. K. (2011), The potential impacts of ocean acidification: scaling from physiology to fisheries. *Fish and Fisheries*, 12: no. doi: 10.1111/j.1467-2979.2011.00423.x
- Leathwick JR, Elith J, Francis MP, Hastie T, Taylor P (2006a) Variation in demersal fish species richness in the oceans surrounding New Zealand: an analysis using boosted regression trees. *Mar Ecol Prog Ser* 321:267-281
- Leathwick JR, Rowden A, Nodder S, Gorman R, Bardsley S, Pinkerton M, Baird SJ, Hadfield M, Currie K, Goh A (2010). Development of a benthic-optimised marine environment classification for waters within the New Zealand EEZ. Final Research Report for Ministry of Fisheries Research Project BEN200601, Objective 5. (Unpublished report held by Ministry of Fisheries, Wellington.)
- Leathwick, J., Francis, M., and Julian, K. (2006b). Development of a demersal fish community map for New Zealand's Exclusive Economic Zone. NIWA Client Report HAM2006-062, prepared for Department of Conservation. National Institute of Water & Atmospheric Research, Hamilton, New Zealand.

- Leathwick, J.R.; Dey, K.L.; Julian, K. (2006c). Development of a marine environmental classification optimised for demersal fish. NIWA Client report HAM2006-063
- Leduc, D., A. A. Rowden, D. A. Bowden, P. K. Probert, C. A. Pilditch, and S. D. Nodder. (2012). Unimodal relationship between biomass and species richness of deep-sea nematodes: implications for the link between productivity and diversity. *Marine Ecology Progress Series* 454:53-64.
- Levin, L.A., Gambi, M.C., Barry, J.P., Genin, A. & Thrush, S. (2011). The Dayton legacy: baselines, benchmarks, climate, disturbance and proof. *Marine Ecology*, 32:261-165.
- Link, J. (1999) Reconstructing Food Webs and Managing Fisheries. *Ecosystem Approaches for Fisheries Management*. Proceedings of the 16th Lowell Wakefield Fisheries Symposium. AK-SG-99-01:571-588.
- Livingston ME (2009). Towards a National Marine Environment Monitoring Programme in New Zealand. A discussion paper submitted to the Biodiversity Research Advisory Group Workshop on Marine Environmental Monitoring, July 2009. (Unpublished report held at MFishMPI, Wellington)
- Livingston, M. (2004). A sampling programme to construct and quantify food-webs in two key areas supporting important fish and invertebrate species in New Zealand. Final Research Report for Ministry of Fisheries Project ENV2002-07, Objective 1. NIWA.
- Lohrer, A.M., Chiaroni, L.D., Thrush, S.F., Hewitt, J.E. (2010). Isolated and interactive effects of two key species on ecosystem function and trophic linkages in New Zealand soft-sediment habitats. *New Zealand Aquatic Environment and Biodiversity Report No. 44*. 69 p.
- Lohrer, A.M., Cummings, V.J., Thrush, S.F. (2012). Altered sea ice thickness and permanence affects benthic ecosystem functioning in coastal Antarctica. *Ecosystems* 10.1007/s10021-012-9610-7.
- Lohrer, A.M., Halliday, N.J., Thrush, S.F., Hewitt, J.E. & Rodil, I.F. (2010). Ecosystem functioning in a disturbance-recovery context: contribution of macrofauna to primary production and nutrient release on intertidal flats. *Journal of Experimental Marine Biology and Ecology*, 390:6-13.
- Lohrer, A.M., Hewitt, J.E., Hailes, S.F., Thrush, S.F., Ahrens, M. & Halliday, J. (2011). Contamination on sandflats and the decoupling of linked ecological functions. *Austral Ecology*, 36:378-388.
- Lohrer, A.M., Rodil, I.F., Townsend, M., Chiaroni, L.D., Hewitt, J.E., Thrush, S.F. (In press) Biogenic habitat transitions influence facilitation in a marine soft-sediment ecosystem. *Ecology*.
- Lohrer, A.M., Townsend, M., Rodil, I.F., Hewitt, J.E., Thrush, S.F. (In press) Detecting shifts in ecosystem functioning: the decoupling of fundamental relationships with increased pollutant stress on sandflats. *Marine Pollution Bulletin*.
- Lorrey, A. *et al.* (in press). Natural drivers of environmental change in New Zealand's marine environment during the last millennium. Final Research Report: ZBD200505 MS 6 Part A,
- Lörz, A.N. (2011a). Pacific Epimeriidae (Amphipoda: Crustacea): Epimeria. *Journal of the Marine Biological Association of the United Kingdom* 91(2): 471-477.
- Lörz, A.-N. (2011b). Biodiversity of an unknown New Zealand habitat: bathyal invertebrate assemblages in the benthic boundary layer. *Marine Biodiversity* 41(2): 299-312.
- Lörz AN, Smith PJ, Linse K, Steinke D. (in press.) High genetic diversity within *Epimeria georgiana* (Amphipoda) from the southern Scotia Arc. *Marine Biodiversity* (PDF)
- Lörz A-N & Coleman O (2009), Living jems: jewel-like creatures from the deep. *Water and Atmosphere* 17(1) 16-17 2009
- Lörz A-N (2009): Synopsis of Amphipoda from two recent Ross Sea voyages with description of a new species of Epimeria (Epimeriidae, Amphipoda, Crustacea). *Zootaxa* (2167): 59-68
- Lörz A-N, Maas E, Linse K, and Coleman C.O. (2009). Do circum-Antarctic species exist in peracarid Amphipoda? A case study in the genus Epimeria Costa, 1851 (Crustacea: Peracarida: Epimeriidae). *Zookeys* (18): 91-128 SI
- Lörz, A.N.; Berkenbusch, K.; Nodder, S.; Ah Yong, S.; Bowden, D.; McMillan, P.; Gordon, D.; Mills, S.; Mackay, K.; (2012a). A review of deep-sea benthic biodiversity associated with trench, canyon and abyssal habitats below 1500 m depth in New Zealand waters. *New Zealand aquatic environment and biodiversity report ; Ministry of Fisheries, Wellington*. 133 p.
- Lörz, A.N.; Linse, K.; Smith, P.; Steinke, D. (2012b). First evidence for underestimated biodiversity of Rhachotropis (Crustacea, Amphipoda) with description of a new species. *PloS ONE* 7(3): e32365.
- Lörz, AN (2010) Deep-sea Rhachotropis (Crustacea: Amphipoda: Eusiridae) from New Zealand and the Ross Sea with key to the Pacific, Indian Ocean and Antarctic species. *ZOOTAXA*, (2482): 22-48
- Lundquist, C.J. Thrush, S.F. Coco, G. Hewitt, J.E (2010). Interactions between distributions and dispersal decrease persistence thresholds of a marine benthic community. *Mar. Ecol. Prog. Ser.* 413: 217-228.
- Maas, E. M.; Voyles, K.M.; Pickmere, S.; Hall, J.A.; Bowden, D.A.; Clark, M.R. (2010). Bacterial and Archaeal diversity and exo-enzyme activity in Ross Sea, Antarctica sediments. Poster presented at SAME 11- Symposium on Aquatic Microbial Ecology, Piran, Slovenia, 30th August - 4th September 2010.
- Maas, E.W.; Law, C.S.; Hall, J.A.; Pickmere, S.; Currie, K.I.; Chang, F.H.; Voyles, K.M.; Caird, D. (2012). Effect of ocean acidification on bacterial abundance, activity and diversity in the Ross Sea. *Aquatic Microbial Ecology* (in review).
- Maas, E.W.; Voyles, K.M.; Caird, D.; Pickmere, S.; Hall, J.A. (2012). Bacterioplankton abundance, activity and diversity in the Ross Sea Antarctica. *Aquatic Microbial Ecology* (in review).
- MacDiarmid A, editor. (2007). A summary of the biodiversity in the New Zealand Marine Ecoregion. Wellington: WWF-New Zealand; The treasures of the sea: Ngā taonga a Tangaroa.
- MacDiarmid, A. Stewart, R (2012). Ross Sea and Balleny Islands biodiversity: routine observations and opportunistic sampling of biota made during a geophysical survey to the Ross Sea in 2006. . *Aquatic Environment and Biodiversity Report No. XX*, 99 p.
- MacDiarmid, A.; McKenzie, A.; Sturman, J.; Beaumont, J.; Mikaloff-Fletcher, S.; Dunne, J. (2012). Assessment of anthropogenic threats to New Zealand marine habitats New Zealand Aquatic Environment and Biodiversity Report No. 93.255 p.
- MacDiarmid, A.B.; Cleaver, P.; Stirling, B. (in press). Historical evidence for exploitation of the marine environment in the Hauraki Gulf and along the Otago/Catkins shelf 1790- 1930 *Aquatic Environment and Biodiversity Report No XX*.,
- MacDiarmid, AB Smith I, Paul L, Francis M, McKenzie A, Parsons D, Hartill B, Stirling B, Cleaver et al (in press) A complete history of the exploitation of an ecologically important inshore finfish species in the Hauraki Gulf, New Zealand: a synthesis of archaeological, historical and fisheries data. *Aquatic Environment and Biodiversity Report XX*, 25 p.
- McKenzie, A; MacDiarmid, AB (in press). Biological parameters and biomass estimates for some commercial fish stocks in the Hauraki Gulf and Otago-Catkins shelf for the period 1930-2006. *Aquatic Environment and Biodiversity Report XX*, 97 p.
- McMillan, P.J.; Francis, M.P.; G.D., J.; Paul, L.J.; Marriott, P.J.; Mackay, E.; Wood, B.A.; Griggs, L.H.; Sui, H.; Wei, F. (2011a). New Zealand fishes. Volume 1: A field guide to common species caught by bottom and midwater fishing. *New Zealand Aquatic Environment and Biodiversity Report* 68.

- McMillan, P.J.; Francis, M.P.; Paul, L.J.; Marriott, P.J.; Mackay, E.; Baird, S.J.; Griggs, L.H.; Sui, H.; Wei, F. (2011b). New Zealand fishes. Volume 2: A field guide to less common species caught by bottom and midwater fishing. New Zealand Aquatic Environment and Biodiversity Report 78.
- McMillan, P.J.; Griggs, L.H.; Francis, M.P.; Marriott, P.J.; Paul, L.J.; Mackay, E.; Wood, B.A.; Sui, H.; Wei, F. (2011c). New Zealand fishes. Volume 3: A field guide to common species caught by surface fishing. New Zealand Aquatic Environment and Biodiversity Report 69.
- McMillan, P.J., Iwamoto, T., Stewart, A., Smith, P.J. (2012). A new species of grenadier, genus *Macrourus* (Teleostei, Gadiformes, Macrouridae) from the southern hemisphere and a revision of the genus. *Zootaxa*, **3165**, 1–24.
- McWethy, D.B., Whitlock, C., Wilmshurst, J.M., McGlone, M.S., Fromont, M., Li, X., Dieffenbacher-Krall, A., Hobbs, W.O., Fritz, S.C., Cook, E.R. (2010). Rapid landscape transformation in South Island, New Zealand, following initial Polynesian settlement. *Proceedings of the national Academy of Sciences* 107: 21343–21348.
- MfE (2007). Environment New Zealand 2007.
- Maxwell, K.; MacDiarmid, A.B. (in press). Oral histories of marine fish and shellfish state and use in the Hauraki Gulf and along the Otago/Catlins coast 1940–2000. Aquatic Environment and Biodiversity Report XX.
- Ministry for Primary Industries (2012). Report from the Fisheries Assessment Plenary, May 2012: stock assessments and yield estimates. Compiled by the Fisheries Science Group, Ministry for Primary Industries, Wellington, New Zealand. 1194 p.
- Mitchell L. Sogin, Hilary G. Morrison, Julie A. Huber, David Mark Welch, Susan M. Huse, Phillip R. Neal, Jesus M. Arrieta, and Gerhard J. Herndl (2006). Microbial diversity in the deep sea and the underexplored “rare biosphere” *Proceedings Natural Academy Science U S A*. 2006 August 8; 103(32): 12115–12120. [www.ncbi.nlm.nih.gov/pmc/articles/PMC1524930/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1524930/)
- Mitchell, J. (2008). Initial environmental evaluation (IEE) NZ IPY-CAML voyage 2008 report. Report prepared for Antarctic Policy Unit, Ministry of Foreign Affairs and Trade. May 2008. 35 p.
- Mitchell, J. MacDiarmid, A. (2006) Voyage Report TAN06-02 EASTERN ROSS SEA VOYAGE Unpublished report for MPI project ZBD2005-03
- Mitchell, J.; Clark, M. (2004). Voyage report TAN04-02. Western Ross Sea voyage 2004. Hydrographic and biodiversity survey RV Tangaroa 27 January to 13 March 2004. Cape Adare, Cape Hallett, Possession Islands and Balleny Islands, Antarctica. NIWA Voyage Report TAN04-02. 102 p. (Unpublished report held in NIWA library, Wellington.)
- Morrison M, Lowe M, Parsons D, Usmar N, McLeod I (2009). A review of land-based effects on coastal fisheries and supporting biodiversity in New Zealand. New Zealand Aquatic Environment and Biodiversity Report No. 37. 100 p.
- Murphy, E.J., Cavanagh, R.D., Hofmann, E. E., Hill, S.L., Constable, A.J., Costa, D.P., Pinkerton, M.H., Johnston, N. M., Trathan, P. N., Klinck, J.M., Wolf-Gladrow, D.A., Daly, K.L., Maury, O. and Doney, S.C. (2012) Developing integrated models of Southern Ocean food webs: including ecological complexity, accounting for uncertainty and the importance of scale. *Progress in Oceanography*, <http://dx.doi.org/10.1016/j.pocean.2012.03.006>.
- Needham, H.C., Pilditch, C.A., Lohrer, A. & Thrush, S. (2010). Habitat dependence in the functional traits of *Austrohelice crassa*, a key bioturbating species. *Marine Ecology Progress Series*, 414:179–193.
- Needham, H.C., Pilditch, C.A., Lohrer, A.M. & Thrush, S.F. (2011). Context-specific bioturbation mediates changes to ecosystem functioning. *Ecosystems*, 14:1096–1109.
- Needham, H.R., Pilditch, C.A., Lohrer, A.M., Thrush, S.F. (In press). Density and habitat dependent effects of crab burrows on sediment erodibility. *Journal of Sea Research*
- Neil, H., et al (in press). Insights into historical marine productivity using ancient fish otoliths. Final Research Report: ZBD200505 MS 6 Part B.
- Neill, K., D’Archino, R., Farr, T., Nelson, W. (2012). Macroalgal diversity associated with soft sediment habitats in New Zealand. NZ Aquatic Environment and Biodiversity Report 87. 127 pp.
- Nelson WA, Gordon DP. (1997). Assessing New Zealand’s marine biological diversity — a challenge for policy makers and systematists. *NZ Sci Rev.*54:58–66.
- Nelson, W., Cummings, V., D’Archino, R., Halliday, J., Marriott, P., Neill, K. (2010). Macroalgae and benthic biodiversity of the Balleny Islands, Southern Ocean. New Zealand Aquatic Environment and Biodiversity Report No. 55. 99 p.
- Nelson, W.A. Villouta, E. Neill, K.F. Williams, G.C.Adams, N.M. Slivsgaard, (2002)R. Marine Macroalgae of Fiordland, New Zealand *Tuhinga* 13: 117–152 Copyright © Te Papa Museum of New Zealand
- Nelson, W.A.; Neill, K.; Farr, T.; Barr, N.; D’Archino, R.; Miller, S.; Stewart, R. (2012). Rhodolith beds in northern New Zealand: characterisation of associated biodiversity and vulnerability to environmental stressors. New Zealand Aquatic Environment and Biodiversity Report 99. 102 pp.
- Nelson, W.A. (2009). Calcified macroalgae - critical to coastal ecosystems and vulnerable to change: A review. *Marine and Freshwater Research* 60:787–801.
- Nodder S, Maas E, Bowden D, Pilditch C. (2011) Physical, Biogeochemical and Microbial Characteristics of Sediment Samples from the Chatham Rise and Challenger Plateau. New Zealand. Draft Aquatic Environment and Biodiversity Report No 70.
- Nodder SD (2008) OS 20/20 Chatham Rise & Challenger Plateau Hydrographic, Biodiversity & Seabed Habitats, NIWA Client Report: WLG2008-27, National Institute of Water & Atmospheric Research, Wellington, New Zealand
- Norkko, A., Andrew, N., Thrush, S., Cummings, V., Schwarz, A-M., Hawes, I., Mercer, S., Budd, R. Gibbs, M., Funnell, G., Hewitt, J., Goring, D. (2002). Ecology and biodiversity of coastal benthic communities in McMurdo Sound, Ross Sea: development of sampling protocols and initial results. Final Research Report for Ministry of Fisheries Research Project ZBD2001/02, Objectives 1, 2 & 3. 119p. (peer)
- Norkko, A., Thrush, S.F., Cummings, V.J., Funnell, G.A., Schwarz, A-M.S., Andrew, N.L., Hawes, I. (2004). Ecological role of Phyllophora antarctica drift accumulations in coastal soft-sediment communities of McMurdo Sound, Antarctica. *Polar Biology* 27: 482–494.
- Norkko, A., Thrush, S.F., Cummings, V.J., Gibbs, M.M., Andrew, N.L., Norkko, J., Schwarz, A.-M. (2007). Trophic structure of coastal Antarctic food webs associated with changes in sea ice and food supply. *Ecology* 88: 2810–2820.
- Norkko, J., Norkko, A., Thrush, S.F., Cummings, V.J. (2005). Growth under environmental extremes: spatial and temporal patterns in nucleic acid ratios in two Antarctic bivalves. *Journal of Experimental Marine Biology and Ecology* 326: 114–156.
- Norkko, J., Norkko, A., Thrush, S.F., Valanko, S. & Suurkuikka, H. (2010). Conditional responses to increasing scales of disturbance, and potential implications for threshold dynamics in soft-sediment communities. *Marine Ecology Progress Series*, 413:253–266.
- O’Driscoll, R.L., Macaulay, G.J., Gauthier, S., Pinkerton, M., Hanchet, S. (2010). Distribution, abundance and acoustic properties of Antarctic silverfish (*Pleuragramma antarcticum*) in the Ross Sea. CCAMLR document WG-FSA-10/P4. 17 p. (Deep-Sea Research II (2010), <http://dx.doi.org/10.1016/j.dsr2.2010.05.018>)
- O’Driscoll R.L. (2009). Preliminary acoustic results from the New Zealand IPY-CAML survey of the Ross Sea region in February–March 2008. Final Research Report for MFishMPI project IPY200701 objective 8. 14 p.

- O'Driscoll R.L.; Macaulay, G.J.; Gauthier, S.; Pinkerton, M.; Hanchet, S. (2009). Preliminary acoustic results from the New Zealand IPY-CAML survey of the Ross Sea region in February-March 2008. CCAMLR SG-ASAM-09-05. 37 p.
- O'Driscoll, R.L.; Hanchet, S.M.; Miller, B.S. (in press). Can acoustic methods be used to monitor grenadier (Macrouridae) abundance in the Ross Sea region? *Journal of Ichthyology*.
- O'Driscoll, R.L.; Macaulay, G.J.; Gauthier, S.; Pinkerton, M.; Hanchet, S. (2011). Distribution, abundance, and acoustic properties of Antarctic silverfish (*Pleuragramma antarcticum*) in the Ross Sea. *Deep Sea Research II* 58: 181–195.
- O'Loughlin MP, Paulay G, Davey N, Michonneau F (2011) The Antarctic region as a marine biodiversity hotspot for echinoderms: Diversity and diversification of sea cucumbers. *Deep Sea Research Part II: Topical Studies in Oceanography* 58:264-275
- Paavo, B., Jonker, R., Thrush, S. & Probert, P.K. 2011. Macrofaunal community patterns of adjacent coastal sediments with wave-reflecting or wave-dissipating characteristics. *Journal of Coastal Research*, 27:515-528.
- Page M.J.; Alcock, N.; Gordon, D.; Kelly-Shanks, M.; Nelson, W.; Neill, K.; Watson, J. (2001). Preliminary assessment of the biodiversity of benthic macrofauna of the western Ross Sea, Antarctica. Final Research Report to the Ministry of Fisheries. 29 p
- Pakhomov, E.A.; Hall, J.; Williams, M.J.M.; Hunt, B.P.V.; Stevens, C.J. (2011) Biology of *Salpa thompsoni* in waters adjacent to the Ross Sea, Southern Ocean, during austral summer 2008. *Polar Biology* 34: 257–271. (DOI 10.1007/s00300-010-0878-9)
- Parker, S.J., A.J. Penney and M.R. Clark (2009). Detection criteria for managing trawl impacts on vulnerable marine ecosystems in high seas fisheries of the South Pacific Ocean. *Marine Ecology Progress Series*, 397: 309–317.
- Parker, S.J., S. Mormede, D. Tracey and M. Carter. (2009). Evaluation of VME taxa monitoring by scientific observers from New Zealand in the Ross Sea Antarctic toothfish longline fishery during the 2008-09 season. Document WG-TASO 09/08. CCAMLR, Hobart, Australia. 13p.
- Parravicini, V., Thrush, S.F., Chiantore, M., Morri, C., Croci, C. & Bianchi, C.N. (2010). The legacy of past disturbance: Chronic angling impairs long-term recovery of marine epibenthic communities from acute date-mussel harvesting. *Biological Conservation*, 143:2435-2440.
- Parsons, D *et al.* (2012). Risks of shifting baselines highlighted by anecdotal accounts of New Zealand's snapper fishery . *New Zealand Journal of Marine and Freshwater Research* 43: 965-983.
- Paul, L. (in press). Trends in the exploitation of finfish from the Hauraki Gulf, 1850–2006. *Aquatic Environment and Biodiversity Report* No. XX, 156 p.
- Paul, L., (2012). A history of the Firth of Thames dredge fishery for mussels: use and abuse of a coastal resource, *New Zealand Aquatic Environment and Biodiversity Report* No. 94, p. 27.
- Peart, R, Serjeant, K. Mulcahy K. (2011). Governing our oceans. Environmental reform for the exclusive Economic Zone. Environmental Defence Society Policy Paper; Available as e-book: <http://www.eds.org.nz/>.
- Pinkerton, M. (in press) Mass balance modeling of a NZ shelf ecosystem since 1200AD. *Aquatic Environment and Biodiversity Report* No. .
- Pinkerton, M., Hanchet, S., Bradford-Grieve, J., Cummings, V., Wilson, P., Williams, M. (2006). Modelling the effects of fishing in the Ross Sea. Final Research Report for Ministry of Fisheries Research Project ANT2004-05. 169 p.
- Pinkerton, M.H., A. Dunn and S.M. Hanchet. (2007). Ecological risk management and the fishery for Antarctic toothfish (*Dissostichus mawsoni*) in the Ross Sea, Antarctica. Pp 22. Document EMM-07-24, CCAMLR, Hobart, Australia.
- Pinkerton, M.H., J. Bradford-Grieve and S.M. Hanchet. (2009a). A balanced model of the food web of the Ross Sea, Antarctica. Submitted to CCAMLR Science. Under review.
- Pinkerton, M.H.; Forman, J.; Bury, S.J.; Brown, J.; Horn, P.; O'Driscoll, R.L. (in press). Diet and trophic niche of Antarctic silverfish (*Pleuragramma antarcticum*) in the Ross Sea, Antarctica. *Journal of Fish Biology*.
- Pinkerton, M.H.; J. Forman; D.W. Stevens; S.J. Bury; J. Brown (2012a). Diet and trophic niche of Macrourus spp. (Gadiformes, Macrouridae) in the Ross Sea region of the Southern Ocean. *Journal of Ichthyology*, Special Issue on Grenadiers (Ed: Alexei Orlov). (in press)
- Pinkerton, M.H.; V. Cummings; J. Forman; J. Brown; S.J. Bury (2009b). Trophic connections in the Ross Sea: information from stomach contents analysis and stable isotopes of carbon and nitrogen. Report to Ministry of Fisheries, project IPY200701 Obj10. Pp 18.
- Probert PK, McKnight DG, Grove SL. (1997) Benthic invertebrate bycatch from a deep-water trawl fishery, Chatham Rise, New Zealand. *Aquat Conserv: Mar Freshw Ecosyst*;7:27–40.
- Ramirez-Llodra E, Tyler PA, Baker MC, Bergstad OA, Clark MR, *et al.* (2011). Man and the Last Great Wilderness: Human Impact on the Deep Sea. *PLoS ONE* 6(8): e22588. doi:10.1371/journal.pone.0022588
- Reid, D.J., Chiaroni, L.D., Hewitt, J.E., Lohrer, A.M., Matthaei, C.D., Phillips, N.R., Scarsbrook, M.R., Smith, B.J., Thrush, S.F., Townsend, C.R., Van Houte-Howes, K.S.S. & Wright-Stow, A.E. 2011. Sedimentation effects on the benthos of streams and estuaries: A cross-ecosystem comparison. *Marine and Freshwater Research*, 62:1201-1213.
- Roberts CD, Stewart AL. (2001). Ross Sea fishes: a collection-based biodiversity research programme. *Seafood New Zealand*, Dec. 9(11): 79-84.
- Rodil, I.F., Lohrer, A.M., Chiaroni, L.D., Hewitt, J.E. & Thrush, S. (2011). Disturbance of sandflats by thin deposits of terrigenous sediment: consequences for primary production and nutrient release. *Ecological Applications*, 21:416-426.
- Rodríguez LO, editor. [UNEP/CBD/SBSTTA/4/INF6.]. Paris: IUBS/DIVERSITAS; (2000). Implementing the GTI: Recommendations from DIVERSITAS Core Programme Element 3, Including an Assessment of Present Knowledge of Key Species Groups.
- Rogers, A.D. & Laffoley, D.d'A. (2011). International Earth system expert workshop on ocean Stresses and impacts. Summary report. IPSO Oxford, 18 pp.
- Rowden AA, Berkenbusch K, Brewin PE, Dalen J, Neill KF, Nelson WA, Oliver MD, Probert PK, Schwarz A-M, Sui PH, Sutherland D (2007). A review of the marine soft-sediment assemblages of New Zealand. *NZ Aquatic Environment and Biodiversity Report*. 184 pp.
- Rowden AA, O'Shea S, Clark MR. (2002) Benthic biodiversity of seamounts on the northwest Chatham Rise. [NZ Min Fish] *Mar Biodiv Biosecurity Rep*.;2:1–21.
- Rowden, A.A. Kroger K. Clark, M. (in press). Biodiversity of macroinvertebrates and fish assemblages of the Balleny Islands seamounts. *AEBR Series*
- Rowden, A.A., Clark, M.R., O'Shea, S., McKnight, D.G. (2003). Benthic biodiversity of seamounts on the southern Kermadec volcanic arc. *Marine Biodiversity Biosecurity Report* No. 3. 23 p.
- Rowden, A.A., Clark, M.R. (2010). Benthic biodiversity of seven seamounts on the southern end of the Kermadec volcanic arc, northeast New Zealand. *New Zealand Aquatic Environment and Biodiversity Report* No. 62. 31 p.
- Rowden, A.A., Clark, M.R., O'Shea, S. (2004). Benthic biodiversity of seamounts on the Northland Plateau. *Marine Biodiversity Biosecurity Report* No. 5. 21 p.
- Rowden, A.A., Oliver, M., Clark, M.R., Mackay, K. (2008). New Zealand's "SEAMOUNT" database: recent updates and its potential use for ecological risk assessment. *New Zealand Aquatic Environment and Biodiversity Report* No. 27. 49 p.

- Rowden, A.A., O'Shea, S., Clark, M.R. (2002). Benthic biodiversity of seamounts on the northwest Chatham Rise. *Marine Biodiversity Biosecurity Report No. 2*. 21 p.
- Rowden, A.A.; Clark, M.R.; Wright, I.C. (2005). Physical characterisation and a biologically focused classification of "seamounts" in the New Zealand region. *New Zealand Journal of Marine and Freshwater Research* 39: 1039-1059.
- Russell LK, Hepburn CD, Hurd CL, Stuart MD. 2008 The expanding range of *Undaria pinnatifida* in southern New Zealand: distribution, dispersal mechanisms and the invasion of wave-exposed environments. *Biol Invasions*.;10:103–115.
- Safi, K.; Robinson, K.; Hall, J.; Schwarz, J.; Maas, E. (2012). Ross Sea deep-ocean and epipelagic microzooplankton during the summer-autumn transition period. *Aquatic Microbial Ecology* 67(2): 123-137.
- Savage C, Thrush SF, Lohrer AM, Hewitt JE (2012) Ecosystem Services Transcend Boundaries: Estuaries Provide Resource Subsidies and Influence Functional Diversity in Coastal Benthic Communities. *PLoS ONE* 7(8): e42708. doi:10.1371/journal.pone.0042708
- Savage, C. (2009). Development of bioindicators for the assimilation of terrestrial nutrient inputs in coastal ecosystems as a tool for watershed management. *New Zealand Aquatic Environment and Biodiversity Report No. 30*. 35 p.
- Schiaparelli S, Alvaro MC, Bohn J, Albertelli G (2010) 'Hitchhiker' polynoid polychaetes in cold deep waters and their potential influence on benthic soft bottom food webs. *Antarctic Science* 22:399-407
- Schiel, D.R. (2011) Biogeographic patterns and long-term changes on New Zealand coastal reefs: Non-trophic cascades from diffuse and local impacts. *Journal of Experimental Marine Biology and Ecology* 400:1-2, 33-51 Online publication date: 1-Apr-2011 Read More: <http://www.esajournals.org/doi/abs/10.1890/03-3107>
- Schwarz, A., Taylor, R., Hewitt, J., Phillips, N., Shima, J., Cole, R., Budd, R. (2006). Impacts of terrestrial runoff on the biodiversity of rocky reefs. *New Zealand Aquatic Environment and Biodiversity Report No 7*. 109 p.
- Schwarz, A-M., Hawes, I., Andrew, N., Mercer, S., Cummings, V., Thrush, S. (2005). Primary production potential of non-geniculate coralline algae at Cape Evans, Ross Sea, Antarctica. *Marine Ecology Progress Series* 294: 131-140.
- Schwarz, A-M., Hawes, I., Andrew, N., Norkko, A., Cummings, V., Thrush, S. (2003). Macroalgal photosynthesis near the southern global limit for growth; Cape Evans, Ross Sea, Antarctica. *Polar Biology* 26: 789-799.
- Sewell, M.A., (2005). Examination of the meroplankton community in the southwestern Ross Sea, Antarctica, using a collapsible plankton net. *Polar Biol.* 28:119–131.
- Sewell, M.A., (2006). The meroplankton community of the northern Ross Sea: a preliminary comparison with the McMurdo Sound region. *Antarct. Sci.* 18:595–602.
- Sewell, M.A., Lavery, S., Baker, C.S. (2006). Whose larva is that? Molecular identification of planktonic larvae of the Ross Sea. *New Zealand Aquatic Environment and Biodiversity Report No. 3*. 57 p.
- Sharp, B.R., S.J. Parker, M.H. Pinkerton (lead authors); also B.B. Breen, V. Cummings, A. Dunn, S.M. Grant, S.M. Hanchet, H.J.R. Keys, S.J. Lockhart, P.O'B. Lyver, R.L. O'Driscoll, M.J.M. Williams, P.R. Wilson. (2010). Bioregionalisation and Spatial Ecosystem Processes in the Ross Sea Region. CCAMLR document WG-EMM-10/30, Hobart, Australia.
- Smith AM. (2009) Bryozoans as southern sentinels of ocean acidification: a major role for a minor phylum. *Mar Freshw Res.*;60:475–482.
- Smith AM, Gordon, D. (2011). Bryozoans of southern New Zealand Field Identification Guide AEBR.
- Smith PJ, Steinke D, McMillan PJ, Stewart AL, McVeagh SM, Diaz De Astarloa JM, Welsford D, Ward RD. (2011). DNA barcoding highlights a cryptic species of grenadier (genus *Macrourus*) in the Southern Ocean. *Journal of Fish Biology*, 78(1):355-65.
- Smith PJ, Steinke, D., McMillan PJ, Stewart AL, Ward RD. (2011). DNA barcoding of morids (Actinopterygii, Moridae) reveals deep divergence in the anti tropical *Halargyreus johnsoni* but little distinction between *Antimora rostrata* and *A. microlepis*. *Mitochondrial DNA*, (doi:10.3109/19401736.2010.532329).
- Smith, F. (2006). Balleny Islands Ecology Research Voyage Report, R.V. Tiama. Unpublished report held at Ministry of Fisheries for project ZBD2005/01.
- Smith, I.W.G. (2005). Retreat and resilience: Fur seals and human settlement in New Zealand. In: Monks, G.G. (ed) *The Exploitation and Cultural Importance of Marine Mammals*. Oxford, Oxbow Books. 6-18.
- Smith, I.W.G. (2011). Estimating the magnitude of pre-European Maori marine harvest in two New Zealand study areas, New Zealand Aquatic Environment and Biodiversity Report, No. 82, p. 72 p)
- Smith, P.J., Steinke, D., McMillan, P.J., McVeagh, S.M., Struthers, C.D. (2008). DNA database for commercial marine fish. *New Zealand Aquatic Environment and Biodiversity Report No. 22*. 62 p.
- Snelder T, Leathwick J, Dey K, Rowden A, Weatherhead M, (2006). Development of an ecological marine classification in the New Zealand region. *Envir Managemt.*;39:12–29.
- Stein D. (2012). Snailfishes (Family Liparidae) of the Ross Sea, Antarctica, and closely adjacent waters. *Zootaxa* 3285. 120 pp. ISBN 978-1-86977-870-5 (Online edition)
- Stevens DW, Dunn MR. (2010). Different food preferences in four sympatric deep-sea Macrourid fishes. *Marine Biology*. DOI 10.1007/s00227-010-1542-1
- Stevens, D.W.; Hurst, R.J.; Bagley, N.W. (2011). Feeding habits of New Zealand fishes: a literature review and summary of research trawl database records 1960 to 2000. *New Zealand Aquatic Environment and Biodiversity Report No*
- Sutherland DL. (2008). Surface-associated diatoms from marine habitats at Cape Evans, Antarctica, including the first record of living *Eunotogramma marginopunctatum*. *Polar Biology* DOI 10.1007/s00300-008-0426-z.
- Thrush SF and Dayton PK (2002) Disturbance to marine benthic habitats by trawling and dredging: implications for marine biodiversity. *Annual Review of Ecology and Systematics* 33:449–473
- Thrush, S.F. and Cummings, V.J. (2011). Massive icebergs, alteration in primary food resources and change in benthic communities at Cape Evans, Antarctica. *Marine Ecology*, 32:289–299.
- Thrush, S.F. & Dayton, P. (2010). What can ecology contribute to Ecosystem-based Management of marine fisheries? *Annual Reviews in Marine Science*, 2:419-441.
- Thrush, S.F. & Lohrer, A.M. (2012). Why bother going outside: the role of observational studies in understanding biodiversity-ecosystem function relationships. In: *Marine biodiversity futures and ecosystem functioning Frameworks, methodologies and integration*, edited by D. M. Paterson, M. Solan and R. Aspinall, Oxford University Press, p198-212.
- Thrush, S.F. D. Lohrer and C. Savage (in press) Carbonate sediments: the positive and negative effects of land-coast interactions on functional diversity, AEBC report;
- Thrush, S.F., Chiantore, M., Asnagi, V., Hewitt, J., Fiorentino, D. & Cattaneo-Vietti, R. (2011). Habitat-diversity relationships in rocky shore algal turf infaunal communities *Marine Ecology Progress Series*, 424:119-132.
- Thrush, S.F., Cummings, V.J. (in press) Massive icebergs, alteration in primary food resources and change in benthic communities at Cape Evans, Antarctica. *Marine Ecology*.
- Thrush, S.F., Cummings, V.J. (2011). Massive icebergs, alteration in primary food resources and change in benthic communities at Cape Evans, Antarctica. *Marine Ecology* 32 (3): 289–299. doi: 10.1111/j.1439-0485.2011.00462.x

- Thrush, S.F., Dayton, P.K., Cattaneo-Vietti, R., Chiantore, M., Cummings, V.J., Andrew, N.L., Hawes, I., Kim, S., Kvitek, R., Schwarz, A.-M. (2006). Broad-scale factors influencing the biodiversity of coastal benthic communities of the Ross Sea. *Deep Sea Research II* 53: 959–971.
- Thrush, S.F., Hewitt, J., Cummings, V.J., Norkko, A. & Chiantore, M. (2010).  $\beta$ -Diversity and species accumulation in Antarctic coastal benthos: Influence of habitat, distance and productivity on ecological connectivity. *PLoS ONE*, 5: E11899.
- Thrush, S.F., Hewitt, J.E. & Lohrer, A.M. (2012). Interaction networks in coastal soft-sediments highlight the potential for change in ecological resilience. *Ecological Applications*, 22:1213–1223.
- Thrush, S.F., Hewitt, J.E., Lohrer A.M., Chiaroni L.D. (In press). When small changes matter: the role of cross-scale interactions between habitat and ecological connectivity in recovery. *Ecological Applications*.
- Thrush, S.F.; Hewitt, J.E.; Cummings, V.J.; Green, M.O.; Funnell, G.A.; Wilkinson, M.R. (2000). The generality of field experiments: interactions between local and broad-scale processes. *Ecology* 81: 399–415.
- Thrush, S.F.; Pridmore, R.D.; Bell, R.G.; Cummings, V.J.; Dayton, P.K.; Ford, R.; Grant, J.; Hewitt, J.E.; Hines, A.H.; Hume, T.M.; Lawrie, S.M.; Legendre, P.; McArdle, B.H.; Morrissey, D.; Schneider, D.C.; Turner, S.J.; Walters, R.; Whitlatch, R.B.; Wilkinson, M.R. (1997). The sandflat habitat: Scaling from experiments to conclusions. *Journal of Experimental Marine Biology and Ecology* 216: 1–9. [did not see this referred to in the text]
- Tittensor, D.P., A.R. Baco, P.E. Brewin, M.R. Clark, M. Consalvey, J. Hall-Spencer, A.A. Rowden, T. Schlacher, K.I. Stocks and A.D. Rogers (2009). Predicting global habitat suitability for stony corals on seamounts. *Journal of Biogeography*, 36: 1111–1128.
- Townsend, M., Thrush, S. & Carabines, M. (2011). Simplifying the complex: an ecosystem principles approach to goods and services management in marine coastal systems. *Marine Ecology Progress Series*, 434:291–301.
- Tracey, D.; Baird, S.J.; Sanders, B.M.; Smith, M.H. (2011). Distribution of protected corals in relation to fishing effort and assessment of accuracy of observer identification. NIWA Client Report No: WLG2011-33 prepared for Department of Conservation, Wellington. 74 p.
- Tracey, D., Carter, M., Parker, S. (2010). Evaluation of VME taxa monitoring by scientific observers. Final Research Report for Ministry of Fisheries Research Project ANT2009/01 Objective 8. 17 p.
- Tracey, D., Rowden, A., Mackay, K., Compton, T. (2011). Habitat-forming coldwater corals show affinity for seamounts in the New Zealand region. *Mar Ecol Prog Ser* 430, 1–22.
- Tracey, D., Susan Jane Baird, S.J. Brian Sanders, B., Smith, M. H. Smith (2011). Identification of Protected Corals: distribution in relation to fishing effort and accuracy of observer identifications (MCSINT 2010/03). Final Report prepared for Marine Conservation Services (MCS), Department of Conservation | Te Papa Atawhai. 74 p.
- Tracey, D.M., Anderson, O.F., Clark, M.R., Oliver, M.D. (2005). A guide to common deepsea invertebrates in New Zealand waters. New Zealand Aquatic Environment and Biodiversity Report No. 1. 160 p.
- Tracey, D.M., Anderson, O.F., Naylor, J. R. (2007). A guide to common deepsea invertebrates in New Zealand waters. New Zealand Aquatic Environment and Biodiversity Report No. 10. 282 p.
- Tracey, D.M.; Anderson, O.F.; Naylor, R.J. (2011). A guide to common deepsea invertebrates in New Zealand waters. New Zealand Aquatic Environment and Biodiversity Report 86. 317 p.
- Tuck, I., Cole, R., Devine, J. (2009). Ecosystem indicators for New Zealand fisheries. New Zealand Aquatic Environment and Biodiversity Report No. 42. 188 p.
- Tuck, I., Drury, J., Kelly, M., Gerring, P. (2010). Designing a programme to monitor the recovery of the benthic community between North Cape and Cape Reinga. New Zealand Aquatic Environment and Biodiversity Report No. 53. 78 p.
- Turner SJ, Thrush SF, Hewitt JE, Cummings VJ, Funnell GA. (1999). Fishing impacts and the degradation or loss of habitat structure. *Fish Managemt Ecol*;6:401–420.
- Varian, S.J. (2005). A summary of the values of the Balleny Islands, Antarctica. *Marine Biodiversity Biosecurity Report* No. 6. 13 p.
- Williams R, Gould B, Christian S. Shipwrecks 2008— an international biosecurity risk? *Surveillance*;35:4–6.
- Williams, A.; Schlacher, T.A.; Rowden, A.A.; Althaus, F.; Clark, M.R.; Bowden, D.A.; Stewart, R.; Bax, N.J.; Consalvey, M.; Kloser, R.J. (2010) Seamount megabenthic assemblages fail to recover from trawling impacts. *Marine Ecology* 31(suppl. 1): 183–199.
- Wing, S.R. (2005) Fiordland Biodiversity Research Cruise Final Research Report ZBD2003-04
- Woelkerling, Wm J.; Nelson, W.A. (2004). A baseline summary and analysis of the taxonomic biodiversity of coralline red algae (Corallinales, Rhodophyta) recorded from the New Zealand region. *Cryptogamie Algologie* 25: 39–106.
- Woodin, S.A., Wetthey, D.S., Hewitt, J.E. & Thrush, S.F. (2012). Small scale terrestrial clay deposits on intertidal sandflats: Behavioral changes and productivity reduction. *Journal of Experimental Marine Biology and Ecology*, 413:184–191.

## 11.6. Appendix

### Technical rationale for the goals and targets of the strategic plan for the period 2011-2020. UNEP/CBD/COP/10/9 18 July 2010.

#### *Strategic goal A. Address the underlying causes of biodiversity loss by mainstreaming biodiversity across government and society*

Strategic actions should be initiated immediately to address, over a longer term, the underlying causes of biodiversity loss. This requires policy coherence and the integration of biodiversity into all national development policies and strategies and economic sectors and at all levels of government. Approaches to achieve this include communication, education and public awareness, appropriate pricing and incentives, and the broader use of planning tools such as strategic environmental assessment. Stakeholders across all sectors of government, society and the economy, including business, will need to be engaged as partners to implement these actions. Consumers and citizens must also be mobilized to contribute to biodiversity conservation and sustainable use, to reduce their ecological footprints and to support action by Governments.

[Note: Targets 1-5 not given here.] Targets 6-11 are directly quoted from the document.

**Target 6: By 2020, overfishing is ended, destructive fishing practices are eliminated, and all fisheries are managed sustainably.] or [By 2020, all exploited fish stocks and other living marine and aquatic resources are harvested sustainably [and restored], and the impact of fisheries on threatened species and vulnerable ecosystems are within safe ecological limits.**

Overexploitation is the main pressure on marine fisheries globally and the World Bank estimates that overexploitation represents a lost profitability of some \$50 billion per year and puts at risk some 27 million jobs and the well-being of more than one billion people. Better fisheries management, which may include a reduction in fishing effort is needed to reduce pressure on ecosystems and to ensure the sustainable use of fish stocks. The specific target should be regarded as a step towards ensuring that all fisheries are sustainable while building upon existing initiatives such as the Code of Conduct for Responsible Fishing. Indicators to measure progress towards this target include the Marine Trophic Index, the proportion of products derived from sustainable sources and trends in abundance and distribution of selected species. Other possible indicators include the proportion of collapsed species, fisheries catch, catch per unit effort, and the proportion of stocks overexploited. Baseline information for several of these indicators is available from the Food and Agriculture Organization of the United Nations.

**Target 7: By 2020, areas under agriculture, aquaculture and forestry are managed sustainably, ensuring conservation of biodiversity.**

The increasing demand for food, fibre and fuel will lead to increasing losses of biodiversity and ecosystem services if management systems do not become increasingly sustainable with regard to the biodiversity. Criteria for sustainable forest management have been adopted by the forest sector and there are many efforts by Governments, indigenous and local communities, NGOs and the private sector to promote good agricultural, aquaculture and forestry practices. The application of the ecosystem approach would also assist with the implementation of this target. While, as yet, there are no universally agreed sustainability criteria, given the diversity of production systems and environmental conditions, each sector and many initiatives have developed their own criteria which could be used pending the development of a more common approach. Similarly, the use of certification and labelling systems or standards could be promoted as part of this target. Relevant indicators for this target include the area of forest, agricultural and aquaculture ecosystems under sustainable management, the proportion of products derived from sustainable sources and trends in genetic diversity of domesticated animals, cultivated plants and fish species of major socioeconomic importance. Existing sustainability certification schemes could provide baseline information for some ecosystems and sectors. UNEP/CBD/COP/10/9 Page 5 /...

**Target 8: By 2020, pollution, including from excess nutrients, has been brought to levels that are not detrimental to ecosystem function and biodiversity.**

Pollution, including nutrient loading is a major and increasing cause of biodiversity loss and ecosystem dysfunction, particularly in wetland, coastal, marine and dryland areas. Humans have already more than doubled the amount of “reactive nitrogen” in the biosphere, and business-as-usual trends would suggest a further increase of the same magnitude by 2050. The better control of sources of pollution, including efficiency in fertilizer use and the better management of animal wastes, coupled with the use of wetlands as natural water treatment plants where appropriate, can be used to bring nutrient levels below levels that are critical for ecosystem functioning, without curtailing the application of fertilizer in areas where it is necessary to meet soil fertility and food security needs. Similarly, the development and application of national water quality guidelines could help to limit pollution and excess nutrients from entering freshwater and marine ecosystems. Relevant indicators include nitrogen deposition and water quality in freshwater ecosystems. Other possible indicators could be the ecological footprint and related concepts, total nutrient use, nutrient loading in freshwater and marine environments, and the incidence of hypoxic zones and algal blooms. Data which could provide baseline information already exists for several of these indicators, including the global aerial deposition of reactive nitrogen and the incidence of marine dead zones (an example of human-induced ecosystem failure).

**Target 9: By 2020, invasive alien species are identified, prioritized and controlled or eradicated and measures are in place to control pathways for the introduction and establishment of invasive alien species.**

Invasive alien species are a major threat to biodiversity and ecosystem services, and increasing trade and travel means that this threat is likely to increase unless additional action is taken. Pathways for the introduction of invasive alien species can be managed through improved border controls and quarantine, including through better coordination with national and regional bodies responsible for plant and animal health. While well-developed and, globally-applicable indicators are lacking, some basic methodologies do exist which can serve as a starting point for further monitoring or provide baseline information. Process indicators for this target could include the number of countries with national invasive species policies, strategies and action plans and the number of countries which have ratified international agreements and standards related to the prevention and control of invasive alien species. One outcome-oriented indicator is trends in invasive alien species while other possible indicators could include the status of alien species invasion, and the Red List Index for impacts of invasive alien species.

**Target 10: By [2020][2015], to have minimized the multiple pressures on coral reefs, and other vulnerable ecosystems impacted by climate change or ocean acidification, so as to maintain their integrity and functioning.**

Given the ecological inertias related to climate change and ocean acidification, it is important to urgently reduce other pressures on vulnerable ecosystems such as coral reefs so as to give vulnerable ecosystems time to cope with the pressures caused by climate change. This can be accomplished by addressing those pressures which are most amenable to rapid positive changes and would include activities such as reducing pollution and overexploitation and harvesting practices which have negative consequences on ecosystems. Indicators for this target include the extent of biomes ecosystems and habitats (% live coral, and coral bleaching), Marine Trophic Index, the incidence of human-induced ecosystem failure, and the health and well-being of communities who depend directly on local ecosystem goods and services, proportion of products derived from sustainable sources. UNEP/CBD/COP/10/9 Page 6 /...

**Strategic goal C: To improve the status of biodiversity by safeguarding ecosystems, species and genetic diversity**

Whilst longer term actions to reduce the underlying causes of biodiversity loss are taking effect, immediate actions, such as protected areas, species recovery programmes, land-use planning approaches, the restoration of degraded ecosystems and other targeted conservation interventions can help conserve biodiversity and critical ecosystems. These might focus on culturally-valued species and key ecosystem services, particularly those of importance to the poor, as well as on threatened species. For example, carefully sited protected areas could prevent the extinction of threatened species by protecting their habitats, allowing for future recovery.

**Target 11: By 2020, at least [15%][20%] of terrestrial, inland-water and [X%] of coastal and marine areas, especially areas of particular importance for biodiversity and ecosystem services, are conserved through comprehensive, ecologically representative and well-connected systems of effectively managed protected areas and other means, and integrated into the wider land- and seascape.**

Currently, some 13 per cent of terrestrial areas and 5 per cent of coastal areas are protected, while very little of the open oceans are protected. Therefore reaching the proposed target implies a modest increase in terrestrial protected areas globally, with an increased focus on representativity and management effectiveness, together

with major efforts to expand marine protected areas. Protected areas should be integrated into the wider land- and seascape, bearing in mind the importance of complementarity and spatial configuration. In doing so, the ecosystem approach should be applied taking into account ecological connectivity and the concept of ecological networks, including connectivity for migratory species. Protected areas should also be established and managed in close collaboration with, and through participatory and equitable processes that recognize and respect the rights of indigenous and local communities, and vulnerable populations. Other means of protection may also include restrictions on activities that impact on biodiversity, which would allow for the safeguarding of sites in areas beyond national jurisdiction in a manner consistent with the jurisdictional scope of the Convention as contained in Article 4. Relevant indicators to measure progress towards this target are the coverage of sites of biodiversity significance covered by protected areas and the connectivity/fragmentation of ecosystems. Other possible indicators include the overlay of protected areas with ecoregions, and the governance and management effectiveness of protected areas. Good baseline information already exists from sources such as the World Database of Protected Areas the Alliance for Zero Extinction, and the IUCN Red List of Threatened Species and the IUCN World Commission on Protected Areas.

## 12. Appendices

### 12.1. *Terms of Reference for the Aquatic Environment Working Group in 2012*

#### Overall purpose

For all New Zealand fisheries in the New Zealand TS and EEZ as well as other important fisheries in which New Zealand engages:

to assess, based on scientific information, the effects of (and risks posed by) fishing, aquaculture, and enhancement on the aquatic environment, including:

- bycatch and unobserved mortality of protected species (e.g. seabirds and marine mammals), fish, and other marine life, and consequent impacts on populations
- effects of bottom fisheries on benthic biodiversity, species, and habitat
- effects on biodiversity, including genetic diversity
- changes to ecosystem structure and function from fishing, including trophic effects
- effects of aquaculture and fishery enhancement on the environment and on fishing

Where appropriate and feasible, such assessments should explore the implications of the effect, including with respect to government standards, other agreed reference points, or other relevant indicators of population or environmental status. Where possible, projections of future status under alternative management scenarios should be made.

AEWG assesses the effects of fishing or environmental status, and may evaluate the consequences of alternative future management scenarios. AEWG does not make management recommendations or decisions (this responsibility lies with MPI fisheries managers and the Minister responsible for Fisheries).

MPI also convenes a Biodiversity Research Advisory Group (BRAG) which has a similar review function to the AEWG. Projects reviewed by BRAG and AEWG have some commonalities in that they relate to aspects of the marine environment. However, the key focus of projects considered by BRAG is on marine issues related to the functionality of the marine ecosystem and its productivity, whereas projects considered by AEWG are more commonly focused on the direct effects of fishing.

#### Preparatory tasks

1. Prior to the beginning of AEWG meetings each year, MPI fisheries scientists will produce a list of issues for which new assessments or evaluations are likely to become available prior to the next scheduled sustainability round or decision process. AEWG Chairs will determine the final timetables and agendas.
2. The Ministry's research planning processes should identify most information needs well in advance but, if urgent issues arise, MPI-Fisheries or standards managers will alert MPI-Fisheries science managers and the Principal Advisor Fisheries Science, at least 3 months prior to the required AEWG meetings to other cases for which assessments or evaluations are urgently needed.

### **Technical objectives**

3. To review any new research information on fisheries impacts, including risks of impacts, and the relative or absolute sensitivity or susceptibility of potentially affected species, populations, habitats, and systems.
4. To estimate appropriate reference points for determining population, system, or environmental status, noting any draft or published Standards.
5. To conduct environmental assessments or evaluations for selected species, populations, habitats, or systems in order to determine their status relative to appropriate reference points and Standards, where such exist.
6. In addition to determining the status of the species, populations, habitats, and systems relative to reference points, and particularly where the status is unknown, AEWG should explore the potential for using existing data and analyses to draw conclusions about likely future trends in fishing effects or status if current fishing methods, effort, catches, and catch limits are maintained, or if fishers or fisheries managers are considering modifying them in other ways.
7. Where appropriate and practical, to conduct or request projections of likely future status using alternative management actions, based on input from AEWG, fisheries plan advisers and fisheries and standards managers, noting any draft or published Standards.
8. For species or populations deemed to be depleted or endangered, to develop ideas for alternative rebuilding scenarios to levels that are likely to ensure long-term viability based on input from AEWG, fisheries managers, noting any draft or published Standards.
9. For species, populations, habitats, or systems for which new assessments are not conducted in the current year, to review and update any existing Fisheries Assessment Plenary report text in order to determine whether the latest reported status summary is still relevant; else to revise the evaluations based on new data or analyses, or other relevant information.

### **Working Group input to annual Aquatic Environment and Biodiversity Review**

10. To include in contributions to the environment analogue of the Fisheries Assessment Plenary Report (the Aquatic Environment and Biodiversity Review, AEBAAR) summaries of information on selected issues that may relate to species, populations, habitats, or systems that may be affected by fishing. These contributions are analogous to Working Group Reports from the Fisheries Assessment Working Groups.
11. To provide information and scientific advice on management considerations (e.g. area boundaries, by-catch issues, effects of fishing on habitat, other sources of mortality, and input controls such as mesh sizes and minimum legal sizes) that may be relevant for setting sustainability measures.
12. To summarise the assessment methods and results, along with estimates of relevant standards, reference points, or other metrics that may be used as benchmarks or to identify risks to the aquatic environment.
13. It is desirable that full agreement among technical experts is achieved on the text of contributions to the AEBAAR. If full agreement among technical experts cannot be reached, the Chair will determine how this will be depicted in the AEBAAR, will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.

14. To advise the Principal Advisor Fisheries Science, about issues of particular importance that may require review by a plenary meeting or summarising in the AEBAR, and issues that are not believed to warrant such review. The general criterion for determining which issues should be discussed by a wider group or summarised in the AEBAR is that new data or analyses have become available that alter the previous assessment of an issue, particularly assessments of population status or projection results. Such information could include:
- New or revised estimates of environmental reference points, recent or current population status, trend, or projections
  - The development of a major trend in bycatch rates or amount
  - Any new studies or data that extend understanding of population, system, or environmental susceptibility to an effect or its recoverability, fishing patterns, or mitigation measures that have a substantial implications for a population, system, or environment or identify risks associated with fishing activity
  - Consistent performance outside accepted reference points or Standards

**Membership and Protocols for all Science Working Groups (paragraph numbers consistent with those in Terms of Reference for Fisheries Assessment Working Groups)**

**Working Group chairs**

17. The Ministry will select and appoint the Chairs for Working Groups. The Chair will be a MPI fisheries scientist who is an active participant in the Working Group, providing technical input, rather than simply being a facilitator. Working Group Chairs will be responsible for:
- \* ensuring that Working Group participants are aware of the Terms of Reference for the working group, and that the Terms of Reference are adhered to by all participants.
  - \* setting the rules of engagement, facilitating constructive questioning, and focussing on relevant issues.
  - \* ensuring that all peer review processes are conducted in accordance with the Research and Science Information Standard for New Zealand Fisheries<sup>110</sup> (the Research Standard), and that research and science information is reviewed by the Working Group against the *P R I O R* principles for science information quality (page 6) and the criteria for peer review (pages 12-16) in the Standard.
  - \* requesting and documenting the affiliations of participants at each Working Group meeting that have the potential to be, or to be perceived to be, a conflict of interest of relevance to the research under review (refer to page 15 of the Research Standard). Chairs are responsible for managing conflicts of interest, and ensuring that fisheries management implications do not jeopardise the objectivity of the review or result in biased interpretation of results.
  - \* ensuring that the quality of information that is intended or likely to inform fisheries management decisions is ranked in accordance with the information ranking guidelines in the Research Standard (page 21-23), and that resulting information quality ranks are appropriately documented in Working Group reports and, where appropriate, in Status of Stock summary tables.
  - \* striving for consensus while ensuring the transparency and integrity of research analyses, results, conclusions and final reports.
  - \* reporting on Working Group recommendations, conclusions and action items; and ensuring follow-up and communication with the MPI Principal Advisor Fisheries Science, relevant MPI fisheries management staff, and other key stakeholders.

---

<sup>110</sup> Link to the Research Standard: <http://www.fish.govt.nz/en-nz/Publications/Research+and+Science+Information+Standard.htm>

## Working Group members

18. Working Groups will consist of the following participants:
  - \* MPI fisheries science chair – required
  - \* Research providers – required (may be the primary researcher, or a designated substitute capable of presenting and discussing the agenda item)
  - \* Other scientists not conducting analytical assessments to act in a peer review capacity
  - \* Representatives of relevant MPI fisheries management teams
  - \* Any interested party who agrees to the standards of participation below.
19. Working Group participants must commit to:
  - \* participating in the discussion
  - \* resolving issues
  - \* following up on agreements and tasks
  - \* maintaining confidentiality of Working Group discussions and deliberations (unless otherwise agreed in advance, and subject to the constraints of the Official Information Act)
  - \* adopting a constructive approach
  - \* avoiding repetition of earlier deliberations, particularly where agreement has already been reached
  - \* facilitating an atmosphere of honesty, openness and trust
  - \* respecting the role of the Chair
  - \* listening to the views of others, and treating them with respect
20. Participants in Working Group meetings will be expected to declare their sector affiliations and contractual relationships to the research under review, and to declare any substantial conflicts of interest related to any particular issue or scientific conclusion.
21. Working Group participants are expected to adhere to the requirements of independence, impartiality and objectivity listed under the Peer Review Criteria in the Research Standard (pages 12-16). It is understood that Working Group participants will often be representing particular sectors and interest groups, and will be expressing the views of those groups. However, when reviewing the quality of science information, representatives are expected to step aside from their sector affiliations, and to ensure that individual and sector views do not result in bias in the science information and conclusions.

### *Information Quality Ranking:*

22. Science Working Groups are required to rank the quality of research and science information that is intended or likely to inform fisheries management decisions, in accordance with the science information quality ranking guidelines in the Research Standard (pages 21-23). This information quality ranking must be documented in Working Group reports and, where appropriate, in Status of Stock summary tables.
  - \* Working Groups are not required to rank all research projects and analyses, but key pieces of information that are expected or likely to inform fisheries management decisions should receive a quality ranking.
  - \* Explanations substantiating the quality rankings must be included in Working Group reports. In particular, the quality shortcomings and concerns for moderate/mixed and low quality information must be documented.
  - \* The Chair, working with participants, will determine which pieces of information require a quality ranking. Not all information resulting from a particular research project would be expected to achieve the same quality rank, and different quality ranks may be assigned to

different components, conclusions or pieces of information resulting from a particular piece of research.

*Working Group papers:*

23. Working group papers will be posted on the MPI-Fisheries website prior to meetings if they are available. As a general guide, Powerpoint presentations and draft or discussion papers should be available at least 2 working days before a meeting, and near-final papers should be available at least 5 working days before a meeting if the Working Group is expected to agree to the paper. However, it is also likely that many papers will be tabled during the meeting due to time constraints. If a paper is not available for sufficient time before the meeting, the Chair may provide for additional time for written comments from Working Group members.
24. Working Group papers are “works in progress” whose role is to facilitate the discussion of the Working Groups. They often contain preliminary results that are receiving peer review for the first time and, as such, may contain errors or preliminary analyses that will be superseded by more rigorous work. **For these reasons, no-one may release the papers or any information contained in these papers to external parties. In general, Working Group papers should never be cited.** Exceptions may be made in rare instances by obtaining permission in writing from the Principal Advisor Fisheries Science, and the authors of the paper.
25. Participants who use Working Group papers inappropriately, or who do not adhere to the standards of participation, may be requested by the Chair to leave a particular meeting or, in more serious instances, to refrain from attending one or more future meetings.
26. Meetings will take place as required, generally January-April and July-November for FAWGs and throughout the year for other working groups (AEWG, BRAG, Marine Amateur Fisheries and Antarctic Working Groups).
27. A quorum will be reached when the Chair, the designated presenter, and three or more other technical experts are present. In the absence of a quorum, the Chair may decide to proceed as a sub-group, with outcomes being taken forward to the next meeting at which a quorum is formed.
28. The Chair is responsible for deciding, with input from the entire Working Group, but focussing primarily on the technical discussion and the views of technical expert members:
  - \* The quality and acceptability of the information and analyses under review
  - \* The way forward to address any deficiencies
  - \* The need for any additional analyses
  - \* Contents of Working Group reports
  - \* Choice of base case models and sensitivity analyses to be presented
  - \* The status of the stocks, or the status/performance in relation to any relevant environmental standards or targets
29. The Chair is responsible for facilitating a consultative and collaborative discussion.
30. Working Group meetings will be run formally, with agendas pre-circulated, and formal records kept of recommendations, conclusions and action items.
31. A record of recommendations, conclusions and action items will be posted on the MPI-Fisheries website after each meeting has taken place.

32. Data upon which analyses presented to the Working Groups are based must be provided to MPI in the appropriate format and level of detail in a timely manner (i.e. the data must be available and accessible to MPI; however, data confidentiality concerns mean that such data are not necessarily available to Working Group members)
33. The outcome of each Working Group round will be evaluated, with a view to identifying opportunities to improve the Working Group process. The Terms of Reference may be updated as part of this review.
34. MPI fisheries scientists and science officers will provide administrative support to the Working Groups.

### **Record-keeping**

35. The overall responsibility for record-keeping rests with the Chair of the Working Group, and includes:
  - \* keeping notes on recommendations, conclusions and follow-up actions for all Working Group meetings, and to ensure that these are available to all members of the Working Group and the Principal Advisor Fisheries Science in a timely manner. If full agreement on the recommendations or conclusions cannot readily be reached amongst technical experts, then the Chair will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.
  - \* compiling a list of generic assessment issues and specific research needs for each Fishstock or species or environmental issue under the purview of the Working Group, for use in subsequent research planning processes.

## **12.2. AEWG Membership 2012**

**Convenors:** Martin Cryer (protected species) and Rich Ford (other issues)

**Members:** Blake Abernethy, Ed Abraham, Owen Anderson, William Arlidge, Chris Baigent, Karen Baird, Suze Baird, Barry Baker, Michael Batson, Michelle Beritzhoff, Jenny Black, Tiffany Bock, Laura Boren, Yoland Bosiger, Paul Breen, Stephen Brouwer, Martin Cawthorn, Simon Childerhouse, Louise Chilvers, Tom Clarke, Malcolm Clark, Deanna Clement, George Clement, Owen Cox, Rohan Currey, Igor Debski, Ian Doonan, Alastair Dunn, Charles Edwards, Ursula Ellenburg, Jack Fenaughty, Chris Francis, Malcolm Francis, Kevin Hackwell, Judi Hewitt, Rosie Hurst, Aaron Irving, Catherine Jones, Dan Kluza, Craig Lawson, Mary Livingston, Dave Lundquist, Greg Lydon, Warrick Lyon, Pamela Mace, Darryl Mackenzie, Rob Mattlin, Tania McPherson, Sarah Meadows, David Middleton, Laura Mitchell, Sophie Mormede, Mark Morrison, Richard O'Driscoll, Tracey Osborne, Milena Palka, Johanna Pierre, Irene Pohl, Kris Ramm, Vicky Reeve, Pat Reid, Yvan Richard, Jim Roberts, Ashley Rowden, Carol Scott, Ben Sharp, Liz Slooten, Paul Starr, Kevin Stokes, Katrina Subedar, John Taunton-Clarke, Alex Thompson, David Thompson, Finlay Thompson, Geoff Tingley, Ian Tuck, Dee Wallace, Barry Weeber, Richard Wells, Francene Wineti, Ray Wood, Bob Zuur.

### **12.3. Terms of Reference for the Biodiversity Research Advisory Group (BRAG) 2012**

#### **Overall purpose**

Since 2000, the objectives of the Biodiversity Research Programme have been drawn directly from MFish commitments to Theme 3 of the New Zealand Biodiversity Strategy. Within this framework, the Biodiversity Medium Term Research Plan has been adapted over time as new issues emerge, to build on synergies with other research programmes and work where biodiversity is under greatest threat from fishing or other anthropogenic activity.

Within the constraints of the overall purpose of the Programme,

“To improve our understanding of New Zealand marine ecosystems in terms of species diversity, marine habitat diversity, and the processes that lead to healthy ecosystem functioning, and the role that biodiversity has for such key processes<sup>111</sup>;

and the NZBS definition of biodiversity (the variability among living organisms from all sources including inter alia, terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are a part; this includes diversity within species, between species and of ecosystem) the science currently commissioned broadly aims to:

- Describe and characterise the distribution and abundance of fauna and flora, as expressed through measures of biodiversity, and improving understanding about the drivers of the spatial and temporal patterns observed;
- determine the functional role of different organisms or groups of organisms in marine ecosystems, and assess the role of marine biodiversity in mitigating the impacts of anthropogenic disturbance on healthy ecosystem functioning;
- identify which components of biodiversity must be protected to ensure the sustainability of a healthy marine ecosystem as well as to meet societal values on biodiversity.

MPI also convenes an Aquatic Environment Working Group (AEWG) which has a similar review function to the BRAG. Projects reviewed by BRAG and AEWG have some commonalities in that they relate to aspects of the marine environment. However, the key focus of projects considered by BRAG is on marine issues related to the functionality of the marine ecosystem and its productivity, whereas projects considered by AEWG are more commonly focused on the direct effects of fishing.

BRAG may identify natural resource management issues that extend beyond fisheries management and make recommendations on priority areas of research that will inform MPI or other government departments of emerging science results that require the attention of managers, policymakers and decision-makers in the marine sector. BRAG does not make management recommendations or decisions (this responsibility lies with the MPI Fisheries Management Group and the Minister of Primary Industry).

#### **Preparatory tasks**

1. Prior to the beginning of BRAG meetings each year, MPI fisheries scientists will produce a list of issues for which new research projects are likely to be required in the forthcoming financial year. The BRAG Chair will determine the final timetables and agendas.

---

<sup>111</sup> See MFish Biodiversity Research Programme 2010: Part 1. Context and Purpose

2. The Ministry's research planning processes should identify most information needs well in advance but, if urgent issues arise, MPI fisheries managers will alert the Aquatic Environment and Biodiversity Science Manager and the Principal Advisor Fisheries Science at least three months prior to the required meetings where possible.

### **BRAG Technical objectives**

3. To review, discuss and convey views on the results of marine biodiversity research projects contracted by Ministry for Primary Industries MPI (formerly Ministry of Fisheries).

It is the responsibility of the BRAG to review, discuss, and convey views on the results of marine biodiversity research projects contracted by MPI and the former Ministry of Fisheries. The review process is an evaluation of how existing research results can be built upon to address emerging research issues and needs. It is essentially an evaluation of "what we already know" and how this can be used to obtain "what we need to know". This information should be used by the BRAG to identify gaps in our knowledge and for developing research plans to address these gaps.

4. Discuss, evaluate, make recommendations and convey views on a 3 to 5 year Medium Term Research Plan.

It is the responsibility of BRAG participants to discuss, evaluate, make recommendations and convey views on a 3 to 5 year Medium Term Research Plan for its particular research area as required. Individual related projects on a species or fishery or research topic need to be integrated into Medium Term Research Plans. The Medium Term Research Plans should encompass research needs and directions for at least the next 3 to 5 years.

The Biodiversity Medium Term Research Plan is aligned to relevant strategic and policy directions such as the "MPI Statement of Intent" and any Strategic Research Plan (Fisheries 2030, Deepwater 10 year research plan) and fisheries plans developed for the appropriate species/fishery or research area, including biodiversity.

The recommendations on project proposals for the next financial year will be submitted via the Chair of BRAG to the Principal Science Advisor Fisheries (MPI).

5. The Biodiversity Research Programme includes research in New Zealand's TS, EEZ, Extended Continental Shelf, the South Pacific Region and the Ross Sea region and has seven scientific work streams as follows:
  - To develop ecosystem-scale understanding of biodiversity in the New Zealand marine environment
  - To classify and characterise the biodiversity, including the description and documentation of biota, associated with nearshore and offshore marine habitats in New Zealand
  - To investigate the role of biodiversity in the functional ecology of nearshore and offshore marine communities.
  - To assess developments in all aspects of biodiversity, including genetic marine biodiversity and identify key topics for research
  - To determine the effects of climate change and increased ocean acidification on marine biodiversity, as well as effects of incursions of non-indigenous species, and other threats and impacts.

- To develop appropriate diversity metrics and other indicators of biodiversity that can be used to monitor change
- To identify threats and impacts to biodiversity and ecosystem functioning beyond natural environmental variation

#### **BRAG input to MPI “Aquatic Environment and Biodiversity Annual Review”**

6. To contribute to and summarise progress on biodiversity research in the Aquatic Environment and Biodiversity Annual Review. This contribution is analogous to Working Group Reports from the Fishery Assessment Working Groups.
7. To summarise the assessment methods and results, along with estimates of relevant standards, references points, or other metrics that may be relevant to biodiversity objectives by MPI, the Biodiversity Strategy and international obligations.
8. It is desirable that full agreement among technical experts is achieved on the text of these contributions. If full agreement among technical experts cannot be reached, the Chair will determine how this will be depicted in the Aquatic Environment and Biodiversity Annual Review, will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.
9. To advise the Principal Science Advisor Fisheries (MPI), about issues of particular importance that may require review by a plenary meeting or summarising in the Aquatic Environment and Biodiversity Annual Review. The general criterion for determining which issues should be discussed by a wider group include:
  - Emerging issues, recent or current biodiversity status assessments, trends, or projections
  - The development of a major trend in the marine environment that will impact on marine productivity or ecosystem resilience to stressors
  - Any new studies or data that impact on international obligations

#### **Membership and Protocols for all Science Working Groups (NOTE: paragraph numbers consistent with those in Terms of Reference for Fisheries Assessment Working Groups)**

##### **Working Group chairs**

17. The Ministry will select and appoint the Chairs for Working Groups. The Chair will be a MPI fisheries scientist who is an active participant in the Working Group, providing technical input, rather than simply being a facilitator. Working Group Chairs will be responsible for:
  - \* ensuring that Working Group participants are aware of the Terms of Reference for the working group, and that the Terms of Reference are adhered to by all participants.
  - \* setting the rules of engagement, facilitating constructive questioning, and focussing on relevant issues.
  - \* ensuring that all peer review processes are conducted in accordance with the Research and Science Information Standard for New Zealand Fisheries<sup>112</sup> (the Research Standard), and that

---

<sup>112</sup> Link to the Research Standard: <http://www.fish.govt.nz/en-nz/Publications/Research+and+Science+Information+Standard.htm>

research and science information is reviewed by the Working Group against the *P R I O R* principles for science information quality (page 6) and the criteria for peer review (pages 12-16) in the Standard.

- \* requesting and documenting the affiliations of participants at each Working Group meeting that have the potential to be, or to be perceived to be, a conflict of interest of relevance to the research under review (refer to page 15 of the Research Standard). Chairs are responsible for managing conflicts of interest, and ensuring that fisheries management implications do not jeopardise the objectivity of the review or result in biased interpretation of results.
- \* ensuring that the quality of information that is intended or likely to inform fisheries management decisions is ranked in accordance with the information ranking guidelines in the Research Standard (page 21-23), and that resulting information quality ranks are appropriately documented in Working Group reports and, where appropriate, in Status of Stock summary tables.
- \* striving for consensus while ensuring the transparency and integrity of research analyses, results, conclusions and final reports.
- \* reporting on Working Group recommendations, conclusions and action items; and ensuring follow-up and communication with the MPI Principal Advisor Fisheries Science, relevant MPI fisheries management staff, and other key stakeholders.

### **Working Group members**

18. Working Groups will consist of the following participants:
    - \* MPI fisheries science chair – required
    - \* Research providers – required (may be the primary researcher, or a designated substitute capable of presenting and discussing the agenda item)
    - \* Other scientists not conducting analytical assessments to act in a peer review capacity
    - \* Representatives of relevant MPI fisheries management teams
    - \* Any interested party who agrees to the standards of participation below.
  19. Working Group participants must commit to:
    - \* participating in the discussion
    - \* resolving issues
    - \* following up on agreements and tasks
    - \* maintaining confidentiality of Working Group discussions and deliberations (unless otherwise agreed in advance, and subject to the constraints of the Official Information Act)
    - \* adopting a constructive approach
    - \* avoiding repetition of earlier deliberations, particularly where agreement has already been reached
    - \* facilitating an atmosphere of honesty, openness and trust
    - \* respecting the role of the Chair
    - \* listening to the views of others, and treating them with respect
  20. Participants in Working Group meetings will be expected to declare their sector affiliations and contractual relationships to the research under review, and to declare any substantial conflicts of interest related to any particular issue or scientific conclusion.
  21. Working Group participants are expected to adhere to the requirements of independence, impartiality and objectivity listed under the Peer Review Criteria in the Research Standard (pages 12-16). It is understood that Working Group participants will often be representing particular sectors and interest groups, and will be expressing the views of those groups. However, when reviewing the quality of science information, representatives are expected to
-

step aside from their sector affiliations, and to ensure that individual and sector views do not result in bias in the science information and conclusions.

*Information Quality Ranking:*

22. Science Working Groups are required to rank the quality of research and science information that is intended or likely to inform fisheries management decisions, in accordance with the science information quality ranking guidelines in the Research Standard (pages 21-23). This information quality ranking must be documented in Working Group reports and, where appropriate, in Status of Stock summary tables.
  - \* Working Groups are not required to rank all research projects and analyses, but key pieces of information that are expected or likely to inform fisheries management decisions should receive a quality ranking.
  - \* Explanations substantiating the quality rankings must be included in Working Group reports. In particular, the quality shortcomings and concerns for moderate/mixed and low quality information must be documented.
  - \* The Chair, working with participants, will determine which pieces of information require a quality ranking. Not all information resulting from a particular research project would be expected to achieve the same quality rank, and different quality ranks may be assigned to different components, conclusions or pieces of information resulting from a particular piece of research.

*Working Group papers:*

23. Working group papers will be posted on the MPI-Fisheries website prior to meetings if they are available. As a general guide, Powerpoint presentations and draft or discussion papers should be available at least 2 working days before a meeting, and near-final papers should be available at least 5 working days before a meeting if the Working Group is expected to agree to the paper. However, it is also likely that many papers will be tabled during the meeting due to time constraints. If a paper is not available for sufficient time before the meeting, the Chair may provide for additional time for written comments from Working Group members.
24. Working Group papers are “works in progress” whose role is to facilitate the discussion of the Working Groups. They often contain preliminary results that are receiving peer review for the first time and, as such, may contain errors or preliminary analyses that will be superseded by more rigorous work. **For these reasons, no-one may release the papers or any information contained in these papers to external parties. In general, Working Group papers should never be cited.** Exceptions may be made in rare instances by obtaining permission in writing from the Principal Advisor Fisheries Science, and the authors of the paper.
25. Participants who use Working Group papers inappropriately, or who do not adhere to the standards of participation, may be requested by the Chair to leave a particular meeting or, in more serious instances, to refrain from attending one or more future meetings.
26. Meetings will take place as required, generally January-April and July-November for FAWGs and throughout the year for other working groups (AEWG, BRAG, Marine Amateur Fisheries and Antarctic Working Groups).
27. A quorum will be reached when the Chair, the designated presenter, and three or more other technical experts are present. In the absence of a quorum, the Chair may decide to proceed as

a sub-group, with outcomes being taken forward to the next meeting at which a quorum is formed.

28. The Chair is responsible for deciding, with input from the entire Working Group, but focussing primarily on the technical discussion and the views of technical expert members:
  - \* The quality and acceptability of the information and analyses under review
  - \* The way forward to address any deficiencies
  - \* The need for any additional analyses
  - \* Contents of Working Group reports
  - \* Choice of base case models and sensitivity analyses to be presented
  - \* The status of the stocks, or the status/performance in relation to any relevant environmental standards or targets
29. The Chair is responsible for facilitating a consultative and collaborative discussion.
30. Working Group meetings will be run formally, with agendas pre-circulated, and formal records kept of recommendations, conclusions and action items.
31. A record of recommendations, conclusions and action items will be posted on the MPI-Fisheries website after each meeting has taken place.
32. Data upon which analyses presented to the Working Groups are based must be provided to MPI in the appropriate format and level of detail in a timely manner (i.e. the data must be available and accessible to MPI; however, data confidentiality concerns mean that such data are not necessarily available to Working Group members)
33. The outcome of each Working Group round will be evaluated, with a view to identifying opportunities to improve the Working Group process. The Terms of Reference may be updated as part of this review.
34. MPI fisheries scientists and science officers will provide administrative support to the Working Groups.

### **Record-keeping**

35. The overall responsibility for record-keeping rests with the Chair of the Working Group, and includes:
  - \* keeping notes on recommendations, conclusions and follow-up actions for all Working Group meetings, and to ensure that these are available to all members of the Working Group and the Principal Advisor Fisheries Science in a timely manner. If full agreement on the recommendations or conclusions cannot readily be reached amongst technical experts, then the Chair will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.
  - \* compiling a list of generic assessment issues and specific research needs for each Fishstock or species or environmental issue under the purview of the Working Group, for use in subsequent research planning processes.

## **12.4. BRAG attendance 2011-2012**

**Convenor:** Mary Livingston (MPI chair),

**Members:** Malcolm Clark, Mark Morrison, Wendy Nelson, Cliff Law, Di Tracey, Dennis Gordon, Anne-Nina Lorz, Stuart Hanchet, Richard O'Driscoll, Jonathon Gardner, Simon Thrush, David Bowden, Matt Pinkerton, Els Maas, Ashley Rowden, Carolyn Lundquist, Judi Hewitt, Drew Lohrer, Alison MacDiarmid, Julie Hall, Vonda Cummings, Kate Neill, Tracy Farr, Di Tracey, Barb Hayden (all NIWA), Richard Wells, Greg Lydon (SeaFIC), Rich Ford (MPI), Shane Lavery, Mark Costello (Auckland University)

## **12.5. Generic Terms of Reference for Research Advisory Groups (Sept 2010)**

### **Overall purpose**

1. The purpose of the Research Advisory Groups (RAGs) is to develop research proposals to meet management information needs and support standards development.

### **Context**

2. To assist RAG members with their work this section outlines the wider process that RAGs will operate within.

#### ***Fisheries Plans will guide the management of fisheries***

3. From 1 July 2011 the Ministry of Fisheries (MFish) will be using Fisheries Plans in the following five areas to guide the management of fisheries:
  - Deepwater
  - Highly Migratory Species
  - Inshore – Finfish
  - Inshore – Freshwater
  - Inshore – Shellfish
4. In each of those five areas there will be:
  - A Fisheries Plan that sets out management objectives over a 5 year period.
  - An Annual Operational Plan that sets out what will be done in a financial year to help meet those objectives, including in the areas of science research, compliance and observer coverage (i.e., the Annual Operational Plan will be where priorities are set each year). Note that external stakeholders will have an opportunity to provide comment on prioritisation through draft Annual Operational Plans.
  - An Annual Review Report that will assess progress made against the management objectives, and help identify gaps to be considered in setting the next set of priorities.

#### ***RAGs will largely be aligned to the Fisheries Plan areas***

5. There will be a RAG for each of the five Fisheries Plan areas above.

6. In addition there will be a RAG for Aquatic Environment (Standards), for research needed to support standards development, and another for Antarctic research. (Note that biodiversity research is dealt with through a separate process that has more of a cross-agency focus.)

***RAGs will develop research proposals to be considered as part of a subsequent prioritisation process***

7. As part of the process for developing the Annual Operational Plans, the identification and prioritisation of science research will broadly occur as follows:
  - i. MFish fisheries managers will identify the fisheries management objectives and information needs that they want the relevant RAG to consider. This will be done in conjunction with MFish scientists, and will draw on the following:
    - The relevant Annual Review Report discussed above
    - Existing research plans
    - Science Assessment Working Groups' feedback arising from research that has been evaluated previously
    - Ad-hoc issues as they arise
    - Initial indications of the available budget
  - ii. The RAGs will then develop proposals for scientific research to meet those management and information needs.
  - iii. MFish fisheries managers will then run a process for prioritising the research proposals that have been developed and updating multi-year research plans, in conjunction with MFish scientists. This will be part of the wider process for developing Annual Operational Plans.
8. In the Aquatic Environment (Standards) and Antarctic areas a similar process will be followed to that above, involving relevant MFish managers.
9. In practice, these processes are likely to iterate between the above steps, e.g., when prioritising research proposals fisheries managers may identify additional questions that they want a RAG to consider.
10. RAGs will only be convened when necessary. If, for example, all of the research for the coming year under review has previously been approved as part of a multi-year funding package for an area, and no additional management needs have emerged, the relevant RAG will not be convened.
11. During 2010-11 RAGs will be used, as required, in all areas except Inshore, given that the three Inshore Fisheries Plans are still being developed through the year. For the Inshore areas a transitional process will be used, with RAGs commencing during 2011-12.

## **Research proposals**

12. RAGs will provide recommendations to fisheries managers on research to meet management needs. This section provides more detail on the research proposals that the RAGs will produce.
13. The RAGs will produce an initial set of project proposals to meet the management and information needs provided to the RAG, for consideration in the subsequent prioritisation process.
14. The proposals may be in the form of multi-year projects where appropriate.

15. While the prioritisation of research is outside the scope of the work of the RAGs, the proposals will include information on potential cost and feasibility to guide decisions on prioritisation. Cost estimates should be specified as ranges so as to not unduly influence subsequent research provider costings.
16. Where the RAG identifies more than one desirable option for scientific research to meet management and information needs, the RAG's proposals will cover those options, their relative pros and cons, their respective potential costs, and the RAG's recommendation as to the preferred option.
17. Once prioritisation decisions have been made on the initial set of research proposals, the RAG may be asked to produce more fully developed project proposals for inclusion in the relevant Annual Operational Plan, and for the purposes of cost recovery consultation and tendering.

## **Membership**

18. Membership of RAGs is expertise-based.
19. Membership will be by invitation from MFish only.
20. A RAG will consist of a core group of one MFish scientist and one manager from the relevant Fisheries Plan or Standards team, with the option to "call in" relevant technical expertise (internal and/or external) as needed.
21. External participants will be paid for their time. This will include preparing for and attending RAG meetings, and any time spent writing proposals.

## **Protocols**

22. All RAG members will commit to:
  - participating in the discussion in an objective and unbiased manner;
  - resolving issues;
  - following up on agreements and tasks;
  - adopting a constructive approach;
  - facilitating an atmosphere of honesty, openness and trust;
  - having respect for the role of the Chair; and
  - listening to the views of others, and treating them with respect.
23. RAG meetings will be run formally with agendas pre-circulated and formal records kept of recommendations, conclusions and action items.
24. Participants who do not adhere to the standards of participation may be requested by the Chair to leave a particular meeting or, in more serious instances, will be excluded from the RAG.

## *Chairpersons*

25. The Chair of each RAG will be a MFish scientist with appropriate expertise.
26. The Chair commits to undertaking the following roles:
  - The Chair is an active participant in RAGs, who also provides technical input, rather than simply being a facilitator.
  - The Chair is responsible for: setting the rules of engagement; promoting full participation by all members; facilitating constructive questioning; focussing on relevant issues;

reporting on RAG recommendations, conclusions and action items, and ensuring follow-up; and communicating with relevant MFish managers.

27. The Chair is responsible for facilitating consultative and collaborative discussions.

*Decision-making*

28. The Chair is responsible for working towards an agreed view of the RAG members on their recommendations to the fisheries manager, but where that proves not to be possible then the Chair is responsible for determining the final recommendation. Minority views should be clearly represented in proposals in those cases.
29. A record of recommendations, conclusions and action items will be circulated by e-mail after each meeting by the Chair.
30. Each RAG round will be evaluated by MFish, with a view to identifying opportunities to improve the process. The Terms of Reference may be updated as part of this review.

*Non-disclosure agreements*

31. Participants may be asked to sign a Non-Disclosure Agreement relating to documents that disclose cost details.

*Conflicts of Interest*

32. New Zealand is a small country and fisheries research is a relatively limited market, even internationally. People with the necessary skills and knowledge to participate in this advisory process may also have close working relationships with industry, research providers and other stakeholders. This will apply to nearly all external members of a RAG.
33. Participants will be asked to declare any “actual, perceived or likely conflicts of interest” before involvement in a RAG is approved, and any new conflicts that arise during the process should be declared immediately. These will be clearly documented by the Chair.
34. Management of conflicts of interest will be determined by the Chair in consultation with Fisheries Managers, and approved by the Deputy Chief Executive, Fisheries Management prior to meetings commencing.

*Frequency of Meetings*

35. Relevant MFish managers, in consultation with the Chair of the RAG, will decide on the frequency and timing of RAG meetings.

*Documents and record-keeping*

36. Unless signalled by the Chair, all RAG documents (papers, agendas, formal records of recommendations, conclusions and action items) will be available to all interested parties through the Ministry of fisheries website ([www.fish.govt.nz](http://www.fish.govt.nz)), except where confidentiality is required for reasons of commercial sensitivity (e.g. cost estimates).
37. RAG documents will be distributed securely.
38. Participants who use RAG papers inappropriately may not be invited to subsequent RAG meetings.

39. The overall responsibility for record-keeping rests with the Chair and includes:

- Records of recommendations, conclusions and follow-up actions for all RAG meetings and to ensure that these are available in a timely manner.
- If full agreement on the recommendations or conclusions cannot readily be reached amongst technical experts, then the Chair will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.

## **12.6. Fisheries 2030**

**Use outcome** – Fisheries resources are used in a manner that provides the greatest overall economic, social, and cultural benefit. This means having:

- An internationally competitive and profitable seafood industry that makes a significant contribution to our economy
- High-quality amateur fisheries that contribute to the social, cultural, and economic well-being of all New Zealanders
- Thriving customary fisheries, managed in accordance with kaitiakitanga, supporting the cultural well-being of iwi and hapū
- Healthy fisheries resources in their aquatic environment that reflect and provide for intrinsic and amenity value.

**Governance conditions** – Fundamental to achieving our goal is the recognition that our approach must be based on sound governance. This means having arrangements that lead to:

- The Treaty partnership being realised through the Crown and Māori clearly defining their respective rights and responsibilities in terms of governance and management of fisheries resources
- The public having confidence and trust in the effectiveness and integrity of the fisheries and aquaculture management regimes
- All stakeholders having rights and responsibilities related to the use and management of fisheries resources that are understood and for which people can be held individually and collectively accountable
- Having an enabling framework that allows stakeholders to create optimal economic, social, and cultural value from their rights and interests
- An accountable, responsive, dynamic, and transparent system of management.

Fisheries 2030 draws on a number of values and principles. These seek to outline the behaviour and approach that should be used to undertake the actions, make decisions, and achieve the goal for New Zealand fisheries.

### **Values**

- **Tikanga:** the Māori way of doing things; correct procedure, custom, habit, lore, method, manner, rule, way, code, meaning, reason, plan, practice, convention. It is derived from the word tika meaning ‘right’ or ‘correct’.
- **Kaitiakitanga:** The root word in kaitiakitanga is tiaki, which includes aspects of guardianship, care, and wise management. Kaitiakitanga is the broad notion applied in different situations.
- **Kotahitanga:** Collective action and unity.
- **Manaakitanga:** Manaakitanga implies a duty to care for others, in the knowledge that at some time others will care for you. This can also be translated in modern Treaty terms as “create no further grievances in the settlement of current claims”.
- **Integrity:** Be honest and straightforward in our dealings with one another. If we agree to do something we will carry it out.

- Respect: Treat each other with courtesy. We will respect each other's right to have different values and hold different opinions.
- Constructive relationship: Strive to build and maintain constructive ways of working with each other, which can endure.
- Achieving results: Focus on producing a solution rather than just discussing the problem.

### **Principles**

- Ecosystem-based approach: We apply an ecosystem-based approach to fisheries management decision-making.
- Conserve biodiversity: Use should not compromise the existence of the full range of genetic diversity within and between species.
- Environmental bottom lines: Biological standards define the limits of extraction and impact on the aquatic environment.
- Precautionary approach: Particular care will be taken to ensure environmental sustainability where information is uncertain, unreliable, or inadequate.
- Address externalities: Those accessing resources and space should address the impacts their activities have on the environment and other users.
- Meet Settlement obligations: Act in ways that are consistent with the Treaty of Waitangi principles and deliver settlement obligations.
- Responsible international citizen: Manage in the context of international rights, obligations, and our strategic interests.
- Inter-generational equity: Current use is achieved in a manner that does not unduly compromise the opportunities for future generations.
- Best available information: Decisions need to be based on the best available and credible biological, economic, social, and cultural information from a range of sources.
- Respect rights and interests: Policies should be formulated and implemented to respect established rights and interests.
- Effective management and services: Use least-cost policy tools to achieve objectives where intervention is necessary and ensure services are delivered efficiently.
- Recover management costs for the reasonable expenses of efficiently provided management and services, from those who benefit from use, and those who cause the risk or adverse effect.
- Dynamic efficiency: Frameworks should be established to allow resources to be allocated to those who value them most.

Fisheries 2030 includes a “plan of action” for the five years from 2009, including: improving the management framework; supporting aquaculture and international objectives; ensuring sustainability of fish stocks; improving fisheries information; building sector leadership and capacity; meeting obligations to Māori; and enabling collective management responsibility. The key components guiding this document are ensuring sustainability of fish stocks and improving fisheries information:

### **Ensuring sustainability of fish stocks**


- Setting and implementing fisheries harvest strategy standards
- Setting and monitoring environmental standards, including for threatened and protected species and seabed impacts
- Enhancing the framework for fisheries management planning, including the use of decision rules to adjust harvest levels over time

### **Improving fisheries information**

- Determining best options for information collection on catch from amateur fisheries, including the implementation of charter boat reporting
- Improving our knowledge of fish stocks and the environmental impacts of fishing through long-term research plans
- Gaining access to increased research and development funding

## 12.7. OUR STRATEGY 2030: Growing and protecting New Zealand

Also available at: <http://www.mpi.govt.nz/Portals/0/Documents/about-maf/strategy.pdf>



## 12.8. Other strategic policy documents

### 12.8.1. Biodiversity Strategy

New Zealand's Biodiversity Strategy was launched in 2000 in response to the decline of New Zealand's indigenous biodiversity — described in the State of New Zealand's Environment report as our “most pervasive environmental issue”. It can be found on the government's biodiversity website at:

(<http://www.biodiversity.govt.nz/picture/doing/nzbs/contents.html>)

The Strategy also reflects New Zealand's commitment, through ratification of the international Convention on Biological Diversity, to help stem the loss of biodiversity worldwide. Strategic Priority 7 of the strategy was “*To manage the marine environment to sustain biodiversity*”. Fishing practices, the effects of activities on land, and biosecurity threats are identified as constituting the areas of greatest risk to marine biodiversity. Pertinent objectives and summarised actions from the strategy are as follows:

**Objective 3.1: Improving our knowledge of coastal and marine ecosystems** (Substantially increase our knowledge of coastal and marine ecosystems and the effects of human activities on them, especially assessing the importance of, and threats facing, marine biodiversity, and establishing environmental monitoring capabilities to assess the effectiveness of measures to avoid, remedy or mitigate impacts on marine biodiversity).

**Objective 3.4: Sustainable marine resource use practices** (Protect biodiversity in coastal and marine waters from the adverse effects of fishing and other coastal and marine resource uses, especially maintaining harvested species at sustainable levels, integrating marine biodiversity protection into an ecosystem approach, applying a precautionary approach, identifying marine species and habitats most sensitive to disturbance, and integrating environmental impact assessments into fisheries management decision making.)

**Objective 3.6: Protecting marine habitats and ecosystems** (Protect a full range of natural marine habitats and ecosystems to effectively conserve marine biodiversity, using a range of appropriate mechanisms, including legal protection, especially establishing a network of areas that protect marine biodiversity.)

**Objective 3.7: Threatened marine and coastal species management** (Protect and enhance populations of marine and coastal species threatened with extinction, and prevent additional species and ecological communities from becoming threatened.)

In addition to its annual reviews (<http://www.biodiversity.govt.nz/news/publications/index.html>), the Biodiversity Strategy was reviewed by Green and Clarkson at the end of its 5-year term. This review was published in 2006 (<http://www.biodiversity.govt.nz/pdfs/nzbs-5-year-review-synthesis-report.pdf>). Most relevant to this synopsis were their findings on Objective 3.4 (Sustainable marine resource use) where they cited “Moderate progress”. “*The policy move towards adopting a more ecosystem approach to fisheries management should be encouraged and strengthened. We acknowledge, however, the difficulties associated with obtaining the necessary information to make this approach effective. There are links to Objective 3.1 and the need for a more coordinated approach to identifying priority areas for marine research.*”

### 12.8.2. Biosecurity Strategy

In its 2003 Biosecurity Strategy, The Ministry of Agriculture and Forestry's Biosecurity NZ defined biosecurity as "*the exclusion, eradication or effective management of risks posed by pests and diseases to the economy, environment and human health*". New Zealand is highly dependent on effective biosecurity measures because our indigenous flora, fauna, biodiversity, and, consequently, our primary production industries, including fisheries are uniquely at risk from invasive species. Information can be found on the Biosecurity New Zealand website at: <http://www.biosecurity.govt.nz/biosec/sys/strategy/biostrategy/biostrategynz>

A complementary Biosecurity Science Strategy for New Zealand was developed in 2007 to address the science expectations of the Biosecurity Strategy. The science strategy identified the need to:

- prioritise science needs;
- minimise biosecurity risks at the earliest stage possible by increasing focus on research that is strategic and proactive;
- improve planning, integration and communication in the delivery of science;
- ensure research outputs can be used effectively to improve biosecurity operations and decision making.

### 12.8.3. Marine Protected Areas Policy

The Marine Protected Areas (MPA) Policy and Implementation Plan was released for consultation in December 2005 jointly by the Ministry of Fisheries and Department of Conservation. It confirmed Government's commitment to ensuring that New Zealand's marine biodiversity was protected, and established MPA Policy as a key component of that commitment. The MPA Policy objective is to protect marine biodiversity by establishing a network of Marine Protected Areas that is comprehensive and representative of New Zealand's marine habitats and ecosystems. The Policy involved a four-stage approach to implementation:

- Stage 1: Development of the approach to classification, formulation of a standard of protection, and mapping of existing protected areas and/or mechanisms. Scientific workshops will be used to assist with the process, and the results will be put on the website for comment
- Stage 2: Development of the MPA inventory, identification of gaps in the MPA network, and prioritisation of new MPAs
- Stage 3: Establishment of new MPAs to meet gaps in the network. This will be undertaken at a regional level and a national process will be followed for offshore MPAs
- Stage 4: Evaluation and monitoring.

Stage 1 and the inventory specified for Stage 2 are complete and regional forums were established for the Subantarctic and West Coast bioregions. In June 2009, these planning forums released consultation documents on implementation of the MPA Policy in their bioregions:

*Consultation Document - Implementation of the Marine Protected Areas Policy in the Territorial Seas of the Subantarctic Biogeographic Region of New Zealand:*

<http://www.biodiversity.govt.nz/pdfs/seas/subantarctics-mpa-policy-consultation-document.pdf>

*Proposed Marine Protected Areas for the South Island's West Coast Te Tai o Poutini: A public consultation document:*

<http://www.westmarine.org.nz/documents/ProposedMPAsWestCoastSubmissiondocumentwebresv2.pdf>

The MPA Classification, Protection Standard, Implementation Guidelines, together with a summary of subsequent consultation processes around implementing the policy can be found on the Government Biodiversity website at:

[http://www.biodiversity.govt.nz/seas/biodiversity/protected/mpa\\_consultation.html](http://www.biodiversity.govt.nz/seas/biodiversity/protected/mpa_consultation.html)

#### 12.8.4. Revised Coastal Policy Statement

The revised New Zealand Coastal Policy Statement (NZCPS) came into force in December 2010, replacing the original 1994 NZCPS. The statement is to be applied, as required by the Resource Management Act 1991 (RMA), by persons exercising functions and powers under that Act. The documentation can be read on the Department of Conservation's website at:

<http://www.doc.govt.nz/publications/conservation/marine-and-coastal/new-zealand-coastal-policy-statement/new-zealand-coastal-policy-statement-2010/>

The NZCPS does not directly apply to fisheries management decision-making, although the Minister of Fisheries is required to have regard to the Statement when making decisions on sustainability measures under section 11 of the Fisheries Act. In addition, this synopsis includes chapters on land use issues and habitats of particular significance for fisheries management for which the main threats are managed under the RMA (e.g., land use practices could increase sedimentation and affect the estuarine nursery grounds of important fishstocks). In other areas, management of effects under the RMA can complement management of the effects of fishing (e.g., complementary management of the habitat and bycatch of a protected species). The following objectives and policies are considered relevant (numbering as per NZCPS, text in parentheses summarises subheadings in the Statement of most relevance to fisheries values):

**Objective 1: To safeguard the integrity, form, functioning and resilience of the coastal environment and sustain its ecosystems, including marine and intertidal areas, estuaries, dunes and land** (especially by maintaining or enhancing natural biological and physical processes in the coastal environment).

**Objective 6: To enable people and communities to provide for their social, economic, and cultural wellbeing and their health and safety, through subdivision, use, and development** (especially by recognising that the protection of habitats of living marine resources contributes to social, economic and cultural wellbeing and that the potential to utilise coastal marine natural resources should not be compromised by activities on land).

**Policy 5: Land or waters managed or held under other Acts** (especially to consider effects on coastal areas held or managed under other Acts with conservation or protection purposes and to avoid, remedy or mitigate adverse effects of activities in relation to those purposes).

**Policy 8: Aquaculture: Recognise the significant existing and potential contribution of aquaculture to the social, economic and cultural well-being of people and communities** (especially by taking account of the social and economic benefits of aquaculture, recognising the need for high water quality, and including provision for aquaculture in the coastal environment).

**Policy 11: Indigenous biodiversity: To protect indigenous biological diversity in the coastal environment** (especially by avoiding, remedying or mitigating adverse effects on: habitats that are important during the vulnerable life stages of indigenous species; ecosystems and habitats that are particularly vulnerable to modification; and habitats of indigenous species that are important for recreational, commercial, traditional or cultural purposes).

**Policy: 21 Enhancement of water quality: Where the quality of water in the coastal environment has deteriorated so that it is having a significant adverse effect on ecosystems, natural habitats, or water based recreational activities, or is restricting existing uses, such as aquaculture, shellfish gathering, and cultural activities, give priority to improving that quality.**

**Policy 22: Sedimentation** (especially with respect to impacts on the coastal environment).

**Policy 23: Discharge of contaminants** (especially with respect to impacts on ecosystems and habitats).

### **12.8.5. Management of Activities in the EEZ**

In August 2007 the Ministry for the Environment (MfE) released a discussion paper “*Improving regulation of environmental effects in New Zealand’s Exclusive Economic Zone*” seeking comment on a preferred legislative option for managing the impacts of activities in the EEZ. The discussion paper stated that environmental effects in the EEZ were, at that time, managed by sector-specific legislation, which creates the following problems:

- gaps and inconsistencies in the operational control of environmental effects
- unclear environmental outcomes against which activities and their effects should be assessed
- uncertainty for investors about the regulatory environment
- uncertainty about how the effects of activities on each other should be managed.

The MfE website (<http://www.mfe.govt.nz/issues/oceans/current-work/index.html>) states that EEZ legislation is a priority for the current government. In response to the Gulf of Mexico oil spill, the Ministry of Economic Development is commissioning an independent study on New Zealand's health, safety and environmental provisions around minerals activities such as deep sea drilling. This report, along with the proposed legislation developed by the last government, will be considered by Ministers before making final policy and timeline decisions for EEZ legislation.

Proposals for EEZ legislation to manage effects other than those caused by fishing do not directly apply to fisheries management decision-making under the Fisheries Act. However, there are issues around the management of cumulative effects (e.g., of more than one activity on benthic communities) and around effects of any proposed new activities in the EEZ on fishing activity already occurring. Some projects already completed or currently underway are likely to be useful for these processes (e.g., detailed maps of fishing effort produced under ENV2001/07 and BEN2006/01 and enhancements of the Marine Environment Classification produced under ZBD2005-02 for demersal fishes and BEN2006/01A for benthic invertebrates).

### **12.8.6. Ministry for Research Science and Technology Roadmaps**

The Ministry for Science Research and Technology (MRST, now a component of the Ministry of Business, Innovation and Employment, MBIE) stated in its 2006 overview “*Science for New Zealand*” that our science system aims to set long-term direction for RS&T, but allows flexibility to alter direction as needs and opportunities change. Recent direction setting has replaced periodic national processes with a range of continuous processes, often focused on particular areas or topics including:

- Government-led strategy processes around particular areas of national need or opportunity. The Biodiversity Strategy and Biosecurity Strategy are recent examples that have led to changes in institutional arrangements, policies, and funding in RS&T.
- More focused processes by research organisations and or user communities around how a particular area of science could better support national needs, or may be needed to retain or build new capability. These may be endorsed by Ministers or implemented directly by research organisations.
- ‘Roadmaps for Science’, led by MRST, aimed at developing and coordinating RS&T directions and bringing a stronger RS&T perspective to other Government strategies. Roadmaps describe New Zealand’s current research activity, interpret Government’s objectives and strategies in the area, and provide guidance to public research investment agencies as well as other participants in the science system.

Roadmaps for Science were published by MRST for Energy Research (December 2006), Nanoscience & Nanotechnologies (February 2007), Biotechnology Research (March 2007), and Environment Research (June 2007). Probably the most relevant of these is that for Environment Research which can be found at MRST's website at:

<http://www.msi.govt.nz/update-me/archive/publications-archive/>

(if you would like to request a copy of this publication, please email [info@msi.govt.nz](mailto:info@msi.govt.nz)).

It is important to note that these roadmaps relate primarily to research funded by the erstwhile Foundation for Research Science and Technology (FRST, now also part of MBIE) and much less to applied, operational research purchased by the Ministry of Fisheries and some other government departments. However, the Environmental Roadmap for Science noted that environmental management decisions increasingly require an understanding of whole system processes and a multi-dimensional approach. More integrated and systems-based approaches can offer environmental managers and decision-makers answers to many of the questions they are facing. A crucial task then becomes one of creating a New Zealand science environment within which systems-based approaches can develop and flourish, acknowledging that small-scale studies remain important to underpin these approaches. MRST identified three overarching themes that require additional focus: systems understanding and integration (e.g., ecosystem aspects of fisheries management); transfer and uptake (including adaptive management to advance scientific understanding); and information systems (including management of databases and collections).

From a suite of six key research areas (global environmental change, land, water and coasts, including the coastal marine area, urban design and hazards, biosecurity, biodiversity, and oceanic systems), MRST identified five key research directions, two of which are of most relevance to fisheries interests:

Direction 4: Over the next few years, the government will give priority to developing more integrated multidisciplinary approaches, and to improving transfer, uptake and information systems in the following areas:

- global environmental change – with a focus on providing the knowledge for integrated ecological, physical and socio-economic modelling of climate change impacts on water and soil resources, land use, biosecurity, biodiversity and potential global impacts;
- land, water and coasts – with a focus on sustainable land and coastal aquatic use, including the impacts of land use on freshwater and the impacts of freshwater, land management and aquatic production on coastal marine environments;
- biosecurity – reflecting the directions set in the Biosecurity Science Research and Technology Strategy.

Direction 5: Over the longer-term, the government will focus on more integrated multidisciplinary approaches, and improved transfer and uptake, and information systems in the biodiversity and oceanic systems areas.

MRST believes that the Environmental Science Roadmap will make a difference by:

- Equipping environmental managers with integrated research results and tools which will help them avoid, remedy or mitigate future environmental problems.
- Enhancing New Zealand's potential as a test bed and world leader for new innovations and business developments in environmental technologies.
- Improved predictions of and responses to natural hazards events.
- Improved responses to climate change.

#### **12.8.7. National Plan of Action to Reduce the Incidental Catch of Seabirds in New Zealand Fisheries**

The National Plan of Action (NPOA) to Reduce the Incidental Catch of Seabirds in New Zealand Fisheries came into action in April 2004. The NPOA-Seabirds provides a framework to inform the management of seabird/fisheries interactions. It also sets a number of time-bound objectives to be reached by partnering with the fishing industry to reduce the incidental catch of seabirds.

The document is available online at:

<http://www.doc.govt.nz/documents/conservation/native-animals/birds/npoa.pdf>

A revised version of this document is currently under consultation, with a draft version available online through the consultations page of the Fisheries website of the Ministry for Primary Industries at:

[http://www.fish.govt.nz/en-nz/Consultations/default.htm?wbc\\_purpose=bas](http://www.fish.govt.nz/en-nz/Consultations/default.htm?wbc_purpose=bas)

*Note that these links are likely to change due to the current revision. The revised document will be available early in 2013.*

#### **12.8.8. New Zealand National Plan of Action for the Conservation and Management of Sharks**

The New Zealand National Plan of Action (NPOA) for the Conservation and Management of Sharks was approved by the Minister of Fisheries on 13 October 2008. The purpose of the NPOA-Sharks is to ensure the conservation and management of sharks and their long-term sustainable use. It also contains a set of actions in order to meet this purpose. The document is available online at:

<http://www.fish.govt.nz/NR/rdonlyres/F0530841-CD61-4C3E-9E50153A281A4180/0/NPOAsharks.pdf>

*Note that the NPOA-Sharks is currently under review with a revised edition due in 2013.*

## **12.9. Appendix of Aquatic Environment and Biodiversity funded and related projects**

The following listing of projects are those relevant to aquatic environment research that have been through research planning and subsequently been funded by the Ministry of Fisheries (MFish), the Ministry for Primary Industries (MPI) or the fishing industry. These projects have been ordered by the research themes:

1. Protected species
2. Non-protected bycatch
3. Benthic impacts
4. Ecosystem effects
5. Biodiversity

Within these themes projects are ordered chronologically (from the most recent to the oldest). A list of references cited within the table is included at the end of this appendix.

Each project or row of the table is described by a project number (used by MFish/MPI), a project title, specific objectives (where there are many objectives and some are clearly not relevant to aquatic environment research they may not be listed), project status and any relevant citations from the project.

Citations listed below can be accessed differently depending upon the type of output. Finalised FARs (Fisheries Assessment Reports) and AEBRs (Aquatic Environment and Biodiversity Reports), historical FARDs (Fisheries Assessment Research Documents) and MMBRs (Marine Biodiversity and Biosecurity Reports), and some FRRs (final Research Reports) can be found at: <http://fs.fish.govt.nz/Page.aspx?pk=61&tk=209>. Increasingly, reports will be available from the MPI website at: <http://www.mpi.govt.nz/news-resources/publications>. For unpublished documents or those not available on either of these websites please contact [Science.Officer@mpi.govt.nz](mailto:Science.Officer@mpi.govt.nz). Every attempt has been made to make this table comprehensive and correct, but if any errors are found please send suggested corrections or additions through to [Science.Officer@mpi.govt.nz](mailto:Science.Officer@mpi.govt.nz).

*AEBAR 2012: Appendices: Past projects*

**PROTECTED SPECIES**

<b>Project Code</b>	<b>Project Title</b>	<b>Specific Objectives</b>	<b>Status</b>	<b>Citation/s</b>
PRO2012-02	Assessment of the risk to marine mammal populations from New Zealand commercial fisheries	<ol style="list-style-type: none"> <li>1. To scope the risk assessment, including producing an agreed list of marine mammal populations (in concert with MAF and DOC).</li> <li>2. To review the literature, compile the required information and evaluate the appropriate level of risk assessment for the marine mammal populations identified in objective 1.</li> <li>3. To conduct a risk assessment for the marine mammal populations identified in objective 1 using, where possible, a risk index reflecting the ratio of fisheries-related mortality to the level of potential biological removal.</li> <li>4. To refine the results of the risk assessment for priority marine mammal populations by incorporating spatially and temporally-explicit abundance, distribution and capture information.</li> </ol>	Approved but not contracted	
PRO2012-07	Cryptic mortality of seabirds in trawl and longline fisheries	<ol style="list-style-type: none"> <li>1. To review available information from international literature and unpublished sources to characterize and inform estimation of cryptic mortality and live releases for at-risk seabirds in New Zealand trawl and longline fisheries</li> <li>2. To review the extent to which fisheries observer data informing current estimates of seabird captures may be used to also estimate cryptic mortalities in different fishery groups in the seabird risk assessment, and identify key assumptions and associated uncertainty in the estimation of cryptic mortalities.</li> <li>3. To identify those species and/or fishery groups for which current uncertainty regarding cryptic mortality contributes most strongly to high risk scores for at-risk seabird species, and recommend options to improve estimation of cryptic mortality for those species / fishery group combinations.</li> </ol>	Approved but not contracted	

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
PRO2012-08	Improved estimation of spatio-temporal overlap with fisheries for at-risk seabird species	1. To generate seabird distribution map layers for seabird species which the existing level 1 risk assessment identifies as being at-risk, but for which no level 2 assessment has been completed. 2. To modify seabird distribution layers used in the current level 2 risk assessment, for those species that the L2 assessment identifies as at-risk and for which: i) spatial distributions used in the current L2 assessment are known to be wrong, or ii) improved spatial distribution layers are readily available (e.g. from new satellite telemetry data). 3. To seasonally disaggregate seabird spatial distribution data layers for those at-risk seabird species with a strongly seasonal abundance and/or distribution in the New Zealand EEZ. 4. To utilize updated spatial/seasonal seabird distribution layers to generate improved estimates of spatio-temporal overlap with fisheries, for integration into the existing level 2 seabird risk assessment framework.	Approved but not contracted	
PRO2012-09	Improvements to key information gaps for highest risk seabird populations TBC	1. To improve estimates of the population size of specified seabirds where this will substantially reduce uncertainty in the risk ratio estimated in the Level 2 seabird risk assessment. 2. To improve estimates of the age at first breeding for specified seabird populations where this will substantially reduce uncertainty in the risk ratio estimated in the Level 2 seabird risk assessment. 3. To improve estimates of the average adult survival rate for specified seabird populations where this will substantially reduce uncertainty in the risk ratio estimated in the Level 2 seabird risk assessment.	Approved but not contracted	
PRO2012-10	Level 3 risk assessment for Antipodean albatross TBC	1. Develop an Antipodean albatross population model 2. Assess the effect of fisheries mortality on population viability 3. As information permits, assess the effect of alternative management strategies	Approved but not contracted	
ENV2011-01	NPOA-sharks science review	1. To collate and summarise information in support of a review of the National Plan of Action for the Conservation and Management of Sharks (NPOA-sharks). 2. To identify research gaps from objective 1 and suggest cost-effective ways these could be addressed.	Completed	Francis & Lyon 2012

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
No project number	A risk assessment of threats to Maui's dolphins	To evaluate of the risks posed to Maui's dolphin to support the review of the TMP.	Completed	Currey <i>et al.</i> 2012
SRP2011-03	Probabilistic modelling of sea lion interactions	1. Estimate the probability that a sea lion suffers mild head trauma following a collision with a SLED grid	Completed	Abraham 2011
SRP2011-04	HSL Modelling	1. Revise Breen-Fu-Gilbert sea lion model	Completed	Breen <i>et al.</i> 2010
DEE2010-03	Development of a methodology to estimate cryptic mortalities to ETP species from DW fishing activity	1. To conduct a review of existing national and international techniques to estimate cryptic mortality of endangered, threatened and protected species caused by deepwater fishing activities 2. To develop one or more approaches to estimating cryptic mortality of endangered, threatened and protected species caused by deepwater fishing activities 3. To field test one or more approaches to estimating cryptic mortality of endangered, threatened and protected species caused by deepwater fishing activities	Ongoing analysis	
No project number	A risk assessment framework for incidental seabird mortality associated with New Zealand fishing in the New Zealand EEZ	To describe the conceptual and methodological framework of this risk assessment approach to guide the completion of similar risk assessments elsewhere.	Completed	Sharp <i>et al.</i> 2011
PRO2010-01	Estimating the nature and extent of incidental captures of seabirds, marine mammals and turtles in New Zealand commercial fisheries	1. To estimate the nature and extent of captures of seabirds, marine mammals and turtles, and the warp strikes of seabirds in New Zealand fisheries for the fishing years 2009/10, 2010/11 and 2011/12.	Ongoing analysis	Thompson <i>et al.</i> 2011
PRO2010-02	Research into key areas of uncertainty or development of mitigation techniques for the revised npoa-seabirds	1.To provide the information necessary to underpin the revised NPOA-seabirds or develop mitigation techniques to reduce risk identified via the revised NPOA-seabirds.	Ongoing analysis	

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
SRP2010-03	Fur Seal interactions with a SED excluder device	1. Fur seal interactions with SED excluder device (Dr J Lyle)	Completed	Lyle 2011
SRP2010-05	Fur seal interaction with an SLED excluder device	<ul style="list-style-type: none"> <li>• Using a series of 10-15 impact tests at a maximum collision speed of 5 or 6 ms<sup>-1</sup>, develop a "HIC map" for the SLED grid to enable the consequences of collisions with different parts of the grid by sea lions of different head masses to be predicted (scaling values (for eq 3) will include -1/3, -2/3, and -3/4)</li> <li>• Using a small number of collision tests, verify that the HIC for a glancing blow can be predicted with sufficient accuracy by resolving vectors</li> <li>• Calculate the maximum possible sensitivity to different boundary conditions using the relative masses of the SLED grid and sea lion heads</li> <li>• Clarify in the final research report that undertaking tests in air (as opposed to underwater) should not affect the results</li> </ul>	Completed	Ponte et al. 2011
IPA2009-09	Sea Lion bioenergetics modelling	<ol style="list-style-type: none"> <li>1. To review and collate data on growth, metabolism, diet and reproductive parameters of NZ sea lions or, if data are inexistent, of other sea lions species</li> <li>2. To analyse the energy density of various NZ sea lion prey items</li> <li>3. To incorporate the data acquired in objectives 1. and 2. into a bioenergetics model to estimate the energy and food requirements of NZ sea lions</li> </ol>	Completed	Meynier 2010
IPA2009-16	Preliminary impact assessment of NZ sea lion interaction with SLEDs	1. Preliminary impact assessment of New Zealand sea lion interactions with SLEDs	Completed	Ponte et al. 2010
IPA2009-19/20	Level 2 seabird risk assessment rerun	1. To examine the risk of incidental mortality from commercial fishing for 64 seabird species in New Zealand trawl and longline fisheries	Completed	Richard et al. 2011
No project number	External review of NZ sea lion bycatch necropsy data and methods	<p>The primary purposes of this review were to determine whether, in the opinion of a group of independent experts:</p> <ul style="list-style-type: none"> <li>- the interpretation of necropsy findings and trauma classification system used by Dr Wendi Roe are valid</li> <li>- sea lions recovered from trawl nets have sustained clinically significant trauma</li> <li>- some or all of the sea lions exiting through SLEDs are likely to survive</li> </ul>	Completed	Roe 2010a
PRO2009-01A	Abundance & distribution of Hector's & maui's dolphins (5 year project)	<ol style="list-style-type: none"> <li>1. To estimate the distribution of the South Coast South Island Hector's dolphin sub-population in both winter and summer.</li> <li>2. The work for this sub-project was subsequently extended to include data collection necessary to estimate abundance.</li> </ol>	Completed	Clement & Mattlin 2010

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
PRO2009-01B	Abundance, distribution, and productivity of Hector's (and Maui's) dolphins	<ol style="list-style-type: none"> <li>1. To estimate the likely precision of abundance estimates from summer aerial surveys for Hector's dolphins along the East Coast South Island (ECSI; from Farewell Spit to Nugget Point) under different levels of sampling intensity and stratification.</li> <li>2. To estimate the likely precision of abundance estimates and the likely quality of distribution information from winter aerial surveys for Hector's dolphins along the ECSI under different levels of sampling intensity and stratification.</li> <li>3. To identify and quantify trade-offs between the precision of abundance estimates and the quality of distribution information as well as between overall precision and likely cost (e.g., based on the number of flying hours required).</li> <li>4. To identify key areas and times for which it would be particularly useful to have information on Hector's dolphin distribution (e.g., where risk may come from overlap with particular fisheries) and quantify trade-offs between the precision of ECSI-wide surveys and collecting such fine-scale information.</li> <li>5. Assess the extent to which two-phase or adaptive approaches would be useful to improve the surveys' utility for assessing dolphin distribution, particularly the seaward limit.</li> </ol>	Ongoing analysis	
PRO2009-04	Development and efficacy of seabird mitigation measures	1. To test the efficacy of a variety of configurations of mitigation techniques at reducing seabird mortality (or appropriate proxies for mortality) in longline fisheries	Completed	No reports specified as required output
ENV2008-03	Bycatch of basking sharks in New Zealand fisheries	<ol style="list-style-type: none"> <li>1. To review the productivity of basking sharks</li> <li>2. To describe the nature and extent of fishery-induced mortality of basking sharks in New Zealand waters and recommend methods of reducing the overall catch.</li> </ol>	Completed	Francis & Smith 2010
PRO2008-01	Risk assessment of protected species bycatch in NZ fisheries	1. To provide an assessment of the risk posed by different fisheries to the viability of New Zealand protected species, and to assign a risk category to all New Zealand fishing operations.	Completed	Waugh <i>et al.</i> 2009

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
SAP2008-14	Sea lion population modelling, additional	1. To assess the likely performance of different bycatch control rules for the SQU6T fishery. 2. To correct and update the Breen-Fu-Gilbert (2008) sea lion model- including assessment of the performance of 200-series and 300-series management control rules. 3. To document the development of the model- including all four objectives of project IPA2006/09 and objective 1 of this project- in a single report suitable for an international review.	Completed	Breen <i>et al.</i> 2010
Deepwater Group	Necropsy of marine mammals captured in New Zealand fisheries in the 2007-08 fishing year	Necropsy of marine mammals captured in New Zealand fisheries in the 2007-08 fishing year	Completed	Roe 2009a
IPA2007-09	Protected species risk assessment	To provide an assessment of the risk posed by different fisheroes to the viability of NZ protected species- and to assign a risk category to all NZ fishing operations	Completed	Waugh <i>et al.</i> 2008
PRO2007-01	Estimating the nature and extent of incidental captures of seabirds in New Zealand commercial fisheries	1. Estimate capture rates per unit effort and total captures of seabirds for the New Zealand EEZ and in selected fisheries by method, area, target fishery, in relation to mitigation methods in use, and, where possible, by seabird species for the fishing year 2006/07, 2007/08 and 2008/09. 2. Examine the incidence of seabird warp strike in trawl fisheries where these data are available from fisheries observers, and estimate the rate of incidents (birds affected per hour) and total number of seabirds affected by fishery, area and method. Examine the factors (fishery, environmental, seasonal, mitigation, area) that influence the probability of warp-strike occurring.	Completed	Abraham 2010; Abraham & Thompson 2009a; 2010; 2011a; b; Thompson & Abraham 2009a; Abraham <i>et al.</i> 2010b
PRO2007-02	Estimating the nature and extent of incidental captures of seabirds in New Zealand commercial fisheries	1. Estimate capture rates per unit effort and total captures of seabirds for the New Zealand EEZ and in selected fisheries by method, area, target fishery, in relation to mitigation methods in use, and, where possible, by seabird species for the fishing year 2006/07, 2007/08 and 2008/09. 2. Examine the incidence of seabird warp strike in trawl fisheries where these data are available from fisheries observers, and estimate the rate of incidents (birds affected per hour) and total number of seabirds affected by fishery, area and method. Examine the factors (fishery, environmental, seasonal, mitigation, area) that influence the probability of warp-strike occurring.	Completed	Abraham <i>et al.</i> 2010a; Thompson & Abraham 2009a; 2009b; 2009c; 2010; 2011; Thompson <i>et al.</i> 2010a; 2010b

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2006-05	The use of electronic monitoring technology in New Zealand longline fisheries	<ol style="list-style-type: none"> <li>1. Trial the deployment of electronic monitoring systems in selected longline fisheries, monitoring incidental take of protected species.</li> <li>2. Evaluate the efficacy of electronic monitoring in allowing enumeration and identification of protected species captures.</li> <li>3. Recommend options for data management and information transfer arising from the deployment of electronic monitoring in selected fisheries.</li> </ol>	Completed	McElderry <i>et al.</i> 2008
IPA2006-02	The efficacy of warp strike mitigation devices: trials in the 2006 squid fishery	<ol style="list-style-type: none"> <li>1. Groom the mitigation trial data and produce a summary of the data (100%)</li> <li>2. Examine strike rates and capture rates on warps and mitigation devices (100%)</li> <li>3. Determine the relative efficacy of mitigation devices tested in the trial (100%)</li> <li>4. Make recommendations regarding future trials (100%)</li> <li>5. Compare seabird warp strike data for 2005 and 2006 (100%)</li> <li>6. Work with SeaFIC and the mitigation trials TAG to produce analyses and outputs (100%)</li> </ol>	Completed	Middleton & Abraham 2007
IPA2006-09	Modelling interactions between trawl fisheries and New Zealand Sea lion interactions	<ol style="list-style-type: none"> <li>1. Model the New Zealand sea lion population and explore alternative management procedures for controlling New Zealand sea lion bycatch in the SQU 6T fishery</li> <li>2. Collate and review all available sea lion biological data- fisheries data- and sea lion bycatch data relevant to a population model and management strategy evaluation for the Auckland Islands sea lion population</li> <li>3. Update and improve the existing Breen and Kim sea lion population model (2003) to incorporate all relevant data and address model uncertainties including but not necessarily limited to those identified by the AEWG</li> <li>4. Fit the revised model to all available data and test sensitivity including but not necessarily limited to runs identified by the AEWG</li> <li>5. Test a range of management procedures (rules) with the model to determine if they meet agreed management criteria</li> </ol>	Completed	Breen 2008

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
IPA2006-13	Identification of Marine Mammals Captured in New Zealand Fisheries	1. To determine, through examination of returned marine mammal carcasses, the species, sex, reproductive status, and age-class of marine mammals returned from New Zealand fisheries. 2. To detail any injuries and, where possible, the cause of mortality of marine mammals returned from New Zealand fisheries, and examine relationships between injuries and body condition, breeding status, and other associated demographic characteristics.	Completed	Roe 2009b
PRO2006-01	Data collection of demographic, distributional and trophic information on selected seabird species to allow estimation of effects of fishing on population viability	1 To gather demographic, distributional and dietary information on selected seabird species to allow assessment of effects of fishing on population viability.	Completed	Sagar & Thompson 2008; Sagar et al. 2009a; b; 2010a; b; c; Baker et al. 2008; 2009; 2010
PRO2006-02	Modelling of the effects of fishing on the population viability of selected seabirds	1. Model the effects of fisheries mortalities on population viability compared with other sources of mortality or trophic effects of fishing 2. Examine the overlap of fishing activity with species distribution at sea for different stages of the breeding and life-cycle and for different sexes, and assess the likely risk to species or populations from fisheries (by target species fisheries, fishing methods, area and season) in the New Zealand EEZ	Completed	Francis & Bell 2010; Francis 2012
PRO2006-04	Estimation of the nature and extent of incidental captures of seabirds in New Zealand commercial fisheries	1. To estimate the nature and extent of captures and warp-strikes of seabirds in New Zealand fisheries for the fishing year 2005/06.	Completed	Baird & Smith 2008
PRO2006-05	Estimating the nature and extent of marine mammal captures in New Zealand commercial fisheries	1. To estimate and report the total numbers, releases and deaths of marine mammals where possible by species, fishery and fishing method, caught in commercial fisheries for the years 1990 to the end of the fishing year 2005/06. 2. To analyse factors affecting the probability of fur seal captures for the years 1990 to the end of the fishing year 2005/06. 3. To classify fishing areas, seasons and fishing methods into different risk categories in relation to the probability of marine mammal incidental captures for the years from 1990 through to the end of the fishing year 2005/06.	Completed	Mormede <i>et al.</i> 2008; Baird 2008a; 2008b; Smith & Baird 2009; Smith & Baird 2011; Baird 2011.

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
PRO2006-07	Characterise non-commercial fisheries interactions	1. To characterise non-commercial fisheries interactions with seabirds and marine mammals 2. Characterise non-commercial fisheries risk to seabirds and marine mammals by area and method Recommend mitigation measures appropriate for uptake in non-commercial fisheries in which seabird or marine mammal captures occur	Completed	Abraham et al. 2010a; Thompson & Abraham 2009a; 2009b; 2009c; 2010; 2011; Thompson et al. 2010a; b; c
ENV2005-06	Estimation of protected species captures in longline fisheries using electronic monitoring	1. To provide estimates of seabird and marine mammal mortalities from longline fisheries in New Zealand using electronic monitoring systems and to recommend deployment and data management options for ongoing use of these systems for estimation of protected species incidental take.	Completed	McElderry et al. 2007
ENV2005-01	Estimation of the nature and extent of incidental captures of seabirds in New Zealand fisheries	1. To estimate the nature and extent of captures of seabirds in selected New Zealand fisheries for the fishing year 2004/05.	Completed	Baird & Smith 2007a; Baird & Gibbert 2010
ENV2005-02	Estimation of the nature and extent of marine mammal captures in New Zealand fisheries	To examine the nature and extent of the captures of marine mammals in New Zealand fisheries, for the whole New Zealand EEZ, by Fishery Management Area and fishing season, and by smaller metric as appropriate for the fishing year 2004/05. 2. Examine alternative methods for estimating sea lion captures and recommend one or more alternative standardised methods for describing and estimating sea lion captures in the SQU 6T fishery.	Completed	Abraham 2008; Baird 2007; Smith & Baird 2007b; Baird & Smith 2007b
ENV2005-04	Identification of marine mammals captured in New Zealand	1. To determine the species- sex- and where possible- age and reproductive status of marine mammals captured in New Zealand fisheries. 2. To necropsy marine mammals captured incidentally to New Zealand fishing operations to determine life-history characteristics and the likely cause of mortality. 3. To determine- through examination of returned marine mammal carcasses- the taxon to species-level- sex- and reproductive status- and age-class of marine mammals captured in New Zealand fisheries. 4. To detail the injuries and where possible the cause of mortality of marine mammals returned from New Zealand fisheries- along with their body condition and breeding status- and other associated demographic characteristics. 5. To detail the protocol used for the necropsy of marine mammals- to provide a standardised procedure for autopsy to determine species- age- sex and associated demographic characteristics for fishery-killed specimens.	Completed	Roe 2007

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2005-09	Data collection to estimate key performance indicators in the Chatham albatross, <i>Diomedea eremita</i> .	<ol style="list-style-type: none"> <li>1. To gather data on key population parameters for Chatham albatross <i>Diomedea eremita</i>- to enable population viability to be assessed- and the responses of key parameters to fisheries mortality and fisheries management activities to mitigate fisheries related risk</li> <li>2. To undertake field research to collect data on population growth rates- adult survival- inter-breeding season survival- mortality due to predation at the colony- fecundity and associated parameters for Chatham Albatross- following the study design project</li> <li>3. To undertake field research to determine the range and extent foraging movements of Chatham albatrosses within New Zealand fishing waters- and examine the nature and extent of any association between Chatham albatrosses and fishing activities.</li> </ol>	Completed	No reports specified as required output
ENV2005-13	Assessment of risk to yellow-eyed penguin <i>Megadyptes antipodes</i> from fisheries incidental mortality	<ol style="list-style-type: none"> <li>1. To review existing data on yellow-eyed penguin <i>M. antipodes</i> population performance and fisheries information and provide an analysis of the potential effect of fishing mortality and other factors on population viability.</li> <li>2. To recommend data collection requirements and protocols for the assessment of the effects of fishing on yellow-eyed penguins.</li> </ol>	Completed	Maunder 2007
ENV2004-04	Characterisation of seabird captures in New Zealand fisheries	<ol style="list-style-type: none"> <li>1. Characterisation of seabird captures in New Zealand fisheries.</li> </ol>	Completed	Mackenzie & Fletcher 2006
ENV2004-05	Modelling of impacts of fishing-related mortality on New Zealand seabird populations	<ol style="list-style-type: none"> <li>1. To examine and identify modelling approaches to analyse seabird demographic impacts that may be occurring as a result of fisheries mortality.</li> <li>2. To compile databases of available demographic and distributional data on selected seabirds affected by fisheries mortality and New Zealand fisheries and estimate key population parameters and seasonal distribution for each species.</li> <li>3. To estimate rates of removals related to fishing activities in New Zealand for selected seabird species, where possible by age class and sex.</li> <li>4. To describe the spatial overlap of seabird distributions at sea, with fisheries where the risk of incidental mortality has been demonstrated to be moderate to high.</li> <li>5. To examine the potential for factors other than fisheries removals within the New Zealand zone to influence the population dynamics of the selected study species.</li> <li>6. To characterise selected seabird populations' abilities to sustain removals related to fishing operations within the New Zealand EEZ, and to recommend, where possible environmental standards for assessing the sustainability of selected fishing operations in relation to impacts on seabird populations.</li> </ol>	Completed	Fletcher <i>et al.</i> 2008

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2004-02	Estimation of New Zealand sea lion incidental captures in New Zealand Fisheries	1. To estimate the level of New Zealand sea lion ( <i>Phocartos hookeri</i> ) incidental capture in New Zealand fisheries	Completed	Smith & Baird 2007a
ENV2004-06	Maui's dolphin study	1. To quantify and compare summer and winter distribution of maui's dolphin	Completed	Slooten <i>et al.</i> 2005
IPA2004-14	Seabird warp strike in the southern squid trawl fishery	1. To document seabird warp strike in the southern squid trawl fishery, 2004-05	Completed	Abraham & Kennedy 2008
ENV2003-05	Review of the Current Threat Status of Associated or Dependent Species	1. To assess the current threat status of selected associated or dependent species.	Completed	Baird <i>et al.</i> 2010
No project number	QMA SQU6T New Zealand sea lion incidental catch and necropsy data for the fishing years 2000-01, 2001-02 and 2002-03	Objectives unknown	Completed	Mattlin 2004
MOF2002-03L	Exploring alternative management procedures for controlling bycatch of Hooker's sea lions in the SQU 6T squid fishery	Objectives unknown	Completed	Breen & Kim 2006
ENV2001-01	Estimation of seabird incidental captures in New Zealand fisheries	1. To estimate the level of seabird incidental capture in New Zealand fisheries. 2. To recommend appropriate levels of observer coverage for estimation of seabird incidental capture in New Zealand fisheries.	Completed	Baird 2004a; b; c; Smith & Baird 2008b
ENV2001-02	Incidental capture of <i>Phocartos hookeri</i> (New Zealand sea lions) in New Zealand commercial fisheries, 2001-02.	1. To estimate and report the total numbers of captures, releases, and deaths of <i>Phocartos hookeri</i> caught in fishing operations, including separate estimates for SQU 6T and other areas, as appropriate, during the 2001102 fishing year, including confidence limits and an investigation of any statistical bias in the estimate.	Completed	Baird 2005a; b; c; Baird & Doonan 2005
ENV2001-03	Estimation of <i>Arctocephalus forsteri</i> (New Zealand fur seal) incidental captures in New Zealand fisheries	1. To estimate the level of <i>Arctocephalus forsteri</i> incidental capture in New Zealand fisheries. 2. To recommend appropriate levels of observer coverage for estimation of <i>Arctocephalus forsteri</i> incidental capture in New Zealand fisheries.	Completed	Smith & Baird 2008a; Baird 2005d; e; f

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2000-01	Protected species bycatch	1. To estimate the total numbers of captures, releases, and deaths of seabirds and marine mammals - by species -caught in fishing operations during the 1999-2000 fishing year.	Completed	Baird 2003
ENV2000-02	Estimation of incidental mortality of New Zealand sea lions in New Zealand fisheries	1. To examine the factors that may influence the level of incidental mortality of New Zealand sea lion in New Zealand fisheries 2. To recommend appropriate levels of observer coverage for estimation of incidental mortality of New Zealand sea lion in New Zealand sea lion fisheries	Completed	Doonan 2001; Bradford 2002; Smith & Baird 2005a; b
ENV99-01	Incidental capture of seabirds, marine mammals and sealions in commercial fisheries in New Zealand waters	Objectives unknown	Completed	Baird 2001; Doonan 2000
ENV98-01	Estimation of nonfish bycatch in commercial fisheries in New Zealand waters, 1997–98	Objectives unknown	Completed	Baird 1999b; Baird & Bradford 1999
No project number	Annual review of bycatch in southern bluefin and related tuna longline fisheries in the New Zealand 200 n. mile Exclusive Economic Zone	Objectives unknown	Completed	Baird <i>et al.</i> 1998
SANF01	Report on the incidental capture of nonfish species during fishing operations in New Zealand waters	Objectives unknown	Completed	Baird 1997
No project number	Nonfish Species and Fisheries Interactions	Objectives unknown	Completed	Baird 1996
No project number	Nonfish Species and Fisheries Interactions	Objectives unknown	Completed	Baird 1995
No project number	Incidental catch of Hooker's sea lion in the southern trawl fishery for squid, summer 1994	Objectives unknown	Completed	Doonan 1995

*AEBAR 2012: Appendices: Past projects*

PROTECTED SPECIES *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
No project number	Analyses of factors which influence seabird bycatch in the Japanese southern bluefin tuna longline fishery in New Zealand waters, 1989-93	1. to assess the influence that 15 monitored environmental and fishery related factors had on seabird bycatch rates, and to gauge the effectiveness of various mitigation measures	Completed	Duckworth 1995
No project number	Nonfish Species and Fisheries Interactions	Objectives unknown	Completed	Baird 1994
No project number	Incidental catch of fur seals in the west coast South Island hoki trawl fishery, 1989-92	Objectives unknown	Completed	Mattlin 1993

*AEBAR 2012: Appendices: Past projects*

NON-PROTECTED BYCATCH

Project Code	Project Title	Specific Objectives	Status	Citation/s
No project number	Incidental catch of non-fish species by setnets in New Zealand waters	Objectives unknown	Completed	Taylor 1992
DAE2010-02	Bycatch monitoring & quantification for scampi bottom trawl	1. To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded in the specified fishery, for the fishing years since the last review, using data from Ministry of Fisheries Observers and commercial fishing returns. 2. To compare estimated rates and amounts of bycatch and discards from this study with previous projects on bycatch in the specified fishery. 3. To compare any trends apparent in bycatch rates in the specified fishery with relevant fishery independent trawl surveys. 4. To provide annual estimates of bycatch for nine Tier 1 species fisheries and incorporate into the Aquatic Environment and Biodiversity Report specified in Objective 3 for SQU, SCI, HAK, HOK, JMA, ORH, OEO, LIN, SBW	Completed	Anderson 2012
ENV2009-02	Bycatch and discards in oreo and orange roughy trawl fisheries	1. To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded, in the trawl fisheries for oreos for the fishing years 2002/03 to 2008/09 using data from Scientific Observers and commercial fishing returns. 2. To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded, in the trawl fisheries for orange roughy for the fishing years 2004/05 to 2008/09 using data from Scientific Observers and commercial fishing returns.	Completed	Anderson 2011
IDG2009-01	Finfish field identification guide	1. To complement the field identification guide under IDG2006/01 with the remaining 120 fish species caught by commercial fishers in New Zealand waters	Completed	McMillan 2011 a;b;c
ENV2008-01	Fish and invertebrate bycatch and discards in southern blue whiting fisheries	1. To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded, in the trawl fisheries for southern blue whiting for the fishing years 2002/03 to 2006/07 using data from Scientific Observers and commercial fishing returns.	Completed	Anderson 2009b
ENV2008-02	Estimation of non-target fish catch and both target and non-target fish discards in hoki, hake and ling trawl fisheries	Estimates of the catch of non-target fish species, and the discards of target and non-target fish species in the hoki ( <i>Macruronus novaezelandiae</i> ), hake ( <i>Merluccius australis</i> ), and ling ( <i>Genypterus blacodes</i> ) trawl fisheries for the fishing years 2003–04 to 2006–07 using data from Scientific Observers and commercial fishing returns	Completed	Ballara <i>et al.</i> 2010
ENV2008-04	Productivity of deepwater sharks	1. To determine the growth rate, age at maturity, longevity and natural mortality rate of shovelnose dogfish ( <i>Deania calcea</i> ) and leafscale gulper shark ( <i>Centrophorus squamosus</i> ).	In the process of publication	Parker & Francis 2012

*AEBAR 2012: Appendices: Past projects*

NON-PROTECTED BYCATCH *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2007-03	Productivity and Trends in Rattail Bycatch Species	1. To estimate growth, longevity, rate of natural mortality, and length at maturity of four key rattail bycatch species in New Zealand trawl fisheries. 2. To examine data from trawl surveys and other data sources for trends in catch rates or indices of relative abundance for species in Objective 1.	Completed	Stevens <i>et al.</i> 2010
DEE2006-03	Monitoring the abundance of deepwater sharks	1. To monitor the abundance of deepwater sharks taken by commercial trawl fisheries	Completed	Blackwell 2010
ENV2006-01	Bycatch and discards in ling longline fisheries	To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded, in the longline fisheries for ling for the fishing years 1998/99 to 2005/06 using data from MFish Observers and commercial fishing returns.	Completed	Anderson 2008
IDG2006-01	Finfish field identification guide	1. To produce a field guide for fish species in New Zealand 2. To produce a field identification guide for all QMS and other fish species commonly caught in commercial and non-commercial fisheries	Completed	McMillan 2011 a;b;c
TUN2006-02	Estimation of non-target fish catches in the tuna longline fishery	1. To estimate the catches, catch rates, and discards of non-target fish in tuna longline fisheries data from the Observer Programme and commercial fishing returns for the 2005/06 fishing year. 2. To describe bycatch trends in tuna longline fisheries using data from this project and the results of previous similar projects.	Completed	Griggs <i>et al.</i> 2008
TUN2004-01	Estimation of non-target fish catches in the tuna	To estimate the catch rates of non-target fish in the 10ngline fisheries for tuna using data from the Observer Programme and commercial fishing returns for the 2002/03, 2003/04 and 2004/05 fishing years. 2. To estimate the quantities of non-target fish caught in the longline fisheries for tuna using data from the Observer Programme and commercial fishing returns for the 2002/03, 2003/04 and 2004/05 fishing years. 3. To estimate the discards of non-target fish caught in the longline fisheries for tuna using data from the Observer Programme and commercial fishing returns for the 2002/03, 2003/04 and 2004/05 fishing years. 4. To describe trends in the non-target fish catches in the tuna longline fisheries using data from this project and the results of previous similar projects.	Completed	Griggs <i>et al.</i> 2007
ENV2005-17	Estimation of non-target fish catch and both target and non-target fish discards in jack mackerel trawl fisheries	1. To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded, in the trawl fisheries for jack mackerel for the fishing years 20011/2002 to 2004/05 using data from Mfish observers and commercial fishing returns.	Completed	Anderson 2007a

*AEBAR 2012: Appendices: Past projects*

NON-PROTECTED BYCATCH *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2005-18	Estimation of non-target fish catch and both target and non-target fish discards in orange roughy trawl fisheries	1. To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded, in the trawl fisheries for orange roughy for the fishing years 1999/2000 to 2003/04 using data from Scientific Observers and commercial fishing returns.	Completed	Anderson 2009a
ENV2003-01	Estimation of non-target catches in the hoki fishery	1. To estimate the catch rates, quantity and discards of non-target fish catches and the discards of target fish catches in trawl fisheries for hoki, using data from the Observer Programme and commercial fishing returns for the 1999/00 to 2002/03 fishing years. 2. To compare and contrast the estimates from the four years of data in Specific Objective 1 above with the 1990/91 through 1998/99 series previously reported.	Completed	Anderson & Smith 2005
ENV2002-01	Estimation of non-target fish catch and both target and non-target fish discards for the tuna longline fishery	1. To estimate the catch rates, quantity and discards of non-target fish, particularly oceanic shark species, broadbill swordfish and marlin species, caught in the longline fisheries for tuna, using data from Scientific Observers and commercial fishing returns for the 2000/01 and 2001/02 fishing years.	Completed	Ayers <i>et al.</i> 2004
ENV2001-04	Non-target fish catch and discards in selected New Zealand fisheries	To generate estimates of the catch of non-target fish species, and the discards of target and non-target fish species in three important New Zealand trawl fisheries: arrow squid ( <i>Nototodarus sloani</i> & <i>N. gouldi</i> ), jack mackerel ( <i>Trachurus declivis</i> , <i>T. novaezelandiae</i> , & <i>T. symmetricus murphyi</i> ) and scampi ( <i>Metanephrops challengeri</i> )	Completed	Anderson 2004
ENV2001-05	To assess the productivity and relative abundance of deepwater sharks	1. To review the relative abundance, distribution and catch composition of the most commonly caught deepwater shark species: shovelnose dogfish ( <i>Deania catcea</i> ), Baxter's dogfish ( <i>Etmopterus baxteri</i> ), Owston's dogfish ( <i>Cenohoscyrnus owstoni</i> ), longnosed velvet dogfish ( <i>Centroscymnus crepidater</i> ), leafscale gulper shark ( <i>Cenophom quamosus</i> ), and the seal shark ( <i>Dalatias ticha</i> ).	Completed	Balckwell & Stevenson 2003
ENV2001-07	Reducing bycatch in scampi trawl fisheries	1. Collate and review the international literature on methods of reducing bycatch in crustacean trawl fisheries. 2. Review and analyse the data from New Zealand studies. 3. Develop recommendations on future approaches to reducing bycatch in the New Zealand scampi fishery, including some general thoughts on the experimental design of field trials.	Completed	Hartill <i>et al.</i> 2006

*AEBAR 2012: Appendices: Past projects*

NON-PROTECTED BYCATCH *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
PAT2000-01	Review of rattail and skate bycatch, and analysis of rattail standardised CPUE from the Ross Sea toothfish fishery in Subarea 88.1, from 1997-1998 to 2001-02	Objectives unknown	Completed	Feaunaughty et al. 2003; Marriot et al. 2003
ENV99-02	Estimation of non-target fish catch and both target and non-target fish discards in selected New Zealand fisheries	<ol style="list-style-type: none"> <li>1. To estimate the quantity of non-target fish species caught in the trawl fisheries for hoki and orange roughy for the fishing years 1990-91 to 1998-99 using data from Scientific Observers, commercial fishing returns and from research trawl surveys.</li> <li>2. To estimate the quantity of target and non-target fish species discarded in the trawl fisheries for hoki and orange roughy for the fishing years 1990-91 to 1998-99 using data from Scientific Observers, commercial fishing returns and from research trawl surveys.</li> <li>3. To explore the effects of various factors on the total catch of non-target fish species and the discards of target and non-target fish species in the trawl fisheries for hoki and orange roughy for the fishing years 1990-91 to 1998-99.</li> <li>4. To recommend appropriate levels of observer coverage for estimation of non-target fish catch and discards of target and non-target fish species in the hoki and orange roughy fisheries.</li> </ol>	Completed	Anderson et al. 2001
ENV98-02	Pelagic shark bycatch in the New Zealand tuna longline fishery	To determine pelagic shark bycatch in the New Zealand tuna longline fishery	Completed	Francis <i>et al.</i> 2001
No project number	Fish bycatch in New Zealand tuna longline fisheries	Objectives unknown	Completed	Francis <i>et al.</i> 1999; 2000
ENV97-01	Estimation of nonfish bycatch in New Zealand fisheries	<ol style="list-style-type: none"> <li>1. Unknown</li> <li>2. To provide weekly within season estimates of total captures, releases, and deaths by sex and area for New Zealand sea lions taken in the southern squid trawl fishery beginning two (2) weeks after the start of the fishery until 15 May 1998. Estimates of the confidence intervals and coefficient of variation of the point estimates must also be provided.</li> <li>3. Unknown</li> </ol>	Completed	Doonan 1998; Baird 1999a; Baird <i>et al.</i> 1999

*AEBAR 2012: Appendices: Past projects*

NON-PROTECTED BYCATCH *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
SCI97-01	Scampi stock assessment for 1998 and an analysis of the fish and invertebrate bycatch of scampi trawlers	1. To summarise catch, effort, observer, and research information for scampi fisheries in QMAs 1,2,3,4 (east and western portions), and 6A in 1998	Completed	Cryer <i>et al.</i> 1999

## BENTHIC IMPACTS

Project Code	Project Title	Specific Objectives	Status	Citation/s
BEN2012-02	Spatial overlap of mobile bottom fishing methods and coastal benthic habitats	1. To use existing information and classifications to describe the distribution of benthic habitats throughout New Zealand's coastal zone (0–200 m depth). 2. To rank the vulnerability to fishing disturbance of habitat classes from Objective 1. 3. To describe the spatial pattern of fishing using bottom trawls, Danish seine nets, and shellfish dredges and assess overlap with each of the habitat classes developed in Objective 1.	Approved but not contracted	
DAE2010-04	Monitoring the trawl footprint for deepwater fisheries	1. To estimate the 2009/10 trawl footprint and map the spatial and temporal distribution of bottom contact trawling throughout the EEZ between 1989/90 and 2009/10. 2. To produce summary statistics, for major deepwater fisheries and the aggregate of all deepwater fisheries, of the spatial extent and frequency of fishing by year, by depth zone, by fishable area, and by habitat class, and to identify any trends or changes.	Ongoing analysis	Black <i>et al.</i> In Press
DEE2010-06	Design a camera / transect study	1. To design and provide indicative costs for a programme to monitor trends in deepwater benthic habitats and communities. 2. To explore the feasibility of using existing trawl and acoustic surveys to capture data relevant to monitoring trends in deepwater benthic habitats and communities.	Ongoing analysis	
BEN2009-02	Monitoring recovery of benthic communities in Spirits Bay	1. To survey Spirits Bay and Tom Bowling Bay benthic invertebrate communities according to the monitoring programme designed in ENV2005/23. 2. To assess changes in benthic communities inside and outside the closed area since 1997.	In the process of publication	Tuck & Hewitt In Press
Internally funded 1	SPRFMO	1. To develop detection criteria for measuring trawl impacts on vulnerable marine ecosystems in high sea fisheries of the South Pacific Ocean	Completed	Parker <i>et al.</i> 2009a
Internally funded 2	SPRFMO	1. To document protection measures implemented by New Zealand for vulnerable marine ecosystems in the South Pacific Ocean	Completed	Penney <i>et al.</i> 2009
Internally funded 3	CCAMLR	1. An Impact Assessment Framework for Bottom Fishing Methods in the CCAMLR Convention Area	Completed	Sharp <i>et al.</i> 2009
Internally funded 4	SPRFMO	1. to develop a bottom Fishery Impact Assessment: Bottom Fishing Activities by New Zealand Vessels Fishing in the High Seas in the SPRFMO Area during 2008 and 2009	Completed	Ministry of Fisheries 2008

BENTHIC IMPACTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
IFA2008-04	Guide for the rapid identification of material in the process of managing Vulnerable Marine Ecosystems	To produce a guide for the rapid identification of material in the process of managing Vulnerable Marine Ecosystems	Completed	Tracey <i>et al.</i> 2008
IFA2007-02	Development of a Draft New Zealand High-Seas Bottom Trawling Benthic Assessment Standard	1. To generate data summaries and maps of New Zealand's recent historic high-seas bottom trawling catch and effort in the proposed convention area of the South Pacific Regional Fisheries Management Organization (SPRFMO). 2. To map vulnerable marine ecosystems (VMEs) in the SPRFMO area. 3. To develop a draft standard for assessment of benthic impacts of high-seas bottom trawling on VMEs in the proposed SPRFMO convention area.	Completed	Parker 2008
BEN2007-01	Assessing the effects of fishing on soft sediment habitat, fauna, and processes	1. To design and test sampling and analytical strategies for broad-scale assessments of habitat and faunal spatial structure and variation across a variety of seafloor habitats. 2. To design and carry out experiments to assess the effects of bottom trawling and dredging on benthic communities and ecological processes important to the sustainability of fishing at scales of relevance to fishery managers.	Ongoing analysis	
BEN2006-01	Mapping the spatial and temporal extent of fishing in the EEZ	1. To update maps and develop GIS layers of fishing effort from project ENV2000/05 to show the spatial and temporal distribution of mobile bottom fishing throughout the EEZ between 1989/90 and 2004/05. 2. To produce summary statistics of major fisheries and the aggregate of all bottom impacting fisheries in terms of the extent and frequency of fishing by year, by depth zone, by fishable area, and, to the extent possible, by habitat type. 3. To identify and document any major trends or changes in fishing effort or fishing behaviour. 4. To identify, discuss the implications of, and make recommendations on data quality and other problems with current reporting systems that complicate characterisation and quantification of bottom fishing effort. 5. To integrate information on the distribution, frequency, and magnitude of fishing disturbance with habitat characteristics throughout the EEZ, using information stored in national databases, expert opinion, and the MEC.	Completed	Baird <i>et al.</i> 2009; 2011; Baird & Wood 2010; Leathwick <i>et al.</i> 2010; 2012

BENTHIC IMPACTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2005-15	Information for managing the Effects of Fishing on Physical Features of the Deep-sea Environment	1. To provide an updated database that identifies all known seamounts in the "New Zealand region", encompassing the area from 24°00' – 57°30'S, 157°00'E – 167°00'W. The database will catalogue relevant data (e.g. physical, biological, location, fishing effort) for individual seamounts. 2. To identify indicators and measures suitable for the assessment of risk pertaining to the effects of fishing disturbance on the benthic biota of seamounts, and review suitable ecological risk assessment methods, that can be derived or utilise information contained within the seamount database.	Completed	Rowden et al. 2008; Clark et al. 2010b
ENV2005-16	Investigate the Effects of Fishing on Physical Features of the Deep-sea Environment	1. To monitor changes in fauna and habitats over time on selected UTFs in the Chatham Rise area that have a range of fishing histories. 2. To continue development of the risk assessment model to predict the effects of fishing, and provide options for the management of UTF ecosystems.	Completed	Clark et al. 2010a; b; c; 2011
ENV2005-17	Estimation of non-target fish catch and both target and non-target fish discards in jack mackerel trawl fisheries	1. To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded, in the trawl fisheries for jack mackerel for the fishing years 2001/2002 to 2004/05 using data from Mfish observers and commercial fishing returns.	Completed	Anderson 2007a
ENV2005-18	Estimation of non-target fish catch and both target and non-target fish discards in orange roughy trawl fisheries	1. To estimate the quantity of non-target fish species caught, and the target and non-target fish species discarded, in the trawl fisheries for orange roughy for the fishing years 1999/2000 to 2003/04 using data from Scientific Observers and commercial fishing returns.	Completed	Anderson 2009a

BENTHIC IMPACTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2005-20	Benthic invertebrate sampling and species identification in trawl fisheries	1. To produce identification guides for benthic invertebrate species encountered in the catches of commercial and research trawlers.	Completed	Tracey et al. 2007; Williams et al. 2010; Clark et al. 2009
ENV2005-23	Monitoring recovery of the benthic community between North Cape and Cape Reinga	1. To design a monitoring programme that will provide the following quantitative estimates: i) Estimates of the nature and extent of past fishing impacts on the benthic community between North Cape and Cape Reinga; ii) Estimates of change over time in areas previously fished but subsequently closed to fishing. Estimated parameters will include indices representing biodiversity, community composition, and biogenic structure; iii) Estimates of change over time in areas environmentally comparable to those assessed in (ii), above, but subject to ongoing fishing impacts; and iv) Estimates of change over time in areas comparable to those above, but not impacted by fishing (if any such areas can be found).	Completed	Tuck et al. 2010
ZBD2005-04	Information on benthic impacts in support of the Foveaux Strait Oyster Fishery Plan	1. To assess the distribution- vulnerability to disturbance- and ecological importance of habitats in Foveaux Strait- and describe the spatial distribution of the Foveaux Strait oyster fishery relative to those habitats. 2. To assemble and collate existing information on the Foveaux Strait system between the Solander Islands and Ruapuke Island or other area to be agreed with MFish. 3. To map- using best available information- substrate type- bathymetry- wave energy- and tidal flow in this area. 4. To assess the extent to which these data can be used to define useful functional categories that might serve as habitat classes. 5. To rank the vulnerability to fishing disturbance of habitat classes developed in Objective 3 using approximate regeneration times. 6. To describe the functional role and ecosystem services provided by each habitat class developed in Objective 3- including an assessment of the relative importance of each to overall ecosystem function and productivity. 7. To describe the spatial pattern and intensity of dredge fishing for Foveaux Strait oysters over the past 10 fishing years and relate this to natural disturbance regimes and habitat classes developed in Objective 3. 8. To carry out a qualitative video survey of benthic habitats in Foveaux Strait- both within the established commercial oyster fishery area and areas outside the fishery area but within OYU 5.	Completed	Michael et al. 2006

BENTHIC IMPACTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2005-15	Information on benthic impacts in support of the Coromandel Scallops Fishery Plan	<ol style="list-style-type: none"> <li>1. To assemble and collate existing information on the coromandel Scallop Fishery between cape Rodney and Town Point or other, wider area to be agreed with Mfish.</li> <li>2. To map, using best available information, substrate type, bathymetry, wave energy, and tidal flow in this area.</li> <li>3. To assess the extent to which data can be used to define useful functional categories that might serves as habitat classes.</li> <li>4. To rank the vulnerability of fishing disturbance of habitat classes developed in Objective 3 using approximate regeneration times.</li> <li>5. To describe the functional role and ecosystem services provided by each habitat class developed in Objective 3, including an assessment of the relative importance of each to overall ecosystem function and productivity.</li> <li>6. To describe the spatial pattern and intensity of dredge and trawl fishing within the Coromandel scallop fishery over the past 15 fishing years and relate this to natural disturbance regimes and habitat classes developed in Objective 3.</li> </ol>	Completed	Tuck et al. 2006a; b
ZBD2005-16	Information on benthic impacts in support of the Southern Blue Whiting Fishery Plan	<ol style="list-style-type: none"> <li>1. To assemble and collate existing information on the Southern Blue Whiting fishery in SBW6A, SBW6B, SBW6I, and SBW6R or other wider area to be agreed with MFish</li> <li>2. To map, using best available information, substratum type, bathymetry, wave energy, tides, and ocean currents in these areas</li> <li>3. To assess the extent to which these data can be used to define useful functional categories that might serve as habitat categories.</li> <li>4. To rank the vulnerability to fishing disturbance of habitat classes developed in Objective 3 using approximate regeneration times.</li> <li>5. To describe the functional role and ecosystem services provided by each habitat class developed in Objective 3, including an assessment of the relative importance of each to overall ecosystem function and productivity.</li> <li>6. To describe the spatial pattern and intensity of trawl fishing within the Southern Blue Whiting fishery over the past 10 fishing years and relate this to natural disturbance regimes and habitat classes developed in Objective 3.</li> </ol>	Completed	Cole et al. 2007
ENV2004-02	Estimation of New Zealand sea lion incidental captures in New Zealand Fisheries	<ol style="list-style-type: none"> <li>1. To estimate the level of New Zealand sea lion (<i>Phocartos hookeri</i>) incidental capture in New Zealand fisheries</li> </ol>	Completed	Smith & Baird 2007a
ENV2003-03	Determining the spatial extent, nature and effect of mobile bottom fishing methods	<ol style="list-style-type: none"> <li>1. To determine the spatial extent, nature and time between disturbances of mobile bottom fishing methods in the Chatham Rise trawl fisheries.</li> </ol>	Completed	Baird et al. 2006

*AEBAR 2012: Appendices: Past projects*

**BENTHIC IMPACTS** *continued*

<b>Project Code</b>	<b>Project Title</b>	<b>Specific Objectives</b>	<b>Status</b>	<b>Citation/s</b>
ENV2002-04	Benthic invertebrate sampling and specific identification in trawl fisheries	1. To quantify and map the benthic invertebrate species incidental catch in commercial and research trawling throughout the New Zealand EEZ	Completed	Tracey et al. 2005
ENV2001-07	Reducing bycatch in scampi trawl fisheries	1. Collate and review the international literature on methods of reducing bycatch in crustacean trawl fisheries. 2. Review and analyse the data from New Zealand studies. 3. Develop recommendations on future approaches to reducing bycatch in the New Zealand scampi fishery, including some general thoughts on the experimental design of field trials.	Completed	Hartill et al. 2006
ENV2001-09	The effects of mobile bottom fishing gear on benthic-pelagic coupling	To describe any effects of fishing that might modify benthic-pelagic coupling (a complex, interlinked suite of processes transferring energy, oxygen, carbon, and nutrients between pelagic and benthic systems), to consider the scale of such possible effects, and to put the summary in a New Zealand context.	Completed	Cryer et al. 2004
ENV2001-15	The effects of bottom impacting trawling on seamounts	1. To design a programme in New Zealand waters previously trawled and now closed to trawling to monitor the rate of regeneration of benthic communities on seamounts.	Completed	Clark & O'Driscoll 2003; Clark & Rowden 2009
OYS2001-01	Foveaux Strait oyster stock assessment	1. To carry out a survey and determine the distribution and absolute abundance of pre-recruit and recruited oysters in both non-commercial and commercial areas of Foveaux Strait. The target coefficient of variation (c.v.) of the estimate of absolute recruited abundance is 20%. 2. To estimate the sustainable yield for the areas of the commercial oyster fishery in Foveaux Strait for the year 2002 oyster season. 3. To identify and count benthic macro-biota collected during the dredge survey.	Completed	Rowden et al. 2007
ENV2000-05	Spatial extent, nature and impact of mobile bottom fishing methods in the New Zealand EEZ	1. To determine the spatial extent, nature and impact of mobile bottom fishing methods within the New Zealand EEZ.	Completed	Cryer and Hartill 2002; Baird et al. 2002
ENV2000-06	Review of technologies and practices to reduce bottom trawl bycatch and seafloor disturbance in New Zealand	Objectives unknown	Completed	Booth et al. 2002; Beentjes & Baird 2004

**BENTHIC IMPACTS** *continued*

<b>Project Code</b>	<b>Project Title</b>	<b>Specific Objectives</b>	<b>Status</b>	<b>Citation/s</b>
ENV98-05	The effects of fishing on the benthic community structure between North Cape and Cape Reinga	1. To determine the effects of fishing on the benthic community structure between North Cape and Cape Reinga.	Completed	Cryer et al. 2000

## ECOSYSTEM EFFECTS

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2012-01	A literature review of Nitrogen levels and adverse ecological effects in embayments in temperate regions.	1. To complete a literature review of Nitrogen levels and adverse ecological impacts from temperate embayments in order to assist aquaculture consenting authorities in determining at what concentration of Nitrogen adverse effects may be expected.	Approved but not contracted	
ZBD2012-06	Ocean status: trends in NZ marine environment and Tier 1 statistic	1. To provide an up to date overview of climatic trends and cycles and how they affect New Zealand oceanographic conditions, and highlight key changes since the previous assessment. 2. To identify candidate oceanographic variables for potential development as part of the proposed Tier 1 Statistic, Atmospheric and Ocean Climate Change	Approved but not contracted	
ANT2011-01	Antarctic fisheries	1. To develop, implement and refine approaches for assessing the stock status of toothfish ( <i>Dissostichus</i> spp.) in the Ross Sea region. 2. To develop, implement and refine approaches for assessing and monitoring the status of non-target fish species, and dependent and related species. 3. To develop, implement and refine approaches for understanding and managing the ecological relationships between the toothfish ( <i>Dissostichus</i> spp.) fishery and the Ross Sea ecosystem.	Ongoing analysis	
DAE2010-01	Taxonomic identification of benthic specimens	1. To identify benthic invertebrates in samples taken during research trawls and by Observers on fishing vessels. 2. To update relevant databases recording the catch of invertebrates in research trawls and commercial fishing.	Ongoing analysis	
DAE2010-03	Ecological risk assessment for deepwater stocks	1. To undertake a qualitative (level 1) risk assessment for tier 3 fishstocks within the deepwater fisheries plan.	Ongoing analysis	
DEE2010-04	Development of a methodology for Environmental Risk Assessments for deepwater fisheries	To review approaches to Ecological Risk Assessments (ERA) and methods available for deepwater fisheries both QMS and non-QMS. 2. To develop and recommend a generic, cost effective, method for ERA in deepwater fisheries by using or modifying methods identified in Objective 1.	Completed	Clark et al. In Press
DEE2010-05	Development of a suite of environmental indicators for deepwater fisheries	1. To review the literature and hold a workshop to recommend a suite of ecosystem and environmental indicators that will contribute to assessing the performance of deepwater fisheries within an environmental context. 2. To examine available data and design a data collection programme to enable future calculation of the indicators identified in Specific Objective 1.	Ongoing analysis	
ENV2010-03	Habitats of particular significance for inshore finfish fisheries management	1. To review the literature to determine the most important juvenile or reproductive (spawning, pupping or egg-laying) areas for inshore finfish target species. 2. To use a gap analysis to prioritize areas for future research concerning the important juvenile or reproductive (spawning, pupping or egg-laying) areas for target inshore finfish fisheries	Ongoing analysis	

ECOSYSTEM EFFECTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2010-05A&B and SEA 2010-15	Habitats of particular significance for fisheries management: shark nursery areas	<ol style="list-style-type: none"> <li>1. Identify, from the literature, important nursery grounds for rig in estuaries around mainland New Zealand.</li> <li>2. Design and carry out a survey of selected estuaries and harbours around New Zealand to quantify the relative importance of nursery ground areas.</li> <li>3. Identify threats to these nursery ground areas and recommend mitigation measures.</li> </ol>	In the process of publication	Francis et al. 2012; Jones et al. In Press
ZBD2010-42	Development of a National Marine Environment Monitoring Programme	<ol style="list-style-type: none"> <li>1. To design a Marine Environment Monitoring Programme (MEMP) to track the physical, chemical and biological changes taking place across New Zealand's marine environment over the long term</li> <li>2. To prepare an online inventory (metadatabase) of repeated (time series) biological and abiotic marine observations/datasets in New Zealand</li> <li>3. To review, evaluate fitness for purpose, and identify gaps in the utility and interoperability of these datasets for inclusion in MEMP from both science and policy perspectives</li> <li>4. To design a MEMP that includes relevant existing data collection and proposed new time series</li> </ol>	Ongoing analysis	
ANT2009-01	Antarctic fisheries	<ol style="list-style-type: none"> <li>1. To explore the biology of fishes captured in the toothfish fishery to underpin future stock assessment and ecosystem modelling research</li> <li>2. To develop and refine stock assessment approaches for toothfish in the Ross Sea</li> <li>3. To assess the status of toothfish stocks in the Ross Sea</li> <li>4. To explore the Ross Sea toothfish fishery at an ecosystem level</li> <li>5. To review and further develop procedures for the ageing of Antarctic toothfish (<i>Dissostichus mawsoni</i>) and Patagonian toothfish (<i>D. eleginoides</i>).</li> <li>6. To review and update the species profiles for toothfish</li> <li>7. To characterise the toothfish fishery in the Ross Sea up to 2009/10</li> <li>8. To further develop toothfish biological and modelling parameters</li> <li>9. To assess the status of the Ross Sea toothfish stock(s) with respect to CCAMLR performance measures</li> <li>10. To further develop approaches to assessing the status of skates in the Ross Sea region with respect to CCAMLR performance measures</li> <li>11. Further develop the SPM approach</li> <li>12. To develop new approaches and refine existing approaches to understanding the impacts of fishing on potential VMEs</li> <li>13. To further develop ecosystem monitoring through the analysis of the diet of toothfish in the north and slope fisheries.</li> <li>14. To refine the draft data collection plan for the Ross Sea region fisheries and undertake associated preliminary reviews of fishery and observer performance against targets immediately post-season</li> </ol>	Completed	Parker & Bowden 2010; Parker et al. 2009c; Tracey et al. 2010

ECOSYSTEM EFFECTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2009-04	Trends in relative mesopelagic biomass using time series of acoustic backscatter data from trawl surveys	1. To evaluate relative changes in abundance of mesopelagic fish and other biological components from acoustic records collected during Chatham Rise and Sub-Antarctic trawl surveys. 2. To explore links between trends in mesopelagic biomass and climate variables and variations, and condition indices of commercial species in the Chatham Rise and Sub-Antarctic areas.	Completed	O'Driscoll et al. 2011
ENV2009-07	Habitats of particular significance for fisheries management: kaipara harbour	1. Collate and review information on the role and spatial distribution of habitats in the Kaipara Harbour that support fisheries production. 2. Assess historical, current, and potential anthropogenic threats to these habitats that could affect fisheries values, including fishing and land-based threats. 3. Design and implement cost-effective habitat mapping and monitoring surveys of habitats of particular significance for fisheries management in the Kaipara Harbour.	Ongoing analysis	
GMU2009-01	Spatial Mixing of GMU1 using Otolith Microchemistry	1. To determine the level of spatial mixing and connectivity of grey mullet ( <i>Mugil cephalus</i> ) populations using otolith microchemistry. 2. To collect and analyse the chemical composition of grey mullet otoliths. 3. To analyse the otoliths collected under Objective 1 to determine if the samples can be spatially separated.	Ongoing analysis	
IPA2009-11	Trophic studies publication of review	1. To publish the comprehensive review of New Zealand-wide trophic studies completed in 2000 that was prepared by NIWA.	Completed	Stevens et al. 2011
ANT2008-03	Ecosystem effects of fishing in the Ross Sea	To evaluate the VMEIO classification accuracy of observers, identify potential causes for taxonomic confusion, and make recommendations for improvements in the classification guide, observer training, and in the data collection protocols	Completed	Parker et al. 2009b; 2010
AQE2008-02	Review of ecological effects of farming shellfish and other species	1. To collate and review information on the ecological effects of farming mussels ( <i>Perna canaliculus</i> ), including offshore mussel farming and spat catching, in the New Zealand marine environment. 2. To collate and review information on the ecological effects of farming oysters in the New Zealand marine environment. 3. To collate and review information on the ecological effects of farming species other than mussels ( <i>Perna canaliculus</i> ), oysters, and finfish, in the New Zealand marine environment.	Completed	Keeley et al. 2009

ECOSYSTEM EFFECTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
TOH2008-01	Distribution and abundance of Toheroa	<ol style="list-style-type: none"> <li>1. To estimate the size structure and absolute abundance of toheroa on Oreti Beach, during February 2009. The target c.v. for the estimate of absolute abundance of legal sized toheroa ( 100 mm shell length) is 20%.</li> <li>2. To describe changes in the size structure and absolute abundance of toheroa on Oreti Beach by comparing the results from this work with those from previous surveys.</li> <li>3. To estimate the size structure and absolute abundance of toheroa on Bluecliffs Beach, during February 2009. The target c.v. for the estimate of absolute abundance of legal sized toheroa ( 100 mm shell length) is 20%.</li> <li>4. To describe changes in the size structure and absolute abundance of toheroa on Bluecliffs Beach by comparing the results from this work with those from previous surveys.</li> </ol>	Completed	Beentjes 2010
IFA2008-08	Inputs to the Ross Sea bioregionalisation	<ol style="list-style-type: none"> <li>1. To produce one or more benthic invertebrate classifications of the Ross Sea region;</li> <li>2. To use fishery catch data to examine spatial distributions of major demersal fish species;</li> <li>3. To prepare other biological or environmental spatial data layers for use in the Ross Sea workshop.</li> </ol>	Completed	Pinkerton et al. 2009a
ANT2007-01	Biology of fishes in the toothfish fishery	<ol style="list-style-type: none"> <li>3. To develop an identification guide for observers of benthic invertebrate species (especially sponges, corals etc) caught in the Ross Sea region fisheries.</li> </ol>	Completed	Parker et al. 2008
BEN2007-01	Assessing the effects of fishing on soft sediment habitat, fauna, and processes	<ol style="list-style-type: none"> <li>1. To design and test sampling and analytical strategies for broad-scale assessments of habitat and faunal spatial structure and variation across a variety of seafloor habitats.</li> <li>2. To design and carry out experiments to assess the effects of bottom trawling and dredging on benthic communities and ecological processes important to the sustainability of fishing at scales of relevance to fishery managers.</li> </ol>	Ongoing analysis	
BEN2007-05	Risk assessment framework for assessing fishing & other anthropogenic effects on coastal fisheries	<ol style="list-style-type: none"> <li>1. To collate existing information on the distribution, intensity, and frequency of anthropogenic disturbances in the coastal zone that could be used in a risk assessment model to estimate their likely aggregate effect on ecosystem function across habitats and over different scales of ecosystem functioning and biological organization.</li> <li>2. To develop a risk assessment framework in conjunction with a variety of stakeholders and environmental scientists.</li> </ol>	Completed	MacDiarmid et al. 2012
ENH2007-01	Stock enhancement of blackfoot paua	<ol style="list-style-type: none"> <li>1. To assess the survival rate of enhanced paua from introduction into the wild through to harvest.</li> <li>2. To assess the genetic diversity of hatchery spawned juvenile paua bred for enhancement purposes.</li> <li>3. To assess interactions between introduced and wild paua populations and to recommend research and monitoring to quantify those impacts that are potentially adverse.</li> </ol>	Ongoing analysis	

ECOSYSTEM EFFECTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2007-04	Climate and Oceanographic Trends Relevant to New Zealand Fisheries	1. To summarise, for fisheries managers, climatic and oceanographic fluctuations and cycles that affect productivity, fish distribution and fish abundance in New Zealand.	Completed	Hurst et al. 2012
ENV2007-06	Trophic Relationships of Commercial Middle Depth Species on the Chatham Rise	1. To quantify the inter-annual variability in the diets of hoki, hake and ling on the Chatham Rise 1992–2007 2. To quantify seasonal dietary cycles for hoki, hake and ling that have been collected from the commercial fleet throughout the year	Completed	Horn & Dunn 2010
HAB2007-01	Biogenic habitats as areas of particular significance for fisheries management	1. To collate and review available information on the location, value, functioning, threats to, and past and current status of biogenic habitats that may be important for fisheries production in the New Zealand marine environment. 2. To identify information gaps, in the New Zealand context, and recommend measures to address those important to an ecosystem approach to fisheries management	Ongoing analysis	
IPA2007-07	Land Based Effects on Coastal Fisheries	1. To review and collate scientific knowledge and research on the impacts of land-based activities on coastal fisheries and biodiversity	Completed	Morrisson et al. 2009
TOH2007-03	Toheroa Abundance	1. To investigate variations in the abundance of toheroa. 2. To investigate sources of mortality of toheroa and factors affecting the recruitment of toheroa	Ongoing analysis	Williams et al. In Press
ENV2006-04	Ecosystem indicators for New Zealand fisheries	1. To carry out a literature review of potential fish-based ecosystem indicators and identify a suite of indicators to be tested in Objective 2 2. To test a suite of fish-based ecosystem indicators (identified by Objective 1) on existing trawl survey time series in New Zealand. The utility of these indicators for monitoring the effects of fishing in New Zealand should also be evaluated	Completed	Tuck et al. 2009
GBD2006-01	DNA database for commercial marine fish and invertebrates	1. To collect DNA sequences for vouchered specimens of commercially important marine fishes and submit the DNA data to the international Barcode of Life Database (BOLD). 2. To collect DNA sequences for vouchered specimens of commercially important marine invertebrates and submit the DNA data to the international Barcode of Life Database (BOLD). Note: The funding was limited to \$60 000 for this Objective. Therefore MFish agreed to omit the invertebrate species (Objective 2) from this project and reduce the number of fish species sequenced from 100 to 80 (up to 5 specimens per species). During the course of the project MFish staff asked NIWA to identify smoked eel product, suspect shark fillets, and possible paua slime with DNA markers, consequently the project was modified to accommodate these requests	Completed	No reports specified as required output

ECOSYSTEM EFFECTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
SAP2006-06	West coast south island review	<ol style="list-style-type: none"> <li>1. To publish a review document summarising oceanic and environmental research information particularly relevant to hoki- but also other fisheries- that spawn off Westland in winter</li> <li>2. Update the draft chapters prepared in 2004 by oceanographers- modellers and scientists towards the overall objective</li> <li>3. Incorporate a section on other west coast spawning fisheries</li> </ol>	Completed	Bradford-Grieve & Livingston 2011
IPA2006-08	Review of the Ecological Effects of Marine Finfish aquaculture: Final Report	<ol style="list-style-type: none"> <li>1. Summarise and review existing information on ecological effects of finfish farming on the marine environment in New Zealand and overseas</li> </ol>	Completed	Forrest et al. 2007
ANT2005-02	Aspects of the biology of fishes in the toothfish fishery	<ol style="list-style-type: none"> <li>1. Estimate length and age at maturity for Antarctic toothfish in the Ross Sea</li> <li>2. Examine TOA length at age by depth and area</li> <li>3. Estimate biological parameters for TOA (M, growth rates corrected for selectivity, h, r)</li> <li>4. Determine stock structure of TOA based on parasite data</li> <li>5. Determine length-weight relationships, diet, reproduction, age and growth of C.dewitti</li> <li>6. ID and speciation of Antarctic skates</li> <li>7. Develop an ID guide for scientific Observers of fish in the Ross Sea fishery</li> <li>8. Identify heavy metal contents of selected fish species in the Ross Sea fishery</li> </ol>	Completed	McMillan et al. 2007; Smith et al. 2007; Sutton et al. 2006
ANT2005-04	Ecosystem modelling of the Ross Sea	<ol style="list-style-type: none"> <li>1. Carry out stable isotope analysis of TOA and 3 key fish prey to determine trophic links</li> <li>2. Determine squid diet by analysis of squid beaks for stable isotope analysis</li> <li>3. Participate in the design of an IPY survey</li> <li>4. Participate in EMM as required</li> </ol>	Completed	Pinkerton et al. 2007b
ENV2005-08	Experimental design of a programme of indicators	<ol style="list-style-type: none"> <li>1. To assess the utility/feasibility of using demographic information to assess the effects of fishing on seabird populations.</li> <li>2. To identify population indicators and to provide sampling protocols and experimental design for selected high to medium priority seabird populations.</li> <li>3. To recommend experimental protocols for sampling of selected seabird populations in New Zealand influenced by fisheries mortality, employing robust-design methodology and including recommendations for inclusions of data into Ministry of Fisheries databases.</li> </ol>	Completed	MacKenzie & Fletcher 2010
SAM2005-02	Effects of climate on commercial fish abundance	To examine the possible effects of climate on fishery yields and abundance indices for commercial fisheries around New Zealand	Completed	Dunn et al. 2009

ECOSYSTEM EFFECTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ANT2004-01	Characterisation of the toothfish fishery	<ol style="list-style-type: none"> <li>1. update descriptive analysis of toothfish fishery in the Ross Sea to 04/05</li> <li>2. analyse age, LF and sex ratio for toothfish and rattails for 04/05</li> <li>3. update and refine the CPUE for TOA in Ross Sea for 04/05</li> <li>4. determine diet of sub-adult TOA in the Ross Sea</li> <li>5. review the TOA parasite collection protocol</li> <li>6. document TOA tagging protocol</li> <li>7. review approaches to monitoring and assessing rattails and skates in the Ross Sea</li> <li>8. descriptive analysis of stake tagging programme in the Ross Sea</li> <li>9. determine factors affecting bycatch of rattails and skates between vessels</li> <li>10. carry out risk assessment for <i>M. whitsoni</i> and <i>A. georgina</i> in the Ross Sea</li> </ol>	Completed	Smith & Notman 2005; Stevens 2006
ANT2004-05	Modelling of the ecosystem effects of fishing in the Ross Sea	<ol style="list-style-type: none"> <li>1. develop an effects of fishing model based around toothfish fishery</li> <li>2. investigate possible consequences of different management strategies</li> <li>3. make recommendations for future research to decrease uncertainty in the model</li> </ol>	Completed	Pinkerton et al. 2005; 2006
HOK2004-01	Hoki Population modelling and stock assessment	<ol style="list-style-type: none"> <li>2. To investigate the prediction of year class strength from environmental variables.</li> </ol>	Completed	Francis et al. 2005
AQE2003-01	Effects of aquaculture and enhancement stock sources on wild fisheries resources and the marine environment.	<ol style="list-style-type: none"> <li>1. To identify, discuss the effects and qualitatively assess the risks of aquaculture and enhancement stocks improved by hatchery technology on New Zealand's wild fisheries resources and the marine environment.</li> <li>2. To identify, discuss the effects and qualitatively assess the risks associated with the translocation of aquaculture and enhancement stocks on New Zealand's wild fisheries resources and the marine environment.</li> <li>3. To make recommendations on priority issues, risks, or research to be undertaken, as a result of information discussed and evaluated in objectives 1-2.</li> </ol>	Completed	Speed 2005
EEL2003-01	Non-fishing mortality of freshwater eels	<ol style="list-style-type: none"> <li>1. To undertake a feasibility study on establishing an estimate of the mortality of eels caused by hydroelectric turbines and other point sources of mortality caused by human activity.</li> </ol>	Completed	Bentjees et al. 2005
MOF2003-01	The implications of marine reserves for fisheries resources and management in the New Zealand context	Objectives unknown	Completed	Speed et al. 2006
ENV2002-03	Beach cast seaweed review	<ol style="list-style-type: none"> <li>1. To collate existing information on the role of beach-cast seaweed in coastal ecosystems to assess the nature and extent of the impacts that the removal of beach cast seaweed may have on the marine environment.</li> <li>2. On the basis of the review in Specific Objective 1 above, to identify key research gaps related to any marine environment impacts that the removal of beach cast seaweed may have.</li> </ol>	Completed	Zemke-White et al. 2005

ECOSYSTEM EFFECTS *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ENV2002-07	Energetics and trophic relationships of important fish and invertebrate species	1. To quantify food webs supporting important fish and invertebrate species	Completed	Livingston 2004
CRA2000-01	Rock lobster stock assessment	Objective 11: To conduct a desktop study to identify and explore data needs associated with managing the effects of rock lobster fishing on the environment.	Completed	Breen 2005
ENV2000-04	Identification of areas of habitat of particular significance for fisheries management within the New Zealand EEZ	1. To review literature and existing data for all significant fish species, including all QMS species, encountered from the 200 1500 m contour within the New Zealand EEZ to: a) determine areas of important juvenile fish habitat; b) determine areas of importance to spawning fish populations; and c) determine areas of importance for shark populations for pupping or egg laying. 2. To review literature and existing data for all significant pelagic fish species (excluding highly migratory species) encountered within the New Zealand EEZ to: a) determine areas of important juvenile fish habitat; b) determine areas of importance to spawning fish populations; and c) determine areas of importance for shark populations for pupping or egg laying 3. To review literature and existing data for all significant marine invertebrate species encountered within the New Zealand EEZ to: a) determine areas of important juvenile habitat; and b) determine areas of importance to spawning populations	Completed	O'Driscoll et al. 2003
MOF2000-02A	Future research requirements for the Ross Sea Antarctic toothfish ( <i>Dissostichus mawsoni</i> ) fishery.	Objectives unknown	Completed	Hanchet 2000
ENV99-03	Identification of areas of habitat of particular significance for fisheries management within the NZ EEZ.	1. To determine areas of habitat of importance to fisheries management within the New Zealand EEZ for selected fish species in selected areas	Completed	Hurst et al. 2000
ENV99-04	A framework for evaluating spatial closures as a fisheries management tool	Unknown	Completed	Bentley et al. 2004
No project number	The fishery for freshwater eels ( <i>Anguilla</i> spp.) in New Zealand	Objectives unknown	Completed	Jellyman 1994

## BIODIVERSITY

Project Code	Project Title	Specific Objectives	Status	Citation/s
SRP2011-02	IDG 2009-01 field guide completion	1. IDG 2009-01 field guide completion	Completed	McMillan 2011 a;b;c
ZBD2010-39	Improved benthic invertebrate species identification in trawl fisheries	1. To revise and update the document "A guide to common deepsea invertebrates in New Zealand waters (second edition)" to allow a third edition of this guide to be printed	Completed	Tracey et al. 2011a
ZBD2010-40	Predictive modelling of the distribution of vulnerable marine ecosystems in the South Pacific Ocean region.	1. To develop & test spatial habitat modelling approaches for predicting distribution patterns of vulnerable marine ecosystems in the convention Area of the South Pacific Regional Fisheries Management Organisation with agreed international partners. 2. To collate datasets and evaluate modelling approaches which are likely to be useful to predict the distribution of vulnerable marine ecosystems in the South Pacific Ocean region.	Ongoing analysis	
ZBD2010-41	Ocean acidification in fisheries habitat	1. To assess the risks of ocean acidification to deep sea corals and deepwater fishery habitat 2. To determine the carbonate mineralogy of selected deep sea corals found in the New Zealand region 3. To assess the distribution of deep sea coral species in the New Zealand region relative to improved knowledge of current and predicted aragonite and calcite saturation horizons, assessment of potential locations vulnerable to deep water upwelling 4. Through a literature search and analysis, determine the most appropriate tools to age and measure the effects of ocean acidification on deep sea habitat-forming corals, and recommend the best approach for future assessments of the direct effects	Ongoing analysis	Tracey et al. 2011b
IPA2009-14	Bryozoan identification guides	1. For each of ~50 species of common bryozoans, provide photos and text to allow for identification. Provide information on distribution and habitat (as far as is known) and further references for each species and on bryozoans as a whole. 2. Submit these data for publication in the Ministry of Fisheries series New Zealand Aquatic Environment and Biodiversity Research.	Completed	Smith & Gordon 2011
ZBD2009-03	To evaluate the vulnerability of New Zealand rhodolith species to environmental stressors and to characterise diversity of rhodolith beds.	1. To characterise the distribution and physical characteristics of two New Zealand rhodolith beds and characterise the associated biodiversity. 2. To measure the growth rates and evaluate the vulnerability of New Zealand species of rhodoliths to environmental stressors.	Completed	Nelson et al. 2012

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2009-10	Multi-species analysis of coastal marine connectivity	<ol style="list-style-type: none"> <li>1. Determine overall patterns of regional connectivity in a broad range of NZ coastal marine organisms to define the geographic units of genetic diversity for protection and the dispersal processes that maintain this diversity.</li> <li>2. Review previous studies of marine connectivity and population genetics in NZ coastal organisms to determine the preliminary range of patterns observed and the principal gaps (taxonomic geographic and ecological) in our understanding.</li> <li>3. In a range of invertebrate and vertebrate marine organisms determine geographic patterns of genetic variation using standardised sampling and molecular techniques.</li> <li>4. Analyse data across past and present studies to reveal both common and unique patterns of connectivity around the NZ coastline and the locations of common barriers to dispersal.</li> </ol>	Completed	Gardner et al. 2010
ZBD2009-13	Ocean acidification impact on key nz molluscs	<ol style="list-style-type: none"> <li>1. Controlled laboratory experiments will be used to determine the effect of pCO<sub>2</sub> levels that are predicted to occur in NZ waters over the next few decades on appropriate life history stages of at least two key NZ mollusc species. A number of response variables will be assessed.</li> <li>2. Implications of these responses to the local and broader ecosystems will be assessed.</li> </ol>	Ongoing analysis	Cummings 2011; Cummings et al. 2011b
ZBD2009-25	Predicting impacts of increasing rates of disturbance on functional diversity in marine benthic ecosystems	<ol style="list-style-type: none"> <li>1. Further develop the landscape ecological model of disturbance/recovery dynamics in marine benthic communities, incorporating habitat connectivity, based on existing model by Lundquist, Thrush, and Hewitt.</li> <li>2. Predict impacts of increasing rates of disturbance on rare species abundance, functional diversity, relative importance of biogenic habitat structure, and ecosystem productivity.</li> <li>3. Use literature and expert knowledge to quantify rare species abundance, biomass, functional diversity, habitat structure, and productivity of various successional community types in the model.</li> <li>4. Field test predictions of the model in appropriate marine benthic communities where historical rates of disturbance are known, and benthic communities have been sampled.</li> </ol>	Ongoing analysis	Lundquist et al. 2010

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2008-01	Biogenic large-habitat-former hotspots in the near-shore coastal zone (50–250 m); quantifying their location, identity, function, threats and protection	<ol style="list-style-type: none"> <li>1. To collect and integrate existing knowledge on biogenic habitat-formers in the &lt;5–150 m depth zone of New Zealand's continental shelf, from sources including structured fisher interviews, primary and grey literature, and other sources as available.</li> <li>2. Using the findings of Objective 1, design and deploy a series of sampling voyages to selected locations, to map and characterise locations of significant biogenic structure (either still existing, or historical), and collect relevant biological samples (both through visual census, and physical collection).</li> <li>3. Process and analyse the samples collected in Objective 2, to provide a hierarchical, quantitative description of the biogenic habitats and associated species encountered.</li> <li>4. Using the findings from Objective 1–3, assess the present status, likely extent, ecological role, and threats to, biogenic habitat formers in the &lt;5–150 m depth zone. This should include a spatial modelling and risk assessment framework. Integrate (as appropriate) with other information sources and/or approaches that may exist by the year 2010/11.</li> </ol>	Ongoing analysis	
ZBD2008-05	Macroalgal diversity associated with soft sediment habitats	<ol style="list-style-type: none"> <li>1. Conduct a targeted collection programme across diverse soft sediment environments to develop a permanent reference collection of representative macroalgae.</li> <li>2. Examine algal distribution in soft sediment habitats in relation to selected environmental variables.</li> <li>3. Prepare an annotated checklist of macroalgae found in soft sediment environments in the New Zealand region.</li> </ol>	Completed	Neill et al. 2012
ZBD2008-07	Carbonate sediments: the positive and negative effects of land-coast interactions on functional diversity	<ol style="list-style-type: none"> <li>1. To quantify shifts in community structure and functional diversity in mollusc dominated habitats along gradients associated with an estuary-coast interface in two locations.</li> <li>2. To characterise the influence of estuary-derived food sources across these gradients for key species.</li> <li>3. To measure changes in growth of key species in relation to changes in food supply and land-derived sediment impacts.</li> <li>4. To quantify carbon and nitrogen uptake and tissue turnover rates of key species in laboratory experiments.</li> </ol>	Completed	Thrush et al. In Press; Savage et al. 2012

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2008-11	Predicting changes in plankton biodiversity and productivity of the EEZ in response to climate change induced ocean acidification	<ol style="list-style-type: none"> <li>1. To document the spatial and inter-annual variability of coccolithophore abundance and biomass- and assess in terms of the phytoplankton abundance-biomass and community composition in sub-tropical and sub-Antarctic water.</li> <li>2. To document the seasonal and inter-annual variability of foraminifera and pteropod abundance and biomass at fixed locations in sub-tropical and sub-Antarctic water by analysis of sediment trap material from time-series data collection.</li> <li>3. To document the spatial and seasonal distribution of the key coccolithophore species- <i>Emiliana huxleyi</i>- using both archived and ongoing ingestion of satellite images of Ocean Colour- and ground-truth the reflectance.</li> <li>4. To determine the sensitivity of- and response of <i>E. huxleyi</i> and other EEZ coccolithophores to pH under a range of realistic atmospheric CO<sub>2</sub> concentrations in perturbation experiments- using monocultures and mixed populations from in situ sampling.</li> <li>5. To document the spatial variability of diazotrophs (nitrogen-fixing organisms) and associated nitrogen fixation rate- and assess in terms of phytoplankton abundance- biomass and community composition in sub-tropical waters north of the STF.</li> <li>7. To determine the sensitivity of- and response of <i>Trichodesmium</i> spp. and other diazotrophs to pH under a range of realistic atmospheric CO<sub>2</sub> concentrations in perturbation experiments using monocultures</li> </ol>	Ongoing analysis	Law et al. 2012; Boyd & Law 2011
ZBD2008-14	What and where should we monitor to detect long-term marine biodiversity and environmental changes-remote sensing, biota, context, inshore offshore workshop	<ol style="list-style-type: none"> <li>1. Identify the key questions to be addressed by long-term monitoring of marine biodiversity and environment.</li> <li>2. Identify appropriate monitoring indices, how they should be spatially distributed and their sampling frequency.</li> <li>3. Identify relevant existing monitoring programmes across the range of New Zealand agencies and science providers and identify gaps.</li> <li>4. Provide those agencies setting environmental goals/ standards or research needs (MoRST, FRST, MFish, DoC, MfE, Commissioner for the Environment) with a thorough situational analysis, including a list of priority monitoring projects/plans.</li> </ol>	Ongoing analysis	Livingston 2009

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2008-15	Continuous plankton recorder project: implementation and identification	<ol style="list-style-type: none"> <li>1. To set up a time series of annual CPR data collection by deployment from a toothfish vessel on the annual summer transit between New Zealand and the Ross Sea.</li> <li>2. To identify phytoplankton and zooplankton according to strict observation protocols determined by the SAHFOS[1] CPR Survey and SO-CPR[2].</li> <li>3. To enter species data, frequency and location along the transect into a spreadsheet that will allow spatial mapping of the plankton density and distribution.</li> <li>4. To analyse the full dataset after 5 years of data collection to: (a) determine trends in the dataset and (b) compare results with Australian datasets available through SO-CPR.</li> <li>5. To evaluate the continuation of the programme</li> </ol>	Ongoing analysis	
ZBD2008-20	Ross sea benthic ecosystem function: predicting consequences of shifts in food supply	<ol style="list-style-type: none"> <li>1. To increase understanding of Ross Sea coastal benthic ecosystem function</li> <li>2. Conduct in situ investigations into responses to and utilisation of primary food sources by key species, at two contrasting coastal Ross Sea locations</li> </ol>	Completed	Cummings & Lohrer 2011; Cummings et al. 2011a; Lohrer et al. 2012
ZBD2008-22	Acidification and ecosystem impacts in NZ and southern ocean waters (data collected during IPY).	<ol style="list-style-type: none"> <li>1. To assess the response of coccolithophorids, and their replacement by non-calcifying organisms during incubation under a range of dissolved CO<sub>2</sub> concentrations.</li> <li>2. To describe and characterise changes in abundance and biodiversity of microbial components of the samples incubated at sea under a range of dissolved CO<sub>2</sub> concentrations.</li> <li>3. To predict the likely impacts of higher acidity on foodwebs and on carbon fixation under scenarios to be encountered in the Southern Ocean under forecasted trends associated with climate change.</li> </ol>	Completed	Maas et al. 2010b
ZBD2008-23	Macroalgae diversity and benthic community structure at the Balleny Islands	<ol style="list-style-type: none"> <li>1. To describe and characterise macroalgae diversity from the Balleny Islands and the Western Ross Sea.</li> <li>2. To describe and quantify benthic community structure from one location at the Balleny Islands</li> <li>3. To complete anatomical and morphological investigations &amp; molecular sequencing required for the identification of macroalgae samples from the Balleny Islands &amp; western Ross Sea coastline to describe &amp; characterise macroalgae diversity in Balleny Islands</li> <li>4. To process and analyse samples collected at the Balleny Islands- to analyse them using ICECUBE methodology- and compare results with those from other ICECUBE sampling locations along the Ross Sea coastline</li> </ol>	Completed	Nelson et al. 2010

*AEBAR 2012: Appendices: Past projects*

**BIODIVERSITY** *continued*

<b>Project Code</b>	<b>Project Title</b>	<b>Specific Objectives</b>	<b>Status</b>	<b>Citation/s</b>
ZBD2008-27	Scoping investigation into New Zealand abyss and trench biodiversity	<ol style="list-style-type: none"> <li>1. Review what is already known of abyssal, canyon and trench faunas in NZ.</li> <li>2. Review what is already known of abyssal, canyon and trench faunas around the world.</li> <li>3. Prioritise science questions and locations for exploration.</li> <li>4. Assess NZ capacity to sample at the required depths; identify sampling equipment needs.</li> <li>5. Design a suitable vessel-based sampling programme</li> </ol>	Completed	Lörz et al. 2012
ZBD2008-50	OS2020 Chatham Rise Biodiversity Hotspots	<ol style="list-style-type: none"> <li>1. To improve understanding of the effects of trawl fishing in New Zealand on the biodiversity of seamounts- knolls and hills.</li> <li>2. To describe differences in benthic biodiversity between northwestern and eastern regions of the Chatham Rise</li> <li>3. To continue the time series of observations in the NW Chatham Rise to demonstrate recovery in terms of biodiversity</li> <li>4. To extend the observations on fished-unfished contrasts and recovery of fauna on protected seamounts to an oceanographically distinct location</li> </ol>	Completed	Clark et al. 2009
IPY2007-02	International polar year census of antarctic marine life post-voyage analysis: Ross Sea - Southern Ocean Biodiversity	<ol style="list-style-type: none"> <li>1. To measure and describe key elements of species distribution- abundance (density or biomass) &amp; biodiversity for the Ross Sea and Southern Ocean for main habitats and key functional ecosystem roles- for major groups- viruses- bacteria- archaea.....</li> <li>2. To report on the diversity of Antarctic Cephalopoda (Octopus and Squid)- including a complete inventory of taxa- &amp; reports on ontogenetic &amp; sexual variation in species- their systematics- diversity- distribution- life histories- &amp; trophic importance.</li> <li>3. To Beak/Biomass Regression Equations</li> <li>4. Life cycle determination</li> </ol>	Completed	Garcia 2010

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
IPY2007-01	International polar year census of antarctic marine life post-voyage analysis: Ross Sea - Southern Ocean Biodiversity	<ol style="list-style-type: none"> <li>1. To measure seabed depth and rugosity using the multibeam system to identify topographic features such as bottom type, iceberg scouring, seamounts etc and to determine areas for targeted benthic faunal sampling.</li> <li>2. To continue the analysis of opportunistic seabird and marine mammal distribution observations from this and previous BioRoss voyages and published records, and in relation to environmental variables.</li> <li>3. To identify and determine near-surface spatial distribution, diversity and abundance of phytoplankton, and zooplankton, based on Continuous Plankton Recorder samples collected during transit to and from the Ross Sea.</li> <li>4. To collect &amp; analyse data collected both underway, &amp; at stations for salinity, temperature nutrient and chlorophyll a data, spot optical measurements with the SeaWiFS.</li> <li>5. To identify and determine the spatial distribution, abundance (biomass), diversity, and size structure of epipelagic, mesopelagic (and possibly bathypelagic) species using acoustics and net sampling.</li> <li>6. To identify and measure diversity, distribution &amp; densities of mesozooplankton, macrozooplankton &amp; meroplankton (as collected by all plankton sampling methods except transit CPR samples).</li> <li>7. To determine diversity, distribution &amp; densities of viral, bacterial, phytoplankton &amp; microzooplankton species in the water column.</li> <li>8. To determine the spatial distribution, abundance (biomass), diversity, and size structure of shelf and slope demersal fish species and associated invertebrate species using a demersal survey.</li> <li>9. To determine the diversity, abundance/density, spatial distribution, and physical habitat associations of benthic assemblages across a body size spectrum from megafauna to bacteria, for shelf, slope, seamounts, and abyssal sites in Ross Sea.</li> <li>10. To describe trophic/ecosystem relationships in the Ross Sea ecosystem (pelagic and benthic, fish and invertebrates).</li> <li>11. Assess molecular taxonomy and population genetics of selected Antarctic fauna and flora to estimate evolutionary divergence within and among ocean basins in circumpolar species. Provide DNA barcoding.</li> </ol>	Completed	<p>Allcock et al. 2009; 2010; Submitted; Alvaro et al. 2011; Bowden et al. 2011a; In Prep; Clark et al. 2010a; Dettai et al. 2011; Eakin et al. 2009; Eleaume et al. 2011; In Prep; Ghiglione et al. 2012; Gordon 2000; Grotti et al. 2008; Hanchet et al. 2008a; 2008b; 2008c; 2008d; Hanchet 2009; 2010; Hanchet et al. In Press; Heimeier et al. 2010; Hemery et al. In prep; Koubbi et al. 2011; Leduc et al. 2012a; b; Linse et al. 2007; Lörz 2009; Lörz 2010a; 2010b; 2010c; Lörz &amp; Coleman 2009; Lörz et al. 2007; 2009; 2012a; b; In Press; In Prep; Maas et al. 2010a; McMillan et al. 2012.; Mitchell 2008; Nielsen et al. 2009; Norkko et al. 2005; O'Driscoll 2009; O'Driscoll et al. 2009; 2010; O'Driscoll et al. In Press; O'Loughlin et al. 2011; Pakhomov et al. 2011; Pinkerton et al. 2007a; Pinkerton et al. 2009a; b; Pinkerton et al. In review; In press; Schiaparelli et al. 2006; 2008; 2010; Smith et al. 2011a; b; Stein 2012; Strugnelli et al. Submitted</p>

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2007-01	Chatham-Challenger Oceans 20/20 Post-Voyage	<ol style="list-style-type: none"> <li>1. To quantify in an ecological manner- the biological composition and function of the seabed at varying scales of resolution- on the Chatham Rise and Challenger Plateau</li> <li>2. To elucidate the relative importance of environmental drivers- including fishing- in determining sea bed community composition and structure.</li> <li>3. To determine if remote-sensed data (e.g. acoustic) and environmentally derived classification schemes (e.g. marine environmental classification system) can be utilized to predict bottom community composition- function and diversity</li> <li>4. To count- measure- and identify to species-level (where possible- otherwise to genus) all macro invertebrates (&gt; 2 mm) and fish collected during Oceans 20/20 voyages.</li> <li>5. To count- measure and identify to species-level (where possible- otherwise to genus or family) all meiofauna (&gt; 2 mm) from multicore samples collected during the Oceans 20/20 voyages.</li> <li>6. To count- measure and identify to species- level (where possible- otherwise to genus or family) all fauna collected by hyper-benthic sled during the Oceans 20/20 voyages.</li> <li>7. To count- measure- and identify to species-level all macrofauna observed on DTIS images collected during the Oceans 20/20 voyages. The number of biogenic features (burrows/mounds) and habitat (spatial) complexity should also be estimated.</li> <li>8. To count- measure- and identify to species-level (where possible- otherwise to genus or family) all macrofauna observed on DTIS video footage collected during the Oceans 20/20 voyages.</li> <li>9. To calculate and compare the performance of a suite of diversity measures (species and taxonomic-based) at varying levels of resolution.</li> <li>10. To estimate particle size composition and organic content of sediment samples. Sediment samples should be aggregated over the top 5 cm of sediment.</li> <li>11. To measure the bacterial biomass (top 2 cm) of the sediment and in the sediment surface water samples- collected during the Oceans 20/20 voyages</li> <li>12. To elucidate the relationships- patterns and contrasts in species composition- assemblages- habitats- biodiversity and biomass (abundance) both within and between stations- strata and areas.</li> <li>13. To define habitats (biotic) encountered during the survey and assess their relative sensitivity to modification by physical disturbance- their recoverability and their importance to ecosystem function / production.</li> <li>14. To quantify the productivity- energy flow (trophic networks) and the energetic coupling (benthic pelagic or otherwise) of the area surveyed areas at various levels of resolution.</li> </ol>	Completed	Bowden 2011; Bowden et al. 2011 ; In press; Bowden & Hewitt 2012; Compton et al. 2012; Coleman and Lörz 2010; Hewitt et al. 2011a; 2011b; Lörz 2011a; 2011b; Nodder et al. 2012; Floerl et al. 2012

*AEBAR 2012: Appendices: Past projects*

		<p>15. To assess the extent to which patterns of species distributions and communities can be predicted using environmental data (including fishing) collected during the Ocean 20/20 voyages or held in other databases.</p> <p>16. To provide an interactive- high resolution mapping facility for displaying &amp; plotting all data collected &amp; derived indices. Includes environmental data- the abundance of species- indices of biomass or diversity- and statistically derived groupings</p> <p>17. To assess the extent to which acoustic- environmental- or other remote-sensed data can provide cost-effective- reliable means of assessing biodiversity at the scale of the Oceans 20/20 surveys.</p> <p>18. To assess the extent to which the 2005 MEC and subsequent variants can provide cost-effective- reliable means of assessing biodiversity at the scale of the Oceans 20/20 surveys.</p> <p>19. Collating all information and analysis from all objectives- devise a series of statistically supported recommendations for surveying marine biodiversity in the future. Including- but may not be limited to- statistical analyses and modelling.</p>		
--	--	---	--	--

**BIODIVERSITY** *continued*

<b>Project Code</b>	<b>Project Title</b>	<b>Specific Objectives</b>	<b>Status</b>	<b>Citation/s</b>
ZBD2006-02	Ongoing NABIS development	<p>As part of NABIS, users will be able to identify spatial information relating to the annual distribution (average distribution over the period of a year) of particular species within the waters around New Zealand and in the terrestrial environment (including off shore islands) of New Zealand. Users will also be able to interrogate metadata and attribute data related to the information layers presented. Users will employ NABIS to identify where a particular species is found, to identify what species are found within an area of interest, and be able to compare the spatial distribution of a particular species with other information layers.</p> <p>2. Some species may have notable changes in their spatial distribution throughout a year. For such species, users of NABIS will be able to view spatial information relating to the seasonal distribution of particular species within the waters around New Zealand and in the terrestrial environment (including offshore islands) of New Zealand. Users will also be able to interrogate metadata and attribute data related to the information layers presented. For species with a seasonal component to their biological distribution, users will employ NABIS to identify where a particular species is found within the waters around New Zealand and in the terrestrial environment (including off shore islands) of New Zealand at a particular time of the year, to identify what species are found within an area of interest at a particular time of year, or be able to compare the distribution of a particular species at a particular time of year, with other information layers.</p> <p>3. To provide analysis of the data used in determining the hotspot distribution.</p>	Completed	Anderson 2007b

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2006-03	Antarctic coastal marine systems	1. Quantify patterns in benthic community structure and function at two coastal Ross Sea locations (Terra Nova Bay and Cape Evans). 2. Quantify benthic community structure and function at selected locations in Terra Nova Bay and Cape Evans.	Completed	Cummings et al. 2003; 2006b; 2008; Thrush & Cummings 2011; Thrush et al. 2010
ZBD2006-04	Chatham/challenger oceans 20/20	1. To collect seabed fauna, sediment samples and photographic images along transects in the Chatham Rise and the Challenger Plateau, as determined by the sampling protocol described in the Voyage Programmes for Voyages 2 and 3 of the project. Multibeam data should be collected opportunistically as time allows. 2. To describe the distribution of broad macro epifauna groups (I.D. level to be determined at sea during Surveys 2 & 3), their relative abundance, the substrate and habitat types, including representative photographic images of each sea-bed habitat and associated fauna along transects in the survey areas. 3. To provide a description of the observed evidence of fishing along transects. 4. To provide indicative measures of alpha biodiversity (richness, number of taxonomic groups) at appropriate scales within and between transects, and between the Chatham Rise and the Challenger Plateau. 5. To determine broad scale variability in sea-bed habitats and associated biodiversity within and between MEC classes at 20 class level. 6. To process and archive biological samples and data into databases and collections for future analysis in meeting the Overall Objectives above.	Completed	Nodder 2008; Nodder et al. 2011
ZBD2005-02	Marine Environment Classification Project	1. Co-fund the Marine Environment Classification Project (being done by NIWA) with the Department of Conservation.	Completed	Snelder et al. 2005; 2006; Leathwick et al. 2006a; b; c
ZBD2005-09	Rocky reef ecosystems - how do they function? Integrating the roles of primary and secondary production, biodiversity and connectivity across coastal habitats	1. To develop a qualitative numerical model of how New Zealand's rocky reef systems are functionally structured 2. To quantify the effects of human predation, and environmental degradation across reef gradients – top-down, or bottom-up functioning? 3. To advance our understanding of how subtidal reef systems are fuelled through primary and secondary production (from a range of sources), the role that biodiversity plays, and how this varies across different reef settings. 4. To quantify how subtidal reef systems are linked with other habitats and ecosystems at broader spatial scales, including the connectivity of MPAs with other habitats and areas.	In the process of publication	MacDiarmid et al. In Press c

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2005-03	Tangaroa ross sea voyage	<p>1. To test the feasibility of obtaining estimates of demersal fish relative abundance using cameras with and without flood lights in areas of high importance for the Ross Sea toothfish fishery (principally 800-1200 m).</p> <p>2. To utilise deepwater camera transects, supported by other direct sampling methods, to characterise the relative abundance, distribution, and diversity of demersal fish species (assuming Objective 1 yields satisfactory results) and of benthic macro-invertebrates, and to examine relationships between demersal fishes and benthic habitats/communities. Camera transects will be deployed opportunistically, with focus on the following high-priority areas (in order of high to low priority) wherever possible:</p> <p>i) Areas of the continental shelf break at depths of high importance for the toothfish fishery (principally 800-1200 m but also 600-800m &amp; 1200-1500 m if time permits),</p> <p>ii) Shallow (50-200 m) water in the immediate vicinity of the Balleny Islands;</p> <p>iii) Deeper water in the vicinity of the Balleny Islands; iv) seamounts around and between Scott Island and the Balleny Islands; and v) at other locations (&lt; 600 m) as opportunity arises (e.g. around Scott Island, western Ross Sea, south-eastern Ross Sea).</p> <p>3. To collect specimens/tissues of selected benthic and pelagic organisms with priority in the vicinity of the Balleny Islands (and to the east/southeast, for pelagic specimens especially Antarctic krill species) and deliver specimens to other projects for stable isotope analysis in order to contribute to understanding of trophic relationships.</p> <p>4. To acquire a continuous acoustic survey of the water column, opportunistically undertake species verification of acoustic marks, integrate the acoustic marks and produce a GIS map of verified and unverified distributions of functionally important mesopelagic species (e.g. krill, Antarctic silverfish).</p> <p>5. To undertake routine identification and abundance estimates of marine mammal and seabird species and deliver raw and GIS summarised data to other related projects in order to generate spatially and temporally explicit population biomass and foraging distribution estimates for top air-breathing predators in the Ross Sea.</p> <p>6. To undertake automated water sampling in order to monitor the identities and spatial and temporal distributions of plankton in the Ross Sea region and to allow ground-truthing of data collection from satellites (e.g. surface seawater temperature, and chlorophyll-a concentration).</p>	In the process of publication	MacDiarmid & Stewart In Press; Mitchell & MacDiarmid 2006

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2005-01	Balleny Islands Ecology Research, Tiama Voyage (2006)	<p>1. To characterise shallow benthic communities across a range of habitat settings around the Balleny Islands, utilising a range of data collection methodologies (including SCUBA-based rock-wall suspension feeder photo quadrats, SCUBA-based linear video transects, and drop camera photography), and to analyse community patterns with reference to possible physical/oceanographic, biological, and/or biogeographic influences on community structure.</p> <p>2. To characterise aspects of the marine food web of the Balleny Islands area, using stable isotope analysis of specimens from important functional groups, and to make inferences about factors affecting ecosystem-scale trophodynamics in the Balleny Islands area and potential implications for the function of the wider ecosystem.</p> <p>3. To characterise the spatial and temporal distributions of higher-level consumer species (birds, seals and whales) and of dominant pelagic prey (i.e. krill swarms) by opportunistically recording all at-sea sightings, and by systematic observation of landbased top predators (birds and seals) while sailing along the coast of the islands.</p> <p>4. To collect and photograph and/or retain fish specimens from shallow benthic environments using a range of fishing methods, including food-baited fish traps, lightbaited fish traps, rotenone sampling, and/or baited lines.</p> <p>5. To continuously collect bathymetric data and water-column acoustic data (i.e. mesopelagic acoustic marks) throughout the voyage, using an acoustic sounder.</p> <p>6. To opportunistically collect a variety of data/materials during shore-based landings, including wherever possible: i) breast feathers from living penguins; ii) tissue samples/feathers/bones from dead seals/penguins/other sea birds; iii) seal scats; iv) visual estimates of adult and juvenile penguin numbers; v) visual assessments of penguin colony status; vi) photographs of penguin colonies; vii) sediment excavations of occupied and abandoned colonies. (Where appropriate these data will contribute to Objective 2).</p>	Terminated	Smith 2006

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2005-05	Long-term effects of climate variation and human impacts on the structure and functioning of New Zealand shelf ecosystems	<ol style="list-style-type: none"> <li>1. To estimate changes in marine productivity via fluctuations in ocean climate and terrestrial nutrient input over the last 1000 years.</li> <li>2. To assess and collate existing archaeological, historical and contemporary data (including catch records and stock assessments) on relevant components of the marine ecosystem to provide a detailed description of change in the shelf marine ecosystem in two areas of contrasting human occupation over last 1000 years.</li> <li>3. To collect additional oral histories from Maori and non-Maori fishers and shellfish gatherers regarding the distribution, sizes and relative abundance (compared to present availability) of key fish and invertebrate stocks in both regions during the first half of the 20th century before the start of widespread modern industrial fishing.</li> <li>4. To build mass-balance ecosystem models (e.g. Ecopath) of the coastal and shelf ecosystem in each area for five critical time periods: now, 60 years BP (before modern industrial fishing), 250 years BP (before European whaling and sealing), 600 y BP (early Maori phase) and 1000 years BP (before human settlement).</li> <li>5. To use qualitative modelling techniques to determine the critical interactions amongst species and other ecosystem components in order to identify those that should be a priority for future research.</li> </ol>	In the process of publication	Carroll et al. In Press; Jackson et al. In Press; Lalas et al. In Press a; b; Lalas & MacDiarmid In Press; Lorrey et al. In Press; MacDiarmid et al. In Press a; b; Maxwell & MacDiarmid In Press; Neil et al. In Press; Paul 2012; Parsons et al. In Press; Pinkerton In Press; Smith 2011
ZBD2004-01	Baseline information on the diversity and function of marine ecosystems	<ol style="list-style-type: none"> <li>1. To quantify, and compare, the macro-invertebrate assemblage composition of a number of seamounts at the southernmost end of the Kermadec volcanic arc.</li> <li>2. To compare the macro-invertebrate diversity of the southernmost end of the Kermadec volcanic arc with that of seamounts already sampled and reported on.</li> </ol>	Completed	Rowden & Clark 2010; Smith et al. 2008
ZBD2004-02	Ecosystem-scale trophic relationships: diet composition and guild structure of middle-depth fish on the Chatham Rise	<ol style="list-style-type: none"> <li>1. To quantitatively characterise the diets of abundant middle-depth fish species on the Chatham Rise, by analysis of fish stomach contents collected from the January 2005, January 2006 and January 2007 Chatham Rise middle-depths trawl surveys.</li> <li>2. To quantitatively characterise Chatham Rise fish diets throughout the year, for a period of 24 months, by analysis of fish stomach contents collected opportunistically aboard industry vessels.</li> <li>3. To describe and examine patterns of diet variation within each fish species as a function of spatial, temporal, and environmental variables, and of fish size.</li> <li>4. To define and characterise trophic guilds for abundant fish species on the Chatham Rise, using multivariate analysis of fish diet data, and to analyse the nature and relative strength of potential trophic interactions between guilds.</li> <li>5. To create and populate a diets database to store all of the dietary information collected under Objectives 1 and 2, and for use in subsequent dietary studies.</li> </ol>	Completed	Connell et al. 2010; Dunn 2009; Dunn et al. 2010a; b; c; Dunn et al. In press; Forman & Dunn 2010; Horn et al. 2010; Stevens & Dunn 2010;

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2004-05	Assessment and definition of the biodiversity of coralline algae of northern New Zealand	1. To assess and define the biodiversity of coralline algae in northern New Zealand. 2. To develop rapid identification tools for coralline algae using molecular sequencing data. 3. To contribute representative material to the national Coralline Algal Collections. 4. To produce ID guides to common coralline algae of northern New Zealand.	Completed	Farr et al. 2009
ZBD2004-10	Development of bioindicators in coastal ecosystems	1. Investigate linkages between land use patterns in catchments and nitrogen loading to recipient estuaries and coastal ecosystems 2. Characterise isotopic signatures of selected bioindicator organisms in relation to different terrestrial nutrient loads; and 3. Validate the use of bioindicators using controlled laboratory and field experiments.	Completed	Savage 2009
ZBD2004-19	Ecological function and critical trophic linkages in New Zealand soft-sediment habitats	1. Define the interactive effects of two functionally important benthic species in maintaining critical trophic linkages in soft-sediment systems from a series of integrated field experiments. 2. Quantify effects of heart urchins ( <i>Echinocardium australe</i> ) on sediment properties- benthic primary production- and macrofaunal diversity through manipulative field experiments in Mahurangi Harbour. 3. Test for interactions between pinnid bivalves ( <i>Atrina zelandica</i> ) and heart urchins ( <i>Echinocardium australe</i> ) in field experiments- and measure their respective and combined contributions to sediment properties- benthic primary production- and macrofauna 4. Determine the dependence of results from objectives 1 and 2 (functional contributions of <i>Echinocardium</i> and <i>Atrina</i> ) in an environmental context by conducting experiments along an estuarine-coastal gradient.	Completed	Lohrer et al. 2010
ZBD2003-02	Biodiversity of Coastal Benthic Communities of the North Western Ross Sea.	1. Quantify patterns in biodiversity and community structure in the coastal Ross Sea region 2. Quantify biodiversity in benthic communities at selected locations in the Ross sea north of Terra Nova Bay 3. Describe ecosystem function at selected locations in the Ross Sea north of Terra Nova Bay.	Completed	Cummings et al. 2003; 2006a; 2010; De Domenico et al. 2006; Guidetti et al. 2006; Norkko et al. 2004
ZBD2003-03	Biodiversity of deepwater invertebrates and fish communities of the north western Ross Sea	1. To describe, and quantify the diversity of, the benthic macroinvertebrates and fish assemblages of the Balleny Islands and adjacent seamounts, and to determine the importance of certain environmental variables influencing assemblage composition.	Completed	Rowden et al. 2012a; In Press; Mitchell & Clark 2004

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2003-04	Fiordland Biodiversity Research Cruise	1. How can ecotone boundaries be defined? 2. If you have an ecotone boundary defining the edge of a commercial exclusion zone how wide is the transition zone across the boundary? 3. If you have an area delineated as a marine protected area or a commercial exclusion zone, does it adequately represent the different habitats or biodiversity of the whole region?	Completed	Wing 2005
ZBD2003-09	Macquarie Ridge Complex Research Review	To review and summarise both biological and physical research carried out on or around the section of the Macquarie Ridge Complex that lies between New Zealand and Macquarie Island	Completed	Grayling 2004
ZBD2002-01	Ecology of Coastal Benthic Communities in Antarctica	Objectives unknown	Completed	Schwarz et al. 2003; 2005; Thrush et al. 2006; Thrush & Cummings 2011; Cummings et al. 2003; Sharp et al. 2010; Sutherland 2008
ZBD2002-02	Whose larvae is that? Molecular identification of planktonic larvae of the Ross Sea.	1. To use molecular sequencing tools in the taxonomic identification of cryptic/invasive marine 2. To provide a molecular description and characterisation of gobies that are introduced ( <i>Arenigobius bifrenatus</i> and <i>Acentrogobius pflaumii</i> ) cryptogenic ( <i>Parioglossus marginalis</i> ) or native (eg. <i>Favonigobius lentiginosus</i> and <i>F. expuissitus</i> ). 3. To describe the molecular diversity of the above species throughout their native and introduced distributions- and characterise a range of the greatest potential invasive gobioid and blennioid species from the Australasian region. 4. To develop molecular criteria to rapidly identify invasive or cryptogenic gobioid and blennioid fish	Completed	Sewell 2005; 2006; Sewell et al. 2006
ZBD2002-06A	Impacts of terrestrial run-off on the biodiversity of rocky reefs	1. Conduct field and laboratory experiments to determine relationships between sediment loading, epifaunal assemblages, and mortality of filter feeding invertebrates. 2. Conduct field and laboratory experiments to identify the influence of sediment on early life stages of key grazers. 3. Determine photosynthetic characteristics and survival of large brown seaweeds and understory algal species in relation to a sediment gradient.	Completed	Schwarz et al. 2006

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2002-12	Molecular identification of cryptogenic/invasive marine species – gobies.	1. To use molecular sequencing tools in the taxonomic identification of cryptic/invasive marine species 2. To provide a molecular description and characterisation of gobies that are introduced ( <i>Arenigobius bifrenatus</i> and <i>Acentrogobius pflaumii</i> ) cryptogenic ( <i>Parioglossus marginalis</i> ) or native (eg. <i>Favonigobius lentiginosus</i> and <i>F. expuisitus</i> ). 3. To describe the molecular diversity of the above species throughout their native and introduced distributions- and characterise a range of the greatest potential invasive gobioid and blennioid species from the Australasian region. 4. To develop molecular criteria to rapidly identify invasive or cryptogenic gobioid and blennioid fish.	Completed	Lavery et al. 2006
ZBD2002-16	Joint New Zealand and Australian Norfolk Ridge	1. To describe the marine biodiversity of the Norfolk Ridge and Lord Howe Rise seamount communities. 2. To survey- sample and document the marine biodiversity and environmental data from seamounts on the Norfolk Ridge and Lord Howe Rise to a depth of at least 1-000m depth. (b) To preserve samples of fishes and invertebrates and hold these in ac...	Completed	Clark & Roberts 2008
ZBD2002-18	Quantitative survey of the intertidal benthos of Farewell Spit Golden Bay	1. To undertake a baseline survey of intertidal macrobenthic organisms at Farewell Spit Nature Reserve and adjacent flats. 2. To undertake an initial field survey of <i>Zostera</i> distribution at Farewell Spit Nature Reserve and adjacent intertidal flats. 3. To undertake a preliminary survey of sediment characteristics of the intertidal flats at Farewell Spit Nature Reserve and adjacent flats.	Completed	Battley et al. 2005
ZBD2001-02	Documentation of New Zealand Seaweed	1. To publish a regional algal flora of Fiordland based on voucher herbarium specimens. 2. To assemble a database of references and to review the current state of knowledge about New Zealand macroalgae.	Completed	Nelson et al. 2002
ZBD2001-03	Ecology and biodiversity of coastal benthic communities in Antarctica.	1. To develop sampling protocols for estimating the relative abundance of algae and benthic invertebrates 2. To quantify patterns in biodiversity and benthic community structure at two locations in McMurdo Sound 3. To analyse Ross Island Sea-Level data.	Completed	Norkko et al 2002
ZBD2001-04	"Deep Sea New Zealand"	To help publish the book "Deep Sea New Zealand"	Completed	Batson 2003
ZBD2001-05	Crustose coralline algae of New Zealand	1. To assess the biodiversity of crustose coralline algae in NZ using modern taxonomic methods and molecular sequence tools. 2. To establish the NZ National Coralline Algal Collection. 3. To produce identification guides to NZ species.	Completed	Harvey et al. 2005; Farr et al. 2009; Broom et al 2008

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2001-06	Biodiversity of New Zealand's soft-sediment communities	<ol style="list-style-type: none"> <li>1. To review the current knowledge of the biodiversity of macroinvertebrates and macrophytes living in and on soft-sediment substrates in New Zealand's harbours- estuaries- beaches and to 1000 m water depth.</li> <li>2. To review existing published and unpublished sources of information on soft-sediment marine assemblages around New Zealand.</li> <li>3. Using the results of Objective 1- identify gaps in the knowledge- hotspots of biodiversity- areas of particular vulnerability- and make recommendations on areas or assemblages that could be the subject of directed research in future years.</li> </ol>	Completed	Rowden et al. 2012b
ZBD2001-10	Additional Research on Biodiversity of Seamounts	<ol style="list-style-type: none"> <li>1. To determine the macro-invertebrate assemblage composition on Cavalli seamount, and adjacent seamount W1, by photographic transects and epibenthic sled sampling.</li> <li>2. To determine the distribution of macro-invertebrate assemblages on the seamounts.</li> <li>3. To compare the macro-invertebrate species diversity of neighbouring seamounts.</li> <li>4. To evaluate and collect samples from suitable macro-invertebrate species for genetic analysis.</li> <li>5. To map bathymetry and habitat characteristics of the seamounts.</li> <li>6. To compare macro-invertebrate assemblage composition of the seamounts with nearby hard bottom low relief (under 100 m) on the slope, if suitable areas can be located.</li> </ol>	Completed	Rowden et. al 2004
MOF2000-01	Bryozoan thickets off Otago Peninsula	Objectives unknown	Completed	Batson & Probert 2000
ZBD2000-01	A review of current knowledge describing the biodiversity of the Ross Sea region	<ol style="list-style-type: none"> <li>1. To review and document existing published and unpublished information describing the biodiversity of the Ross Sea region.</li> <li>2. To identify and document Ross Sea region marine communities that are under high pressure or likely to come under high pressure from human activities in the near future.</li> </ol>	Completed	Bradford-Grieve & Fenwick 2001a; 2001b; Fenwick & Bradford-Grieve 2002a; 2002b; Bradford-Grieve & Fenwick 2002; Varian 2005
ZBD2000-02	Exploration and description of the biodiversity, in particular the benthic macrofauna, of the western Ross Sea	<ol style="list-style-type: none"> <li>1. To utilise sampling opportunities provided by the presence of RV Tangaroa in the western Ross Sea in February / March 2001 to make collections of (primarily) benthic organisms as a contribution to the understanding of biodiversity in the region.</li> <li>2. To identify and document the organisms collected and provide for their proper storage in national collections.</li> <li>3. To describe the logistic constraints of working in the Ross Sea region, and make recommendations for future research to improve understanding of biodiversity in the Ross Sea.</li> </ol>	Completed	Page et al. 2001

BIODIVERSITY *continued*

Project Code	Project Title	Specific Objectives	Status	Citation/s
ZBD2000-03	The spatial extent and nature of the bryozoan communities at Separation Point, Tasman Bay	1. To assess the present state and extent of bryozoan communities around Separation Point. 2. To characterise the bryozoan communities around Separation Point.	Completed	Grange et al. 2003
ZBD2000-04	Supplementary Research on Biodiversity of Seamounts	1. To determine the biodiversity of seamounts of the southern Kermadec volcanic arc (Rumble V, Rumble 111, Brothers). 2. To describe the distribution of fauna, with an emphasis on mapping the nature and extent, of biodiversity associated with hydrothermal vents. 3. To compare the biodiversity of the three seamounts, and adjacent slope. 4. To collect samples from near the vent sources (if possible, as these are thought to be very localised) to measure chemical and thermal aspects of the environment	Completed	Rowden et al. 2002 and 2003; Clark & O'Driscoll 2003
ZBD2000-06	"The Living Reef: The Ecology of New Zealand's Rocky Reefs"	1. Funding to support the publication of this book.	Completed	Andrew & Francis (Eds.) 2003
ZBD2000-08	A review of current knowledge describing New Zealand's Deepwater Benthic Biodiversity	1. To review and document existing published and unpublished reports and data describing New Zealand's deepwater benthic biodiversity. 2. To make recommendations on representative communities and potentially impacted communities that could be the subject of directed research.	Completed	Key 2002
ZBD2000-09	Antarctic fish taxonomy	1. Ross Sea fishes processing and identification	Completed	Roberts & Stewart & 2001

## References

- Abraham, E., 2008. Evaluating methods for estimating the incidental capture of New Zealand sea lions, New Zealand Aquatic Environment and Biodiversity Report No. 15, 25 p.
- Abraham, E., 2010. Warp strike in New Zealand trawl fisheries, 2004-05 to 2008-09, New Zealand Aquatic Environment and Biodiversity Report No. 60, 29 p.
- Abraham, E., 2011. Probability of Mild Traumatic Brain Injury for sea lions interacting with SLEDs, Unpublished Final Research Report for Ministry of Fisheries project SRP2011-03 (held by the Ministry of Fisheries, Wellington), 21 p.
- Abraham, E., Berkenbusch, K., Richard, Y., 2010. The capture of seabirds and marine mammals in New Zealand non-commercial fisheries, New Zealand Aquatic Environment and Biodiversity Review. No. 64, 52 p.
- Abraham, E., Kennedy, A., 2008. Seabird warp strike in the southern squid trawl fishery, 2004-05, New Zealand Aquatic Environment and Biodiversity Report No. 16, 39 p.
- Abraham, E., Thompson, F., 2009a. Capture of protected species in New Zealand trawl and longline fisheries, 1998-99 to 2006-07, New Zealand Aquatic Environment and Biodiversity Report No. 32, 197 p.
- Abraham, E., Thompson, F., 2009b. Warp strike in New Zealand trawl fisheries, 2004-05 to 2006-07, New Zealand Aquatic Environment and Biodiversity Report No. 33, 21 p.
- Abraham, E., Thompson, F., 2010. Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998-99 to 2008-09, New Zealand Aquatic Environment and Biodiversity Report No. 80., 172 p p.
- Abraham, E., Thompson, F., 2011a. Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002-03 to 2008-09, New Zealand Aquatic Environment and Biodiversity Report No. 79., 74 p.
- Abraham, E., Thompson, F., 2011b. Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998-99 to 2008-09, Unpublished Final Research Report for the Ministry of Fisheries project PRO2007-01 170 p p.
- Abraham, E., Thompson, F., Oliver, M., 2010. Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998-99 to 2007-08, New Zealand Aquatic Environment and Biodiversity Report No. 45, 148 p.
- Allcock, A., Barratt, I., Eleaume, M., Linse, K., Norman, M., Smith, P., Steinke, D., Stevens, D., Strugnell, J., 2010. Cryptic speciation and the circumpolarity debate: A case study on endemic Southern Ocean octopuses using the COI barcode of life. *Deep Sea Research* doi:10.1016/j.dsr2.2010.05.016.
- Allcock, A., Norman, M., Smith, P., Steinke, D., Stevens, D., Strugnell, J., 2009. Cryptic speciation and the circumpolarity debate: a case study on endemic Southern Ocean octopuses using the COI barcode of life, CAML Symposium, Genoap.
- Allcock, A., Others, Submitted. Bipolarity in marine invertebrates: myth or marvel. p.
- Alvaro, M., Blazewicz-Paszkwycz, M., Davey, N., Schiaparelli, S., 2011. Skin-digging tanaids: the unusual parasitic behaviour of *Exspinia typica* in Antarctic waters and worldwide deep basins. *Antarctic Science* 23: : 343-348.
- Anderson, O., 2004. Fish discards and non-target fish catch in the trawl fisheries for arrow squid, jack mackerel, and scampi in New Zealand waters, New Zealand Fisheries Assessment Report 2004/10, 61 p.
- Anderson, O., 2007a. Fish discards and non-target fish catch in the New Zealand jack mackerel trawl fishery, 2001-02 to 2004-05., New Zealand Aquatic Environment and Biodiversity Report No. 8, 36 p.
- Anderson, O., 2007b. NABIS biological distributions, Unpublished Final Research Report for Ministry of Fisheries Project ZBD2006-02, held by the Ministry of Fisheries. , 20 p.
- Anderson, O., 2008. Fish and invertebrate bycatch and discards in ling longline fisheries, 1998-2006, New Zealand Aquatic Environment and Biodiversity Report No. 23, 43 p.
- Anderson, O., 2009a. Fish discards and non-target fish catch in the New Zealand orange roughy trawl fishery: 1999-2000 to 2004-05, New Zealand Aquatic Environment and Biodiversity Report No. 39, 40 p.
- Anderson, O., 2009b. Fish and invertebrate bycatch and discards in southern blue whiting fisheries, 2002-07, New Zealand Aquatic Environment and Biodiversity Report. No. 43, 42 p.
- Anderson, O., 2011. Fish and invertebrate bycatch and discards in orange roughy and oreo fisheries from 1990-91 until 2008-09, New Zealand Aquatic Environment and Biodiversity Report No. 67, 60 p.
- Anderson, O., 2012. Fish and invertebrate bycatch and discards in New Zealand scampi fisheries from 1990-91 until 2009-10., New Zealand Aquatic Environment and Biodiversity Report No. 100, 65 p.
- Anderson, O., Gilbert, D., Clark, M., 2001. Fish discards and non-target catch in the trawl fisheries for orange roughy and hoki in New Zealand waters for the fishing years 1990-91 to 1998-99, New Zealand Fisheries Assessment Report 2001/16, 57 p.
- Anderson, O., Smith, M., 2005. Fish discards and non-target fish catch in the New Zealand hoki trawl fishery, 1999/2000 to 2002/03., New Zealand Fisheries Assessment Report 2005/03, 37 p.
- Andrew, N., Francis, M., 2003. The living Reef: the ecology of New Zealand's Living Reef. Crag Potton Publishing, Nelson, 283 p.
- Ayers, D., Francis, M., Griggs, L., Baird, S., 2004. Fish bycatch in New Zealand tuna longline fisheries, 2000-01 and 2001-02, New Zealand Fisheries Assessment Report 2004/46, 47 p.
- Baird, S., 1994. Nonfish Species and Fisheries Interactions Working Group Report. , N.Z. Fisheries Assessment Working Group Report 94/1, N.Z. Ministry of Agriculture and Fisheries, Wellington, 54 p.
- Baird, S., 1995. Nonfish Species and Fisheries Interactions Working Group Report. , N.Z. Fisheries Assessment Working Group Report - April 1995. 95/1, N.Z. Ministry of Agriculture and Fisheries, Wellington, 24 p.
- Baird, S., 1996. Nonfish Species and Fisheries Interactions Working Group Report. , N.Z. Fisheries Assessment Working Group Report - May 1996. 96/1, N.Z. Ministry of Agriculture and Fisheries, Wellington, 34 p.
- Baird, S., 1997. Report on the incidental capture of nonfish species during fishing operations in New Zealand waters, Unpublished Final Research Report for Ministry of Fisheries SANF01 contract, 15 p plus appendices on New Zealand fur seal-trawl fishery interaction (54 p) and seabird-tuna longline fishery interaction (34 p) p.
- Baird, S., 1999a. Determination of factors which affect nonfish bycatch in some New Zealand fisheries, Unpublished Final Research Report for Ministry of Fisheries Project ENV9701 – Objective 3, 83 p.
- Baird, S., 1999b. Estimation of nonfish bycatch in commercial fisheries in New Zealand waters, 1997-98, Unpublished Final Research Report for Objective 1 of Ministry of Fisheries Project ENV9801, 57 p.

- Baird, S., 2001. Estimation of the incidental capture of seabird and marine mammal species in commercial fisheries in New Zealand waters, 1998-99, New Zealand Fisheries Assessment Report 2001/14, 43 p.
- Baird, S., 2003. New Zealand breeding seabirds: human-induced mortality — a review, Unpublished Report prepared for the Ministry of Fisheries Project ENV2000/09, 74 p.
- Baird, S., 2004a. Estimation of the incidental capture of seabird and marine mammal species in commercial fisheries in New Zealand waters, 1999-2000, New Zealand Fisheries Assessment Report 2004/41, 56 p.
- Baird, S., 2004b. Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2000-01, New Zealand Fisheries Assessment Report 2004/58, 63 p.
- Baird, S., 2004c. Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2001-02, New Zealand Fisheries Assessment Report 2004/60, 51 p.
- Baird, S., 2005a. Incidental capture of *Phocartos hookeri* (New Zealand sea lions) in New Zealand commercial fisheries, 2002-03, New Zealand Fisheries Assessment Report 2005/8, 17 p.
- Baird, S., 2005b. Incidental capture of *Phocartos hookeri* (New Zealand sea lions) in New Zealand commercial fisheries, 2002-03, New Zealand Fisheries Assessment Report 2005/9, 13 p.
- Baird, S., 2005c. Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2002/03, New Zealand Fisheries Assessment Report 2005/02, 50 p.
- Baird, S., 2005d. Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in commercial fisheries in New Zealand waters, 2000-01, New Zealand Fisheries Assessment Report 2005-11, 34 p.
- Baird, S., 2005e. Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in commercial fisheries in New Zealand waters, 2002-03, New Zealand Fisheries Assessment Report 2005/12, 35 p.
- Baird, S., 2005f. Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in commercial fisheries in New Zealand waters, 2001-02, New Zealand Fisheries Assessment Report 2005/12, 33 p.
- Baird, S., 2007. Incidental capture of cetaceans in commercial fisheries in New Zealand waters 2003-04 and 2004-05, New Zealand Aquatic Environment and Biodiversity Report. No. 13, 27 p.
- Baird, S., 2008a. Incidental capture of cetaceans in commercial fisheries in New Zealand waters, 1994-95 to 2005-06, New Zealand Aquatic Environment and Marine Biodiversity Report No. 21, 29 p.
- Baird, S., 2008b. Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in longline fisheries in New Zealand waters, 1994-95 to 2005-06, New Zealand Aquatic Environment and Biodiversity Report No. 20, 21 p.
- Baird, S., 2011. New Zealand fur seals - summary of current knowledge, New Zealand Aquatic Environment and Biodiversity Report No. 72, 51 p.
- Baird, S., Bagley, N., Wood, B., Dunn, A., Beentjes, M., 2002. The spatial extent and nature of mobile bottom fishing methods within the New Zealand EEZ, 1989-90 to 1998-99, Unpublished Final Research Report for Objective 1 of project ENV2000/05, p.
- Baird, S., Bradford, E., 1999. Factors that may influence the bycatch of nonfish species in some New Zealand fisheries, Unpublished final Research Report completed for Objective 3 of Ministry of Fisheries Project ENV9801, 106 p.
- Baird, S., Bradford, E., 2000a. Factors that may have influenced bycatch of New Zealand fur seals (*Arctocephalus forsteri*) in the west coast South Island hoki fishery, NIWA Technical Report 92, 35 p.
- Baird, S., Bradford, E., 2000b. Factors that may have influenced the capture of seabirds in New Zealand tuna longline fisheries, NIWA Technical Report 93, 61 p.
- Baird, S., Doonan, L., 2005. *Phocartos hookeri* (New Zealand sea lions): incidental captures in New Zealand commercial fisheries during 2000-2001 and in-season estimates of captures during squid trawling in SQU 6T in 2002, New Zealand Fisheries Assessment Report 2005/17, 18 p.
- Baird, S., Francis, M., Griggs, L., Dean, H., 1998. Annual review of bycatch in southern bluefin and related tuna longline fisheries in the New Zealand 200 n. mile Exclusive Economic Zone, CCSBT-ERS/9806/31 (Report prepared for the Third Meeting of the Ecologically Related Species Working Group, Tokyo, 9-13 June 1998)p.
- Baird, S., Gilbert, D., 2010. Initial assessment of risk posed by trawl and longline fisheries to selected seabird taxa breeding in New Zealand waters, New Zealand Aquatic Environment and Biodiversity Report No. 50, 98 p.
- Baird, S., Griggs, L., 2005. Estimation of within-season chartered southern bluefin tuna (*Thunnus maccoyii*) longline seabird incidental captures, 2003, New Zealand Fisheries Assessment Report 2005/01, 15 p.
- Baird, S., Sanders, B., Dean, H., Griggs, L., 1999. Estimation of nonfish bycatch in commercial fisheries in New Zealand waters, 1990-91 to 1993-94, Final Research Report for Ministry of Fisheries Project ENV9701 - Objective 1, 63 p.
- Baird, S., Smith, M., 2007a. Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2003-04 and 2004-05, New Zealand Aquatic Environment and Biodiversity Report No. 9, 108 p.
- Baird, S., Smith, M., 2007b. Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in commercial fisheries in New Zealand waters, 2003-04 and 2004-05, New Zealand Aquatic Environment and Biodiversity Report. No. 14, 98 p.
- Baird, S., Smith, M., 2008. Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2005-06, New Zealand Aquatic Environment and Biodiversity Report, No. 18, 124 p.
- Baird, S., Wood, B., 2010. Extent of coverage of 15 environmental classes within the New Zealand EEZ by commercial trawling with seafloor contact, Unpublished Final Research Report for Objective 5 of project BEN2006/01, p.
- Baird, S., Wood, B., Bagley, N., 2009. The extent of trawling on or near the seafloor in relation to benthic-optimised marine environment classes within the New Zealand EEZ, Unpublished Final Research Report for Objective 5 of project BEN2006/01, p.
- Baird, S., Wood, B., Bagley, N., 2011. Nature and extent of commercial fishing effort on or near the seafloor within the New Zealand 200 n. Mile Exclusive Economic Zone, 1989-90 to 2004-05, New Zealand Aquatic Environment and Biodiversity Report No. 73, 143 p.
- Baird, S., Wood, B., Clark, M., Bagley, N., McKenzie, A., 2006. Description of the spatial extent and nature of disturbances by bottom trawls in Chatham Rise and Southern Plateau fisheries, Unpublished Final Research Report for project ENV2003/03, 139 p.
- Baker, B., Cunningham, R., Hedley, G., King, S., 2008. Data collection of demographic, distributional and trophic information on the Westland petrel to allow estimation of effects of fishing on population viability, Unpublished Final Research Report to the Ministry of Fisheries, 47 p.
- Baker, B., Hedley, G., Cunningham, R., 2010. Data collection of demographic, distributional, and trophic information on the flesh-footed shearwater to allow estimation of effects of fishing on population viability: 2009— 10 Field Season, Unpublished Final Research Report for the Ministry of Fisheries 62 p.

- Baker, G., Jensz, K., Cunningham, R., 2009. Data collection of demographic, distributional and trophic information on the white-capped albatross to allow estimation of effects of fishing on population viability — 2009 Field Season, Unpublished Research Report for the Ministry of Fisheries, 15 p.
- Ballara, S., Anderson, O., 2009. Fish discards and non-target fish catch in the trawl fisheries for arrow squid and scampi in New Zealand waters, New Zealand Aquatic Environment and Biodiversity Report No. 38, 102 p.
- Ballara, S., O'Driscoll, R., Anderson, O., 2010. Fish discards and non-target fish catch in the trawl fisheries for hoki, hake, and ling in New Zealand waters, New Zealand Aquatic Environment and Biodiversity Report. No. 48, 100 p.
- Batson, P., 2003. Deep New Zealand: Blue Water, Black Abyss. Canterbury University Press, Christchurch, 240 p.
- Batson, P., Probert, P.K., 2000. Bryozoan thickets off Otago Peninsula, New Zealand Fisheries Assessment Report 2000/46, 31 p.
- Battley, P., Melville, D., Schuckard, R., Ballance, P., 2005. Quantitative survey of the intertidal benthos of Farewell Spit, Golden Bay., Marine Biodiversity Biosecurity Report No. 7, 119 p.
- Beentjes, M., 2010. Toheroa survey of Oreti Beach, 2009, and review of historical surveys, New Zealand Fisheries Assessment Report 2010/6, 40 p.
- Beentjes, M., Baird, S., 2004. Review of dredge fishing technologies and practice for application in New Zealand, New Zealand Fisheries Assessment Report 2004/37, 40 p.
- Beentjes, M., Boubee, J., Jellyman, D., E, G., 2005. Non-fishing mortality of freshwater eels (*Anguilla* spp.). New Zealand Fisheries Assessment Report 2005/34, 38 p.
- Bentley, N., Davies, C., McNeill, S., Davies, N., 2004. A framework for evaluating spatial closures as a fisheries management tool., New Zealand Fisheries Assessment Report 2004/25, 25 p.
- Black, J., Wood, R., Berthelsen, T., In Press. Monitoring New Zealand's trawl footprint for deepwater fisheries: 1989-1990 to 2009-2010, New Zealand Aquatic Environment and Biodiversity Report No. XX., xx p.
- Blackwell, R., 2010. Distribution and abundance of deepwater sharks in New Zealand waters, 2000-01 to 2005-06, New Zealand Aquatic Environment and Biodiversity Report No. 57, 51 p.
- Blackwell, R., Stevenson, M., 2003. Review of the distribution and abundance of deepwater sharks in New Zealand waters, New Zealand Fisheries Assessment Report 2003/40, 48 p.
- Booth, J., Baird, S., Stevenson, M., Bagley, N., Wood, B., 2002. Review of technologies and practices to reduce bottom trawl bycatch and seafloor disturbance in New Zealand, Unpublished Fisheries Research Assessment Report prepared for the Ministry of Fisheries as completion of ENV2000/06, 61 p.
- Bowden, D., 2011. Benthic invertebrate samples and data from the Ocean Survey 20/20 voyages to the Chatham Rise and Challenger Plateau, 2007, New Zealand Aquatic Environment and Biodiversity Report No. 65, 40 p.
- Bowden, D., Schiaparelli, S., Clark, M., Rickard, G., 2011a. A lost world? Archaic crinoid-dominated assemblages on an Antarctic seamount. Deep Sea Research Part II: Topical Studies in Oceanography 58: 119-127.
- Bowden, D., Compton, T., Snelder, T., Hewitt, J., 2011b. Evaluation of the New Zealand Marine Environment Classifications using Ocean Survey 20/20 data from Chatham Rise and Challenger Plateau. , New Zealand Aquatic Environment and Biodiversity Report No. 77, 27 p.
- Bowden, D., Hanchet, S., Marriott, P., In Prep. Population estimates of Ross Sea demersal fish: a comparison between video and trawl methods. (Submitted as short note to Fisheries Research).
- Bowden, D., Hewitt, J., 2012. Recommendations for surveys of marine benthic biodiversity: outcomes from the Chatham-Challenger Ocean Survey 20/20 Post-Voyage Analyses Project New Zealand Aquatic Environment and Biodiversity Report, No. 91, 34 p.
- Bowden, D., Hewitt, J., Verdier, A.-L., Pallentin, A., In Press. The potential of multibeam echosounder data for predicting benthic invertebrate assemblages across Chatham Rise and Challenger Plateau, Draft Final Research Report Ministry of Fisheries, New Zealand, Wellington.
- Boyd, P., Law, C., 2011. An Ocean Climate Change Atlas for New Zealand waters: A primer for a major new web-based tool to help predict how oceanic species will be affected by climate change, NIWA Information Series No. 79 ISSN 1174-264X, , 24 p.
- Bradford, E., 2002. Estimation of the variance of mean catch rates and total catches of non-target species in New Zealand fisheries., New Zealand Fisheries Assessment Report 2002/54, 60 p.
- Bradford, E., 2003. Factors that might influence the catch and discards of non-target fish species on tuna long-lines, New Zealand Fisheries Assessment Report 2003/57, 75 p.
- Bradford-Grieve, J., Fenwick, G., 2001a. A review of the current knowledge describing the biodiversity of the Balleny Islands, Unpublished Final Research Report to the Ministry of Fisheries.
- Bradford-Grieve, J., Fenwick, G., 2001b. A review of the current knowledge describing the biodiversity of the Ross Sea region, Unpublished Final Research Report to the Ministry of Fisheries, xpp.
- Bradford-Grieve, J., Fenwick, G. 2002. A review of the current knowledge describing the biodiversity of the Balleny Islands. FRR ZBD2000/01, xx p.
- Bradford-Grieve, J., Livingston, M.E., 2011. Spawning fisheries and the productivity of the marine environment off the west coast of the South Island, New Zealand, New Zealand Aquatic Environment and Biodiversity Report. 84, 136 p.
- Breen, P., 2005. Managing the effects of fishing on the environment what does it mean for the rock lobster (*Jasus edwardsii*) fishery?, New Zealand Fisheries Assessment Report 2005/73, 45 p.
- Breen, P., 2008. Sea lion data for use in the population model for Project IPA200609. , Unpublished Final Research Report for Ministry of Fisheries, 30 March 2008 18 pp. plus Confidential Appendix of 16 p.
- Breen, P., Fu, D., Gilbert, D., 2010. Sea lion population modelling and management procedure evaluations, Unpublished Report for Project SAP2008/14, Objective 2: Presented to AEWG March 22 2010, Wellington, New Zealand.
- Breen, P., Fu, D., Gilbert, D., 2010. Sea lion population modelling and management procedure evaluations, Unpublished Report for Project SAP2008/14, Objective 2. Presented to AEWG March 22 2010, Wellington, New Zealand.
- Breen, P., Kim, S., 2006. Exploring alternative management procedures for controlling bycatch of Hooker's sea lions in the SQU 6T squid fishery, Unpublished Final Research Report For Ministry of Fisheries Project M0F2002/03L, Objective 3. Revision 5, 3 February 2006, 88 p.
- Broom, J., Hart, D., Farr, T., Nelson, W., Neill, K., Harvey, A., Woelkerling, W., 2008. Utility of psbA and nSSU for phylogenetic reconstruction in the Corallinales based on New Zealand taxa. Molecular Phylogenetics and Evolution 46: 958-973.
- Carroll, E., Jackson, J., Paton, D., Smith, T., In Press. Estimating nineteenth and twentieth century right whale catches and removals around east Australia and New Zealand, New Zealand Aquatic Environment and Biodiversity Report, xx p.

- Clark, M., Dunn, M., McMillan, P., Pinkerton, M., Stewart, A., Hanchet, S., 2010a. Latitudinal variation of demersal fish assemblages in the western Ross Sea. *Antarctic Science* 22 (6): 782-792.
- Clark, M., Watling, L., Rowden, A., Guinotte, J., Smith, C., 2010b. A global seamount classification to aid the scientific design of marine protected area networks. *Journal of Ocean and Coastal Management* 54: 19-36.
- Clark, M., Bowden, D., Baird, S., Stewart, R., 2010c. Effects of fishing on the benthic biodiversity of seamounts of the "Graveyard" complex, northern Chatham Rise, New Zealand Aquatic Environment and Biodiversity Report, No. 46, 40 p.
- Clark, M., O'Driscoll, R., 2003. Deepwater fisheries and aspects of their impact on seamount habitat in New Zealand. *Journal of Northwest Atlantic Fishery Science* 31: 441-458.
- Clark, M., Roberts, C., 2008. Fish and invertebrate biodiversity on the Norfolk Ridge and Lord Howe Rise, Tasman Sea (NORFANZ voyage, 2003), New Zealand Aquatic Environment and Biodiversity Report, No. 28, 131 p.
- Clark, M., Rowden, A., 2009. Effect of deepwater trawling on the macro-invertebrate assemblages of seamounts on the Chatham Rise, New Zealand. *Deep Sea Research I* 56: 1540-1554.
- Clark, M., Tracey, D., Pallentin, A., Schnabel, K., Anderson, O., Bowden, D., 2009. Voyage report of a survey of "seamounts" on the northwest and southeast Chatham Rise (TAN0905), Unpublished report available from NIWA, Wellington, 49 p.
- Clark, M., Williams, A., Rowden, A., Hobday, A., Consalvey, M., 2011. Development of seamount risk assessment: application to the ERAEF approach to Chatham Rise seamount features., New Zealand Aquatic Environment and Biodiversity Report No. 74, 18 p.
- Clement, D., Mattlin, R., 2010. Abundance, distribution and productivity of Hector's (and Maui's) dolphins, Unpublished Final Research report for the Ministry of Fisheries.
- Cole, R., Stevenson, M., Hanchet, S., Gorman, R., Rickard, G., Morrissey, D., Handley, S., 2007. Information on benthic impacts in support of the Southern Blue Whiting Fishery Plan, Unpublished Final Research Report for Ministry of Fisheries Research Project ZBD2005-16, Objectives 1-6p.
- Coleman, C., Lörz, A., 2010. A new species of Camacho (Crustacea, Amphipoda, Aoridae) from the Chatham Rise, New Zealand. *Zeitschrift für Zoologische Systematik und Evolutionsforschung* 86 (1): 33-40.
- Compton, T., Bowden, D., Pitcher, R., Hewitt, J., Ellis, N., 2012. Biophysical patterns in benthic assemblage composition across contrasting continental margins off New Zealand. *Journal of Biogeography*: doi:10.1111/j.1365-2699.2012.02761.x.
- Connell, A., Dunn, M., Forman, J., 2010. Diet and dietary variation of New Zealand hoki *Macruronus novaezelandiae*. *New Zealand Journal of Marine and Freshwater Research* 44 (4): 289 - 308.
- Cryer, M., Coburn, R., Hartill, B., O'Shea, S., Kendrick, T., Doonan, I., 1999. Scampi stock assessment for 1998 and an analysis of the fish and invertebrate bycatch of scampi trawlers, New Zealand Fisheries Assessment Research Document 99/4, 74 p.
- Cryer, M., Hartill, B., 2002. Relative impacts on benthic invertebrate community structure of trawl fisheries for scampi and finfish in the Bay of Plenty, Unpublished Final Research Report for Ministry of Fisheries Research Project ENV2000/05 Objective 2, 28 p.
- Cryer, M., Nodder, S., Thrush, S., Lohrer, A., Gorman, R., Vopel, K., Baird, S., 2004. The effects of trawling and dredging on benthic-pelagic coupling processes in the New Zealand EEZ., Draft New Zealand Fisheries Assessment Report 2004/xx., 66 p.
- Cryer, M., O'Shea, S., Gordon, D., Kelly, M., Drury, J., Morrison, M., Hill, A., Saunders, H., Shankar, U., Wilkinson, M., Foster, G., 2000. Distribution and structure of benthic invertebrate communities between North Cape and Cape Reinga Unpublished Final Research Report for Ministry of Fisheries Research Project ENV1998-05 Objectives 1-4p.
- Cummings, V., 2011. Ocean acidification: impacts on key NZ molluscs, Unpublished Progress Report for ZBD2009-13, 6 p.
- Cummings, V., Lohrer, D., 2011a. Ross Sea ecosystem function: predicting consequences of shifts in food supply, Unpublished Final Research Report for Ministry of fisheries Project ZBD2008-20, held by the Ministry of Fisheries, 53 p.
- Cummings, V., Hewitt, J., Van Rooyen, A., Currie, K., Beard, S., Thrush, S., Norkko, J., Barr, N., Heath, P., Halliday, J., Sedcole, R., Gomez, A., McGraw, C., Metcalf, V., 2011b. Ocean acidification at high latitudes: potential effects on functioning of the Antarctic bivalve *Laternula elliptica*. *PLoS ONE* 6 (1): e16069. doi:10.1371/journal.pone.0016069.
- Cummings, V., Thrush, S., Andrew, N., Norkko, A., Funnell, G., Budd, R., Hewitt, J., Gibbs, M., Mercer, S., Marriott, P., Anderson, O., 2003. Ecology and biodiversity of coastal benthic communities in McMurdo Sound, Ross Sea: emerging results, Unpublished Final Research Report for the Ministry of Fisheries, xx p.
- Cummings, V., Thrush, S., Chiantore, M., Hewitt, J., Cattaneo-Vietti, R., 2010. Macrobenthic communities of the north western Ross Sea shelf: links to depth, sediment characteristics and latitude. *Antarctic Science* 22 (6): 793-804.
- Cummings, V., Thrush, S., Marriott, P., Funnell, G., Norkko, A., Budd, R., 2008. Antarctic coastal marine ecosystems (ICECUBE), Unpublished Final Research Report for Ministry of Fisheries Research Project, 73 p.
- Cummings, V., Thrush, S., Norkko, A., Andrew, N., Hewitt, J., Funnell, G., Schwarz, A.-M., 2006a. Accounting for local scale variability in benthos: implications for future assessments of latitudinal trends in the coastal Ross Sea. *Antarctic Science* 18 (4): 633-644.
- Cummings, V., Thrush, S., Schwarz, A.-M., Funnell, G., Budd, R., 2006b. Ecology of coastal benthic communities of the northwestern Ross Sea., New Zealand Aquatic Environment and Biodiversity Report, No. 6, 67 p.
- Currey, R., Boren, L., Sharp, B., Peterson, D., 2012. A risk assessment of threats to Maui's dolphins, Ministry for Primary Industries and Department of Conservation, Wellington, 51 p.
- De Domenico, F., Chiantore, M., Buongiovanni, S., Paola Ferranti, M., Ghione, S., Thrush, S., Cummings, V., Hewitt, J., Kroeger, K., Cattaneo-Vietti, R., 2006. Latitude versus local effects on echinoderm assemblages along the Victoria Land coast, Ross Sea, Antarctica. *Antarctic Science* 18 (4): 655-662.
- Dettai, (+45 authors), 2011. DNA barcoding and molecular systematics of the benthic and demersal organisms of the CEAMARC survey. *Polar Science* 5 298-312.
- Doonan, I., 1995. Incidental catch of Hooker's sea lion in the southern trawl fishery for squid, summer 1994, N.Z. Fisheries Assessment Research Document 95/22, 13 p.
- Doonan, I., 1998. Estimation of sealion captures in southern fisheries in 1998. Final Research Report for Ministry of Fisheries Research Project ENV1997/01 Objective 2, 7 p.
- Doonan, I., 2001. Estimation of Hooker's sea lion, *Phocarctos hookeri*, captures in the southern squid trawl fisheries, 2001., New Zealand Fisheries Assessment Report 2001/67, 10 p.
- Doonan, I.J., 2000. Estimation of Hooker's sea lion, *Phocarctos hookeri*, captures in the southern squid trawl fisheries in 2000, New Zealand Fisheries Assessment Report 2000/41, 11 p.

- Duckworth, K., 1995. Analyses of factors which influence seabird bycatch in the Japanese southern bluefin tuna longline fishery in New Zealand waters, 1989-93, New Zealand Fisheries Assessment Report 1995/26, 59 p.
- Dunn, M., 2009 Feeding habits of the ommastrephid squid *Nototodarus sloanii* on the Chatham Rise, New Zealand. Journal of marine and Freshwater Research 43: 1103-1113
- Dunn, M., Connell, A., Forman, J., Stevens, D., Horn, P., 2010a. Diet of Two Large Sympatric Teleosts, the Ling (*Genypterus blacodes*) and Hake (*Merluccius australis*). PLoS ONE 5 (10): e13647. doi:10.1371/journal.pone.0013647
- Dunn, M., Griggs, L., Forman, J., PL, H., 2010b. Feeding habits and niche separation among the deep-sea chimaeroid fishes *Harriotta raleighana*, *Hydrolagus bemisi* and *Hydrolagus novaezealandiae* Marine Ecology Progress Series 407: 209-225.
- Dunn, M., Szabo, A., McVeagh, M., Smith, P., 2010c. The diet of deepwater sharks and the benefits of using DNA identification of prey. Deep-Sea Research Part 1 57 (7): 923-930.
- Dunn, M., Hurst, R., Renwick, J., Francis, R., Devine, J., McKenzie, A., 2009. Fish abundance and climate trends in New Zealand, New Zealand Aquatic Environment and Biodiversity Report. No. 31, 75 p.
- Dunn, M., PL, H., Connell, A., Stevens, D., Forman, J., Pinkerton, M., Griggs, L., Notman, P., B, W., In Press. Ecosystem-scale trophic relationships: diet composition and guild structure of middle-depth fish on the Chatham Rise., New Zealand Aquatic Environment and Biodiversity Report, xx p.
- Eakin, R., Eastman, J., Near, T., 2009. A new species and a molecular phylogenetic analysis of the Antarctic Fish Genus Pogonophryne (Nototheniidae: Artedidraconidae). Copeia No. 4: 705-713.
- Eléaume, M., Hemery, V., Bowden, D., Roux, M., 2011. A large new species of the genus *Ptilocrinus* (Echinodermata, Crinoidea, Hyocrinidae) from Antarctic seamounts. Polar Biology, DOI: 10.1007/s00300-011-0993-2.
- Eleaume, M., Hemery, V., Roussel, S., S, S., Steinke, D., Strugnell, J., Hibberd, T., Bohn, J., Riddle, M., Ameziane, N., In Prep. Southern Ocean crinoid diversity patterns and processes. Polar Biology
- Farr, T., Broom, J., Hart, D., Neill, K., Nelson, W., 2009. Common coralline algae of northern New Zealand: an identification guide, NIWA Information Series No. 70, 120 p.
- Fenaughty, J., Stevens, D., Hanchet, S., 2003. Diet of the Antarctic toothfish (*Dissostichus mawsoni*) from the Ross Sea, Antarctica (Subarea 88.1). CCAMLR Science 10: 113-123.
- Fenwick, G., Bradford-Grieve, J., 2002a. Human pressures on Ross Sea region marine communities: recommendations for future research, Unpublished Final Research Report for the Ministry of Fisheries, xx p.
- Fenwick, G., Bradford-Grieve, J., 2002b. Recommendations for future directed research to describe the biodiversity of the Ross Sea region, Unpublished Final Research Report for the Ministry of Fisheries, xx p.
- Fletcher, D., Mackenzie, D., Dillingham, P., 2008. Modelling of impacts of fishing-related mortality on NZ seabird populations, Unpublished Final Research Report on ENV2004/05, xx p.
- Floerl, O., J, H., Bowden, D., 2012. Chatham-Challenger OS 20/20 Post Voyage Analyses: (ZBD2007-01) Objective 9- Patterns in species composition. , New Zealand Aquatic Environment and Biodiversity Report No. 97, 40 p.
- Forman, J., Dunn, MR, 2010. The influence of ontogeny and environment on the diet of lookdown dory, *Cyttus traversi* New Zealand Journal of Marine and Freshwater Research 44 (4): 329-342.
- Forrest, B., Keeley, N., Gillespie, P., Hopkins, G., Knight, B., Govier, D., 2007. Review of the Ecological Effects of Marine Finfish Aquaculture: Final Report, Unpublished Cawthron Report for the Ministry of Fisheries, 80 p.
- Francis, M., Griggs, L., Baird, S., 2001. Pelagic shark bycatch in the New Zealand tuna longline fishery. Marine and Freshwater Research 52: 165-178.
- Francis, M., Griggs, L., Baird, S., 2004. Fish bycatch in New Zealand tuna longline fisheries, 1998-99 to 1999-2000, New Zealand Fisheries Assessment Report 2004/22, 62 p.
- Francis, M., Griggs, L., Baird, S., Murray, T., Dean, H., 1999. Fish bycatch in New Zealand tuna longline fisheries, NIWA Technical Report 55, 70 p.
- Francis, M., Griggs, L., Baird, S., Murray, T., Dean, H., 2000. Fish bycatch in New Zealand tuna longline fisheries, 1988-89 to 1997-98, NIWA Technical Report 76, 79 p.
- Francis, M., Lyon, W., 2012. Review of research and monitoring studies on New Zealand sharks, skates, rays and chimaeras, 2008-2012., New Zealand Aquatic Environment and Biodiversity Report No. 102, 70 p.
- Francis, M., Lyon, W., Jones, E., Notman, P., Parkinson, D., Getzlaff, C., 2012. Rig nursery grounds in New Zealand: a review and survey, New Zealand Aquatic Environment and Biodiversity Report No. 95p.
- Francis, M., Smith, M., 2010. Basking shark (*Cetorhinus maximus*) bycatch in New Zealand fisheries, 1994-95 to 2007-08, New Zealand Aquatic Environment and Biodiversity Report No. 49, 57 p.
- Francis, R., 2012. Fisheries Risks to the Population Viability of White-capped Albatross *Thalassarche steadi*, New Zealand Aquatic Environment and Biodiversity Report. No. 104, 24 p.
- Francis, R., Bell, E., 2010. Fisheries risks to the population viability of black petrel (*Procellaria parkinsoni*), New Zealand Aquatic Environment and Biodiversity Report No. 51, 57 p.
- Francis, R., Hadfield, M., Bradford-Grieve, J., Renwick, J., Sutton, P., 2005. Environmental predictors of hoki year-class strengths: an update, New Zealand Fisheries Assessment Report 2005/58p.
- Garcia, A., 2010. Comparative study of the morphology and anatomy of octopuses of the family Octopodidae. Auckland University of Technology, 247 p.
- Gardner, J., Bell, J., Constable, H., Hannan, D., Ritchie, P., Zuccarello, G., 2010. Multi-species coastal marine connectivity: a literature review with recommendations for further research, New Zealand Aquatic Environment and Biodiversity Report. No. 58, 47 p.
- Ghiglione, J.-F., Galand, P., Pommier, T., Pedros-Alio, C., Maas, E.W., Bakker, K., Bertilson, S., Kirchman, D., Lovejoy, C., Yager, P., Murray, A., 2012. Pole to pole biogeography of surface and deep marine bacterial communities. Proceedings of the national Academy of Sciences of the United States of America www.pnas.org/cgi/doi/10.1073/pnas.1208160109.
- Gordon, D., 2000. The Pacific Ocean and global OBIS: a New Zealand perspective. Oceanography 13: 41-47.
- Grange, K., Tovey, A., Hill, A., 2003. The spatial extent and nature of the bryozoan communities at Separation Point, Tasman Bay, Marine Biodiversity Biosecurity Report No. 4, 22 p.
- Grayling, S. 2004. A review of scientific studies conducted on the Macquarie Ridge. Unpublished Final Research Report (ZBD2003-09) for the Ministry of Fisheries. 33p.
- Griggs, L., Baird, S., Francis, M., 2007. Fish bycatch in the New Zealand tuna longline fisheries 2002-03 to 2004-05, New Zealand Fisheries Assessment Report 2007/18, 58 p.
- Griggs, L., Baird, S., Francis, M., 2008. Fish bycatch in New Zealand tuna longline fisheries in 2005-06., New Zealand Fisheries Assessment Report 2008/27, 47 p.

- Grotti, M., Soggia, F., Lagomarsino, C., Dalla Riva, S., Goessler, W., Francesconi, K., 2008. Natural variability and distribution of trace elements in marine organisms from Antarctic coastal environments. *Antarctic Science* 20 (1): 39-51.
- Guidetti, M., Marcato, S., Chiantore, M., Patarnello, T., Albertelli, G., Cattaneo-Vietti, R., 2006. Exchange between populations of *Adamussium colbecki* (Mollusca: Bivalvia) in the Ross Sea. *Antarctic Science* 18 (4): 645-653.
- Hanchet, S., 2000. Future research requirements for the Ross Sea Antarctic toothfish (*Dissostichus mawsoni*) fishery. , Unpublished Final Research Report for MFish Research Project MOF2000/02A Objective 8, 10 p.
- Hanchet, S., Fu, D., Dunn, A., 2008a. Indicative estimates of biomass and yield of Whitson's grenadier (*M. whitsoni*) on the continental slope of the Ross Sea in Subareas 88.1 and 88.2. , WG-FSA-08/32p.
- Hanchet, S., Mitchell, J., Bowden, D., Clark, M., Hall, J., O'Driscoll, R., 2008b. Ocean survey 20/20: New Zealand IPY-CAML Final Voyage Report, NIWA Client Report: WLG2008-74, October 2008, 193 p.
- Hanchet, S., Mitchell, J., Bowden, D., Clark, M., Hall, J., O'Driscoll, R., Pinkerton, M., Robertson, D., 2008c. Preliminary report of the New Zealand RV Tangaroa IPY-CAML survey of the Ross Sea region, Antarctica in February-March 2008, Unpublished NIWA report to CCAMLR working group on ecosystem monitoring and management, WG-EMM-08/18, 15 p.
- Hanchet, S., Stevenson, M., Jones, C., Marriott, P., McMillan, P., O'Driscoll, R., Stevens, D., Stewart, A., Wood, B., 2008d. Biomass estimates and size distributions of demersal finfish on the Ross Sea shelf and slope from the New Zealand IPY-CAML survey, February-March 2008., WG-FSA-08/31p.
- Hanchet, S., Stewart, A., McMillan, P., Clark, M., O'Driscoll, R., Stevenson, M., (In Press). Diversity, relative abundance, new locality records, and updated fish fauna of the Ross Sea, Antarctica. *Antarctic Science*.
- Hanchet, S.C., 2009. New Zealand IPY-CAML Progress Report, 76 p.
- Hanchet, S.C., 2010. New Zealand IPY-CAML Progress Report, 76 p.
- Hartill, B., Cryer, N., MacDiarmid, A., 2006. Reducing bycatch in New Zealand's scampi trawl fisheries, New Zealand Aquatic Environment and Biodiversity Report No. 4, 53 p p.
- Harvey, A., Woelkerling, W., Farr, T., Neill, K., Nelson, W., 2005. Coralline algae of central New Zealand: an identification guide to common 'crustose' species, NIWA Information Series No. 57p.
- Heimeier, D., Lavery, S., Sewell, MA, 2010. Using DNA barcoding and phylogenetics to identify Antarctic invertebrate larvae: Lessons from a large scale study. *Marine Genomics* 3: 165-177.
- Hemery, L., Eléaume, M., Roussel, V., Ameziane, N., Wilson, N., In Prep. Circumpolar distribution of a complex of cryptic species.
- Hewitt, J., Julian, K., Bone, E., 2011a. Chatham-Challenger Ocean Survey 20/20 Post-Voyage Analyses: Objective 10 - Biotic habitats and their sensitivity to physical disturbance., New Zealand Aquatic Environment and Biodiversity Report No. 81, 36 p.
- Hewitt, J., Lundquist, C., Bowden, D., 2011b. Chatham-Challenger Ocean Survey 20/20 Post Voyage Analyses: Diversity Metrics, New Zealand Aquatic Environment and Biodiversity Report No. 83, 64 p.
- Horn, P., Dunn, M., 2010. Inter-annual variability in the diets of hoki, hake, and ling on the Chatham Rise from 1990 to 2009, New Zealand Aquatic Environment and Biodiversity Report No. 54, 57 p.
- Horn, P., Forman, J., Dunn, MR., 2010. Feeding habits of alfonsino *Beryx splendens*. *Journal of Fish Biology* 76: 2382-2400
- Hurst, R., Renwick, J., Sutton, P., Uddstrom, M., Kennan, S., Law, C., Rickard, G., Korpela, A., Stewart, C., J. E., 2012. Climate and ocean trends of potential relevance to fisheries in the New Zealand Region, 2010, New Zealand Aquatic Environment and Biodiversity Report No. 90, 202 p.
- Hurst, R., Stevenson, M., Bagley, N., Griggs, L., Morrison, M., Francis, M., 2000. Areas of importance for spawning, pupping or egg-laying and juveniles of new Zealand coastal fish, Unpublished Final Research Report for Ministry of Fisheries Research Project ENV1999/03 Objective 1, 56 p.
- Jackson, J., Carroll, E., Smith, T., Patenaude, N., Baker, C., In Press. Taking stock – the historical demography of the New Zealand right whale (the Tohoro) 1820-2008, New Zealand Aquatic Environment and Biodiversity Report, xx p.
- Jellyman, D., 1994. The fishery for freshwater eels (*Anguilla* spp.) in New Zealand New Zealand Fisheries Assessment Research Document 94/14, 25 p.
- Jones, E., Francis, M., Paterson, C., Rush, N., Morrison, M., In Press. Habitats of particular significance for fisheries management: identification of threats and stressors to rig nursery areas., New Zealand Aquatic Environment and Biodiversity Report, xx p.
- Key, J., 2002. A review of current knowledge describing New Zealand's deepwater benthic biodiversity, Marine Biodiversity Biosecurity Report No. 1, 25 p.
- Koubbi, P., Masato, M., Duhamel, G., Goarant, A., Hulley, P.-A., O'Driscoll, R., Takashi, I., Pruvost, P., Tavenier, E., Hosie, G., 2011. Ecological importance of micronektonic fish for the ecoregionalisation of the Indo-Pacific sector of the Southern Ocean: role of myctophids. . *Deep Sea Research II* 58: 170-180.
- Lalas, C., MacDiarmid, A., Abraham, E., In Press a. Estimates for annual consumption by a recovered population of New Zealand fur Seals., New Zealand Aquatic Environment and Biodiversity Report xx p.
- Lalas, C., MacDiarmid, A., Abraham, E., In Press b. Estimates for annual consumption rates by New Zealand sea lions., New Zealand Aquatic Environment and Biodiversity Report xx p.
- Lalas, C., MacDiarmid, A., In Press. Rapid recolonisation of south-eastern New Zealand by New Zealand fur seals *Arctocephalus forsteri*, New Zealand Aquatic Environment and Biodiversity Report xx p.
- Lavery, S., Clements, K., Hickey, A., 2006. Molecular identification of cryptogenic/invasive gobies in New Zealand., New Zealand Aquatic Environment and Biodiversity Report No. 5, 48 p.
- Leathwick, J., Dey, K., Julian, K., 2006a. Development of a marine environmental classification optimised for demersal fish, NIWA Client report HAM2006-063p.
- Leathwick, J., Elith, J., Francis, M., Hastie, T., Taylor, P., 2006b. Variation in demersal fish species richness in the oceans surrounding New Zealand: an analysis using boosted regression trees. *Marine Ecology Progress Series* 321: 267-281.
- Leathwick, J., Francis, M., Julian, K., 2006c. Development of a demersal fish community map for New Zealand's Exclusive Economic Zone, NIWA Client Report HAM2006-062, prepared for Department of Conservation, National Institute of Water & Atmospheric Research, Hamilton, New Zealand. p.
- Leathwick, J., Rowden, A., Nodder, S., Gorman, R., Bardsley, S., Pinkerton, M., Baird, S., Hadfield, M., Currie, K., Goh, A., 2012. A Benthic-optimised Marine Environment Classification (BOMEc) for New Zealand waters, New Zealand Aquatic Environment and Biodiversity Report No 89, 54 p.

- Leathwick JR, Rowden A, Nodder S, Gorman R, Bardsley S, Pinkerton M, Baird SJ, Hadfield M, Currie K, Goh A 2010. Development of a benthic-optimised marine environment classification for waters within the New Zealand EEZ. Final Research Report for Ministry of Fisheries Research Project BEN200601, Objective 5.
- Leduc, D., Rowden, A., Bowden, D., Probert, P., Pilditch, C., 2012a. Direct evidence for a unimodal relationship between productivity and deep-sea benthic diversity. *Marine Ecology Progress Series* 454: 53-64.
- Leduc, D., Rowden, A., Bowden, D., Probert, P., Pilditch, C., Nodder, S., 2012b. Unimodal relationship between biomass and species richness of deep-sea nematodes: implications for the link between productivity and diversity. *Marine Ecology Progress Series* 454: 53-64.
- Linse, K., Cope, T., Lörz, A., Sands, C., 2007. Is the Scotia Sea a centre of Antarctic marine diversification? Some evidence of cryptic speciation in the circum-Antarctic bivalve *Lissarca notorcadensis* (Arcoidea : Philobryidae). *Polar Biology* 30: 1059-1068.
- Livingston, M., 2004. A sampling programme to construct and quantify food-webs in two key areas supporting important fish and invertebrate species in New Zealand, Unpublished Final Research Report for Ministry of Fisheries Project ENV2002-07, Objective 1 NIWA, xx p.
- Livingston, M., 2009. Towards a National Marine Environment Monitoring Programme in New Zealand: A discussion paper submitted to the Biodiversity Research Advisory Group Workshop on Marine Environmental Monitoring, Unpublished report held at MPI, Wellington, xx p.
- Livingston, M., Clark, M., Baird, S., 2003. Trends in incidental catch of major fisheries on the Chatham Rise for Orfishing years 1989-90 to 1998-99., New Zealand Fisheries Assessment Report 2003/52, 74 p.
- Lohrer, A., Chiaroni, L., Thrush, S.F., Hewitt, J., 2010. Isolated and interactive effects of two key species on ecosystem function and trophic linkages in New Zealand soft-sediment habitats, New Zealand Aquatic Environment and Biodiversity Report No. 44, 69 p.
- Lohrer, A., Cummings, V., Thrush, S., 2012. Altered sea ice thickness and permanence affects benthic ecosystem functioning in coastal Antarctica. *Ecosystems* 10.1007/s10021-10012-19610-10027.
- Lorrey, A., Goff, J., McFadgen, B., Chagué-Goff, C., Neil, H., MacDiarmid, A., In Press. A synthesis of climatic and geophysical driver activity in New Zealand and environmental changes during the last 1000 years, New Zealand Aquatic Environment and Biodiversity Report, xx p.
- Lörz, A., 2009. Synopsis of Amphipoda from two recent Ross Sea voyages with description of a new species of *Epimeria* (Epimeriidae, Amphipoda, Crustacea). *Zootaxa* 2167: 59-68.
- Lörz, A., 2010a. Deep-sea Rhachotropis (Crustacea: Amphipoda: Eusiridae) from New Zealand and the Ross Sea with key to the Pacific, Indian Ocean and Antarctic species. *Zootaxa* 2482: 22-48.
- Lörz, A., 2010b. New Zealand Amphipoda, Conference presentation, XIVth International Colloquium on Amphipoda, Seville, Spain p.
- Lörz, A.-N., 2010c. Deep-sea Rhachotropis (Crustacea: Amphipoda: Eusiridae) from New Zealand and the Ross Sea with key to the Pacific, Indian Ocean and Antarctic species. *Zootaxa* 2482 22-48.
- Lörz, A., 2011a. Biodiversity of unknown New Zealand habitat: bathyal invertebrate assemblages in the benthic boundary layer. *Marine Biodiversity* 41 (2): 287-297.
- Lörz, A., 2011b. Pacific Epimeriidae (Amphipoda, Crustacea): Epimeria. *Journal of the Marine Biological Association of the United Kingdom* 91 (2): 471-477.
- Lörz, A., Maas, E., Linse, K., Coleman, C., 2009 Do circum-Antarctic species exist in peracarid Amphipoda? A case study in the genus *Epimeria* Costa, 1851 (Crustacea, Peracarida, Epimeriidae). In: Bruce N (Ed) *Advances in the taxonomy and biogeography of Crustacea in the Southern Hemisphere*. *ZooKeys* 18: 91-128.
- Lörz, A., Maas, E., Linse, K., Fenwick, G., 2007. *Epimeria schiaparelli* sp. nov., an amphipod crustacean (family Epimeriidae) from the Ross Sea, Antarctica, with molecular characterisation of the species complex. *Zootaxa* 1402: 23-37.
- Lörz, A., Smith, P., Linse, K., Steinke, D., 2012a. First Molecular Evidence for Underestimated Biodiversity of Rhachotropis (Crustacea, Amphipoda), with Description of a New Species. *PLoS ONE* 7 (3): e32365. doi:32310.31371/journal.pone.0032365.
- Lörz, A.-N., Berkenbusch, K., Nodder, S., Ahyong, S., Bowden, D., McMillan, P., Gordon, D., Mills, L., Mackay, K., 2012b. A review of deep-sea biodiversity associated with trench, canyon and abyssal habitats deeper than 1500 m in New Zealand waters New Zealand Aquatic Environment and Biodiversity Report, No. 92, 133 p.
- Lörz, A., Smith, P., Linse, K., Steinke, D., In Prep. Molecular insights to deep-sea Rhachotropis from New Zealand and the Ross Sea with description of a new species.
- Lörz, A., Smith, P., Linse, K., Steinke, D., In press. High genetic diversity within *Epimeria georgiana* (Amphipoda) from the southern Scotia Arc. *Marine Biodiversity*.
- Lörz, A.-N., Coleman, O., 2009. Living jems: jewel-like creatures from the deep. *Water and Atmosphere* 17 (1): 16-17.
- Lundquist, C., Thrush, S., Coco, G., Hewitt, J., 2010. Interactions between distributions and dispersal decrease persistence thresholds of a marine benthic community. *Marine Ecology Progress Series* 413: 217-228.
- Lyle, J., 2011. SRP2010-03: Fur seal interactions with SED excluder device, Unpublished Final Research Report for Ministry of Fisheries, 26 July 2011, 20 p.
- Maas, E., Voyles, K., Pickmere, S., Hall, J., Bowden, D., Clark, M., 2010a. Bacterial and Archaeal diversity and exo-enzyme activity in Ross Sea, Antarctica sediments. , Poster presented at SAME 11- Symposium on Aquatic Microbial Ecology, Piran, Slovenia, 30th August - 4th September 2010.p.
- Maas, E., Law, C., Hall, J., Chang, H., Bury, S., Robinson, K., Thompson, K., Voyles, M., 2010b. Impacts of ocean acidification on planktonic ecosystems of the southern ocean, Unpublished Final Research Report for Ministry of Fisheries Project ZBD2008-22, held by the Ministry of Fisheries, 44 p.
- MacDiarmid, A., Cleaver, P., Stirling, B., In Press a. Historical evidence for the state and exploitation of the marine environment in the Hauraki Gulf and along the Otago-Catkins shelf 1769 - 1950, New Zealand Aquatic Environment and Biodiversity Report xx p.
- MacDiarmid, A., Smith, I., Paul, L., Francis, M., McKenzie, A., Parsons, D., Hartill, B., Stirling, B., Cleaver, P., In Press b. A complete history of the exploitation of an ecologically important inshore finfish species in the Hauraki Gulf, New Zealand: a synthesis of archaeological, historical and fisheries data, New Zealand Aquatic Environment and Biodiversity Report Aquatic Environment and Biodiversity Report xx p.
- MacDiarmid, A., Beaumont, J., MacDiarmid, A., Morrison, M., McKenzie, A., Abraham, E., Taylor, R., Gillanders, B., Bury, S., Cowles, A., Parsons, D., Cole, R., Pinkerton, M., Walker, J., In Press c. Rocky reef ecosystems – how do they function, New Zealand Aquatic Environment and Biodiversity Report xx p.

- MacDiarmid, A., McKenzie, A., Sturman, J., Beaumont, J., Mikaloff-Fletcher, S., Dunne, J., 2012. Assessment of anthropogenic threats to New Zealand marine habitats, Aquatic Environment and Biodiversity Report No. 93, 255 p.
- MacDiarmid, A., Stewart, R., In Press. Routine and opportunistic sampling of Ross Sea and Balleny Islands Biodiversity, New Zealand Aquatic Environment and Biodiversity Report xx p.
- MacKenzie, D., Fletcher, D., 2006. Characterisation of seabird captures in NZ fisheries, Unpublished Final Research Report prepared for the Ministry of Fisheries, Proteus Wildlife Research Consultants, 99 p.
- MacKenzie, D., Fletcher, D., 2010. Designing a programme of indicators of population performance with respect to captures of seabirds in New Zealand fisheries, Unpublished Final Research Report for Ministry of fisheries project ENV2005-08, held by the Ministry of Fisheries.
- Marriot, P., Horn, P., McMillan, P., 2003. Species identification and age estimation for the ridge-scaled Macrourid (*Macrourus whitsoni*) the Ross Sea. CCAMLR Science 10: 37-51.
- Mattlin, R., 1993. Incidental catch of fur seals in the west coast South Island hoki trawl fishery, 1989-92, New Zealand Fisheries Assessment Report 1993/19p.
- Mattlin, R., 2004. QMA SQU6T New Zealand sea lion incidental catch and necropsy data for the fishing years 2000-01, 2001-02 and 2002-03, Unpublished Report prepared for the NZ Ministry of Fisheries, Wellington, NZ. July 2004, 21 p.
- Mattlin, R.H., 1993. Incidental catch of fur seals in the west coast South Island hoki trawl fishery, 1989-92, New Zealand Fisheries Assessment Research Document 93/19., 17 p.
- Maunder, M., 2007. Assessment of risk of yellow-eyed penguins *Megadyptes antipodes* from fisheries incidental mortality in New Zealand fisheries and definition of information requirements for managing fisheries related risk Unpublished Final Research Report for the Ministry of Fisheries, 29 p.
- Maxwell, K., MacDiarmid, A., In Press. Oral histories of Maori marine resource state and use in the Hauraki Gulf and along the Otago-Catkins coast 1940-2008, Darft Aquatic Environment and Biodiversity Report xx p.
- McElderry, H., Schrader, J., Anderson, S., 2007. Electronic Monitoring to Assess Protected Species Interactions in New Zealand Longline Fisheries: A Pilot Study Unpublished report prepared for the New Zealand Ministry of Fisheries, Wellington, NZ by Archipelago Marine Research Ltd., Victoria, British Columbia, Canada., 45 p.
- McElderry, H., Schrader, J., Anderson, S., 2008. Electronic monitoring to assess protected species interactions in New Zealand longline fisheries: a pilot study, New Zealand Aquatic Environment and Biodiversity Report No. 24, 39 p.
- McMillan, P., Griggs, L., Francis, M., Marriott, P., Paul, L., Mackay, E., Wood, B., Sui, H., Wei, F., 2011a. New Zealand fishes. 1: A guide to common species caught by trawling, New Zealand Aquatic Environment and Biodiversity Report. No. 68, 331 p.
- McMillan, P., Griggs, L., Francis, M., Marriott, P., Paul, L., Mackay, E., Wood, B., Sui, H., Wei, F., 2011b. New Zealand fishes. 2: A guide to less common species caught by trawling, New Zealand Aquatic Environment and Biodiversity Report. No. 78, 184 p.
- McMillan, P., Griggs, L., Francis, M., Marriott, P., Paul, L., Mackay, E., Wood, B., Sui, H., Wei, F., 2011c. New Zealand fishes 3: A guide to common species caught by surface fishing methods, New Zealand Aquatic Environment and Biodiversity Report No. 69, 147 p.
- McMillan, P., Iwamoto, T., Stewart, A., Smith, P., 2012. A new species of grenadier, genus *Macrourus* (Teleostei, Gadiformes, Macrouridae) from the Antarctic and a revision of the genus. Zootaxa 3165: 1-24.
- McMillan, P., Marriott, P., Hanchet, S., Fenaughty, J., Mackay, E., Sui, H., 2007. Field identification guide to the main fishes caught in the Ross Sea long-line fishery., Unpublished Final Research Report to the Ministry of Fisheries. Project ANT2005/02, objective 7. , 14 p.
- Meynier, L., 2010. New Zealand sea lion bio-energetic modelling, Unpublished Final Report for Project IPA2009-09. Presented to AEWG May 17 2011, Wellington, New Zealand, 34 p.
- Michael, K., Kroger, K., Richardson, K., Hill, N., 2006. Summary of information in support of the Foveaux Strait Oyster Fishery Plan: the Foveaux Strait ecosystem and effects of oyster dredging, Unpublished Final Research Report for Ministry of Fisheries project ZBD2005-04, xx p.
- Middleton, D., Abraham, E., 2007. The efficacy of warp strike mitigation devices: trials in the 2006 squid fishery, Unpublished Final Research Report for the Ministry of Fisheries for project IPA2006-02, xx p.
- Ministry of Fisheries, 2008. Bottom Fishery Impact Assessment: Bottom Fishing Activities by New Zealand Vessels Fishing in the High Seas in the SPRFMO Area during 2008 and 2009, Available from <http://www.southpacificrmo.org/benthic-impact-assessments/>, Wellington, 102 p.
- Mitchell, J., 2008. Initial environmental evaluation (IEE) NZ IPY-CAML voyage 2008 report, Unpublished Report prepared for Antarctic Policy Unit, Ministry of Foreign Affairs and Trade., 35 p.
- Mitchell, J., Clark, M., 2004. Voyage report TAN04-02. Western Ross Sea voyage 2004. Hydrographic and biodiversity survey RV Tangaroa 27 January to 13 March 2004. Cape Adare, Cape Hallett, Possession Islands and Balleny Islands, Antarctica. , NIWA Voyage Report TAN04-02, Unpublished report held in NIWA library, Wellington, 102 p.
- Mitchell, J., MacDiarmid, A., 2006. Voyage Report TAN06-02 Eastern Ross Sea Voyage, Unpublished report for MFish project ZBD2005-03, xx p.
- Mormede, S., Baird, S., Smith, M., 2008 Factors that may influence the probability of fur seal capture in selected New Zealand fisheries, New Zealand Aquatic Environment and Biodiversity Report. No. 19, 42 p.
- Morrison, M., D, P., 2008. Distribution and abundance of toheroa (*Paphies ventricosa*) on Ninety Mile Beach, March 2006, New Zealand Fisheries Assessment Report 2008/26, 27 p.
- Morrison, L., Lowe, M., Parsons, D., Usmar, N., McLeod, I., 2009. A review of land-based effects on coastal fisheries and supporting biodiversity in New Zealand, New Zealand Aquatic Environment and Biodiversity Report No. 37, 100 p.
- Murray, T., Taylor, P., Greaves, J., Bartle, J., Molloy, J., 1992. Seabird bycatch by Southern Fishery longline vessels in New Zealand waters, New Zealand Fisheries Assessment Research Document 92/22, 21 p.
- Naylor, J., Webber, W., Booth, J., 2005. A guide to common offshore crabs in New Zealand waters., New Zealand Aquatic Environment and Biodiversity Report No. 2, 47 p.
- Neil, H., MacDiarmid, A., Marriot, P., Paul, L., Horn, P., In Press. Insights into historical New Zealand marine shelf productivity using ancient fish otoliths, Unpublished Final Research Report for Project ZBD200505 MS6 - Part B, Ministry of Fisheries, Wellington, New Zealand., xx p.
- Neill, K., D'Archino, R., Farr, T., Nelson, W., 2012. Macroalgal diversity associated with soft sediment habitats in New Zealand, New Zealand Aquatic Environment and Biodiversity Report No. 87, 130 p.
- Nelson, W., Cummings, V., D'Archino, R., Halliday, J., Marriott, P., Neill, K., 2010. Macroalgae and benthic biodiversity of the Balleny Islands, Southern Ocean, New Zealand Aquatic Environment and Biodiversity Report No. 55, 99 p.

- Nelson, W., Neill, K., Farr, T., Barr, N., D'Archino, R., Miller, S., Stewart, R., 2012. Rhodolith Beds in Northern New Zealand: Characterisation of Associated Biodiversity and Vulnerability to Environmental Stressors., New Zealand Aquatic Environment and Biodiversity Report No. 99, 102 p.
- Nelson, W., Villouta, E., Neill, K., Williams, G., Adams, N., Slivsgaard, R., 2002. Marine Macroalgae of Fiordland, New Zealand *Tuhinga* 13: 117-152
- Nielsen, J., Lavery, S., Lörz, A., 2009. Synopsis of a new collection of sea spiders (Arthropoda: Pycnogonida) from the Ross Sea, Antarctica. *Polar Biology* 32: 1147-1155.
- Nodder, S., 2008. OS 20/20 Chatham Rise & Challenger Plateau Hydrographic, Biodiversity & Seabed Habitats, NIWA Client Report: WLG2008-27, National Institute of Water & Atmospheric Research, Wellington, New Zealand.
- Nodder, S., Bowden, D., A, P., Mackay, K., 2012. Seafloor habitats and benthos of a continental ridge: Chatham Rise, New Zealand. , in: Harris, P., Baker, E. (Eds.), *Seafloor geomorphology as benthic habitat: GEOHAB atlas of geomorphic features and benthic habitats*. Elsevier, London, pp. 763-776.
- Nodder, S., Maas, E., Bowden, D., Pilditch, C., 2011. Physical, biogeochemical, and microbial characteristics of sediment samples from the Chatham Rise and Challenger Plateau., *Aquatic Environment and Biodiversity Report No. 70p*.
- Norkko, A., Andrew, N., Thrush, S., Cummings, V., Schwarz, A.-M., Hawes, I., Mercer, S., Budd, R., Gibbs, M., Funnell, G., Hewitt, J., Goring, D., 2002. Ecology and biodiversity of coastal benthic communities in McMurdo Sound, Ross Sea: development of sampling protocols and initial results, Unpublished Final Research Report for Ministry of Fisheries Research Project ZBD2001/02, Objectives 1, 2 & 3. , 119 p.
- Norkko, A., Thrush, S., Cummings, V., Funnell, G., Schwarz, A.-M., Andrew, N., Hawes, I., 2004. Ecological role of Phyllophora antarctica drift accumulations in coastal soft-sediment communities of McMurdo Sound, Antarctica. *Polar Biology* 27: 482-494.
- Norkko, A., Thrush, S., Cummings, V., Gibbs, M., Andrew, N., Norkko, J., Schwarz, A.-M., 2007. Trophic structure of coastal Antarctic food webs associated with changes in sea ice and food supply. *Ecology* 88: 2810-2820.
- Norkko, J., Norkko, A., Thrush, S., Cummings, V., 2005. Growth under environmental extremes: spatial and temporal patterns in nucleic acid ratios in two Antarctic bivalves. *Journal of Experimental Marine Biology and Ecology* 326: 114-156
- O'Driscoll, R., Hurst, R., Dunn, M., Gauthier, S., Ballara, S., 2011. Trends in relative mesopelagic biomass using time series of acoustic backscatter data from trawl surveys, New Zealand Aquatic Environment and Biodiversity Report. No. 96, 99 p.
- O'Driscoll, R., Macaulay, G., Gauthier, S., Pinkerton, M., Hanchet, S., 2010. Distribution, abundance and acoustic properties of Antarctic silverfish (*Pleuragramma antarcticum*) in the Ross Sea. CCAMLR document WG-FSA-10/P4 58 (1-2): 181-195.
- O'Driscoll, R., 2009 Preliminary acoustic results from the New Zealand IPY-CAML survey of the Ross Sea region in February-March 2008, Unpublished Final Research Report for MFish project IPY200701 objective 8, 14 p.
- O'Driscoll, R., Booth, J., Bagley, N., Anderson, O., Griggs, L., Stevenson, M., Francis, M., 2003. Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand deepwater fish, pelagic fish, and invertebrates., NIWA Technical Report 119, 377 p.
- O'Driscoll, R., Hanchet, S., Miller, B., In press. Can acoustic methods be used to monitor grenadier (Macrouridae) abundance in the Ross Sea region. *Journal of Ichthyology*.
- O'Driscoll, R., Macaulay, G., Gauthier, S., Pinkerton, M., Hanchet, S., 2009. Preliminary acoustic results from the New Zealand IPY-CAML survey of the Ross Sea region in February-March 2008, CCAMLR SG-ASAM-09-05, 37 p.
- O'Loughlin, M., Paulay, G., Davey, N., Michonneau, F., 2011. The Antarctic region as a marine biodiversity hotspot for echinoderms: Diversity and diversification of sea cucumbers. *Deep Sea Research Part II: Topical Studies in Oceanography* 58: 264-275.
- Page, M., Alcock, N., Gordon, D., Kelly-Shanks, M., Nelson, W., Neill, K., Watson, J., 2001. Preliminary assessment of the biodiversity of benthic macrofauna of the western Ross Sea, Antarctica, Unpublished Final Research Report to the Ministry of Fisheries, 29 p.
- Pakhomov, E., Hall, J., Williams, M., Hunt, B., Stevens, C., 2011. Biology of *Salpa thompsoni* in waters adjacent to the Ross Sea, Southern Ocean, during austral summer 2008. *Polar Biology* 34: 257-271.
- Parker, S., 2008. Development of a New Zealand High Seas Bottom Trawling Benthic Assessment Standard for Evaluation of Fishing Impacts to Vulnerable Marine Ecosystems in the South Pacific Ocean, Unpublished Final Research Report for Ministry of Fisheries Research Projects IFA2007-02 Objectives 3 and 4p.
- Parker, S., Bowden, D., 2010. Identifying taxonomic groups vulnerable to bottom longline fishing gear in the Ross Sea region. *CCAMLR Science* 17: 105-127.
- Parker, S., Francis, M., 2012. Productivity of two species of deepwater sharks, *Deania calcea* and *Centrophorus squamosus* in New Zealand, *Aquatic Environment and Biodiversity Report No. 103*, 48 p.
- Parker, S., Penney, A., Clark, M., 2009a. Detection criteria for measuring trawl impacts on vulnerable marine ecosystems in high sea fisheries of the South Pacific Ocean. *Marine Ecology Progress Series* 397: 309-317.
- Parker, S., Mormede, S., Tracey, D., Carter, M., 2009b. Evaluation of VME taxa monitoring by scientific observers from five vessels in the Ross Sea Antarctic toothfish longline fishery during the 2008-09 season, Unpublished Final Research Report for the Ministry of Fisheries, xx p.
- Parker, S., Tracey, D., Mackay, E., Mills, S., Marriott, P., Anderson, O., Schnabel, K., Bowden, D., Lockhart, S., 2009c. CCAMLR VME taxa Identification guide Version 2009., Available at <http://www.ccamlr.org/pu/e/sc/obs/vme-guide.pdf>, Hobart, Tasmania, Australia, 4 p.
- Parker, S., Tracey, D., Mackay, E., Mills, S., Marriott, P., Anderson, O., Schnabel, K., Bowden, D., Kelly, M., 2008. Classification guide for potentially vulnerable invertebrate taxa in the Ross Sea long-line fishery, Unpublished Final Research Report for Ministry of Fisheries Research Project ANT2007-01 (Objective 3), 7 p.
- Parker, S., Wood, B., Hanchet, S., Dunn, A., 2010. A bathymetric data framework for fisheries management in the Ross Sea region, Unpublished Final Research Report for the Ministry of Fisheries, 14 p.
- Parsons, D., Morrison, M., MacDiarmid, A., Stirling, B., Cleaver, P., Smith, I., Butcher, M., In press. Long-term effects of climate variation and human impacts on the structure and functioning of New Zealand's Shelf Ecosystems, New Zealand Aquatic Environment and Biodiversity Report xx p.
- Paul, L., 2012. A history of the Firth of Thames dredge fishery for mussels: use and abuse of a coastal resource, New Zealand Aquatic Environment and Biodiversity Report No. 94, 27 p.
- Penney, A., Parker, S., Brown, J., 2009. Protection measures implemented by New Zealand for vulnerable marine ecosystems in the South Pacific Ocean. *Marine Ecology Progress Series* 397: 341-354.

- Pinkerton, M., In Press. Trophic modelling of a New Zealand marine shelf ecosystem since AD 1000 New Zealand Aquatic Environment and Biodiversity Report xx p.
- Pinkerton, M., Bradford-Grieve, J., Hanchet, S., In Review. A balanced model of the food web of the Ross Sea, Antarctica. Submitted to CCAMLR Science.
- Pinkerton, M., Hanchet, S., Bradford-Grieve, J., Cummings, V., Wilson, P., Williams, M., 2006. Modelling the effects of fishing in the Ross Sea, Unpublished Final Research Report for Ministry of Fisheries Research Project ANT2004-05, 169 p.
- Pinkerton, M., Cummings, V., Forman, J., Brown, J., Bury, S., 2009b. Trophic connections in the Ross Sea: information from stomach contents analysis and stable isotopes of carbon and nitrogen., Unpublished Report to Ministry of Fisheries, project IPY200701 Obj10, 18 p.
- Pinkerton, M., Dunn, A., Hanchet, S., 2007a. Ecological risk management and the fishery for Antarctic toothfish (*Dissostichus mawsoni*) in the Ross Sea, Antarctica, Document EMM-07-24, CCAMLR, Hobart, Australia, 22 p.
- Pinkerton, M., Bury, S., Vorster, I., Hanchet, S., 2007b. Stable isotope analysis of Southern Ocean fish tissue samples: preliminary results, Unpublished Final Research Report to the Ministry of Fisheries. Project ANT2005/04, objective 1, 32 p.
- Pinkerton, M., Forman, J., Bury, S., Brown, J., Horn, P., O'Driscoll, R., In press. Diet and trophic niche of Antarctic silverfish (*Pleuragramma antarcticum*) in the Ross Sea, Antarctica. Journal of Fish Biology.
- Pinkerton, M., Hanchet, S., Bradford Grieve, J., 2005. Modelling the effects of fishing in the Ross Sea, Unpublished Final Research Report for MFish Research Project ANT2004/05, 130 p.
- Pinkerton, M., Mormede, S., Bowden, D., 2009a. Inputs to the Ross Sea Bioregionalisation/Spatial Management planning workshop, Unpublished Final Research Report to the Ministry of Fisheries, 40 p.
- Ponte, G., van den Berg, A., Anderson, R., 2010. Impact characteristics of the New Zealand Fisheries sea lion exclusion device stainless steel grid, IPA2009-16 Report prepared for NZ Ministry of Fisheries (Project IPA2009-16), Oct. 2010, 24 p.
- Ponte, G., van den Berg, A., Anderson, R., 2011. Further analysis of the impact characteristics of the New Zealand Fisheries sea lion exclusion device stainless steel grid, Unpublished Report prepared for NZ Ministry of Fisheries (Project SRP2010-05), 24 p.
- Richard, Y., Abraham, E., Filippi, D., 2011. Assessment of the risk to seabird populations from New Zealand commercial fisheries, Unpublished Final Research Report held by the Ministry of Fisheries, xx p.
- Roberts, C., Stewart, A., 2001. Ross Sea fishes: a collection-based biodiversity research programme. Seafood New Zealand 9 (11): 79-84.
- Roe, W., 2007. Necropsy of marine mammals captured in New Zealand fisheries in the 2005-06 fishing year, New Zealand Aquatic Environment and Biodiversity Report No. 11, 24 p.
- Roe, W., 2009a. Necropsy of marine mammals captured in New Zealand fisheries in the 2007-08 fishing year, New Zealand Aquatic Environment and Biodiversity Report. No. 34, 35 p.
- Roe, W., 2009b. Necropsy of marine mammals captured in New Zealand fisheries in the 2006-07 fishing year, New Zealand Aquatic Environment and Biodiversity Report No. 29, 32 p.
- Roe, W., 2010a External review of NZ sea lion bycatch necropsy data and methods, Unpublished Report prepared for the NZ Ministry of Fisheries, Wellington, 8 p.
- Roe, W., 2010b. Necropsy of marine mammals captured in New Zealand fisheries in the 2008-09 fishing year, New Zealand Aquatic Environment and Biodiversity Report No. 47, 22 p.
- Roe, W., Meynier, L., 2012. Review of Necropsy Records for Bycaught NZ sea lions (*Phocarcos hookeri*), 2000-2008, New Zealand Aquatic Environment and Biodiversity Report No. 98, 46 p.
- Rowden, A., Clark, M., 2010. Benthic biodiversity of seven seamounts on the southern end of the Kermadec volcanic arc, northeast New Zealand, New Zealand Aquatic Environment and Biodiversity Report No. 62, 31 p.
- Rowden, A., Clark, M., O'Shea, S., 2004. Benthic biodiversity of seamounts on the Northland Plateau, Marine Biodiversity Biosecurity Report No. 5, 21 p.
- Rowden, A., Clark, M., O'Shea, S., McKnight, D., 2003. Benthic biodiversity of seamounts on the southern Kermadec volcanic arc, Marine Biodiversity Biosecurity Report. No. 3, 23 p.
- Rowden, A., Cranfield, H., Mann, P., Wood, A.L., 2007. Benthic macrofauna bycatch of oyster dredging in Foveaux Strait., New Zealand Fisheries Assessment Report 2007/10, 27 p.
- Rowden, A., Kroeger, K., Clark, M., 2012a. Biodiversity of macroinvertebrates and fish assemblages of the northwestern Ross Sea shelf, New Zealand Aquatic Environment and Biodiversity Report No. 101, 111 p.
- Rowden, A., Berkenbusch, K., Brewin, P., Dalen, J., Neill, K., Nelson, W., Oliver, M., Probert, P., Schwarz, A.-M., Sui, P., Sutherland, D., 2012b. A review of the marine soft-sediment assemblages of New Zealand, New Zealand Aquatic Environment and Biodiversity Report No. 92, 165 p.
- Rowden, A., Kroeger, K., Clark, M., In Press. Biodiversity of macroinvertebrates and fish assemblages of the Balleny Islands seamounts, New Zealand Aquatic Environment and Biodiversity Report, xx p.
- Rowden, A., Oliver, M., Clark, M., Mackay, K., 2008. New Zealand's "SEAMOUNT" database: recent updates and its potential use for ecological risk assessment, New Zealand Aquatic Environment and Biodiversity Report. No. 27, 49 p.
- Rowden, A., O'Shea, S., Clark, M., 2002. Benthic biodiversity of seamounts on the northwest Chatham Rise. , Marine Biodiversity and Biosecurity Report. 2, 1-21 p.
- Sagar, P., Charteris, M., Carroll, J., Sagar, P., 2009a. Population assessment of Salvin's albatrosses at the Snares Western Chain, 29 September - 17 October 2008, Unpublished Final Research Report for the Ministry of Fisheries, 4 p.
- Sagar, P., Fraser, M., Hunt, S., Scofield, P., Robertson, C., 2009b. Population dynamics of the Chatham Albatross at The Pyramid, 9 November- 7 December 2008, Unpublished Final Research Report for the Ministry of Fisheries, 5 p.
- Sagar, P., Carroll, J., Charteris, M., 2010a. Population assessment of Salvin's albatrosses at the Snares Western Chain, 29 September - 14 October 2009, Unpublished Final Research Report for Ministry of Fisheries, 4 p.
- Sagar, P., Fraser, M., Cameron, N., Scofield, P., Robertson, C., 2010b. Population assessment of Northern Buller's Albatross and Northern Giant Petrels at the Forty-Fours, Chatham Islands, 1 - 8 December 2009, Unpublished Final Research Report to the Ministry of Fisheries. Project PRO2006-05, 11 p.
- Sagar, P., Fraser, M., Palmer, D., Deppe, L., Scofield, P., Robertson, C., 2010c. Population dynamics of the Chatham Albatross at The Pyramid, 20 November- 14 December 2009, Unpublished Final Research Report for the Ministry of Fisheries, 6 p.
- Sagar, P., Thompson, D., 2008. Data collection of demographic, distributional, and trophic information on selected petrels, New Zealand Aquatic Environment and Biodiversity Report No. 17, 12 p.

- Savage, C., 2009. Development of bioindicators for the assimilation of terrestrial nutrient inputs in coastal ecosystems as a tool for watershed management, New Zealand Aquatic Environment and Biodiversity Report No. 30, 35 p.
- Savage, C., Thrush, S., Lohrer, A., Hewitt, J., 2012. Ecosystem Services Transcend Boundaries: Estuaries Provide Resource Subsidies and Influence Functional Diversity in Coastal Benthic Communities. PLoS ONE 7 (8): e42708. doi:10.1371/journal.pone.0042708.
- Schiaparelli, S., Alvaro, M., Bohn, J., Albertelli, G., 2010 'Hitchhiker' polynoid polychaetes in cold deep waters and their potential influence on benthic soft bottom food webs. Antarctic Science 22: 399-407.
- Schiaparelli, S., Lörz, A., Cattaneo-Vietti, R., 2006. Diversity and distribution of mollusc assemblages on the Victoria Land coast and the Balleny Islands, Ross Sea, Antarctica. Antarctic Science 18 (4): 615-631.
- Schiaparelli, S., Olivero, M., Taviani, M., Griffiths, H., Lörz, A., Albertelli, G., 2008. Circumpolar distribution of the pycnogonid-ectoparasitic gastropod *Dickdella labioflecta* (Dell, 1990) (Mollusca: Zerotulidae). Antarctic Science 20: 497-498.
- Schwarz, A., Taylor, R., Hewitt, J., Phillips, N., Shima, J., Cole, R., Budd, R., 2006. Impacts of terrestrial runoff on the biodiversity of rocky reefs, New Zealand Aquatic Environment and Biodiversity Report No. 7, 109 p.
- Schwarz, A.-M., Hawes, I., Andrew, N., Mercer, S., Cummings, V., Thrush, S., 2005. Primary production potential of non-geniculate coralline algae at Cape Evans, Ross Sea, Antarctica. Marine Ecology Progress Series 294: 131-140.
- Schwarz, A.-M., Hawes, I., Andrew, N., Norkko, A., Cummings, V., Thrush, S., 2003. Macroalgal photosynthesis near the southern global limit for growth; Cape Evans, Ross Sea, Antarctica. Polar Biology 26: 789-799.
- Sewell, M.A., 2005. Examination of the meroplankton community in the southwestern Ross Sea, Antarctica, using a collapsible plankton net. Polar Biology. 28:119-131.
- Sewell, M.A., 2006. The meroplankton community of the northern Ross Sea: a preliminary comparison with the McMurdo Sound region. Antarctic Science. 18:595-602.
- Sewell, M., Lavery, S., Baker, C., 2006 Whose larva is that? Molecular identification of larvae of the Ross Sea, New Zealand Aquatic Environment and Biodiversity Report No. 3, 57 p.
- Sharp, B., Parker, S., Pinkerton, M., (lead authors), also Breen, B., Cummings, V., Dunn, A., Grant, S., Hanchet, S., Keys, H., Lockhart, S., Lyver, P.B., O'Driscoll, R., Williams, M., Wilson, P., 2010. Bioregionalisation and Spatial Ecosystem Processes in the Ross Sea Region, CCAMLR document WG-EMM-10/30, Hobart, Australia.
- Sharp, B., Parker, S., Smith, N., 2009. An Impact Assessment Framework for Bottom Fishing Methods in the CCAMLR Convention Area. CCAMLR Science 16: 195-210.
- Sharp, B., Waugh, S., Walker, N., 2011. A risk assessment framework for incidental seabird mortality associated with New Zealand fishing in the New Zealand EEZ., Unpublished report held by the Ministry of Fisheries, Wellington., 39 p.
- Slooten, E., Dawson, S., Rayment, W., Childerhouse, S., 2005. Distribution of Maui's dolphin, *Cephalorhynchus hectori maui*, New Zealand Fisheries Assessment Report. 2005/28, 21 p.
- Smith, A., Gordon, D., 2011. Bryozoans of southern New Zealand: a field identification guide, New Zealand Aquatic Environment and Biodiversity Report. No. 75, 65 p.
- Smith, F., 2006. Balleny Islands Ecology Research Voyage Report, R.V. Tiama, Unpublished report held at Ministry of Fisheries for project ZBD2005/01p.
- Smith, I., 2011. Estimating the magnitude of pre-European Maori marine harvest in two New Zealand study areas, New Zealand Aquatic Environment and Biodiversity Report, No. 82, 72 p.
- Smith, M., Baird, S., 2005a. Factors that may influence the level of incidental mortality of New Zealand sea lions (*Phocarctos hookeri*) in the squid (*Nototodar* spp.) trawl fishery in SQU 6T, New Zealand Fisheries Assessment Report 2005/20, 35 p.
- Smith, M., Baird, S., 2005b. Representativeness of past observer coverage, and future coverage required for estimation of New Zealand sea lion (*Phocarctos hookeri*) captures in the SQU 6T fishery., New Zealand Fisheries Assessment Report 39 p.
- Smith, M., Baird, S., 2007a. Estimation of incidental captures of New Zealand sea lions (*Phocarctos hookeri*) in New Zealand fisheries in 2003-04, with particular reference to the SQU 6T squid trawl fishery, New Zealand Fisheries Assessment Report 2007/7, 32 p.
- Smith, M., Baird, S., 2007b. Estimation of incidental captures of New Zealand sea lions (*Phocarctos hookeri*) in New Zealand fisheries in 2004-05, with particular reference to the SQU 6T squid trawl fishery., New Zealand Aquatic Environment and Biodiversity Report No. 12, 31 p.
- Smith, M., Baird, S., 2008a. Observer coverage required for the prediction of incidental capture of New Zealand fur seals in New Zealand commercial fisheries, New Zealand Aquatic Environment and Biodiversity Report. No. 26, 76 p.
- Smith, M., Baird, S., 2008b. Observer coverage required for the prediction of incidental capture of seabirds in New Zealand commercial fisheries, New Zealand Aquatic Environment and Biodiversity Report No. 25, 107 p.
- Smith, M., Baird, S., 2009. Model-based estimation of New Zealand fur seal (*Arctocephalus forsteri*) incidental captures and strike rates for trawl fishing in New Zealand waters for the years 1994-95 to 2005-06, New Zealand Aquatic Environment and Biodiversity Report No. 40, 90 p.
- Smith, M., Baird, S., 2011. Predicted incidental captures of New Zealand sea lions (*Phocarctos hookeri*) in the Auckland Islands SQU 6T squid trawl fishery for 1995 to 2006., New Zealand Aquatic Environment and Biodiversity Report. No. 71, 40 p.
- Smith, P., Notman, P., 2005. Inventory of biological samples collected from Antarctic toothfish *Dissostichus mawsoni* over summer 2005, Unpublished Final Research Report for Ministry of Fisheries Project ANT200401, Objective 5p.
- Smith, P., Roberts, C., Stewart, A., McVeagh, M., Struthers, C., 2007. Identification and speciation of Antarctic skates, Unpublished Final Research Report to the Ministry of Fisheries ANT2005/02, objective 5, 27 p.
- Smith, P., Steinke, D., McMillan, P., McVeagh, S., Struthers, C., 2008. DNA database for commercial marine fish, New Zealand Aquatic Environment and Biodiversity Report No. 22, 62 p.
- Smith, P., Steinke, D., McMillan, P., Stewart, A., McVeagh, S., Diaz De Astarloa, J., Welsford, D., Ward, R., 2011a. DNA barcoding highlights a cryptic species of grenadier (genus *Macrourus*) in the Southern Ocean Journal of Fish Biology 78 (1): 355-365.
- Smith, P., Steinke, D., McMillan, P., Stewart, A., Ward, R., 2011b. DNA barcoding of morids (Actinopterygii, Moridae) reveals deep divergence in the anti tropical *Halargyreus johnsoni* but little distinction between *Antimora rostrata* and *A. microlepis* Mitochondrial DNA (doi:10.3109/19401736.2010.532329).
- Snelder, T.H., Leathwick, J.R., Dey, K.L., Rowden, A.A., Weatherhead, M.A., Fenwick, G.D., Francis, M.P., Gorman, R.M., Grieve, J.M., Hadfield, M.G., Hewitt, J.E., Richardson, K.M., Uddstrom, M.J., Zeldis, J.R., 2005. The New Zealand Marine Environmental Classification, Unpublished Report for the Ministry for the Environment, 80 p.

- Snelder, T., Leathwick, J., Dey, K., Rowden, A., Weatherhead, M., 2006. Development of an ecological marine classification in the New Zealand region. *Environmental Management* 39: 12-29.
- Speed, S., 2005. Review of the effects of aquaculture and enhancement stock sources on wild fisheries resources and the marine environment- report on Objective 1 & 2, Unpublished Final Research Report for the Ministry of Fisheries, 128 p.
- Speed, S., McLary, D., Jones, E., 2006. The implications of marine reserves for fisheries resources and management in the New Zealand context, Unpublished Final Research Report for the Ministry of fisheries, 72 p.
- Stein, D., 2012. Snailfishes (Family Liparidae) of the Ross Sea, Antarctica, and closely adjacent waters. *Zootaxa* 3285 ( ISBN 978-1-86977-870-5 (Online edition)): 120.
- Stevens, D., 2006. Stomach contents of sub-adult Antarctic toothfish (*Dissostichus mawsoni*) from the western Ross Sea, Antarctica, Unpublished Final Research Report to Ministry of Fisheries, New Zealand for project ANT2004-01, Objective 4., xx p.
- Stevens, D., Dunn, M., 2010. Different food preferences in four sympatric deep-sea Macrourid fishes. *Marine Biology* DOI 10.1007/s00227-010-1542-1.
- Stevens, D., Hurst, R., Bagley, N., 2011. Feeding habits of New Zealand fishes: a literature review and summary of research trawl database records 1960 to 2000, New Zealand Aquatic Environment and Biodiversity Report, No. 85, 218 p.
- Stevens, D., Smith, M., Grimes, P., Devine, J., Sutton, C., MacGibbon, D., Maolagain, C., 2010. Age, growth, and maturity of four New Zealand rattail species, Aquatic Environment and Biodiversity Report. No. 59, 39 p.
- Strugnell, J., Watts, P., Smith, P., Allcock, A., Submitted. Persistent genetic signatures of historic climatic events in an Antarctic octopus.p.
- Sutherland, D., 2008. Surface-associated diatoms from marine habitats at Cape Evans, Antarctica, including the first record of living *Eunotogramma marginopunctatum*. *Polar Biology* DOI 10.1007/s00300-008-0426-z.
- Sutton, C., Manning, M., Stevens, D., Marriott, P., 2006. Biological parameters for icefish (*Chionobathyscus dewitti*) in the Ross Sea, Antarctica, Unpublished Final Research Report to the Ministry of Fisheries. Project ANT2005/02, objective 5, 27 p.
- Taylor, P., 1992. Incidental catch of non-fish species by setnets in New Zealand waters, New Zealand Fisheries Assessment Research Document 92/21, 23 p.
- Thatje, S., Lörz, A., 2005. First record of lithodid crabs from Antarctic waters off the Balleny Islands. *Polar Biology* 28 (4): 334-337.
- Thompson, F., Abraham, E., 2009a. Six monthly summary of the capture of protected species in New Zealand commercial fisheries, summer 2007-08, New Zealand Aquatic Environment and Biodiversity Report. No. 35, 22 p.
- Thompson, F., Abraham, E., 2009b. Dolphin bycatch in New Zealand trawl fisheries, 1995-96 to 2006-07, New Zealand Aquatic Environment and Biodiversity Report No. 36, 24 p.
- Thompson, F., Abraham, E., 2009c. Estimation of the capture of New Zealand sea lions (*Phocarctos hookeri*) in trawl fisheries from 1995-96 to 2006-07, New Zealand Aquatic Environment and Biodiversity Report No. 41, 31 p.
- Thompson, F., Abraham, E., 2010. Estimation of fur seal (*Arctocephalus forsteri*) bycatch in New Zealand trawl fisheries, 2002-03 to 2008-09, New Zealand Aquatic Environment and Biodiversity Report No. 61, 37 p.
- Thompson, F., Abraham, E., 2011. Estimation of the capture of New Zealand sea lions (*Phocarctos hookeri*) in trawl fisheries, from 1995-96 to 2008-09, New Zealand Aquatic Environment and Biodiversity Report No. 66, 25 p.
- Thompson, F., Abraham, E., Berkenbusch, K., 2010a. Common dolphin (*Delphinus delphis*) bycatch in New Zealand mackerel trawl fisheries, 1995-96 to 2008-09, New Zealand Aquatic Environment and Biodiversity Report No. 63, 20 p.
- Thompson, F., Abraham, E., Berkenbusch, K., 2011. Marine mammal bycatch in New Zealand trawl fisheries, 1995-96 to 2009-10, Draft Final Research Report for Ministry of Fisheries project PRO2010-01 (Unpublished report held by the Ministry of Fisheries, Wellington), 80 p.
- Thompson, F., Abraham, E., Oliver, MD, 2010b. Estimation of fur seal bycatch in New Zealand trawl fisheries, 2002-03 to 2007-08, New Zealand Aquatic Environment and Biodiversity Report. No. 56, 29 p.
- Thompson, F., Oliver, M., Abraham, E., 2010c. Estimation of the capture of New Zealand sea lions (*Phocarctos hookeri*) in trawl fisheries, from 1995-96 to 2007-08, New Zealand Aquatic Environment and Biodiversity Report No. 52, 25 p.
- Thrush, S., Cummings, V., 2011. Massive icebergs, alteration in primary food resources and change in benthic communities at Cape Evans, Antarctica. *Marine Ecology*.
- Thrush, S., Dayton, P., Cattaneo-Vietti, R., Chiantore, M., Cummings, V., Andrew, N., Hawes, I., Kim, S., Kvitek, R., Schwarz, A.-M., 2006. Broad-scale factors influencing the biodiversity of coastal benthic communities of the Ross Sea, Deep Sea Research II. 53, 959-971 p.
- Thrush, S., Hewitt, J., Cummings, V., Norkko, A., Chiantore, M., 2010.  $\beta$ -diversity and species accumulation in Antarctic coastal benthic communities: the role of habitat, distance and productivity on ecological connectivity. , *PLoS ONE* <http://dx.plos.org/10.1371/journal.pone.0011899p>.
- Thrush, S., Lohrer, D., Savage, C., In Press. Carbonate sediments: the positive and negative effects of land-coast interactions of functional diversity, Aquatic Environment and Biodiversity Report, xx p.
- Tracey, D., Anderson, O., Clark, M., Oliver, M., (Comps.), 2005. A guide to common deepsea invertebrates in New Zealand waters, New Zealand Aquatic Environment and Biodiversity Report No. 1, 160 p.
- Tracey, D., Anderson, O., Naylor, J.C., 2011a. A guide to common deepsea invertebrates in New Zealand waters, New Zealand Aquatic Environment and Biodiversity Report No. 86, 317 p.
- Tracey, D., Rowden, A., Mackay, K., Compton, T., 2011b. Habitat-forming cold-water corals show affinity for seamounts in the New Zealand region. . *Marine Ecology Progress Series* 430: 1-22.
- Tracey, D., Anderson, O., Naylor, J.R., (Comps.), 2007. A guide to common deepsea invertebrates in New Zealand waters, New Zealand Aquatic Environment and Biodiversity Report No.10., 282 p.
- Tracey, D., Carter, M., Parker, S., 2010. Evaluation of VME taxa monitoring by scientific observers, Unpublished Final Research Report for Ministry of Fisheries Research Project ANT2009/01 Objective 8, 17 p.
- Tracey, D., Parker, S., Mackay, E., Anderson, O., Ramm, K., 2008. Classification guide for potentially vulnerable invertebrate taxa in the SPRFMO Area, Identification guide for Ministry of fisheries 1p.
- Tuck, I., Cole, R., Devine, J., 2009. Ecosystem indicators for New Zealand fisheries, New Zealand Aquatic Environment and Biodiversity Report No. 42, 188 p.
- Tuck, I., Drury, J., Kelly, M., Gerring, P., 2010. Designing a programme to monitor the recovery of the benthic community between North Cape and Cape Reinga, New Zealand Aquatic Environment and Biodiversity Report No. 53, 78 p.
- Tuck, I., Hewitt, J., In Press. Monitoring change in benthic communities in Spirits bay, Aquatic Environment and Biodiversity Report, xx p.

## *AEBAR 2012: Appendices*

- Tuck, I., Parkinson, D., Dey, K., Oldman, K., Sadhwa, W., 2006. Information on benthic impacts in support of the coromandel Scallops fishery plan, Unpublished Ministry of Fisheries Final Research Report 64 p.
- Tuck, I., Parkinson, K., Dey, J., Oldman, S., Wadwha, 2006. Information on benthic impacts in support of the Coromandel scallop fisheries plan. info for SCA-CS fish plan, Unpublished Final Research Report for Ministry of Fisheries Research Project ZBD2005-15p.
- Varian, S., 2005. A summary of the values of the Balleny Islands, Antarctica, Marine Biodiversity Biosecurity Report No. 6, 13 p.
- Waugh, S., Filippi, D., Walker, N., Kirby, D., 2008. Updated preliminary results of an ecological risk assessment for seabirds and marine mammals with risk of fisheries interactions Western and Central pacific Fisheries Comission (WCPFC) Working Group - Fisheries Stock Assessment. 08/51p.
- Waugh, S., Filippi, D., Abraham, E., 2009. Ecological Risk Assessment for Seabirds in New Zealand fisheries, Unpublished Final Research Report for the Ministry of Fisheries 58 p.
- Williams, A., Schlacher, T., Rowden, A., Althaus, F., Clark, M., Bowden, D., Stewart, R., Bax, N., Consalvey, M., Kloser, R., 2010. Seamount megabenthic assemblages fail to recover from trawling impacts. *Marine Ecology* 31(suppl. 1): 183-199.
- Wing, S., 2005. Fiordland Biodiversity Research Cruise, Unpublished Final Research Report for ZBD2003-04, xx p.
- Zemke-White, W.L., Speed, S.R., McClary, D.J., 2005. Beach-cast seaweed: a review, New Zealand Fisheries Assessment Report 2005/44, 47 p.

ISBN: 978-0-478-40503-3 (print)  
ISBN: 978-0-478-40504-0 (online)

New Zealand Government

Ministry for Primary Industries  
Manatū Ahu Matua  
Pastoral House  
25 The Terrace  
PO Box 2526, Wellington, 6140  
New Zealand  
[www.mpi.govt.nz](http://www.mpi.govt.nz)