



Review and summary of the time series of input data
available for the assessment of southern blue whiting
(*Micromesistius australis*) stocks up to and including the
2013 season

New Zealand Fisheries Assessment Report 2015/56

A. Dunn,
S.M. Hanchet

ISSN 1179-5352 (online)
ISBN 978-1-77665-060-6 (online)

September 2015



Requests for further copies should be directed to:

Publications Logistics Officer
Ministry for Primary Industries
PO Box 2526
WELLINGTON 6140

Email: brand@mpi.govt.nz
Telephone: 0800 00 83 33
Facsimile: 04-894 0300

This publication is also available on the Ministry for Primary Industries websites at:
<http://www.mpi.govt.nz/news-resources/publications.aspx>
<http://fs.fish.govt.nz> go to Document library/Research reports

© Crown Copyright - Ministry for Primary Industries

Table of Contents

| | |
|---|-----------|
| Executive Summary | 1 |
| 1. Introduction | 2 |
| 2. Fishery summary | 4 |
| 2.1 Commercial fisheries..... | 4 |
| 2.2 Illegal catches | 7 |
| 2.3 Other sources of fishing mortality..... | 7 |
| 2.4 The 2013 season..... | 8 |
| 3. Biology | 12 |
| 3.1 Stock structure | 12 |
| 3.2 Biological parameters..... | 12 |
| 3.2.1 Age and growth | 12 |
| 3.2.2 <i>Spawning and length and age at maturity</i> | 15 |
| 3.2.3 Natural mortality | 16 |
| 4. Research surveys and other estimates of abundance | 16 |
| 4.1 Acoustic research surveys | 16 |
| 4.1.1 Auckland Islands | 17 |
| 4.1.2 Bounty Platform | 17 |
| 4.1.3 Campbell Island Rise..... | 18 |
| 4.1.4 Pukaki Rise | 19 |
| 4.2 Trawl research surveys | 20 |
| 4.2.1 Auckland Island Shelf | 21 |
| 4.2.2 Campbell Island Rise..... | 22 |
| 4.2.3 Pukaki Rise | 23 |
| 4.3 CPUE analyses..... | 24 |
| 4.3.1 Bounty Platform | 25 |
| 4.3.2 Campbell Island Rise..... | 26 |
| 5. Length and age composition of the fishery | 27 |
| 5.1 Methods | 27 |
| 5.2 Auckland Islands..... | 28 |
| 5.3 Bounty Platform..... | 30 |
| 5.4 Campbell Island Rise | 33 |
| 5.5 Pukaki Rise..... | 37 |
| 5.6 Other areas (SBW1) | 39 |
| 6. Discussion | 40 |
| 7. Acknowledgments | 41 |
| 8. References | 42 |

EXECUTIVE SUMMARY

Dunn, A.; Hanchet, S.M. (2015). Review and summary of the time series of input data available for the assessment of southern blue whiting (*Micromesistius australis*) stocks up to and including the 2013 season.

New Zealand Fisheries Assessment Report 2015/56. 45 p.

This document updates and summarises the observational and research data carried out on southern blue whiting up to and including the 2013 season into a single document. We have included the time series of relative abundance from acoustic surveys for each of the four main stocks (both from the wide area R.V. *Tangaroa* surveys and the local aggregation surveys), CPUE indices for Bounty Platform and Campbell Island Rise, and trawl survey indices for the Auckland Islands Shelf, Campbell Island Rise, and Pukaki Rise, as well as updated time series of length-at-age and catch-at-age.

The main source of information on southern blue whiting stock size remains the acoustic indices from wide area surveys on the Campbell Island Rise and local aggregations surveys on the Bounty Platform. Aggregations surveys for southern blue whiting on the Pukaki Rise have so far been largely unsuccessful with high variability in snapshots between snapshots and years. Estimates of southern blue whiting abundance from the sub-Antarctic trawl surveys on the Auckland Islands Shelf, Campbell Island Rise, and Pukaki Rise were available, but do not appear to be a useful source of information for monitoring these fisheries.

The catch on the Bounty Platform in recent years has been dominated by the strong 2002 year class, with no further recruitment except in 2011 when the 2007 year class entered the fishery. On the Campbell Island Rise, there was strong evidence of several year classes of moderate strength, with both acoustic indices and commercial-catch-at-age proportions suggesting strong recruitment in 2006 and 2009. The catch on the Pukaki Rise has been dominated recently by fish comprising the 2004–2006 year classes but there has been recent recruitment of the 2009 year class, which may prove to be moderately strong. Very few length measurements or otoliths were collected from the Auckland Islands in recent years, and the length data were not sufficient to infer recruitment patterns in this fishery.

1. INTRODUCTION

Southern blue whiting are almost entirely restricted in distribution to sub-Antarctic waters. They are dispersed throughout the Campbell Plateau and Bounty Platform for much of the year, but during August and September they aggregate to spawn near the Campbell Islands, on Pukaki Rise, on Bounty Platform, and near Auckland Islands over depths of 250–600 m (Figure 1). During most years fish in the spawning fishery range between 35 and 50 cm fork length (FL), although occasionally smaller size classes of males (29–32 cm FL) are observed in the catch.

Commercial fishing has concentrated on the Campbell Island Rise and, to a lesser extent, the Bounty Platform. The Pukaki Rise and Auckland Islands have been important fisheries in the past, but have recently had much lower annual catches than the Campbell Island Rise and Bounty Platform fisheries.

Stock assessments for southern blue whiting on the Campbell Island Rise and the Bounty Platform have been conducted at approximately biennial intervals using age-structured stock assessment models. Model inputs have included time series of acoustic survey indices, commercial catch-at-age composition data, and in earlier years, CPUE indices.




Figure 1: Relative total density of the commercial catch of southern blue whiting by location, TCEPR data 1990–2013.

A large amount of research has been carried out on southern blue whiting, including work on stock structure (Hanchet 1998, Hanchet 1999), age and growth (Hanchet & Uozumi 1996), catch-at-age (e.g., Hanchet et al. 2003, Hanchet 2005), acoustic surveys (e.g., O'Driscoll et al. 2009), CPUE analyses (e.g., Hanchet & Blackwell 2003, Hanchet et al. 2005), trawl surveys (Hanchet & Stevenson 2006), and stock

assessments (e.g., Dunn & Hanchet 2011a, Dunn & Hanchet 2011b). The objective of this report is to summarise and document the time series of input data which could potentially be used for stock assessment. Further, we revise and update the commercial catch-at-age proportions for the Bounty Platform, Campbell Island Rise, and Pukaki Rise. This report updates the summary provided by Cole et al. (2013) and Dunn & Hanchet (2014).

The data described in this report were based on data extracts from the Ministry for Primary Industries Catch-Effort database in December 2013 (REPLOG 9257), the Ministry for Primary Industries Observer database in December 2013, and the Ministry for Primary Industries Trawl Survey database in November 2013.

This report is in partial fulfilment of the Ministry for Primary Industries Project DEE201002SBWC: To carry out stock assessments of southern blue whiting (*Micromesistius australis*), including estimating biomass and sustainable yields.

2. FISHERY SUMMARY

2.1 Commercial fisheries

The southern blue whiting fishery was developed by Soviet vessels during the early 1970s, with early reported landings peaking at almost 50 000 t in 1973 (Table 1). Early reports recorded that southern blue whiting spawned in most years on the Bounty Platform (Shpak 1978) and in some years on the Campbell Plateau (Shpak & Kuchina 1983), and that feeding aggregations could be caught on the Pukaki Rise, southeast of the Campbell Island Rise, and on the Auckland Islands Shelf (Shpak 1978). Some fishing probably took place on each of the grounds, but the proportion of catch from each ground was not accurately recorded before 1978. Hence the amount of catch for each ground cannot accurately be determined before 1978.

Landings were chiefly taken by the Soviet foreign licensed fleet during the 1970s and early 1980s. The entire Campbell Plateau (Campbell Island Rise and Pukaki Rise) was fished year-round between 1978 and 1984, but highest catches were usually made during spawning, typically during September. In some seasons (notably 1979, 1982, and 1983) vessels also targeted spawning fish on the Bounty Platform in August and September (Table 1).

As a result of the increase in hoki quota in 1985 and 1986, the Japanese surimi fleet increased its presence in New Zealand waters and some vessels stayed on after the hoki fishery to fish for southern blue whiting. After that, many of the Japanese and Soviet (replaced latterly by Ukrainian) vessels which fish for hoki on the west coast of the South Island during July and August each year move in late August or early September to the southern blue whiting spawning grounds. Between 1986 and 1989, fishing was confined to the spawning grounds on the northern Campbell Island Rise. From 1990 onwards, vessels also started fishing spawning aggregations on the Bounty Platform, the Pukaki Rise, and the southern Campbell Island Rise. Fishing effort increased markedly between 1990 and 1992, culminating in a catch of over 75 000 t in 1992 (Table 1). The increased catch came mainly from the Bounty Platform. In 1993, a fishery developed for the first time on the Auckland Islands spawning grounds but fishing has continued there at a low level since then. Total annual landings over the past five years have ranged between 25 000 t and 40 000 t, most of which has been taken from the Campbell Island Rise grounds. However, a strong year class on the Bounty Platform led to a rapid increase in catches there from 2008 to 2010. Relatively large catches also occurred at the Pukaki Rise in 2009, 2010, and 2012.

On the Bounty Platform and Campbell Island Rise the TACC has usually been almost fully caught in recent years, except that in 2012, the catch on Campbell Island Rise was only two thirds of the TACC. The under-catch on the Campbell Island Rise was due to an oversupply in the market for southern blue whiting, resulting in much reduced effort and hence a reduced catch.

The total catch also approached the level of the TACC at Pukaki Rise in 2009 and 2010, but since then the TACC has been considerably under-caught. At Auckland Islands, the catch limits have generally been under-caught in most years since their introduction. On the Bounty Platform, the amount of fishing effort in any season has depended largely on the timing of the west coast hoki fishery. If there is a delayed hoki season, then the vessels remain longer on the hoki grounds and consequently may miss the peak fishing season on the Bounty Platform. On the Pukaki Rise and Auckland Islands Shelf, operators find it difficult to justify expending time to locate fishable aggregations, given the small allocation available in these areas and the relatively low value of the product.

In general, the fleet has gradually changed from one being dominated by Japanese surimi and Soviet 'Head and Gut' vessels to one dominated by vessels from Ukraine and the Dominican Republic which process fish to a dressed product. In 2013, most catch was taken by vessels from the Ukraine (46%) with the remainder from New Zealand (11%), Japan (15%), and the Dominican Republic (26%). Before 2003, about 70% of the product was surimi and about 30% was dressed, but since 2005 the proportion surimi has declined to 40% in 2006 and 12–16% since 2009 (Figure 2).

Catch quotas, allocated to individual operators, were introduced for the first time in the 1992–93 fishing year. The catch limit of 32 000 t, with stock-specific sub-limits, was retained for the next three years (Table 1). The stock-specific sub-limits were revised for the 1995–96 fishing year, and the total catch limit increased to 58 000 t in 1996–97 for three years (Table 1). In 1997–98, a separate catch limit of 1640 t was set for the Auckland Islands fishery for the first time (Table 1).

The southern stocks of southern blue whiting were introduced to the Quota Management System on 1 November 1999 with the following TACCs: Auckland Islands (SBW 6A) 1640 t, Bounty Platform (SBW 6B) 15 400 t, Campbell Island Rise (SBW 6I) 35 460 t, and Pukaki Rise (SBW 6R) 5500 t (Table 1). At the same time, the fishing year was changed to 1 April to 31 March to reflect the timing of the main fishing season. Southern blue whiting has been managed using a Current Annual Yield (CAY) strategy (Annala et al. 2004), which has contributed to the fluctuating catch limits and TACCs (Table 1). A nominal TACC of 8 t (SBW 1) was set for the rest of the EEZ. Less than 20 t per year has been reported from SBW 1 since 2000–01.

Details of recent stock-specific changes over time are given in Table 1. Once in the QMS the TACC for the Bounty Platform was gradually reduced to 3500 t by 2003, reflecting a period of poor recruitment to the stock. The TACC remained at that level until 2008 when the strong 2002 year class entered the fishery, and the TACC was increased to 9800 t and then 14 700 t. From 1 April 2011, the TACC for the Bounty Platform stock was reduced 6860 t, and for the 2014 season the industry shelved 2832 t under a voluntary agreement resulting in a catch limit of 4028 t. Once in the QMS the TACC for the Campbell Island Rise was gradually reduced to 20 000 t by 2006, reflecting a period of poor to average recruitment to the stock. The TACC remained at that level until 2009 when the strong 2006 year class entered the fishery, and the TACC was increased to 23 000 t in 2010 and then to 29 400 t for 2011, 2012, and 2013. Catch limits for Pukaki Rise and Auckland Islands have remained unchanged since 1997–98.

Table 1: Estimated catches and catch limits (TACCs) (t) of southern blue whiting for 1971 to 2012–13, and by area for 1978 to 2012–13 (source: QMRs and MHRs; ‘–’ denotes no catch limit in place).

| Fishing year ¹ | Bounty Platform (SBW 6B) | | Campbell Island Rise (SBW 6I) | | Pukaki Rise (SBW 6R) | | Auckland Island (SBW 6A) | | Total (All areas) | |
|---------------------------|--------------------------|--------------------|-------------------------------|--------|----------------------|-------|--------------------------|--------------------|--------------------|--------|
| | Catch | Limit ⁴ | Catch | Limit | Catch | Limit | Catch | Limit ² | Catch ³ | Limit |
| 1971 | – | – | – | – | – | – | – | – | 10 400 | – |
| 1972 | – | – | – | – | – | – | – | – | 25 800 | – |
| 1973 | – | – | – | – | – | – | – | – | 48 500 | – |
| 1974 | – | – | – | – | – | – | – | – | 42 200 | – |
| 1975 | – | – | – | – | – | – | – | – | 2 378 | – |
| 1976 | – | – | – | – | – | – | – | – | 17 089 | – |
| 1977 | – | – | – | – | – | – | – | – | 26 435 | – |
| 1978 | 0 | – | 6 403 | – | 79 | – | 15 | – | 6 497 | – |
| 1978–79 | 1 211 | – | 25 305 | – | 601 | – | 1 019 | – | 28 136 | – |
| 1979–80 | 16 | – | 12 828 | – | 5 602 | – | 187 | – | 18 633 | – |
| 1980–81 | 8 | – | 5 989 | – | 2 380 | – | 89 | – | 8 466 | – |
| 1981–82 | 8 325 | – | 7 915 | – | 1 250 | – | 105 | – | 17 595 | – |
| 1982–83 | 3 864 | – | 12 803 | – | 7 388 | – | 184 | – | 24 239 | – |
| 1983–84 | 348 | – | 10 777 | – | 2 150 | – | 99 | – | 13 374 | – |
| 1984–85 | 0 | – | 7 490 | – | 1 724 | – | 121 | – | 9 335 | – |
| 1985–86 | 0 | – | 15 252 | – | 552 | – | 15 | – | 15 819 | – |
| 1986–87 | 0 | – | 12 804 | – | 845 | – | 61 | – | 13 710 | – |
| 1987–88 | 18 | – | 17 422 | – | 157 | – | 4 | – | 17 601 | – |
| 1988–89 | 8 | – | 26 611 | – | 1 219 | – | 1 | – | 27 839 | – |
| 1989–90 | 4 430 | – | 16 542 | – | 1 393 | – | 2 | – | 22 367 | – |
| 1990–91 | 10 897 | – | 21 314 | – | 4 652 | – | 7 | – | 36 870 | – |
| 1991–92 | 58 928 | – | 14 208 | – | 3 046 | – | 73 | – | 76 255 | – |
| 1992–93 | 11 908 | 15 000 | 9 316 | 11 000 | 5 341 | 6 000 | 1 143 | – | 27 708 | 32 000 |
| 1993–94 | 3 877 | 15 000 | 11 668 | 11 000 | 2 306 | 6 000 | 709 | – | 18 560 | 32 000 |
| 1994–95 | 6 386 | 15 000 | 9 492 | 11 000 | 1 158 | 6 000 | 441 | – | 17 477 | 32 000 |
| 1995–96 | 6 508 | 8 000 | 14 959 | 21 000 | 772 | 3 000 | 40 | – | 22 279 | 32 000 |
| 1996–97 | 1 761 | 20 200 | 15 685 | 30 100 | 1 806 | 7 700 | 895 | – | 20 147 | 58 000 |
| 1997–98 | 5 647 | 15 400 | 24 273 | 35 460 | 1 245 | 5 500 | 0 | 1 640 | 31 165 | 58 000 |
| 1998–00 | 8 741 | 15 400 | 30 386 | 35 460 | 1 049 | 5 500 | 750 | 1 640 | 40 926 | 58 000 |
| 2000–01 | 3 997 | 8 000 | 18 049 | 20 000 | 2 864 | 5 500 | 19 | 1 640 | 24 804 | 35 148 |
| 2001–02 | 2 262 | 8 000 | 29 999 | 30 000 | 230 | 5 500 | 10 | 1 640 | 31 114 | 45 148 |
| 2002–03 | 7 565 | 8 000 | 33 445 | 30 000 | 508 | 5 500 | 262 | 1 640 | 41 795 | 45 148 |
| 2003–04 | 3 812 | 3 500 | 23 718 | 25 000 | 163 | 5 500 | 116 | 1 640 | 27 812 | 35 648 |
| 2004–05 | 1 477 | 3 500 | 19 799 | 25 000 | 240 | 5 500 | 95 | 1 640 | 21 620 | 35 648 |
| 2005–06 | 3 962 | 3 500 | 26 190 | 25 000 | 58 | 5 500 | 66 | 1 640 | 30 278 | 35 648 |
| 2006–07 | 4 395 | 3 500 | 19 763 | 20 000 | 1 115 | 5 500 | 84 | 1 640 | 25 363 | 30 648 |
| 2007–08 | 3 799 | 3 500 | 20 996 | 20 000 | 513 | 5 500 | 278 | 1 640 | 25 587 | 30 648 |
| 2008–09 | 9 863 | 9 800 | 20 483 | 20 000 | 1 377 | 5 500 | 143 | 1 640 | 31 887 | 36 948 |
| 2009–10 | 15 468 | 14 700 | 19 040 | 20 000 | 4 853 | 5 500 | 174 | 1 640 | 39 540 | 41 848 |
| 2010–11 | 13 913 | 14 700 | 20 224 | 23 000 | 4 433 | 5 500 | 131 | 1 640 | 38 708 | 44 848 |
| 2011–12 | 6 660 | 6 860 | 30 982 | 29 400 | 701 | 5 500 | 92 | 1 640 | 38 440 | 43 400 |
| 2012–13 | 6 827 | 6 860 | 21 321 | 29 400 | 1 702 | 5 500 | 49 | 1 640 | 29 906 | 43 400 |
| 2013–14 | 4 278 | 4 028 | 28 607 | 29 400 | 14 | 5 500 | 47 | 1 640 | 32 950 | 43 400 |

1. Fishing years defined as 1 April to 30 September for 1978; 1 October to 30 September for 1978–79 to 1997–98; 1 October 1998 to 31 March 2000 for 1998–2000; 1 April to 31 March for 2000–01 to current.
2. Before 1997–98, there were no separate catch limits for Auckland Islands.
3. Totals include SBW 1 (i.e., all EEZ areas outside QMA 6). SBW 1 has a TACC of 8 t and reported annual catches since 2000–01 have ranged from 1 t in 2007–08 to 21 t in 2008–09.
4. In 2013, although the TACC remained at 6860 t, the catch limit was limited to 4028 t as 2832 t was shelved under a voluntary agreement.




Figure 2: Estimated total processed catch for southern blue whiting by processed state, for all areas, between July and October, 1990–2013.

2.2 Illegal catches

The level of illegal and unreported catch for southern blue whiting has been reported as “thought to be low” (Ministry for Primary Industries 2013). However, there have been several instances of area misreporting and illegal discards.

In 2002–03, the operators of one vessel were convicted for area misreporting — the vessel had caught about 204 t of southern blue whiting on the Campbell Island Rise (SBW6I) and reported this against quota for the Pukaki Rise (SBW 6R); another 480 t caught on the Campbell Island Rise had been reported against quota for the Auckland Islands Shelf (SBW 6A). In addition, in 2004, the operators of a vessel were convicted of dumping southern blue whiting at sea — crew members estimated that between 40 and 310 tonnes were illegally discarded during a two and a half week period of fishing on the Campbell Island Rise (Ministry for Primary Industries 2013).

In addition, some catch was alleged to have been misreported between SBW 6R, SBW 6B, and SBW 6I. The then Ministry of Fisheries (now Ministry for Primary Industries (MPI), hereafter referred to as the Ministry) noted that in August and September 2002, there was some evidence that a vessel caught 81 t of southern blue whiting in SBW 6R and misreported it as catch from SBW 6B; and also misreported 108 t from SBW 6I as being from SBW 6R. The Ministry noted that in 2004, there was some evidence that 64.5 t of southern blue whiting was caught in SBW 6I and misreported as being caught in SBW 6B (G. Backhouse, Senior Fisheries Investigator, MPI, pers. comm.).

2.3 Other sources of fishing mortality

Scientific observers reported discards of undersize fish and accidental loss from torn or burst codends, particularly during the early years of the fishery. Discarding in the southern blue whiting fishery has been estimated by Clark et al. (2000), and Anderson (2004, 2009). Anderson (2004) quantified total annual discard estimates (including estimates of fish lost from the net at the surface) as ranging between 0.4% and 2.0% of the estimated catch for all southern blue whiting fisheries. Anderson (2009) reviewed fish and invertebrate bycatch, and discards in the southern blue whiting fishery based on observer data

from 2002 to 2007. He estimated that 0.23% of the catch was discarded from observed vessels. The low levels of discarding occurred primarily because the fishery targeted spawning aggregations.

In August 2010, the fishing vessel *Oyang 70* sank while fishing for southern blue whiting on the Bounty Platform. It was fishing an area between 48° 00' S and 48° 20' S, and 179° 20' E and 180° 00' E between 15 and 17 August 2010, before sinking on 18 August 2010. The Ministry of Fisheries estimated that it had taken a catch of between 120 t and 190 t that was lost with the vessel (G. Backhouse, Senior Fisheries Investigator, MPI, pers. comm.).

2.4 The 2013 season

The location of trawls made during the 2013 season (mid-August to mid-October) is shown in Figure 3. Most of the catch was taken by vessels flagged to Ukraine, Dominican Republic, New Zealand, and Japan and most fishing was carried out on the Bounty and Campbell stocks (Table 2). In 2013, the first vessels arrived on the Bounty Platform on 11 August and gradually worked their way south and east making small to moderate catches. From 15 to 20 August vessels were fishing in the south west and made high catches (typically at least 400 t per day) (Figure 4). Fish were spawning from 22–26 August (Figure 5), at which time the vessels left the area.

In 2013, vessels started fishing the Campbell Island Rise on 15 August and continued fishing until 1 October (Figure 4). As in recent years, the fleet fished the northern spawning aggregations on the Campbell Island Rise ground, but unlike previous years, only lightly fished the east and south. Daily proportions spawning at Campbell Island showed two distinct spawning peaks with the first spawning peak from 1–7 September and the second from 20–25 September (Figure 5).

In 2013, four vessels fished the Pukaki Rise between 17 August and 29 September (Figure 4) but only recorded a total of four target tows (Table 2) and almost no catch was taken (14 t, see Table 1). The timing of spawning could not be determined as coverage was so low.

Six vessels reported catches of southern blue whiting on the Auckland Islands Shelf in 2013, but no targeted fishing was carried out (Table 2) and no fishing was carried out during the spawning season. Daily catches were small and southern blue whiting were taken only as bycatch of fishing for other species.

Table 2: Number of tows and vessels for vessels targeting southern blue whiting by area, 1990–2013 (source: TCEPR data).

| Year | Auckland Islands | | Bounty Platform | | Campbell Island | | Pukaki Rise | | Other | |
|------|------------------|---------|-----------------|---------|-----------------|---------|-------------|---------|-------|---------|
| | Tows | Vessels | Tows | Vessels | Tows | Vessels | Tows | Vessels | Tows | Vessels |
| 1990 | 3 | 1 | 263 | 25 | 1 030 | 35 | 191 | 32 | 3 | 8 |
| 1991 | 1 | 1 | 661 | 31 | 1 228 | 33 | 262 | 24 | 3 | 8 |
| 1992 | 7 | 2 | 1 732 | 49 | 1 530 | 47 | 374 | 40 | 13 | 27 |
| 1993 | 20 | 4 | 433 | 21 | 423 | 20 | 393 | 23 | 6 | 12 |
| 1994 | 43 | 7 | 178 | 9 | 480 | 15 | 81 | 11 | 4 | 4 |
| 1995 | 15 | 5 | 155 | 10 | 285 | 12 | 71 | 9 | 6 | 12 |
| 1996 | 6 | 3 | 67 | 5 | 474 | 11 | 10 | 4 | 1 | 1 |
| 1997 | 18 | 5 | 37 | 5 | 650 | 18 | 46 | 8 | 1 | 2 |
| 1998 | 14 | 5 | 118 | 12 | 959 | 24 | 34 | 11 | 3 | 8 |
| 1999 | 14 | 3 | 288 | 14 | 790 | 21 | 26 | 7 | 2 | 2 |
| 2000 | 1 | 1 | 99 | 6 | 447 | 16 | 57 | 8 | – | – |
| 2001 | 2 | 2 | 32 | 5 | 650 | 14 | 12 | 7 | 1 | 1 |
| 2002 | 6 | 2 | 94 | 6 | 862 | 18 | 15 | 5 | – | – |
| 2003 | – | – | 24 | 3 | 599 | 15 | 4 | 3 | – | – |
| 2004 | 1 | 1 | 32 | 3 | 690 | 16 | 3 | 3 | – | – |
| 2005 | 1 | 1 | 100 | 5 | 755 | 17 | 3 | 2 | – | – |
| 2006 | – | – | 94 | 5 | 521 | 13 | 19 | 1 | – | – |
| 2007 | – | – | 51 | 4 | 544 | 13 | 20 | 3 | – | – |
| 2008 | – | – | 200 | 8 | 557 | 14 | 57 | 4 | – | – |
| 2009 | 12 | 2 | 401 | 13 | 627 | 14 | 158 | 9 | 2 | 3 |
| 2010 | – | – | 394 | 13 | 550 | 12 | 170 | 10 | – | – |
| 2011 | – | – | 175 | 8 | 976 | 14 | 72 | 8 | – | – |
| 2012 | – | – | 173 | 8 | 592 | 11 | 128 | 9 | – | – |
| 2013 | – | – | 77 | 5 | 693 | 10 | 4 | 4 | – | – |




Figure 3: Commercial trawls made during the 2013 season targeting southern blue whiting (late August to early October, black points) and the location of historical target tows 1990–2012 (grey points).




Figure 4: Daily target southern blue whiting catch on the Bounty Platform (blue), Pukaki Rise (black), and Campbell Island Rise (red) in 2013 between July and October.




Figure 5: Daily proportion of females spawning (proportions at stage 4) for target southern blue whiting tows on the Bounty Platform (blue), Pukaki Rise (black), and Campbell Island Rise (red) in 2013 between August and October.

3. BIOLOGY

3.1 Stock structure

Stock structure of southern blue whiting was reviewed by Hanchet (1998, 1999) who examined data on distribution and abundance, reproduction, growth, and morphometrics. There appear to be four main spawning grounds: Bounty Platform, Pukaki Rise, Auckland Islands Shelf, and Campbell Island Rise. There are also consistent differences in the size and age distributions of fish, in the recruitment strength, and in the timing of spawning between these four areas. Multiple discriminant analysis of data collected in October 1989 and 1990 showed that fish from Bounty Platform, Pukaki Rise, and Campbell Island Rise could be distinguished on the basis of their morphometric measurements. This constitutes strong evidence that fish in these areas return to spawn on the grounds where they first recruited. There have been no genetic studies, but given the close proximity of the areas, it is unlikely that there would be detectable genetic differences in the fish among these four areas.

For stock assessment, it is assumed that there are four stocks of southern blue whiting with fidelity within stocks: the Bounty Platform stock, the Pukaki Rise stock, the Auckland Islands stock, and the Campbell Island Rise stock. Southern blue whiting are also managed as four separate stocks.

3.2 Biological parameters

3.2.1 Age and growth

Early growth has been well documented with fish reaching a length of about 20 cm FL after one year and 30 cm FL after two years Hanchet & Uozumi (1996). Growth slows down after five years and virtually ceases after ten years. Ages have been validated up to at least 15 years by following strong year classes, but ring counts from otoliths suggest individual fish may reach 25 years (Hanchet & Uozumi 1996).

An important feature of the biology of southern blue whiting is very high recruitment variability and associated density dependent growth (Hanchet et al. 2003). For example, the very strong 1991 year class on the Campbell Island Rise grew at a much slower rate (smaller length and weight at age) than previous year classes (see Figure 6 and Table 3). A similar large reduction in growth rate occurred on the Bounty Platform with the strong 2002 year class (Figure 7 and Table 4), with the subsequent two year classes also growing at a similar slower rate. For this reason, mean length at age is input into the assessment models as a year specific matrix of lengths at age rather than a vector of length at age based on the von Bertalanffy growth parameters.

Mean length at age estimates for each area (based directly on the annual age-length key) were presented by Hanchet et al. (2003). These estimates have been updated and recalculated using catch-at-age software (Bull & Dunn 2002). In this approach the raw age-length key data are scaled up so that the mean length at age for the plus group is based on the scaled length frequency distribution of fish in the plus group. The results are presented in Figure 6 and Figure 7 for the Campbell Island Rise and the Bounty Platform respectively.

Table 3: Overall mean weight at age (g) and the stronger year classes (1991, 2007, and 2009) mean weight at age (g) for southern blue whiting on the Campbell Island Rise.

| Sex | Year class | Age (y) | | | | | | | | |
|--------|------------|---------|-----|-----|-----|-----|-----|-----|-----|-----|
| | | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| Male | 1991 | 147 | 207 | 274 | 334 | 404 | 434 | 482 | 516 | 556 |
| | 2007 | 141 | 193 | 244 | 329 | 418 | – | – | – | – |
| | 2009 | 133 | 193 | 275 | – | – | – | – | – | – |
| | Overall | 175 | 262 | 369 | 437 | 511 | 560 | 609 | 633 | 674 |
| Female | 1991 | 144 | 243 | 317 | 395 | 503 | 540 | 599 | 644 | 683 |
| | 2007 | 140 | 218 | 282 | 382 | 482 | – | – | – | – |
| | 2009 | 127 | 210 | 312 | – | – | – | – | – | – |
| | Overall | 166 | 307 | 413 | 503 | 595 | 672 | 745 | 776 | 821 |

Table 4: Overall mean weight at age (g) and the stronger year classes (2002 and 2007) mean weight at age (g) for southern blue whiting on the Bounty Platform.

| Sex | Year class | Age (y) | | | | | | | | |
|--------|------------|---------|-----|-----|-----|-----|-----|-----|-----|-----|
| | | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| Male | 2002 | – | 173 | 238 | 293 | 344 | 361 | 396 | 413 | 424 |
| | 2007 | 127 | 190 | 277 | 339 | 365 | – | – | – | – |
| | Overall | 126 | 208 | 299 | 373 | 433 | 533 | 583 | 628 | 671 |
| Female | 2002 | – | 185 | 260 | 337 | 400 | 424 | 475 | 517 | 545 |
| | 2007 | 132 | 270 | 301 | 371 | 418 | – | – | – | – |
| | Overall | 128 | 218 | 328 | 463 | 516 | 639 | 749 | 831 | 881 |




Figure 6: Observed mean length-at-age (ages 2–8) for the Campbell Island stock by sex and year class, 1978–2011.




Figure 7: Observed mean length-at-age (ages 2–8) for the Bounty Platform stock by sex and year class, 1990–2011.

3.2.2 Spawning and length and age at maturity

Southern blue whiting are highly synchronised batch spawners. Four spawning areas have been identified, on Bounty Platform, Pukaki Rise, Auckland Islands Shelf, and Campbell Island Rise. The Campbell Island Rise has two separate spawning grounds to the north and south. Fish appear to recruit first to the southern ground but thereafter spawn on the northern ground (Hanchet 1998). Spawning on

Bounty Platform begins in mid-August and finishes by mid-September. Spawning begins 3–4 weeks later in the other areas, finishing in late September/early October. Spawning appears to mainly occur at night, in mid-water, over depths of 400–500 m on Campbell Island Rise but shallower elsewhere.

The age and length of maturity, and recruitment to the fishery, varies between areas and between years. In some years a small proportion of males mature at age 2, but the majority do not mature until age 3 or 4, usually at a length of 33–40 cm FL. The majority of females also mature at age 3 or 4 at a length of 35–42 cm FL. Ageing studies have shown that this species has very high recruitment variability (Hanchet et al. 2003).

3.2.3 Natural mortality

Natural mortality (M) was estimated using the equation $\log_e(100)/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unexploited stock. Using a maximum age of 22 years, M was estimated as 0.21 y^{-1} , and a value of 0.2 y^{-1} has been assumed in assessments (Table 5). Recent Campbell Island stock assessments have estimated M within the model. MCMC estimates of 0.18–0.22 have been obtained from recent assessments of the Campbell Island stock when M was estimated in the model (e.g., Dunn & Hanchet 2011b).

Table 5: Estimates of biological parameters estimated for the Campbell Island Rise stock, and assumed for all stocks.

| Estimate | Parameter | Male | Female | Source |
|---------------------------------------|-----------|---------|---------|----------------|
| Natural mortality (y^{-1}) | M | 0.2 | 0.2 | Hanchet (1992) |
| Weight = a (length) ^b | a | 0.00515 | 0.00407 | |
| Weight in g, length in cm fork length | b | 3.092 | 3.152 | Hanchet (1991) |

4. RESEARCH SURVEYS AND OTHER ESTIMATES OF ABUNDANCE

4.1 Acoustic research surveys

A programme to estimate southern blue whiting spawning stock biomass on each fishing ground using acoustic techniques began in 1993. The Bounty Platform, Pukaki Rise, and Campbell Island Rise were each surveyed annually between 1993 and 1995. After the first three annual surveys, these areas were surveyed less regularly. The Bounty Platform grounds were surveyed in 1997, 1999, and most recently in 2001. The Pukaki area was surveyed in 1997 and 2000. The only on-going series of research surveys is on the Campbell Island Rise grounds, which have been surveyed in 1998, 2000, 2002, 2004, 2006, 2009, and 2011. All these surveys have been carried out from R.V. *Tangaroa* using towed transducers and have been wide-area surveys intended to survey spawning southern blue whiting and pre-recruits. The results of these surveys have been an important input into southern blue whiting stock assessments for the last decade (e.g., Dunn & Hanchet 2011a, Dunn & Hanchet 2011b). Various designs for acoustic surveys of southern blue whiting were investigated using simulation studies by Dunn & Hanchet (1998) and Dunn et al. (2001), whilst Hanchet et al. (2000a) examined diel variation in southern blue whiting density estimates.

The primary objective of the RV *Tangaroa* wide area surveys has been to estimate the biomass of the adult spawning stock. A secondary objective has been to provide estimates of pre-recruit fish in each of the areas and so the surveys have been extended into shallower water where the younger fish live. When adult southern blue whiting are actively spawning, the marks are easily identified because they are very dense and have characteristic features (McClatchie et al. (2000); Hanchet et al. (2000b)). However, the pre-spawning and post-spawning adult marks are somewhat more diffuse and the adult fish distribution at this time often overlaps with the pre-recruits. The original analysis separated southern blue whiting marks into categories of adult, immature (mainly 2 and 3 years old) and juvenile (mainly 1 year old).

In some areas and years the marks classified as adults also contained some immature 2 and 3 year old fish, whilst juveniles were often a mix of 1 and 2 year old fish. This problem was addressed by Hanchet et al. (2000b) who carried out a re-analysis of the early R.V. *Tangaroa* acoustic survey and decomposed the estimates into age 1, 2, 3, and 4+ fish. This decomposition of acoustic marks was based on the age composition of targeted research trawls for juvenile and immature marks and on the age composition of commercial trawls for adults. Decomposed estimates of abundance were used as input for stock assessments of all three areas until 2012. However, this introduced the concern that the commercial catch-at-age data were being used twice inside the model — once to decompose the adult acoustic indices and then again as a separate input to the stock assessment model. So for the 2013 assessment of the Bounty Platform the model was revised to use the biomass estimates of the southern blue whiting categories (adult, immature, juvenile) instead.

There have also been changes in the target strength–fish length (TS–FL) relationship over time. The original estimate was based on the target strength used for the closely related blue whiting (*M. poutassou*) in the northern hemisphere (Monstad et al. 1992), which had itself been derived from measurements of juvenile cod (Nakken & Olsen 1977). A TS–FL relationship for southern blue whiting was first developed by Dunford & Macaulay (2006), which had a much steeper slope and gave higher adult TS than the previous relationship. More recently, O’Driscoll et al. (2013) developed a revised TS–FL relationship using an autonomous optical acoustic system. This new relationship has a shallower slope and lower intercept and is more similar to a recent value estimated for blue whiting (Pedersen et al. 2011). O’Driscoll et al. (2013) also suggested that part of the reason for the discrepancy between the estimates of Dunford & Macaulay (2006) and the *in situ* estimates of O’Driscoll et al. (2013) was due to an inappropriate application of the Kirchhoff-approximation model at small swimbladder sizes.

The time series of acoustic estimates of the RV *Tangaroa* wide area surveys have been revised on several occasions to reflect these changes in target strength and other changes to the acoustic analysis. Grimes et al. (2007) updated the decomposed estimates by: (i) incorporating the new target strength–fish length relationship of Dunford & Macaulay (2006), (ii) using the revised sound absorption coefficient of Doonan et al. (2003), (iii) including corrections and changes to strata areas, and (iv) estimating CVs of the decomposed estimates by age. Estimates of biomass of the southern blue whiting categories were recalculated in 2010 by P. Grimes (NIWA unpublished data) who revised the estimates by making the changes identified in (i)–(iii) above, and these revised estimates have been summarised in Fu et al. (2013). Most recently, Fu et al. (2013) revised estimates of biomass of the southern blue whiting categories using the new TS–FL relationship of O’Driscoll et al. (2013).

For the purposes of this report, we present the acoustic estimates for almost all areas and surveys using the most recent TS–FL relationship for blue whiting (O’Driscoll et al. 2013). Where necessary, acoustic indices in earlier reports have been updated (Richard O’Driscoll, NIWA, pers. comm.). An update for the single survey of the Auckland Islands Shelf (which used the earlier TS–FL relationship of Monstad et al. 1992) was not available.

4.1.1 Auckland Islands

A single survey of the Auckland Islands Shelf was carried out in 1995 using R.V. *Tangaroa*. This provided a spawning stock biomass estimate of 7800 t (CV = 0.34) based on the original TS–FL relationship for blue whiting (Ingerson & Hanchet 1996).

4.1.2 Bounty Platform

Two time series of acoustic indices are available for the Bounty Platform stock (Fu et al. 2013). The first was a wide-area time series of juvenile, immature, and adult southern blue whiting using the R.V. *Tangaroa* for the period 1993 to 2001 (Table 6).

A time series of aggregation or local area acoustic surveys using industry vessels (usually from only one vessel in each year) was initiated in 2004 and has continued in most years to 2013 (Table 7). These surveys have had mixed levels of success. Acoustic data collected in 2005 could not be used because of acoustic interference from the scanning sonar used by the vessel for searching for fish marks and inadequate survey design. There was also concern that the surveys in 2006 and 2009 did not sample the entire aggregation because on several transects the fish marks extended beyond the area being surveyed (O'Driscoll 2011c). Surveys in 2010, 2011, 2012, and 2013 were thought to have had reasonably good coverage, and to have surveyed the aggregations successfully (O'Driscoll 2011a, O'Driscoll et al. 2012, O'Driscoll et al. 2014).

Table 6: R.V. *Tangaroa* juvenile, immature, and mature acoustic biomass estimates (t) and CVs for the Bounty Platform using the O'Driscoll et al. (2013) target strength (Fu et al. 2013).

| Year | Juvenile | | Immature | | Adult | |
|------|----------|------|----------|------|---------|------|
| | Biomass | CV | Biomass | CV | Biomass | CV |
| 1993 | 6 449 | 0.27 | 15 269 | 0.33 | 43 338 | 0.58 |
| 1994 | 38 | 0.27 | 7 263 | 0.29 | 17 991 | 0.25 |
| 1995 | 25 961 | 0.37 | 0 | – | 17 945 | 0.23 |
| 1997 | 56 | 0.62 | 3 265 | 0.54 | 27 594 | 0.37 |
| 1999 | 674 | 0.57 | 344 | 0.37 | 21 956 | 0.75 |
| 2001 | 2 141 | 0.28 | 668 | 0.12 | 11 784 | 0.35 |

Table 7: The local area acoustic biomass estimates (t) and CVs for the Bounty Platform with the O'Driscoll et al. (2013) target strength and the proportion of catch taken at the time of the survey (TCEPR data).

| Year | Biomass | CV | Proportion ² | Source |
|-------------------|---------|------|-------------------------|--------------------------|
| 2004 | 8 572 | 0.69 | 0.73 | (Fu et al. 2013) |
| 2005 ¹ | – | – | | (Fu et al. 2013) |
| 2006 | 11 949 | 0.12 | 0.78 | (Fu et al. 2013) |
| 2007 | 79 285 | 0.19 | 0.93 | (Fu et al. 2013) |
| 2008 | 75 899 | 0.34 | 0.68 | (Fu et al. 2013) |
| 2009 | 16 640 | 0.21 | 0.29 | (Fu et al. 2013) |
| 2010 | 18 074 | 0.36 | 0.35 | (Fu et al. 2013) |
| 2011 | 20 990 | 0.28 | 0.89 | (Fu et al. 2013) |
| 2012 | 16 333 | 0.07 | 0.84 | (Fu et al. 2013) |
| 2013 | 28 533 | 0.27 | 0.76 | (O'Driscoll et al. 2014) |

1. In 2005, a local area aggregation survey was carried out, but the acoustic data could not be used because of acoustic interference from the scanning sonar used by the vessel.
2. The local area aggregation surveys were carried out by fishing vessels in between fishing activities, and hence the resulting estimates of biomass represented a different time period during the fishing season in each year.

4.1.3 Campbell Island Rise

As of 2012, eleven acoustic surveys of the Campbell Island Rise spawning grounds have been completed using the R.V. *Tangaroa*, and results of recent surveys were reported by O'Driscoll et al. (2005), O'Driscoll et al. (2007), Gauthier et al. (2011), O'Driscoll et al. (2012), and O'Driscoll et al. (2014) are summarised in Table 8.

The first industry survey of the Campbell stock (Table 9) was carried out from F.V. *Aoraki* in September 2003 (O'Driscoll & Hanchet 2004). This demonstrated that industry vessels with hull-mounted acoustic systems could also be used to collect acoustic data on southern blue whiting in good weather (less than 25 knots of wind). However, subsequent surveys from industry vessels on the Campbell Island grounds have not been successful and hence have not provided estimates useful for stock assessment.

Table 8: R.V. *Tangaroa* juvenile, immature, and mature acoustic biomass estimates (t) and CV for the Campbell Island Rise 1993–2013 using the target strength of O’Driscoll et al. (2013).

| Year | Juvenile | | Immature | | Mature | | Total | Source |
|------|----------|------|----------|------|---------|------|---------|--------------------------|
| | Biomass | CV | Biomass | CV | Biomass | CV | Biomass | |
| 1993 | 0 | 0.00 | 35 208 | 0.25 | 16 060 | 0.24 | 51 268 | (Fu et al. 2013) |
| 1994 | 0 | 0.00 | 5 523 | 0.38 | 72 168 | 0.34 | 77 691 | (Fu et al. 2013) |
| 1995 | 0 | 0.00 | 15 507 | 0.29 | 53 608 | 0.30 | 69 114 | (Fu et al. 2013) |
| 1998 | 322 | 0.45 | 6 759 | 0.20 | 91 639 | 0.14 | 98 720 | (Fu et al. 2013) |
| 2000 | 423 | 0.39 | 1 864 | 0.24 | 71 749 | 0.17 | 74 035 | (Fu et al. 2013) |
| 2002 | 1 969 | 0.39 | 247 | 0.76 | 66 034 | 0.68 | 68 250 | (Fu et al. 2013) |
| 2004 | 639 | 0.67 | 5 617 | 0.16 | 42 236 | 0.35 | 48 492 | (Fu et al. 2013) |
| 2006 | 504 | 0.38 | 3 423 | 0.24 | 43 843 | 0.32 | 47 770 | (Fu et al. 2013) |
| 2009 | 0 | – | 24 479 | 0.26 | 99 521 | 0.27 | 124 000 | (Fu et al. 2013) |
| 2011 | 0 | – | 14 454 | 0.17 | 53 299 | 0.22 | 67 753 | (Fu et al. 2013) |
| 2013 | 0 | – | 8 004 | 0.55 | 65 801 | 0.25 | 73 805 | (O’Driscoll et al. 2014) |

Table 9: The local area acoustic biomass estimates (including zero transects) and CVs for the Campbell Island Rise using the target strength of O’Driscoll et al. (2013).

| Year | Vessel | Area | No. transects | Area (km ²) | Biomass (t) | CV |
|------|-----------------------------|-----------|---------------|-------------------------|-------------|------|
| 2003 | <i>Aoraki</i> | Northeast | 6 | 303 | 5 442 | 0.13 |
| | | | 7 | 407 | 7 518 | 0.17 |
| | | | 5 | 579 | 10 359 | 0.24 |
| | | | 5 | 342 | 18 162 | 0.55 |
| | | | 7 | 332 | 15 529 | 0.84 |
| | | | 5 | 330 | 13 586 | 0.57 |
| | | | 7 | 276 | 25 594 | 0.49 |
| | | | 5 | 393 | 22 722 | 0.60 |
| | | | 6 | 330 | 9958 | 0.39 |
| 2006 | <i>Professor Alexandrov</i> | Northeast | 4 | 225 | 4 145 | 0.72 |
| | | | 8 | 255 | 6 940 | 0.17 |
| 2010 | <i>Meridian I</i> | Northeast | 7 | 171 | 32 736 | 0.40 |
| | | | 7 | 168 | 27 891 | 0.67 |
| | | | 6 | 42 | 7 924 | 0.79 |
| | | | 7 | 31 | 5 518 | 0.65 |
| | | Northeast | 7 | 89 | 10 408 | 0.31 |
| | | | 7 | 67 | 11 918 | 0.58 |

4.1.4 Pukaki Rise

A total of five acoustic surveys of the Pukaki Rise spawning grounds were completed using the R.V. *Tangaroa* between 1993 and 2000 (Table 10).

Large aggregations of spawning southern blue whiting were detected by vessels fishing on the Pukaki Rise in 2009 (O’Driscoll 2011c). Three vessels opportunistically collected acoustic data on these aggregations. The acoustic biomass estimates from the snapshots at the Pukaki Rise in 2009 ranged from 200 t (CV 29%) to 11 000 t (CV 38%) (Table 11). The latter estimate was of a similar magnitude to the abundance of sub-adult and adult southern blue whiting estimated from previous wide-area acoustic surveys of the area (Table 10).

Acoustic surveys on Pukaki Rise in September 2010, using the industry vessels *Professor Alexandrov* and *Meridian I*, were reported by O’Driscoll (2011c) (Table 11). The estimated acoustic biomass from the survey by *Meridian I* was 1085 t (CV 17%). Acoustic data collected from the *Professor Alexandrov*

were of lower quality due to bubble aeration and were less extensive than the former, and its echosounder has never been scientifically calibrated, so they were not used to estimate biomass. O'Driscoll (2011b) re-iterated the problems with trying to use aggregation-based surveys on the Pukaki Rise and recommended the use of wide-area surveys instead. Updated estimates of the surveys using the new target strength-fish length relationship are not available.

Table 10: R.V. *Tangaroa* juvenile, immature, sub-adult, and adult acoustic biomass estimates (t) and CVs for the Pukaki Rise using the target strength of O'Driscoll et al. (2013) and reported by Fu et al. (2013).

| Year | Juvenile | | Immature | | Sub-adult | | Adult | |
|------|----------|----|----------|------|-----------|------|---------|------|
| | Biomass | CV | Biomass | CV | Biomass | CV | Biomass | CV |
| 1993 | 0 | – | 9 558 | 0.25 | | | 26 298 | 0.32 |
| 1994 | 0 | – | 125 | 1.00 | 3 591 | 0.48 | 21 506 | 0.44 |
| 1995 | 0 | – | 0 | – | | | 6 552 | 0.18 |
| 1997 | 0 | – | 1 866 | 0.12 | | | 16 862 | 0.34 |
| 2000 | 0 | – | 1 868 | 0.62 | 8 363 | 0.74 | 6 960 | 0.37 |

Table 11: The local area acoustic biomass estimates (ignoring zero transects) for the Pukaki Rise 2009–2012 (O'Driscoll 2011c). Updated estimates of the surveys using the new target strength-fish length relationship are not available.

| Year | Vessel | No. transects | Area (km ²) | Biomass | CV | Source |
|------|--------------------|---------------|-------------------------|---------|------|--------------------|
| 2009 | <i>Meridian 1</i> | 4 | 50 | 188 | 0.29 | (O'Driscoll 2011c) |
| | | 5 | 283 | 9 459 | 0.30 | (O'Driscoll 2011c) |
| | | 5 | 71 | 6 272 | 0.41 | (O'Driscoll 2011c) |
| | <i>Buryachenko</i> | 6 | 60 | 2 361 | 0.12 | (O'Driscoll 2011c) |
| | | 7 | 117 | 7 903 | 0.26 | (O'Driscoll 2011c) |
| | | 6 | 19 | 11 321 | 0.38 | (O'Driscoll 2011c) |
| 2010 | <i>Meridian 1</i> | 10 | 364 | 1 085 | 0.17 | (O'Driscoll 2011c) |

4.2 Trawl research surveys

Trawl surveys of the sub-Antarctic targeting hoki, hake, and ling have been carried out using R.V. *Tangaroa* since 1991 (e.g., O'Driscoll & Bagley 2009). Although southern blue whiting are not a target species of this survey, they are often caught in moderate numbers, particularly on the Pukaki Rise and Campbell Island Rise, and it was considered possible that the surveys could be used to monitor their abundance. Hanchet & Stevenson (2006) reanalysed biomass estimates and scaled length frequency distributions for southern blue whiting from the sub-Antarctic summer and autumn survey series for each of three sub-areas, Pukaki Rise, Campbell Island Rise, and Auckland Island Shelf. They defined the three areas as follows: Pukaki Rise (strata 11, 12); Campbell Island Rise (10, 13, 14, 15, and 9S); Auckland Island Shelf (3, 4, 5, 6, 7, 8, 9N) (Figure 8).




Figure 8: Survey area and stratum boundaries used for R.V. *Tangaroa* sub-Antarctic trawl surveys since 1996. Stratum 9 was split into 9N and 9S at 51°S for the analysis.

4.2.1 Auckland Island Shelf

The CVs of the biomass estimates for the Auckland Islands Shelf were typically 45–80%, making them too imprecise for monitoring abundance (Table 12). There was little consistency in biomass estimates between the summer and autumn series and between adjacent surveys. Hanchet & Stevenson (2006) concluded that because of the erratic biomass estimates and very high CVs, it is extremely unlikely that the trawl survey indices were monitoring abundance on the Auckland Islands Shelf.

Table 12: R.V. *Tangaroa* trawl survey biomass estimates (t), CVs, and number of stations (N) for selected strata for the Auckland Islands Shelf (data for 1991–2005 from Hanchet & Stevenson (2006), data for 2006 onwards N. Bagley, NIWA, pers. comm.).

| Year | Summer | | | | Year | Autumn | | | |
|------|---------|---------|------|----|------|---------|---------|------|----|
| | Trip | Biomass | CV | N | | Trip | Biomass | CV | N |
| 1991 | TAN9105 | 565 | 0.75 | 58 | 1992 | TAN9211 | 125 | 0.98 | 60 |
| 1992 | TAN9204 | 40 | 0.31 | 31 | 1993 | TAN9310 | 3 458 | 0.60 | 51 |
| 1993 | TAN9304 | 159 | 0.89 | 44 | 1996 | TAN9605 | 447 | 0.33 | 40 |
| 2000 | TAN0012 | 135 | 0.61 | 38 | 1998 | TAN9805 | 746 | 0.69 | 25 |
| 2001 | TAN0118 | 527 | 0.68 | 36 | | | | | |
| 2002 | TAN0219 | 68 | 0.76 | 38 | | | | | |
| 2003 | TAN0317 | 281 | 0.85 | 27 | | | | | |
| 2004 | TAN0414 | 28 | 0.69 | 30 | | | | | |
| 2005 | TAN0515 | 3 972 | 0.39 | 98 | | | | | |
| 2006 | TAN0617 | 1 146 | 0.81 | 40 | | | | | |
| 2007 | TAN0714 | 1 686 | 0.45 | 41 | | | | | |
| 2008 | TAN0813 | 275 | 0.55 | 36 | | | | | |
| 2009 | TAN0911 | 1 432 | 0.60 | 39 | | | | | |
| 2011 | TAN1117 | 3 628 | 0.61 | 45 | | | | | |
| 2012 | TAN1215 | 1 724 | 0.45 | 42 | | | | | |

4.2.2 Campbell Island Rise

The CVs of the biomass estimates for the Campbell Island Rise were mostly between 25 and 40%, making them only marginally useful for monitoring abundance (Table 13). There was some consistency in biomass estimates between the summer and autumn series and also between adjacent surveys. However, Hanchet & Stevenson (2006) noted that although the trend in the trawl survey abundance indices on the Campbell Island Rise was generally similar to estimates of biomass from the population model, the trawl survey underestimated biomass at low stock sizes and overestimated biomass at high stock sizes. Hanchet & Stevenson (2006) suggested that increasing the number of trawl stations would improve the precision of the surveys, but they could not determine if this would also remove this bias.

Dunn & Hanchet (2011a) included observations of biomass from the sub-Antarctic trawl survey and the associated age frequencies in an assessment model for the Campbell Island Rise. They drew a similar conclusion to Hanchet & Stevenson (2006), with the fits suggesting some consistency in the pattern of biomass estimates between the summer series, but the observations underestimating biomass at low stock sizes and overestimating biomass at high stock sizes.

Table 13: R.V. *Tangaroa* trawl survey biomass estimates (t), CVs, and number of stations (N) for selected strata for the Campbell Island Rise (data for 1991–2005 from Hanchet & Stevenson (2006), data for 2006 onwards N. Bagley, NIWA, pers. comm.).

| Year | Summer | | | | Year | Autumn | | | |
|------|---------|-------------|------|----|-------------------|---------|-------------|------|----|
| | Trip | Biomass (t) | CV | N | | Trip | Biomass (t) | CV | N |
| 1991 | TAN9105 | 2 328 | 0.53 | 52 | 1992 | TAN9211 | 5 013 | 0.31 | 54 |
| 1992 | TAN9204 | 5 942 | 0.58 | 39 | 1993 | TAN9310 | 2 472 | 0.25 | 52 |
| 1993 | TAN9304 | 1 714 | 0.29 | 34 | 1996 ¹ | TAN9605 | 31 203 | 0.36 | 19 |
| 2000 | TAN0012 | 10 738 | 0.14 | 23 | 1998 | TAN9805 | 10 321 | 0.37 | 17 |
| 2001 | TAN0118 | 6 393 | 0.40 | 23 | | | | | |
| 2002 | TAN0219 | 3 198 | 0.45 | 21 | | | | | |
| 2003 | TAN0317 | 1 047 | 0.56 | 19 | | | | | |
| 2004 | TAN0414 | 778 | 0.26 | 21 | | | | | |
| 2005 | TAN0515 | 1 502 | 0.27 | 17 | | | | | |
| 2006 | TAN0617 | 4 729 | 0.73 | 16 | | | | | |
| 2007 | TAN0714 | 2 631 | 0.53 | 19 | | | | | |
| 2008 | TAN0813 | 5 870 | 0.29 | 17 | | | | | |
| 2009 | TAN0911 | 4 884 | 0.31 | 15 | | | | | |
| 2011 | TAN1117 | 1 610 | 0.25 | 15 | | | | | |
| 2012 | TAN1215 | 13 444 | 0.51 | 23 | | | | | |

¹. Only 1 station for TAN9605 was in stratum 0009S. This was supplemented with a second station taken from 0009N to allow the stratum biomass and variance to be calculated. The contribution of stratum 0009S to the total biomass was about 64 t, and hence the impact of this correction was negligible.

4.2.3 Pukaki Rise

The CVs of the biomass estimates for the Pukaki Rise were quite variable between years but mainly in the range 20–45%, making them only marginally useful for monitoring abundance (Table 14). There was some consistency in biomass estimates between the summer and autumn series and also between adjacent surveys.

Hanchet & Stevenson (2006) concluded that given the reduction in station density over time and poor agreement of the indices with either modelled biomass or catch history, it was unlikely that the trawl survey indices were monitoring abundance on the Pukaki Rise. After reviewing the work, The Middle Depths Working Group recommended that the number of stations in the core Pukaki Rise stratum be increased slightly during the surveys and this has been undertaken in some recent surveys, where time allowed (e.g., O'Driscoll & Bagley 2009). Recent biomass estimates have fluctuated considerably and the conclusions of Hanchet & Stevenson (2006) remain valid.

Table 14: R.V. *Tangaroa* trawl survey biomass estimates (t), CVs, and number of stations (N) for selected strata for the Pukaki Rise (data for 1991–2005 from Hanchet & Stevenson (2006), data for 2006 onwards N. Bagley, NIWA, pers. comm.).

| Year | Summer | | | | Year | Autumn | | | |
|------|---------|-------------|------|----|------|---------|-------------|------|----|
| | Trip | Biomass (t) | CV | N | | Trip | Biomass (t) | CV | N |
| 1991 | TAN9105 | 3 037 | 0.31 | 30 | 1992 | TAN9211 | 2 368 | 0.31 | 29 |
| 1992 | TAN9204 | 2 894 | 0.60 | 17 | 1993 | TAN9310 | 3 550 | 0.24 | 20 |
| 1993 | TAN9304 | 3 684 | 0.44 | 16 | 1996 | TAN9605 | 13 698 | 0.65 | 15 |
| 2000 | TAN0012 | 6 659 | 0.33 | 10 | 1998 | TAN9805 | 11 102 | 0.31 | 10 |
| 2001 | TAN0118 | 2 995 | 0.26 | 14 | | | | | |
| 2002 | TAN0219 | 3 251 | 0.63 | 12 | | | | | |
| 2003 | TAN0317 | 1 731 | 0.35 | 12 | | | | | |
| 2004 | TAN0414 | 2 537 | 0.47 | 10 | | | | | |
| 2005 | TAN0515 | 1 109 | 0.18 | 10 | | | | | |
| 2006 | TAN0617 | 911 | 0.43 | 10 | | | | | |
| 2007 | TAN0714 | 3 747 | 0.28 | 12 | | | | | |
| 2008 | TAN0813 | 9 078 | 0.14 | 14 | | | | | |
| 2009 | TAN0911 | 45 657 | 0.85 | 12 | | | | | |
| 2011 | TAN1117 | 2 106 | 0.21 | 12 | | | | | |
| 2012 | TAN1215 | 6 295 | 0.47 | 12 | | | | | |

4.3 CPUE analyses

Standardised CPUE analyses were carried out for the southern blue whiting spawning fisheries on the Campbell Island Rise from 1986 to 2002, and on the Bounty Platform from 1990 to 2002 by Hanchet & Blackwell (2003). Indices were calculated using lognormal linear models of catch-per-tow, catch-per-hour, and catch-per-day for all vessels, and catch-per-tow for subsets of vessels based on processing type (surimi or dressed), and by relative experience in each fishery. The authors summarised the data and the method of calculating the indices, and then compared the CPUE indices with the results of recent stock assessments. Unstandardised CPUE indices are given in Table 15.

Table 15: Unstandardised target median catch per unit effort indices (t/hour) for the Auckland Islands, Bounty Platform, Campbell Island Rise, and Pukaki Rise fisheries, July–October 1990–2013 (source: TCEPR data).

| Year | Auckland Islands | Bounty Platform | Campbell Island Rise | Pukaki Rise |
|------|------------------|-----------------|----------------------|-------------|
| 1990 | 0.0 | 5.8 | 3.3 | 1.1 |
| 1991 | 0.0 | 4.2 | 4.5 | 4.9 |
| 1992 | 0.0 | 10.0 | 1.0 | 0.9 |
| 1993 | 8.1 | 4.8 | 4.0 | 1.8 |
| 1994 | 0.5 | 0.9 | 5.5 | 3.2 |
| 1995 | 1.5 | 6.1 | 5.8 | 1.9 |
| 1996 | 0.5 | 7.5 | 6.7 | 13.3 |
| 1997 | 5.7 | 6.4 | 5.7 | 4.5 |
| 1998 | 4.7 | 6.5 | 6.3 | 2.7 |
| 1999 | 1.6 | 6.9 | 10.2 | 14.3 |
| 2000 | 0.0 | 3.0 | 6.6 | 3.7 |
| 2001 | 0.0 | 6.0 | 7.2 | 0.4 |
| 2002 | 0.5 | 7.2 | 6.5 | 0.0 |
| 2003 | – | 14.3 | 8.0 | 0.0 |
| 2004 | 1.4 | 5.4 | 7.4 | 1.2 |
| 2005 | 5.6 | 9.0 | 7.2 | 0.7 |
| 2006 | – | 7.9 | 9.8 | 5.5 |
| 2007 | – | 12.9 | 9.7 | 2.4 |
| 2008 | – | 16.6 | 9.4 | 13.6 |
| 2009 | 0.2 | 12.9 | 8.1 | 12.0 |
| 2010 | – | 12.6 | 10.2 | 8.8 |
| 2011 | – | 16.7 | 9.5 | 1.4 |
| 2012 | – | 12.3 | 10.0 | 2.4 |
| 2013 | – | 16.1 | 10.5 | 1.0 |

4.3.1 Bounty Platform

The Bounty Platform analysis was based on a data set of 3288 non-zero records from 1990 to 2002 (Hanchet & Blackwell 2003). The CPUE indices fluctuated considerably, peaking in 1992, 1996–1998, and again in 2002 (Table 16). The indices were similar between most of the CPUE models until 1997, but after 1997 they became more erratic between years and inconsistent amongst vessel subsets. The authors noted that there were other problems with the model assumptions, and that the model structure may be inadequate to reliably determine the indices and their standard errors. Trends in CPUE for the Bounty Platform fishery were consistent with trends in biomass from the 2002 NIWA assessment model of Hanchet (2002), apart from the first two years and last two years. The CPUE indices were rejected as indices of abundance by the Middle Depths Working Group and have not been used for stock assessments.

Table 16: Relative year effects and standard errors (s.e.) for the all vessels catch-per-tow model 1990 to 2002 for the Bounty Platform fishery (Hanchet & Blackwell 2003).

| Year | Standardised CPUE | |
|------|-------------------|------|
| | Index | s.e. |
| 1990 | 1.00 | — |
| 1991 | 1.20 | 0.12 |
| 1992 | 1.69 | 0.15 |
| 1993 | 0.89 | 0.10 |
| 1994 | 0.35 | 0.06 |
| 1995 | 0.57 | 0.09 |
| 1996 | 1.06 | 0.20 |
| 1997 | 0.98 | 0.25 |
| 1998 | 1.06 | 0.16 |
| 1999 | 0.68 | 0.08 |
| 2000 | 0.75 | 0.12 |
| 2001 | 0.98 | 0.25 |
| 2002 | 1.52 | 0.24 |

4.3.2 Campbell Island Rise

The original Campbell Island Rise analysis was based on 11 853 non-zero records from 1986 to 2002. CPUE indices decreased slowly to a minimum in 1992, increased to a peak in 1996, followed by a slight decline to 2002 (Hanchet & Blackwell 2003). This trend was consistent among alternative measures of effort and among subsets of surimi and dressed vessels. *Vessel* was the most important variable, together with *day in season*, *end time of tow*, and *sub-area*. Model diagnostics indicate a poor fit to the data, and the models were unable to fit very high or very low catch rates.

The trends in CPUE for the Campbell Island Rise fishery were consistent with the trends in the 2003 assessment model (Hanchet & Blackwell 2003). They followed the increase from 1993 to 1996 associated with the strong 1991 year class, and then followed the decline in relative abundance as this year class was fished down. Exploratory stock assessment model runs including the CPUE indices gave very similar results to those excluding the CPUE indices. The authors concluded that the CPUE indices for the Campbell Island Rise were monitoring the stock abundance and could be used in future stock assessments. However, they also cautioned that there can be considerable variability in the CPUE indices for individual years, and several years' data may be necessary before any trends become apparent.

The standardised CPUE analysis (Table 17) was updated to 2005 by (Hanchet et al. 2006). They found that there was some divergence in the CPUE indices between the various models in the years 2002 to 2005. The Working Group was unable to agree on which indices were monitoring abundance. As such the CPUE indices were rejected as indices of abundance by the Middle Depths Working Group and have not been used for stock assessments.

Table 17: Relative year effects and standard errors (s.e.) for all vessels catch-per-hour and catch-per-tow models, and raw mean CPUE for the Campbell Island fishery, 1986 to 2005 (source: Hanchet et al. 2006).

| Year | Catch per hour model | | | Catch per tow model | | |
|------|----------------------|------|-------------|---------------------|------|--------------|
| | Year index | s.e. | CPUE (t/hr) | Year index | s.e. | CPUE (t/tow) |
| 1986 | 1.00 | — | 9.7 | 1.00 | — | 14.9 |
| 1987 | 0.79 | 0.06 | 7.7 | 0.91 | 0.06 | 15.4 |
| 1988 | 0.59 | 0.05 | 6.7 | 0.88 | 0.06 | 19.9 |
| 1989 | 0.68 | 0.07 | 8.7 | 1.40 | 0.12 | 27.2 |
| 1990 | 0.52 | 0.05 | 7 | 1.04 | 0.09 | 17.7 |
| 1991 | 0.44 | 0.05 | 7.2 | 1.31 | 0.13 | 18.3 |
| 1992 | 0.29 | 0.03 | 4.3 | 0.60 | 0.06 | 11.7 |
| 1993 | 0.69 | 0.09 | 9.4 | 1.05 | 0.13 | 24 |
| 1994 | 0.69 | 0.10 | 9.2 | 1.19 | 0.14 | 25.8 |
| 1995 | 0.93 | 0.14 | 11.3 | 1.26 | 0.17 | 46.2 |
| 1996 | 1.88 | 0.27 | 14 | 2.34 | 0.29 | 42 |
| 1997 | 1.67 | 0.23 | 10.3 | 2.34 | 0.29 | 32.1 |
| 1998 | 1.17 | 0.15 | 11.5 | 1.79 | 0.21 | 28.3 |
| 1999 | 1.91 | 0.26 | 17.3 | 2.57 | 0.30 | 36 |
| 2000 | 1.23 | 0.17 | 10.8 | 1.87 | 0.23 | 32.7 |
| 2001 | 1.00 | 0.13 | 11.1 | 1.77 | 0.21 | 36.1 |
| 2002 | 1.02 | 0.14 | 11.1 | 1.88 | 0.22 | 33.2 |
| 2003 | 0.82 | 0.11 | 10.3 | 2.11 | 0.25 | 36.6 |
| 2004 | 0.92 | 0.12 | 12.1 | 1.95 | 0.23 | 28.9 |
| 2005 | 0.95 | 0.13 | 13.5 | 2.51 | 0.30 | 33.6 |

5. LENGTH AND AGE COMPOSITION OF THE FISHERY

5.1 Methods

Historical time series of catch-at-age data are available for each of the stocks, and these form an important input into the southern blue whiting stock assessments. A summary of the number of length measurements and otoliths read, on which these catch-at-age estimates were based, is tabulated for each area, in Tables 18–22. The raw LF data were examined graphically for variability in length composition by time of season and/or locality within each of the main areas, and divided into appropriate strata. The length frequency data for each tow was first scaled up to the catch from that tow, and these were then scaled up to the catch for each of the strata, and then the strata were combined to form a single length frequency for that area/year combination.

Age-length keys were year and area specific. In most years, 400–500 otoliths were read for each area/year combination. Before 2002, the catch-at-age was estimated by combining the scaled length frequency data with the age-length key using the old NIWA catch-at-age software. Catch-at-age data for each stock were reanalysed in 2002 using the new NIWA catch-at-age software (Bull & Dunn 2002) and the revised catch-at-age series for the Bounty Platform and Campbell Island Rise were summarised by Hanchet (2005) and Hanchet et al. (2003) respectively. This software produces CVs that incorporate the variance from both the length-frequency data and the age-length key using bootstrapping, and is an improvement over earlier algebraic calculations. Some of the age-length keys used for the analysis were also slightly modified during that re-analysis. Where necessary, ‘proxy’ ages were assumed for those length intervals with no corresponding age — typically only smaller fish lengths (less than about 30 cm) that were assigned age 1 or 2 depending on their size. We therefore ensured that an age was available for every length interval below 50 cm for males and 52 cm for females, for which length frequency observations were available. Any larger fish were put into an ‘unassigned’ category, which were placed in the plus group at age 11 for modelling.

5.2 Auckland Islands

The Auckland Islands has been fished only sporadically since 1990 (Table 18). Some targeting of aggregations of southern blue whiting during the spawning season occurred between 1993 and 1998, but since then most of the southern blue whiting catch has been taken as bycatch of fisheries targeting hoki, hake, ling, and squid during other months of the year (Hanchet & Dunn 2009). Almost 90% of the catch but less than 40% of the tows have been made in the months July to October.

Small numbers of fish were measured in most years from a small amount of effort around the Auckland Islands (Figure 9, Table 18), but few otoliths were collected and these have not been read. Catch-at-age data are available only for 1993 to 1998 (Figure 10). The catch at that time was dominated by the strong 1991 year class.

Table 18: Total number of tows and TCEPR estimated catch of southern blue whiting (including non-target), observed tows and estimated catch (including non-target), number of measured and aged males and females, Auckland Islands 1990–2013 (source: TCEPR and Observer data, 1990–2013).

| Year | Catch | | | Observed | | | | Measured | | Aged | |
|------|---------|------|-------|----------|------|-----|------|----------|--------|------|--------|
| | Vessels | Tows | t | Vessels | Tows | t | %t | Male | Female | Male | Female |
| 1990 | 7 | 90 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 1991 | 6 | 130 | 5 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 1992 | 11 | 90 | 73 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 1993 | 5 | 82 | 1 133 | 2 | 5 | 457 | 40.4 | 495 | 264 | 28 | 37 |
| 1994 | 12 | 315 | 1 056 | 1 | 7 | 324 | 30.6 | 601 | 563 | 57 | 79 |
| 1995 | 5 | 15 | 408 | 4 | 10 | 345 | 84.4 | 732 | 974 | 46 | 94 |
| 1996 | 3 | 6 | 54 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 1997 | 7 | 59 | 935 | 3 | 11 | 517 | 55.4 | 1 019 | 827 | 126 | 114 |
| 1998 | 11 | 121 | 520 | 1 | 6 | 238 | 45.8 | 649 | 550 | 80 | 38 |
| 1999 | 10 | 174 | 214 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2000 | 11 | 273 | 7 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2001 | 9 | 219 | 0 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2002 | 15 | 499 | 45 | 2 | 3 | 3 | 6.0 | 100 | 89 | 20 | 25 |
| 2003 | 14 | 466 | 14 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2004 | 11 | 314 | 27 | 1 | 1 | 4 | 16.7 | 12 | 28 | 0 | 0 |
| 2005 | 13 | 445 | 43 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2006 | 12 | 226 | 35 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2007 | 11 | 488 | 240 | 2 | 5 | 4 | 1.8 | 107 | 77 | 0 | 0 |
| 2008 | 7 | 186 | 67 | 1 | 11 | 16 | 24.4 | 307 | 220 | 0 | 0 |
| 2009 | 8 | 295 | 93 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2010 | 9 | 257 | 39 | 1 | 3 | 5 | 13.5 | 175 | 175 | 0 | 0 |
| 2011 | 9 | 297 | 29 | 1 | 1 | 1 | 2.0 | 9 | 11 | 0 | 0 |
| 2012 | 4 | 268 | 13 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2013 | 6 | 227 | 16 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |




Figure 9: Commercial catch proportions-at-length for the Auckland Islands stock by sex, 2007–2011.




Figure 10: Commercial catch proportions-at-age for the Auckland Islands stock by sex and year class, 1990–2013. Symbol area is proportional to the proportions-at-age within the sampling event.

5.3 Bounty Platform

The Bounty Platform has been fished consistently since 1990 (Table 19), and in each year all of the catch and almost all of the tows have been made between July and October. Catch-at-age data are available for the entire period 1990 to 2013, although the numbers of fish measured and aged were low in some years and zero in 2003 (Table 19). Examination of the raw data showed that the length composition was relatively constant through the season and across the area and so in most years all the length frequency data were placed into a single stratum. The catch in recent years has been dominated by a single mode of fish (the 2002 year class), which can be tracked from 2005, when it first entered the fishery at about 30 cm as 3 year olds, to 2013, when it still dominated the fishery as 11 year olds (Figure 11). In the past few years, a second, weaker 2007 year class has appeared in the length and age frequency. Previously, the catch over the 30 year period has been dominated by several other strong year classes — in particular those from 1986, 1988, 1991, 1992, and 1994 (Figure 12).

Table 19: Total number of tows and TCEPR estimated catch of southern blue whiting (including non-target), observed tows and estimated catch (including non-target), number of measured and aged males and females, Bounty Platform, 1990–2013 (source: TCEPR and Observer data, 1990–2013).

| Year | Catch | | | Observed | | | | Measured | | Aged | |
|------|---------|-------|--------|----------|------|--------|------|----------|--------|------|--------|
| | Vessels | Tows | t | Vessels | Tows | t | %t | Male | Female | Male | Female |
| 1990 | 26 | 269 | 4 438 | 5 | 23 | 391 | 8.8 | 2 569 | 1 690 | 135 | 118 |
| 1991 | 31 | 662 | 11 185 | 3 | 16 | 458 | 4.1 | 1 613 | 1 140 | 85 | 56 |
| 1992 | 49 | 1 732 | 58 696 | 10 | 161 | 10 086 | 17.2 | 12 726 | 12 189 | 318 | 282 |
| 1993 | 21 | 433 | 11 788 | 6 | 72 | 5 037 | 42.7 | 4 901 | 7 065 | 213 | 319 |
| 1994 | 9 | 202 | 3 877 | 4 | 40 | 2 836 | 73.1 | 4 202 | 3 126 | 255 | 253 |
| 1995 | 10 | 156 | 6 473 | 5 | 65 | 5 816 | 89.9 | 5 992 | 4 299 | 215 | 189 |
| 1996 | 5 | 67 | 5 113 | 2 | 22 | 2 511 | 49.1 | 2 171 | 2 465 | 201 | 280 |
| 1997 | 5 | 37 | 2 043 | 3 | 8 | 689 | 33.7 | 692 | 884 | 151 | 293 |
| 1998 | 12 | 119 | 5 824 | 6 | 69 | 5 627 | 96.6 | 7 574 | 6 743 | 211 | 261 |
| 1999 | 14 | 289 | 10 573 | 5 | 73 | 4 765 | 45.1 | 6 145 | 6 217 | 195 | 383 |
| 2000 | 6 | 99 | 3 851 | 3 | 27 | 2 716 | 70.5 | 1 858 | 3 323 | 110 | 288 |
| 2001 | 5 | 32 | 1 554 | 2 | 12 | 1 060 | 68.2 | 836 | 1 133 | 218 | 283 |
| 2002 | 8 | 182 | 6 209 | 1 | 8 | 1 116 | 18.0 | 590 | 671 | 62 | 87 |
| 2003 | 3 | 24 | 3 603 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2004 | 4 | 234 | 1 478 | 1 | 3 | 379 | 25.7 | 202 | 292 | 80 | 111 |
| 2005 | 8 | 284 | 3 769 | 4 | 40 | 2 818 | 74.8 | 3 212 | 3 256 | 159 | 261 |
| 2006 | 6 | 145 | 4 256 | 3 | 62 | 3 375 | 79.3 | 5 658 | 4 231 | 232 | 268 |
| 2007 | 5 | 103 | 3 602 | 3 | 27 | 3 458 | 96.0 | 2 118 | 2 124 | 110 | 190 |
| 2008 | 9 | 209 | 9 582 | 5 | 91 | 6 489 | 67.7 | 6 085 | 9 713 | 130 | 276 |
| 2009 | 14 | 426 | 14 958 | 4 | 104 | 5 269 | 35.2 | 7 637 | 8 526 | 130 | 292 |
| 2010 | 14 | 601 | 13 783 | 4 | 57 | 3 810 | 27.6 | 4 918 | 3 836 | 193 | 203 |
| 2011 | 9 | 241 | 6 468 | 4 | 49 | 3 876 | 59.9 | 4 121 | 4 205 | 140 | 200 |
| 2012 | 8 | 174 | 6 855 | 5 | 62 | 4 049 | 59.1 | 6 024 | 5 280 | 179 | 216 |
| 2013 | 5 | 77 | 3 860 | 5 | 30 | 2 638 | 68.3 | 2 356 | 2 224 | 186 | 228 |




Figure 11: Commercial catch proportions-at-length for the Bounty Platform stock by sex, 2006–2013.




Figure 12: Commercial catch proportions-at-age for the Bounty Platform stock by sex and year class, 1990–2013. Symbol area proportional to the proportions-at-age within the sampling event.

5.4 Campbell Island Rise

The Campbell Island Rise has been fished since 1979, although we have restricted much of the data presented here to that collected since 1990 (Table 20). Almost all of the catch and the tows during these years have been made in the months July to October.

Previous examination of the raw data has shown that the length composition is often different between the northern and southern Campbell Island Rise (Hanchet 1998). Therefore, the analysis was carried out by dividing the area into two strata (at 52° 30' S) for each year. The commercial catch at Campbell Island is currently dominated by a single mode of fish at about 38–40 cm, which comprises the 2006 and 2007 year classes and a mode of smaller fish comprising the 2009 year class (Figure 13–Figure 15). Previously, the catch since 1990 has been dominated by several other strong year classes – in particular those from 1988, 1991, and 2001 (see Figure 14 and Figure 15).

The time series of numbers-at-age (and CVs) from 1979 to 1989 are given in Hanchet et al. (2003). As described in earlier reports (e.g., Hanchet 1991, Hanchet & Ingerson 1995) the data for the early years (1979–1985) came from single vessels fishing during the spawning season and are probably less reliable than the more recent data, which have all been from multiple vessels. This tends to be reflected in the mean weighted CV, which ranged from 0.2–0.5 in the early years but 0.1–0.2 in more recent years (Hanchet et al. 2003).

Table 20: Total number of tows and TCEPR estimated catch of southern blue whiting (including non-target), observed tows and estimated catch (including non-target), number of measured males and females, Campbell Island Rise, 1990–2013 (source: TCEPR and Observer data, 1990–2013).

| Year | Catch | | | Observed | | | Measured | | Aged | | |
|------|---------|-------|--------|----------|------|--------|----------|--------|--------|------|--------|
| | Vessels | Tows | t | Vessels | Tows | t | %t | Male | Female | Male | Female |
| 1990 | 36 | 1 079 | 16 559 | 7 | 94 | 2 508 | 15.1 | 10 459 | 7 677 | 346 | 282 |
| 1991 | 35 | 1 242 | 21 934 | 3 | 52 | 1 107 | 5.0 | 3 852 | 4 864 | 281 | 413 |
| 1992 | 48 | 1 533 | 13 454 | 10 | 121 | 1 911 | 14.2 | 9 839 | 8 287 | 330 | 287 |
| 1993 | 20 | 423 | 8 757 | 5 | 55 | 2 722 | 31.1 | 4 456 | 4 623 | 247 | 321 |
| 1994 | 15 | 480 | 11 405 | 4 | 80 | 5 622 | 49.3 | 8 458 | 4 717 | 416 | 346 |
| 1995 | 12 | 285 | 9 989 | 5 | 76 | 7 726 | 77.3 | 5 807 | 7 301 | 212 | 358 |
| 1996 | 11 | 474 | 16 744 | 4 | 97 | 5 406 | 32.3 | 7 802 | 10 270 | 182 | 347 |
| 1997 | 18 | 650 | 19 145 | 6 | 185 | 9 476 | 49.5 | 16 756 | 16 254 | 239 | 255 |
| 1998 | 24 | 960 | 24 162 | 8 | 254 | 12 740 | 52.7 | 26 603 | 23 237 | 259 | 361 |
| 1999 | 21 | 790 | 27 206 | 9 | 175 | 11 308 | 41.6 | 15 024 | 15 522 | 227 | 190 |
| 2000 | 18 | 556 | 14 470 | 10 | 168 | 9 695 | 67.0 | 15 098 | 14 289 | 210 | 289 |
| 2001 | 16 | 919 | 24 410 | 10 | 321 | 19 144 | 78.4 | 27 994 | 25 500 | 135 | 270 |
| 2002 | 20 | 1 013 | 29 148 | 7 | 185 | 9 863 | 33.8 | 15 990 | 16 212 | 178 | 319 |
| 2003 | 16 | 636 | 22 695 | 5 | 124 | 2 922 | 12.9 | 9 259 | 10 979 | 236 | 383 |
| 2004 | 16 | 726 | 19 513 | 7 | 132 | 7 263 | 37.2 | 12 083 | 11 958 | 276 | 439 |
| 2005 | 17 | 757 | 25 200 | 6 | 187 | 9 041 | 35.9 | 14 184 | 18 757 | 147 | 262 |
| 2006 | 13 | 524 | 18 905 | 4 | 110 | 7 653 | 40.5 | 11 779 | 7 700 | 206 | 294 |
| 2007 | 13 | 549 | 20 437 | 6 | 119 | 8 345 | 40.8 | 10 291 | 11 504 | 182 | 234 |
| 2008 | 14 | 557 | 19 723 | 6 | 171 | 9 658 | 49.0 | 15 112 | 14 513 | 225 | 252 |
| 2009 | 14 | 629 | 18 299 | 3 | 53 | 3 145 | 17.2 | 4 506 | 3 856 | 123 | 311 |
| 2010 | 13 | 553 | 19 415 | 7 | 175 | 8 460 | 43.6 | 14 406 | 13 810 | 214 | 260 |
| 2011 | 14 | 976 | 29 204 | 8 | 246 | 9 739 | 35.2 | 19 884 | 24 570 | 207 | 254 |
| 2012 | 11 | 592 | 20 156 | 10 | 287 | 11 391 | 56.5 | 25 428 | 25 472 | 235 | 260 |
| 2013 | 10 | 694 | 25 624 | 9 | 299 | 13 460 | 52.5 | 23 435 | 23 870 | 191 | 174 |




Figure 13: Commercial catch proportions at length for the Campbell Island Rise stock by sex, 2006–2013.




Figure 14: Commercial catch proportions-at-age for the Campbell Island Rise stock by sex and year class, 1979–1994. Symbol area proportional to the proportions-at-age within the sampling event.




Figure 15: Commercial catch proportions-at-age for the Campbell Island Rise stock by sex and year class, 1995–2013. Symbol area proportional to the proportions-at-age within the sampling event.

5.5 Pukaki Rise

The Pukaki Rise has been fished only sporadically since 1990 with most of the catch taken between 1991 and 1993 and again in 2009 and 2010 (Table 21). While most of the catch has been made in the months July to October, less than half of the effort occurs during this period. The remaining effort has typically targeted hoki and other middle depth species (Hanchet & Dunn 2009).

Catch-at-age data are available for the period 1989 to 2000 and again for 2007, 2009, 2010, and 2012 although the numbers of fish measured and aged were low in some years (Table 21). Examination of the raw data showed that the length composition was relatively constant through the season and across the area and so the length frequency data were analysed as a single stratum. The catch on the Pukaki Rise has been dominated recently by fish comprising the 2004–2006 year classes but there has been recent recruitment of the 2009 year class, which may prove to be moderately strong (Figure 16). The catch over the 20 year period has been dominated by several other moderate year classes – in particular those from 1985, 1986, 1990, and 1991 (Figure 17).

Table 21: Total number of tows and TCEPR estimated catch of southern blue whiting (including non-target), observed tows and estimated catch (including non-target), number of measured males and females, Pukaki Rise 1990–2013 (source: TCEPR and Observer data, 1990–2013).

| Year | Catch | | | Observed | | | | Measured | | Aged | |
|------|---------|------|-------|----------|------|-------|-------|----------|--------|------|--------|
| | Vessels | Tows | t | Vessels | Tows | t | %t | Male | Female | Male | Female |
| 1990 | 35 | 464 | 1 295 | 6 | 20 | 204 | 15.7 | 2 624 | 1 050 | 182 | 197 |
| 1991 | 27 | 512 | 4 697 | 4 | 24 | 771 | 16.4 | 1 983 | 2 265 | 191 | 282 |
| 1992 | 44 | 614 | 2 866 | 5 | 23 | 227 | 7.9 | 1 611 | 1 391 | 233 | 243 |
| 1993 | 23 | 396 | 5 341 | 6 | 43 | 2 004 | 37.5 | 3 496 | 3 237 | 234 | 345 |
| 1994 | 14 | 195 | 1 918 | 4 | 22 | 1 191 | 62.1 | 1 831 | 1 940 | 222 | 188 |
| 1995 | 10 | 82 | 1 364 | 4 | 12 | 725 | 53.2 | 885 | 1 136 | 240 | 274 |
| 1996 | 5 | 11 | 299 | 1 | 1 | 112 | 37.5 | 72 | 113 | 0 | 0 |
| 1997 | 11 | 118 | 2 109 | 4 | 24 | 1 609 | 76.3 | 1 720 | 2 312 | 184 | 305 |
| 1998 | 15 | 115 | 1 219 | 7 | 18 | 1 248 | 102.4 | 1 686 | 1 756 | 174 | 168 |
| 1999 | 10 | 67 | 955 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2000 | 15 | 131 | 2 402 | 3 | 15 | 1 475 | 61.4 | 1 236 | 1 703 | 172 | 229 |
| 2001 | 15 | 68 | 284 | 1 | 2 | 45 | 15.9 | 153 | 157 | 0 | 0 |
| 2002 | 13 | 207 | 111 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2003 | 12 | 113 | 19 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2004 | 11 | 178 | 53 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2005 | 11 | 83 | 44 | 1 | 1 | 4 | 8.3 | 85 | 69 | 0 | 0 |
| 2006 | 8 | 47 | 1 048 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |
| 2007 | 12 | 200 | 391 | 1 | 4 | 103 | 26.4 | 382 | 287 | 39 | 48 |
| 2008 | 8 | 113 | 1 306 | 1 | 1 | 4 | 0.3 | 63 | 117 | 0 | 0 |
| 2009 | 16 | 393 | 4 777 | 4 | 48 | 1 078 | 22.6 | 3 016 | 3 953 | 164 | 261 |
| 2010 | 14 | 470 | 4 168 | 4 | 51 | 1 505 | 36.1 | 3 319 | 4 085 | 170 | 235 |
| 2011 | 10 | 471 | 625 | 4 | 6 | 96 | 15.4 | 482 | 359 | 0 | 0 |
| 2012 | 10 | 251 | 1 421 | 7 | 76 | 1 120 | 78.7 | 5 048 | 6 986 | 166 | 221 |
| 2013 | 7 | 36 | 9.9 | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 |




Figure 16: Commercial catch proportions at length for the Pukaki Rise stock by sex, 2007–2012.




Figure 17: Commercial catch proportions at age for the Pukaki Rise stock by sex and year class, 1990–2013. Symbol area proportional to the proportions-at-age within the sampling event.

5.6 Other areas (SBW1)

The remaining catch has been taken as bycatch of fisheries for hoki and other middle depths species from the Snares Shelf and southern Chatham Rise. Most of the large catches reported from this area are likely to be due to positional errors in the TCEPR database (Table 22). We plan to develop additional error checking algorithms to examine these errors in more detail prior to the 2015 assessment.

Table 22: Total number of tows and TCEPR estimated catch of southern blue whiting (including non-target), observed tows and estimated catch (including non-target), number of measured and aged males and females, SBW 1 1990–2013 (source: TCEPR and Observer data, 1990–2013). Most of the large catches reported from this area are likely to be due to positional errors in the TCEPR database.

| Year | TCEPR | | | Observer | | | Measured | | Aged | |
|------|---------|-------|-------|----------|------|-------|----------|--------|------|--------|
| | Vessels | Tows | Catch | Vessels | Tows | Catch | Male | Female | Male | Female |
| 1990 | 20 | 498 | 144 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1991 | 27 | 899 | 70 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 39 | 1 441 | 658 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 21 | 655 | 711 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 19 | 1 128 | 305 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 14 | 642 | 693 | 1 | 1 | 20 | 139 | 19 | 4 | 2 |
| 1996 | 7 | 405 | 45 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 13 | 823 | 163 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 23 | 1 082 | 93 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 26 | 1 732 | 14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 26 | 1 803 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 29 | 1 660 | 52 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2002 | 29 | 1 948 | 4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2003 | 23 | 1 187 | 8 | 1 | 1 | 0 | 54 | 1 | 0 | 0 |
| 2004 | 23 | 1 394 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2005 | 22 | 1 388 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2006 | 22 | 1 230 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2007 | 19 | 1 402 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2008 | 22 | 1 609 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2009 | 22 | 1 243 | 226 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2010 | 23 | 1 569 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2011 | 20 | 909 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2012 | 20 | 1 051 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2013 | 19 | 862 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

6. DISCUSSION

This document updates and summarises the observational and research data for southern blue whiting into one document. We have included here time series of relative abundance from acoustic surveys for each of the four main stocks from the wide area R.V. *Tangaroa* surveys as well as from local area aggregation industry vessel surveys, CPUE indices for Bounty Platform and Campbell Island Rise, and trawl survey indices for the Auckland Islands Shelf, Campbell Island Rise, and Pukaki Rise, as well as updated time series of length-at-age and catch-at-age.

R.V. *Tangaroa* acoustic surveys were carried out on the three main stocks from 1993 until around 2000, when because of the low catch limits on the Bounty and Pukaki stocks, the returns from the fishery were too low to be able to afford funding additional R.V. *Tangaroa* acoustic surveys and the time series of acoustic surveys was discontinued. Local area aggregation surveys from industry vessels on the Bounty Platform since 2004, and more recently on the Pukaki Rise, have provided the only biomass information on these stocks. However, there has been very large inter-snapshot and inter-annual variability in these biomass estimates making it difficult to use them for assessment and management purposes. On the Bounty Platform there was a seven-fold increase in biomass between 2006 and 2007 followed by a similar sized decline in biomass between 2008 and 2009. On the Pukaki Rise, acoustic snapshots ranged from 188 to 11 321 t in 2009 but declined to 1 085 t in 2010. Without a wide-area survey to provide a ground truthing of the aggregation results there will be on-going uncertainty about the status of these stocks (O'Driscoll 2011c, O'Driscoll et al. 2012).

Industry acoustic surveys on the Campbell Island Rise have also been unsuccessful to date, in part because of the much larger area which needs to be surveyed and the rougher weather conditions. Wide area acoustic surveys using the R.V. *Tangaroa* have been the preferred option for monitoring the Campbell Island stock because of the ability to use a towed acoustic array and the estimates of immature (age 2 and 3 year old) fish provided from this survey.

Estimates of abundance from the sub-Antarctic trawl surveys on the Auckland Islands Shelf, Campbell Island Rise, and Pukaki Rise were available for the period 1991 to 2011. While the surveys were not designed to monitor southern blue whiting, the biomass estimates for the latter two areas had moderate CVs, showed some consistency between years, and the trends showed some correspondence with biomass trajectories from stock assessments (Hanchet & Stevenson 2006). Dunn & Hanchet (2011b) investigated fitting the sub-Antarctic summer trawl survey time series in the Campbell assessment model, but found that although there was some consistency in biomass estimates between the summer series and the model estimates, the trawl survey underestimated biomass at low stock sizes and overestimated biomass at high stock. They concluded that the time series is not particularly useful for monitoring abundance. The utility of the trawl survey biomass indices for the Auckland Islands and the Pukaki Rise stocks is not known.

CPUE indices for the Bounty Platform and Campbell Island Rise are available for the periods 1990–2002 and 1986–2005 respectively. Although most fishing is carried out on highly aggregated spawning concentrations of southern blue whiting, there was moderate agreement between some of the CPUE indices and the biomass trajectories from modelling the stocks (Hanchet et al. 2003, Hanchet 2005). However, the MDWG was unable to agree on a time series to use and rejected these indices for stock assessment modelling (Ministry for Primary Industries 2012).

The time series of catch-at-age and length-at-age were updated for this report. Catch at Bounty Platform was still dominated by the 2002 year class. The 2007 year class has been observed in catches since 2010 and appears to be above average, but year classes from 2003 to 2006 and from 2008 onwards appear to be very weak.

There continues to be strong evidence of several year classes of moderate strength at the Campbell Island Rise, with both acoustic indices and commercial-at-age proportions suggesting strong recruitment in 2006, 2007, and 2009. The catch on the Pukaki Rise has been dominated recently by fish comprising the 2004–2006 year classes but there has been recent recruitment of the 2009 year class, which may prove to be moderately strong. Very little fishing has been carried out around the Auckland Islands in recent years and the data are insufficient to infer when or if strong recruitment has entered the fishery.

Catches from areas outside the four main fisheries remain low and typically have been less than 20 t annually.

7. ACKNOWLEDGMENTS

The authors are grateful to the scientific observers for the collection of the length frequency data and otoliths. Michael Stevenson and Colin Sutton read and processed the southern blue whiting otoliths. We thank Peter Horn for reviewing a draft of this manuscript. This project was funded by the Ministry for Primary Industries under Project DEE201002SBWC.

8. REFERENCES

- Anderson, O. (2004). Fish discards and non-target fish catch in the fisheries for southern blue whiting and oreos. *New Zealand Aquatic Environment and Biodiversity Report 2004/09*. 40 p.
- Anderson, O. (2009). Fish and invertebrate bycatch and discards in southern blue whiting fisheries, 2002–2007. *New Zealand Aquatic Environment and Biodiversity Report 43*. 28 p.
- Annala, J.H.; Sullivan, K.J.; Smith, N.W.M.; Griffiths, M.H.; Todd, P.R.; Mace, P.M.; Connell, A.M. (2004). Report from the Fishery Assessment Plenary, May 2004: stock assessments and yield estimates. 690 p. Ministry of Fisheries. (Unpublished report held in NIWA library, Wellington.)
- Bull, B.; Dunn, A. (2002). Catch-at-age: User manual v1.06.2002/09/12. NIWA Internal Report 114. 23 p. National Institute of Water and Atmospheric Research. (Unpublished report held in NIWA library, Wellington.)
- Clark, M.R.; Anderson, O.F.; Gilbert, D.J. (2000). Discards in trawl fisheries for southern blue whiting, orange roughy, hoki, and oreos in New Zealand waters. *NIWA Technical Report 71*. 73 p.
- Cole, R.; Dunn, A.; Hanchet, S.M. (2013). Review and summary of the time series of input data available for the assessment of southern blue whiting (*Micromesistius australis*) stocks. *New Zealand Fisheries Assessment Report 2013/69*. 41 p.
- Doonan, I.J.; Coombs, R.F.; McClatchie, S. (2003). The absorption of sound in seawater in relation to the estimation of deep-water fish biomass. *ICES Journal of Marine Science* 60(5): 1047–1055.
- Dunford, A.J.; Macaulay, G.J. (2006). Progress in determining southern blue whiting (*Micromesistius australis*) target strength: results of swimbladder modelling. *ICES Journal of Marine Science* 63: 952–955.
- Dunn, A.; Grimes, P.J.; Hanchet, S.M. (2001). Comparative evaluation of two-phase and adaptive cluster sampling designs for acoustic surveys of southern blue whiting (*Micromesistius australis*) on the Campbell Rise. Final Research Report to the Ministry of Fisheries for Project SBW1999/01, Objective 1. 15 p. National Institute of Water and Atmospheric Research. (Unpublished report held by the Ministry for Primary Industries.)
- Dunn, A.; Hanchet, S.M. (1998). Two-phase acoustic survey designs for southern blue whiting on the Bounty Platform and the Pukaki Rise. *NIWA Technical Report 28*. 29 p.
- Dunn, A.; Hanchet, S.M. (2011a). Southern blue whiting (*Micromesistius australis*) stock assessment for the Bounty Platform. *New Zealand Fisheries Assessment Report 2011/26*. 31 p.
- Dunn, A.; Hanchet, S.M. (2011b). Southern blue whiting (*Micromesistius australis*) stock assessment for the Campbell Island Rise, 2009–10. *New Zealand Fisheries Assessment Report 2011/40*. 37 p.
- Dunn, A.; Hanchet, S.M. (2014). Review and summary of the time series of input data available for the assessment of southern blue whiting (*Micromesistius australis*) stocks, 2013. *New Zealand Fisheries Assessment Report 2014/37*. 44 p.
- Fu, D.; Hanchet, S.M.; O'Driscoll, R.L. (2013). Estimates of biomass and c.v.s of southern blue whiting from previous acoustic surveys from 1993 to 2012 using a new target strength – fish length relationship. Final Research Report to the Ministry of Fisheries for Project DEE201002SBWB. 52 p. National Institute of Water and Atmospheric Research. (Unpublished report held by the Ministry for Primary Industries.)

- Gauthier, S.; Fu, D.; O'Driscoll, R.L.; Dunford, A. (2011). Acoustic estimates of southern blue whiting from the Campbell Island Rise, August–September 2009. *New Zealand Fisheries Assessment Report*. 40 p.
- Grimes, P.J.; Fu, D.; Hanchet, S.M. (2007). Estimates of biomass and c.v.s of decomposed age classes of southern blue whiting from previous acoustic surveys from 1993 to 2004 using a new target strength–fish length relationship. Final Research Report to the Ministry of Fisheries for Project SBW2005/01. 7 p. National Institute of Water and Atmospheric Research. (Unpublished report held by the Ministry for Primary Industries.)
- Hanchet, S.M. (1991). Southern blue whiting (*Micromesistius australis*) fishery assessment for the 1991–92 fishing year. New Zealand Fisheries Assessment Research Document 91/7. 32 p. MAF (Fisheries). (Unpublished report held in NIWA library, Wellington.)
- Hanchet, S.M. (1992). Southern blue whiting (*Micromesistius australis*) fishery assessment for the 1992–93 fishing year. New Zealand Fisheries Assessment Research Document 92/19. 28 p. MAF (Fisheries). (Unpublished report held in NIWA library, Wellington.)
- Hanchet, S.M. (1998). A review of southern blue whiting (*Micromesistius australis*) stock structure. New Zealand Fisheries Assessment Research Document 98/8. 29 p. Ministry of Fisheries. (Unpublished report held in NIWA library, Wellington.)
- Hanchet, S.M. (1999). Stock structure of southern blue whiting (*Micromesistius australis*) in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* 34(2): 599–610.
- Hanchet, S.M. (2002). Southern blue whiting (*Micromesistius australis*) stock assessment for the Bounty Platform for 2002 and 2003. *New Zealand Fisheries Assessment Report* 2002/53. 23 p. Ministry of Fisheries.
- Hanchet, S.M. (2005). Southern blue whiting (*Micromesistius australis*) stock assessment for the Bounty Platform for 2004–2005. *New Zealand Fisheries Assessment Report* 2005/45. 36 p.
- Hanchet, S.M.; Blackwell, R.G. (2003). Development and evaluation of catch-per-unit-effort indices for southern blue whiting (*Micromesistius australis*) on the Campbell Island Rise (1986–2002) and the Bounty Platform (1990–2002). Final Research Report to the Ministry of Fisheries for Project SBW2001/01 Objective 3. National Institute of Water and Atmospheric Research. (Unpublished report held by the Ministry for Primary Industries.)
- Hanchet, S.M.; Blackwell, R.G.; Dunn, A. (2005). Development and evaluation of catch per unit effort indices for southern blue whiting (*Micromesistius australis*) on the Campbell Island Rise, New Zealand. *ICES Journal of Marine Science* 62(6): 1131–1138.
- Hanchet, S.M.; Blackwell, R.G.; Stevenson, M.L. (2006). Southern blue whiting (*Micromesistius australis*) stock assessment for the Campbell Island Rise for 2006. *New Zealand Fisheries Assessment Report* 2006/41. 45 p.
- Hanchet, S.M.; Bull, B.; Bryan, C. (2000a). Diel variation in fish density estimates during acoustic surveys of southern blue whiting. *New Zealand Fisheries Assessment Report* 2000/16. 22 p.
- Hanchet, S.M.; Dunn, A. (2009). Southern blue whiting (*Micromesistius australis*) stock assessment for the Bounty Platform for 2007–08. Final Research Report to the Ministry of Fisheries for Projects SBW2006/01 and SBW2007/01, Objectives 1 and 2. 34 p. National Institute of Water and Atmospheric Research. (Unpublished report held by the Ministry for Primary Industries.)

- Hanchet, S.M.; Dunn, A.; Stevenson, M.L. (2003). Southern blue whiting (*Micromesistius australis*) stock assessment for the Campbell Island Rise for 2003. *New Zealand Fisheries Assessment Report 2003/59*. 42 p.
- Hanchet, S.M.; Ingerson, J.K.V. (1995). Southern blue whiting (*Micromesistius australis*) fishery assessment for the 1995–96 fishing year. New Zealand Fisheries Assessment Research Document 95/20. 37 p. Ministry of Fisheries. (Unpublished report held in NIWA library, Wellington.)
- Hanchet, S.M.; Richards, L.; Bradford, E. (2000b). Decomposition of acoustic biomass estimates of southern blue whiting (*Micromesistius australis*) using length and age frequency data. *New Zealand Fisheries Assessment Report 2000/43*. 37 p.
- Hanchet, S.M.; Stevenson, M.L. (2006). Biomass and scaled length frequency distribution of southern blue whiting by stock from the *Tangaroa* sub-Antarctic trawl surveys. Final Research Report to the Ministry of Fisheries for Project SBW2005/01, Objective 4. 22 p. National Institute of Water and Atmospheric Research. (Unpublished report held by the Ministry for Primary Industries.)
- Hanchet, S.M.; Uozumi, Y. (1996). Age validation and growth of southern blue whiting, *Micromesistius australis* Norman, in New Zealand. *New Zealand Journal of Marine and Freshwater Research* 30: 57–67.
- Ingerson, J.K.V.; Hanchet, S.M. (1996). Acoustic biomass estimates of southern blue whiting (*Micromesistius australis*) from the Bounty Platform, Pukaki Rise, Auckland Islands Shelf, and Campbell Island Rise, August–September 1995. New Zealand Fisheries Assessment Research Document 96/18. 29 p. Ministry of Fisheries. (Unpublished report held in NIWA library, Wellington.)
- McClatchie, S.; Thorne, R.; Grimes, P.J.; Hanchet, S.M. (2000). Ground truth and target identification for fisheries acoustics. *Fisheries Research* 47: 173–191.
- Ministry for Primary Industries (2012). Report from the Fishery Assessment Plenary, May 2012: stock assessments and yield estimates. 1194 p. Fisheries Science Group, Ministry for Primary Industries.
- Ministry for Primary Industries (2013). Report from the Fishery Assessment Plenary, May 2013: stock assessments and yield estimates. 1357 p. Fisheries Science Group, Ministry for Primary Industries.
- Monstad, T.; Borkin, I.; Ermolchev, V. (1992). Report of the joint Norwegian-Russian acoustic survey on blue whiting, spring 1992. *ICES CM 1992/H:6, Pelagic Fish Committee*: 26 p.
- Nakken, O.; Olsen, K. (1977). Target strength measurements of fish. *Rapports et Procès-Verbaux des Réunions du Conseil International pour l'Exploration de la Mer* 170: 52–69.
- O'Driscoll, R.L. (2011a). Acoustic biomass estimate of southern blue whiting on the Bounty Platform in 2010. NIWA Client Report: WLG2011-1. 27 p. National Institute of Water and Atmospheric Research. Unpublished Report to the Deepwater Group Ltd
- O'Driscoll, R.L. (2011b). Acoustic biomass estimates of southern blue whiting on the Pukaki Rise and Campbell Island Rise in 2010 NIWA Client Report: WLG2011-3. 37 p. National Institute of Water and Atmospheric Research. Unpublished Report to the Deepwater Group Ltd.
- O'Driscoll, R.L. (2011c). Industry acoustic surveys of spawning southern blue whiting on the Bounty Platform and Pukaki Rise 2004–2009. *New Zealand Fisheries Assessment Report 2011/17*. 52 p.
- O'Driscoll, R.L. (2013). Acoustic estimates of southern blue whiting on the Bounty Platform in 2012. *New Zealand Fisheries Assessment Report 2013/4*. 26 p.

- O'Driscoll, R.L.; Bagley, N.W. (2009). Trawl survey of hoki, hake, and ling in the Southland and Sub-Antarctic areas, November-December 2008 (TAN0813). *New Zealand Fisheries Assessment Report 2009/56*. 67 p.
- O'Driscoll, R.L.; Dunford, A.J.; Fu, D. (2012). Acoustic estimates of southern blue whiting from the Campbell Island Rise, August–September 2011 (TAN1112). *New Zealand Fisheries Assessment Report 2012/18*. 56 p.
- O'Driscoll, R.L.; Dunford, A.J.; Ladroit, Y. (2014). Acoustic estimates of southern blue whiting from the Campbell Island Rise, August-September 2013 (TAN1309). *New Zealand Fisheries Assessment Report 2014/22*. 46 p.
- O'Driscoll, R.L.; Dunford, A.J.; Macaulay, G.J. (2009). Acoustic biomass estimates of southern blue whiting on the Bounty Platform and Pukaki Rise from industry vessels in 2009. NIWA Client Report WL2009-81. 47 p. National Institute of Water and Atmospheric Research. Unpublished Report to the Deepwater Group Ltd
- O'Driscoll, R.L.; Grimes, P.J.; Hanchet, S.M.; Dunford, A. (2005). Acoustic estimates of southern blue whiting from the Campbell Island Rise, August–September 2004. *New Zealand Fisheries Assessment Report 2005/41*. 29 p.
- O'Driscoll, R.L.; Hanchet, S.M. (2004). Acoustic survey of spawning southern blue whiting on the Campbell Island Rise from FV *Aoraki* in September 2003. *New Zealand Fisheries Assessment Report 2004/27*. 31 p.
- O'Driscoll, R.L.; Hanchet, S.M.; Gauthier, S.; Grimes, P.J. (2007). Acoustic estimates of southern blue whiting from the Campbell Island Rise, August-September 2006. *New Zealand Fisheries Assessment Report 2007/20*. 34 p.
- O'Driscoll, R.L.; Oeffner, J.; Dunford, A.J. (2013). *In situ* target strength estimates of optically verified southern blue whiting (*Micromesistius australis*). *ICES Journal of Marine Science* 70: 431–439.
- Pedersen, G.; Rune Godø, O.; Ona, E.; Macaulay, G.J. (2011). A revised target strength–length estimate for blue whiting (*Micromesistius poutassou*): implications for biomass estimates. *ICES Journal of Marine Science* 68(10): 2222–2228.
- Shpak, V.M. (1978). The results of biological investigations of the southern putassu *Micromesistius australis* (Norman, 1937) on the New Zealand plateau and perspectives of its fishery. Unpublished TINRO manuscript. TINRO. (English translation held in NIWA library, Wellington.)
- Shpak, V.M.; Kuchina, V.V. (1983). Dynamics of abundance of southern putassu. *Biologiya Morya* 2: 35–39. (English translation held in NIWA library, Wellington.).